

BMJ

Using conjoint analysis to elicit preferences for health care

Mandy Ryan and Shelley Farrar

BMJ 2000;320:1530-1533
doi:10.1136/bmj.320.7248.1530

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/320/7248/1530>

These include:

Data supplement

"Table"
<http://bmj.com/cgi/content/full/320/7248/1530/DC1>

References

27 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/320/7248/1530#otherarticles>

Rapid responses

2 rapid responses have been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/320/7248/1530#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/320/7248/1530>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top left of the article

Topic collections

Articles on similar topics can be found in the following collections

[Clinical trials \(epidemiology\)](#) (3855 articles)
[Dentistry and oral medicine](#) (1102 articles)
[Communication](#) (227 articles)
[Internet](#) (965 articles)

Notes

To order reprints follow the "Request Permissions" link in the navigation box

To subscribe to *BMJ* go to:
<http://resources.bmj.com/bmj/subscribers>

Using conjoint analysis to elicit preferences for health care

Mandy Ryan, Shelley Farrar

Health Economics
Research Unit,
University of
Aberdeen,
Aberdeen
AB25 2ZD
Mandy Ryan
MRC senior fellow
Shelley Farrar
research fellow

Correspondence to:
M Ryan
m.ryan@abdn.ac.uk

BMJ 2000;320:1530-3

Traditionally the extent of patients' involvement in medical decision making has been minimal. This has been true at both the micro level of the patient consultation with a doctor and the macro level of planning and developing healthcare services. Since 1989, however, greater involvement of patients and the community in these processes has been advocated.¹⁻⁵ In principle, the elicitation of patients' and the community's values represents a big step forward in terms of enhancing the benefits from the provision of health care. For the exercise to be worth while, however, the information obtained must be useful and scientifically defensible.⁶ During the 1990s, conjoint analysis was developed to elicit patients' and the community's views on health care.

Methods

This paper explains conjoint analysis, provides examples of applications in health care which were obtained from a systematic review of databases between 1989 and 1999 (Medline, Embase, HealthSTAR, PsychLIT, EconLIT), and uses a study in orthodontic care to show the uses and pitfalls of the technique.

Conjoint analysis

The survey method of data collection and analysis known as conjoint analysis was developed in mathematical psychology and has a strong theoretical basis.⁷⁻⁹ It has been successfully used in market research,¹⁰ transport economics,¹¹ and environmental economics¹²⁻¹³ and was recommended to the UK Treasury for valuing quality in the provision of public services.¹⁴ Within these areas it has been well received by policymakers.¹²⁻¹⁴ The technique is gaining widespread use in health care and has been applied successfully in several areas, including eliciting patients' and the community's preferences in the delivery of health services¹⁵⁻²²; establishing consultants' preferences in priority setting²³; developing outcome measures²⁴⁻²⁵; determining optimal treatments for patients²⁶⁻²⁷; evaluating alternatives within randomised controlled trials²⁸⁻²⁹; and establishing patients' preferences in the doctor-patient relationship.³⁰⁻³¹

The technique is based on the premises that any good or service can be described by its characteristics (or attributes) and that the extent to which an individual values a good or service depends on the

Summary points

Conjoint analysis is a rigorous method of eliciting preferences

It allows estimation of the relative importance of different aspects of care, the trade-offs between these aspects, and the total satisfaction or utility that respondents derive from healthcare services

The technique can help with decision making for some of the issues facing the NHS

Though further applications of conjoint analysis are encouraged, methodological issues need further consideration

levels of these characteristics. The technique can be used to

- Show how people are willing to trade between characteristics; this is useful when deciding on the optimal way to provide a service within limited resources¹⁶⁻²⁰⁻²²
- Produce overall benefit scores for alternative ways of providing health care; this allows the ranking of health services against one another when setting priorities²¹⁻²²
- Estimate the relative importance of different characteristics of a service; this allows the policymaker to observe the individual impact of each characteristic on overall benefit¹⁵⁻³¹
- Estimate whether an attribute is important; this may be particularly useful when assessing the outcomes of trials. Trials generally include more than one outcome measure. Without further information it is not possible to determine whether these different outcome measures are important to patients.²⁸⁻²⁹

Undertaking a conjoint analysis study

Stage 1: Identifying the characteristics

The characteristics can be identified by various methods. If a policy question is being addressed, the characteristics will be predefined. For use alongside a randomised controlled trial, the characteristics may be defined by the different components of the arms of the trial.²⁸⁻²⁹ Where the characteristics are not predefined, literature reviews, group discussions, and individual interviews will be necessary.¹⁹



A table comparing scenarios is available on the *BMJ* website

Stage 2: Assigning levels to the characteristics

Levels assigned to the characteristics may be cardinal (for example, waiting time, where two weeks is twice as long as one week), ordinal (for example, though “severe pain” is worse than “moderate pain,” we do not know by how much), or categorical (for example, where there is no natural ordering for specialist nurse, general practitioner, or consultant). Pragmatically, the levels must be plausible and actionable, thus encouraging respondents to take the exercise seriously.

Stage 3: Choice of scenarios

Scenarios are then drawn up that describe all possible service (or outcome) configurations given the characteristics and levels chosen. The number of scenarios increases with the number of characteristics and levels. Rarely can all the scenarios generated be included in the questionnaire, and experimental designs are used to reduce the number to a manageable level.^{32 33}

Stage 4: Establishing preferences

Preferences for scenarios included in the questionnaire are elicited by using one of three methods: ranking, rating, or discrete choices. With ranking, respondents are asked to list the scenarios in order of preference. This method has not as yet been used in health care. The rating method requires the respondents to assign a score, of say 1 to 5, to each of the scenarios.^{21 25} For the discrete choice method, respondents are presented with a number of choices and, for each, asked to choose their preferred one. Possible responses include stating that either A or B is preferred,¹⁵⁻³¹ or responding on a five point scale where 1 equals definitely prefer A and 5 equals definitely prefer B (see figure). Given that choices more closely resemble real life decisions, the discrete choice approach has been preferred in health care.

Stage 5: Data analysis

Regression techniques are used to analyse responses. The appropriate method is determined by the type of data collected.³⁴ For discrete choice data, a benefit function is estimated, which can be specified as

$$\Delta B = \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \dots + \beta_n X_n$$

where ΔB is the change in benefit in moving from service A to B, X_j ($j = 1, 2, \dots, n$) are the differences in the attribute levels between A and B, and β_j ($j = 1, 2, \dots, n$) are the coefficients of the model to be estimated.

An application to orthodontic services

This application considers the trade-offs that individuals were willing to make between location of treatment

Table 1 Attributes and levels included in conjoint analysis study

Attributes	Levels	Regression coding
Waiting time (for first appointment) (WAIT)	4 months, 8 months, 12 months, 16 months	4, 8, 12, 16
Location of first appointment (for diagnosis) (LOC1)	Hospital, local	0=local, 1=hospital
Location of second appointment (and subsequent appointments for fixing the appliance) (LOC2)	Hospital, local	0=local, 1=hospital

and waiting time in the provision of orthodontic services.

Subjects and methods

A total of 160 patients attending three orthodontic clinics in the Grampian area of Scotland were asked to take part in the study.

The policy question identified the characteristics to be included in the conjoint analysis study. These and their levels are shown in table 1, together with their coding for the regression analysis. Sixteen scenarios were possible. Fifteen discrete choices were constructed by comparing the current service to all alternatives. Each respondent was presented with the 15 choices and asked, for each, which they preferred (see figure).

A check for internal consistency was included and dominant preferences (respondents not willing to trade a reduction in one attribute for an improvement in another) were identified. Respondents who were inconsistent or had dominant preferences were dropped from further analysis.

The following benefit equation was estimated for consistent traders:

$$\Delta B = \beta_1 LOC1 + \beta_2 LOC2 + \beta_3 WAIT$$

where ΔB is the change in benefit in moving from the current service to an alternative and the explanatory variables are the difference in the attributes of the two clinics, as defined in table 1. Given the nature of the dependent variable, the random effects ordered probit model was used to analyse the data.^{34 35}

Coefficients β_1 to β_3 show the relative importance of the different attributes, with the associated P statistic indicating whether the attribute has a statistically significant effect on choices. When interpreting these coefficients, it is important to be aware of the unit of measurement: β_1 and β_2 show the change in benefit for a unit change in location from central to local, and β_3 for a week's change in waiting time. β_1/β_3 indicates how much waiting time individuals are willing to give up to have their first appointment at a local clinic, and β_2/β_3 shows how much waiting time an individual would be willing to give up to have the second (and all subsequent appointments) at the local clinic. Benefit scores were estimated for different ways of providing orthodontic services.

	(a) Current			OR	(b) Alternative			Which option would you choose? (please tick one box for each choice)				
	First appointment	Second appointment	Waiting time		First appointment	Second appointment	Waiting time	Definitely (a) current	Probably (a) current	No preference	Probably (b) alternative	Definitely (b) alternative
Choice 1	Hospital	Hospital	8 months	OR	Local	Local	12 months	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Choice 2	Hospital	Hospital	8 months	OR	Hospital	Hospital	16 months	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Choice 3	Hospital	Hospital	8 months	OR	Local	Local	16 months	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Example of conjoint analysis question

Table 2 Results from regression analysis (1081 observations in 73 individuals)

Variable	Coefficient	P value
Waiting time*	-0.59	<0.001
Location of first appointment (0=local, 1=central)	-0.77	<0.001
Location of second appointment (0=local, 1=central)	-0.91	<0.001
Log likelihood	-1220	
χ^2	319.49	
McFadden R ²	0.12	

*Extra months willing to wait for a local first appointment=1.3; extra months willing to wait for a local second appointment=1.54.

Results

In all, 157 questionnaires were completed. Dropping inconsistent responders, those with dominant preferences, and missing values left 73 individuals and 1081 observations in the sample.

Table 2 presents the regression results for this group. The positive and significant signs on both location coefficients show that respondents prefer a local clinic to a hospital. Moving from a hospital clinic to a local clinic will increase benefit by 0.77 for the first appointment and 0.91 for the second appointment. The negative and significant sign of waiting time indicates that the higher the waiting time, the lower the benefit. The negative coefficient of 0.59 means that a unit increase in waiting time (for instance, from four months to five months) will reduce the benefit score by 0.59. Individuals are willing to wait an extra 1.3 months (0.77/0.59) to have a local clinic for their first appointment and an extra 1.5 months (0.91/0.59) to have a local clinic for their second appointment. From the equation used in the study, the benefit in moving from the current situation to having local first and second appointments and waiting 12 months was estimated to be -2.28. (Benefit scores were estimated for all the choice situations presented to individuals in the questionnaire; more detailed results are available on the *BMJ* website.) Generally, positive benefit values indicate an improvement over the current service and negative values indicate that patients would be less satisfied with that service configuration. Attempts should be made to reach the best configuration within available resources.

Other applications

The orthodontic study shows one way in which conjoint analysis can be applied, but the potential for the technique is much broader. Consider the issues faced by Dumfries and Galloway Health Board, recently highlighted in the *BMJ*.³⁶ This region faces problems common in the NHS. These relate to issues of new build, optimal information systems, relations between central and periphery hospitals, and patients' rising expectations. Conjoint analysis could be used to aid decision making within these areas. For example:

- Do patients want more personal and familiar local hospital care, where the level of specialisation may be lower, or more specialised, centrally located services?²²
- On what basis should priority be given to competing new build projects?²³

Although the technique can be applied to a wide range of policy questions within the NHS, several methodological issues remain. Some of these are outlined below within the context of the orthodontics study.

Methodological issues

In the orthodontics study, although the attributes were defined by the policy question, defining their levels was more difficult. Variations around the status quo were chosen in this study. In the only study appraising this issue, conjoint analysis has been shown to be relatively insensitive to characteristic levels (M Ryan and S Wordsworth, International Health Economics Association conference, 1999).

Secondly, the attributes and levels gave rise to 16 possible scenarios and, from these, 120 possible choices. Obviously it would not be possible to ask individuals their preferences among 120 choices, so the current situation was compared with the remaining 15 scenarios. An alternative approach is to randomly pair the 16 scenarios into choices.²³ Work is needed to assess the sensitivity of results to the method of setting discrete choices.

Thirdly, in data analysis, there is the issue of what to do with inconsistent responders and those exhibiting dominant preferences. For simplicity, inconsistent responders and "non-traders" were dropped in the application to orthodontic services, following the procedure in previous applications of conjoint analysis in health care. Future work, however, should consider the extent to which inconsistent responders are behaving rationally (and therefore should be included in the analysis) and should explore issues related to non-traders. The bias that exclusion may create in the results also needs considering.

Fourthly, there is a question of how to model the benefit function. In the orthodontics study, a linear additive relation was assumed. Research from outside health care has shown that alternatives to the linear additive model seldom result in a significantly better fit.³⁷ Future work applying conjoint analysis in health care should explore this issue.

Conclusions

This paper proposes conjoint analysis as a rigorous survey technique for eliciting the views of patients and people in the community on health care. The application presented here, which was well received by policy-makers, shows the potential uses and pitfalls of the technique. Conjoint analysis has been successfully applied elsewhere in health care and has great potential as an instrument for establishing the preferences of patients and the community (as well as those of clinicians and policymakers). Though further applications of conjoint analysis are encouraged in the NHS, methodological issues arising from its use should continue to be considered.

We are grateful to all respondents who completed questionnaires, to Dr Kate Dawson for help in setting up the study, and to health promotion assistants for their help in collecting the data.

Funding: Financial support from the Medical Research Council and the Chief Scientist Office of the Scottish Office Department of Health are acknowledged. The views expressed in the paper are those of the authors.

Contributors: MR has been developing and applying conjoint analysis in health care under the auspices of an MRC Senior Fellowship. For the empirical work presented here MR proposed that conjoint analysis be used to evaluate trade-offs between location and waiting time in the provision of orthodontic care. MR and SF developed the study design and

questionnaire and analysed the data. MR drafted the paper and SF provided comments. MR is the guarantor for the paper.

Competing interests: None declared.

- 1 Secretaries of State for Health, Wales, Northern Ireland, and Scotland. *Working for patients*. London: HMSO, 1989.
- 2 NHS Management Executive. *Local voices: the views of local people in purchasing for health*. London: Department of Health, 1992.
- 3 Secretaries of State for Health, Wales, Northern Ireland, and Scotland. *Promoting better health*. London: HMSO, 1989.
- 4 Secretary of State for Health. *The health of the nation*. London: HMSO, 1989.
- 5 Scottish Office Department of Health. *The patient's charter: what users think*. Edinburgh: HMSO, 1992.
- 6 Cleary P. The increasing importance of patient surveys. *BMJ* 1999;319:720-1.
- 7 Luce D, Tukey J. Simultaneous conjoint measurement: a new type of fundamental measurement. *J Math Psychol* 1964;1:1-27.
- 8 Anderson N. Functional measurement and psycho-physical judgement. *Psychol Rev* 1977;77:153-70.
- 9 McFadden D. *Conditional logit analysis of qualitative choice behaviour*. Berkeley, CA: University of California at Berkeley, 1973. (Working paper No 199/BART 10.)
- 10 Cattin P, Wittink D. Commercial use of conjoint analysis: a survey. *J Marketing* 1982;46:44-53.
- 11 [Conjoint analysis.] *J Transport Econ Policy* 1988;22(1).
- 12 Opaluch J, Swallow S, Weaver T, Wessells C, Wichelns D. Evaluating impacts from noxious facilities: including public preferences in current siting mechanisms. *J Environ Econ Manage* 1993;24:41-59.
- 13 Adamowicz W, Louviere J, Williams M. Combining revealed preference and stated preference methods for valuing environmental amenities. *J Environ Econ Manage* 1994;6:271-92.
- 14 Cave M, Birmingham D, Buxton M, Hanney S, Pollitt C, Scanlan M, et al. *The valuation of changes in quality in the public services*. London: HMSO, 1993.
- 15 Bryan S, Buxton M, Sheldon R, Grant A. Magnetic resonance imaging for the investigation of knee injuries: an investigation of preference. *Health Econ* 1998;7:595-604.
- 16 Propper C. Contingent valuation of time spent on NHS waiting list. *Econ J* 1991;100:193-9.
- 17 Propper C. The disutility of time spent on the United Kingdom's National Health Service waiting lists. *J Hum Resources* 1995;30:677-700.
- 18 Ratcliffe J, Buxton M. Patients' preferences regarding the process and outcomes of life saving technology: an application of conjoint analysis to liver transplantation. *Int J Technol Assess Health Care* 1999;15:340-51.
- 19 Ryan M. Using conjoint analysis to go beyond health outcomes: an application to in vitro fertilisation. *Soc Sci Med* 1999;8:535-46.
- 20 Van der Pol M, Cairns J. Establishing preferences for blood transfusion support: an application of conjoint analysis. *J Health Services Res Manage* 1998;3:70-6.
- 21 Ryan M, McIntosh E, Shackley P. Using conjoint analysis to assess consumer preferences in primary care: an application to the patient health card. *Health Expectations* 1998;1:117-29.
- 22 Ryan M, McIntosh E, Dean T, Old P. Trade-offs between location and waiting times in the provision of health care: the case of elective surgery on the Isle of Wight. *J Public Health Med* (in press).
- 23 Farrar S, Ryan M, Ross D, Ludbrook A. Using discrete choice modelling in priority setting: an application to clinical service developments. *Soc Sci Med* 2000;50:63-75.
- 24 Hakim Z, Pathak D. Modelling the EuroQol data: a comparison of discrete choice conjoint and conditional preference modelling. *Health Econ* 1999;8:103-16.
- 25 Harwood R, Rogers A, Dickinson E, Ebrahim S. Measuring handicap: the London handicap scale, a new outcome measure for chronic disease. *Qual Health Care* 1994;3:11-6.
- 26 Maas A, Stalpers L. Assessing utilities by means of conjoint measurement: an application in medical decision analysis. *Med Decision Making* 1992;12:288-97.
- 27 Verhoef C, Maas A, Stalpers L, Verbeek A, Wobbes T, Daal W. The feasibility of additive conjoint measurement in measuring utilities in breast cancer patients. *Health Policy* 1991;17:39-50.
- 28 Ryan M, Hughes J. Using conjoint analysis to value surgical versus medical management of miscarriage. *Health Econ* 1997;6:261-73.
- 29 San Miguel F, Ryan M, McIntosh E. Applying conjoint analysis in economic evaluations: an application to menorrhagia. *Appl Econ* (in press).
- 30 Scott A, Vick S. Patients, doctors and contracts: an application of principal-agent theory to the doctor-patient relationship. *Scot J Political Econ* 1999;46:111-34.
- 31 Vick S, Scott A. What makes a perfect agent? A pilot study of patients' preferences in the doctor patient relationship. *J Health Econ* 1998;17:587-606.
- 32 Kocur G, Alder T, Hyman W, Aunet B. *Guide to forecasting travel demand with direct utility assessment*. Washington, DC: US Department of Transportation, 1982.
- 33 Bradley M. *User's manual for the SPEED version 2.1. Stated preference experiment editor and designer*. Hague: Hague Consulting Group, 1991.
- 34 Greene W. *Econometric analysis*. Englewood Cliffs, NJ: Prentice Hall, 1993.
- 35 Greene W. *LIMDEP version 7.0*. New York: Econometric Software, 1995.
- 36 Smith R. The NHS in Dumfries and Galloway: straining but optimistic. *BMJ* 1999;319:1123-7.
- 37 Emery D, Barron F. Axiomatic and numerical conjoint measurement: an evaluation of diagnostic efficacy. *Psychometrika* 1979;44:195-210.

(Accepted 13 March 2000)

Modernising the NHS

Challenges to the health services: the professions

Isobel Allen

"There is the challenge for the professions to strip out unnecessary demarcations, introduce more flexible training and working practices and ensure that doctors do not use time dealing with patients who could be treated safely by other health care staff." This was the third of five challenges to the health service set out by the prime minister on 22 March, followed up by the secretary of state for health on 23 March with the announcement of six modernisation action teams to focus on the challenges. This announcement expanded on the reference to more flexibility and removing demarcations by adding "in the context of major expansion of the health care workforce."

The challenge to the professions has been made. What are the most important things that would make a difference?

Preparing to act on the challenge

The first thing is for the government to recognise the extent to which members of the professions are ready to take on the challenge. The profile of the medical

Summary points

Professionals are ready to change—and have already done so

NHS organisations need to listen to their staff, who know where systems go wrong

More flexible working and training are long overdue, together with proper training for multidisciplinary working

Innovation needs to be encouraged—and entrepreneurial clinicians supported

This is the fourth in a series of seven articles

Policy Studies Institute, London NW1 3SR

Isobel Allen
professor of health and social policy

i.allen@psi.org.uk

BMJ 2000;320:1533-5

profession has changed radically in the past 10 years. Women now account for more than 50% of doctors leaving medical schools¹ and represent an increasing proportion of doctors under the age of 40, particularly in general practice. Younger doctors are much more