

# Using "small" molecules to facilitate exchange of bicarbonate and chloride anions across liposomal membranes.

Jeffery T. Davis,<sup>1</sup> Philip A. Gale,<sup>2</sup> Oluyomi A. Okunola,<sup>1</sup> Pilar Prados,<sup>3</sup> Jose Carlos Iglesias-Sánchez,<sup>3</sup> Tomás Torroba,<sup>4</sup> and Roberto Quesada<sup>4</sup>

<sup>1</sup>Department of Chemistry and Biochemistry, University of Maryland, College Park, MD 20742, USA. <sup>2</sup>School of Chemistry, University of Southampton, Southampton, SO17 1BJ, United Kingdom. <sup>3</sup>Departamento de Química Orgánica, Universidad Autónoma de Madrid, 28049 Madrid, Spain. <sup>4</sup>Departamento de Química, Universidad de Burgos, 09001 Burgos, Spain. Correspondence should be addressed to J.T.D. (jdavis@umd.edu), P.A.G. (philip.gale@soton.ac.uk) or R.Q. (rquesada@ubu.es)

Please cite this paper as:

*Nature*, 2009, 1, pp 138–144

The publisher's version of this paper is available here:

<http://www.nature.com/nchem/journal/v1/n2/abs/nchem.178.html>

Related articles by Prof Philip A. Gale can be found below:

N. Busschaert and P.A. Gale, (2013) "[Small molecule lipid bilayer anion transporters for biological applications](#)" *Angew. Chem. Int. Ed.* 2013, 52, 1374-1382; DOI: 10.1002/anie.201207535

S.J. Moore, C.J.E. Haynes, J. González, J.L. Sutton, S.J. Brooks, M.E. Light, J. Herniman, G. J. Langley, Vanessa Soto-Cerrato, R. Pérez-Tomás, I. Marques, P.J. Costa, V. Félix and P.A. Gale, (2013) "[Chloride, carboxylate and carbonate transport by ortho-phenylene diamine-based bis-ureas.](#)" *Chem. Sci.*, 4, 103-117; DOI:10.1039/C2SC21112B.

N. Busschaert, I.L. Kirby, S. Young, S.J. Coles, P.N. Horton, M.E. Light and P.A. Gale, (2012) "[Squaramides as Potent Transmembrane Anion Transporters.](#)" *Angew. Chem. Int. Ed.*, 51, 4426-4430; DOI: 10.1002/anie.201200729

N. Busschaert, M. Wenzel, M.E. Light, P. Iglesias-Hernández, R. Pérez-Tomás and P.A. Gale, (2011) "[Structure-activity relationships in tripodal anion transporters: the effect of fluorination](#)" *J. Am. Chem. Soc.*, 133, 14136-14148; DOI: 10.1021/ja205884y.

P.A. Gale, (2011) "[From anion receptors to transporters](#)", *Acc. Chem. Res.*, 44, 216-226; DOI: 10.1021/ar100134p.

N.J. Andrews, C.J.E. Haynes, M.E. Light, S.J. Moore, C.C. Tong, J.T. Davis, W.A. Harrell Jr. and P.A. Gale, (2011) "[Structurally simple lipid bilayer transport agents for chloride and bicarbonate](#)", *Chem. Sci.*, 2, 256-260; DOI:10.1039/C0SC00503G.

## **Using "small" molecules to facilitate exchange of bicarbonate and chloride anions across liposomal membranes.**

Jeffery T. Davis,<sup>1</sup> Philip A. Gale,<sup>2</sup> Oluyomi A. Okunola,<sup>1</sup> Pilar Prados,<sup>3</sup> Jose Carlos Iglesias-Sánchez,<sup>3</sup> Tomás Torroba,<sup>4</sup> and Roberto Quesada<sup>4</sup>

<sup>1</sup>Department of Chemistry and Biochemistry, University of Maryland, College Park, MD 20742, USA. <sup>2</sup>School of Chemistry, University of Southampton, Southampton, SO17 1BJ, United Kingdom. <sup>3</sup>Departamento de Química Orgánica, Universidad Autónoma de Madrid, 28049 Madrid, Spain. <sup>4</sup>Departamento de Química, Universidad de Burgos, 09001 Burgos, Spain. Correspondence should be addressed to J.T.D. (jdavis@umd.edu), P.A.G. (philip.gale@soton.ac.uk) or R.Q. (rquesada@ubu.es)

**Bicarbonate is involved in a variety of biological processes including respiration, regulation of intracellular pH and fertilization. In this study we show that both simple synthetic anion receptors and the natural product prodigiosin are capable of facilitating chloride/bicarbonate exchange (antiport) across liposomal membranes. While it is well known that proteins regulate Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchange in cells, our results suggest that small molecules may also be able to regulate the concentration of these anionic species in biological systems. We also describe new NMR assays that were developed for monitoring transmembrane transport of H<sup>13</sup>CO<sub>3</sub><sup>-</sup>.**

Bicarbonate is an anion of paramount importance in living systems. It is a substrate in photosynthesis,<sup>1</sup> a component of the buffering system to regulate intra- and extracellular pH levels,<sup>2</sup> is generated during cellular respiration from CO<sub>2</sub>,<sup>3</sup> and acts as a cellular signal that activates sperm for fertilization.<sup>4</sup> Indeed, under physiological conditions most dissolved inorganic carbon exits as HCO<sub>3</sub><sup>-</sup>. Bicarbonate cannot diffuse freely across cell membranes and facilitated bicarbonate transport is promoted by membrane bound proteins that function mainly through Na<sup>+</sup>/HCO<sub>3</sub><sup>-</sup> co-transport or Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchange mechanisms.<sup>5</sup> Dysregulation of bicarbonate transport can lead to conditions such as cystic fibrosis, heart disease and infertility.<sup>6-8</sup> The lack of structural data for these proteins means that little is known regarding the anion binding sites that modulate their affinity and selectivity.<sup>7,8</sup> Despite the obvious importance of

transmembrane bicarbonate transport, there have been no reports, to the best of our knowledge, of "small" molecules that are capable of promoting bicarbonate transport across lipid membranes, in contradistinction to the growing body of work on chloride and HCl transport across phospholipid vesicle membranes.<sup>9-15</sup> Phospholipid vesicles have been extensively investigated because of their usefulness as models for biological membranes. Unilamellar vesicles of a specific size are easily produced with control of the entrapped solution. These liposomes can be suspended in an external medium of different composition and release of the encapsulated substrates or influx of substances from the external milieu to the interior of the vesicles, as facilitated by some molecular transporter, can be readily monitored by using fluorescence, NMR or ion selective electrode techniques. Below, we describe studies that demonstrate, for the first time, that synthetic compounds and natural products are able to facilitate transmembrane transport of bicarbonate anion across liposomal membranes.




Chart 1. 4,6-Dihydroxyisophthalamides **1-3** and prodigiosin **4**.

We recently reported the transmembrane chloride transport activity of 4,6-dihydroxyisophthalamide **1**.<sup>16</sup> Isophthalamides have amide NH groups that are able to form hydrogen bonds with anions.<sup>17</sup> In the case of **1**, conformational control of the anion binding cleft by means of intramolecular hydrogen bonds between the 4,6 dihydroxy units and the neighbouring amide carbonyls resulted in an improved anion affinity in solution and in optimal activity for transmembrane transport of chloride anion. For this present study, we also synthesized the closely related isophthalamides **2** and **3**, functionalized with different alkyl substituents on the amide groups (Chart 1). We hypothesized that the bidentate isophthalamide unit might also be effective at binding and, more importantly, transporting trigonal planar oxoanions, such as bicarbonate across lipid membranes.


In addition to synthetic compounds **1-3** we also report below on the bicarbonate transport activity of prodigiosin **4**,<sup>18</sup> a natural product produced by microorganisms such as *Streptomyces* and *Serratia*.<sup>19,20</sup> This tripyrrolic metabolite has been found to have potent immunosuppressive and anticancer activities. Prodigiosin appears to cause apoptosis of cancer cells selectively,<sup>21</sup> and the structural analogue obatoclax is currently in clinical trials for the treatment of cancer.<sup>22</sup> The origin of the biological activity has yet to be unambiguously established, although there is evidence that this class of compound facilitates the co-transport of HCl<sup>23-28</sup> and/or the anion exchange of chloride<sup>29,30</sup> across lipid bilayer membranes. Indeed, the antibiotic

activity of prodigiosin-like molecules has been related to their activity as transmembrane Cl<sup>-</sup> carriers.<sup>28</sup> As for isophthalamides **1-3**, the structure of prodigiosin **4**, with its three convergent pyrrole units, seemed to be well suited for coordinating to a trigonal planar anion such as bicarbonate.

We decided to investigate whether these potent chloride transporters (**1-4**) could also facilitate the transmembrane transport of the bicarbonate anion. In this work, we report 1) the first examples of transmembrane chloride/bicarbonate exchange facilitated by "small" molecules and 2) new NMR methods to monitor the transport of bicarbonate into lipid vesicles. We believe that these studies will set the stage for further development of highly-selective synthetic transporters for bicarbonate anion and perhaps, in the longer term, lead to new approaches for treating diseases caused by defective bicarbonate transport.

## RESULTS

Initially, we wished to compare the chloride transport ability of synthetic receptors **1-3** vs. prodigiosin **4**. The transmembrane anion transport abilities of compounds **1-4** were first evaluated by monitoring chloride efflux from unilamellar POPC (1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine) vesicles using a chloride selective electrode.<sup>31</sup> Initial studies were conducted using nitrate in the extravesicular solution. Nitrate is more hydrophobic than bicarbonate and is frequently used in experiments to assess chloride transport efficiency. The liposomes were loaded with a sodium chloride solution and suspended in a sodium nitrate solution. The carrier molecules, dissolved in a small amount (10  $\mu$ L) of DMSO, were then added to the extravesicular solution and the chloride efflux was monitored over 300 s. At the end of the experiment the vesicles were lysed by the addition of detergent and the final value was normalized to equal complete chloride efflux. Isophthalamides **1-3** proved to be potent chloride transporters using this assay (Figure 1a). Carrier loadings as low as 0.1 % molar carrier to lipid were capable of almost complete chloride efflux within 300 s, with the isopentyl-substituted isophthalamide **2** being the most active synthetic transporter under these conditions. A 0.005% molar carrier to lipid concentration of the natural product prodigiosin **4** showed similar activity to the 0.1% molar carrier to lipid concentration of the synthetic systems (Figure 1).




**Figure 1.** (a) Chloride efflux promoted upon addition of **1** (■), **2** (▲), **3** (●) (0.1 % molar carrier to lipid) and **4** (◆) (0.005 % molar carrier to lipid) to unilamellar POPC vesicles loaded with 488 mM NaCl 5 mM phosphate buffer pH 7.2 dispersed in 488 mM NaNO<sub>3</sub> 5 mM phosphate buffer pH 7.2. At t = 300 s the vesicles were lysed by addition of detergent and the final reading at t = 420 s was considered to equal 100% chloride efflux. (b) Chloride efflux studies upon addition of compounds **1-3** (0.5 % molar carrier to lipid) and **4** (0.04 % molar carrier to lipid) to vesicles composed of POPC. The vesicles contained NaCl (488 mM) and were immersed in Na<sub>2</sub>SO<sub>4</sub> (166 mM), pH 7.2 solution; at 300 s the vesicles were lysed to obtain 100% chloride efflux.

In the assay depicted in Figure 1a, the anion transport activity can occur either *via* H<sup>+</sup>/Cl<sup>-</sup> co-transport or by a Cl<sup>-</sup>/NO<sub>3</sub><sup>-</sup> exchange mechanism. To distinguish between these alternative mechanisms, we carried out the Cl<sup>-</sup> electrode transport assay while varying the anion present in the external medium. If the transport activity is the result of an anion exchange mechanism changing the external anion should impact the transport rate, whilst a H<sup>+</sup>/Cl<sup>-</sup> co-transport mechanism should not be strongly affected by the identity of the external anion to any significant degree. As depicted in Figure 1b, sulfate was used as an external anion and the transport assay was repeated by suspending the chloride-loaded vesicles in a sulfate containing external medium. As the sulfate dianion carries a higher charge and is significantly more hydrophilic than the nitrate anion, transport activity by compounds **1-4** should be greatly reduced if an anion exchange mechanism is operative. Indeed, under these conditions, with sulfate as the external anion, no chloride efflux from the liposomes was detected upon addition of **1-4**, strongly supporting a chloride/nitrate exchange (antiport) mechanism for mediating anion transport across the vesicle bilayer and additionally

demonstrating that the carriers do not release chloride by simply disrupting the lipid bilayer.


While both nitrate and bicarbonate have similar sizes and shapes, bicarbonate is significantly more hydrated than nitrate and one might expect that it would be significantly more challenging to transport bicarbonate than nitrate across a non-polar lipid bilayer.<sup>32,33</sup> Prompted by the  $\text{Cl}^-/\text{NO}_3^-$  anion exchange activity shown by **1-4**, we designed an experiment to determine whether these compounds could also facilitate bicarbonate/chloride exchange across phospholipid barriers. Chloride loaded vesicles were suspended in a sulfate containing medium. After two minutes, a solution of bicarbonate was added and chloride efflux was monitored over an additional five minutes. At the end of the experiment the vesicles were lysed to calibrate the experimental data to 100% chloride release. The results shown in Figure 2 confirmed that, as shown above in Figure 1a, negligible chloride efflux was detected in the presence of sulfate as the external anion. Addition of bicarbonate to the extravesicular solution switched on the chloride efflux promoted by **1-4**, indicating that these compounds enable chloride/bicarbonate antiport across liposomal membranes. The rate of chloride efflux was dependent on the amount of carrier added to the experiment (see supplementary information Figures S9-S12).



**Figure 2.** Chloride efflux promoted upon addition of **1** (■), **2** (▲), **3** (●) (1 % molar carrier to lipid) and **4** (◆) (0.04 % molar carrier to lipid) to unilamellar POPC vesicles loaded with 451 mM NaCl 20 mM phosphate buffer pH 7.2 dispersed in 150 mM Na<sub>2</sub>SO<sub>4</sub> 20 mM phosphate buffer pH 7.2. At t = 120 s a solution of NaHCO<sub>3</sub> was added to give a 40 mM external concentration. At t = 420 s the vesicles were lysed by addition of detergent and the final reading at t = 540 s was considered to equal 100% chloride efflux. (a) In the presence of the carrier compounds **1-4** chloride was not released from the vesicles when suspended in a sulfate solution. (b) Upon introduction of bicarbonate to the solution, chloride efflux began as one component of the chloride/bicarbonate antiport mechanism.

Under the assay conditions addition of bicarbonate induced small changes (~0.2 units) in the pH of the external medium. We therefore carried out control experiments in the presence of compounds **1-4** to rule out the possibility that chloride efflux was driven by a pH gradient. Indeed, addition of NaOH to the external medium resulted in no significant chloride efflux. Furthermore, addition of bicarbonate solutions to a suspension of vesicles without the presence of transporters **1-4** resulted in no chloride efflux.

We used  $^{13}\text{C}$  NMR spectroscopy to verify that transporters **1-4** facilitate transmembrane  $\text{HCO}_3^-/\text{Cl}^-$  anion exchange across phospholipid vesicles. We developed experiments that use paramagnetic  $\text{Mn}^{2+}$  to bleach the  $^{13}\text{C}$  NMR signal for extravesicular  $\text{H}^{13}\text{CO}_3^-$ , allowing for ready discrimination of extravesicular and intravesicular  $\text{H}^{13}\text{CO}_3^-$ . We based these paramagnetic NMR protocols on previous experiments that 1) monitored transmembrane chloride transport in liposomes by  $^{35}\text{Cl}$  NMR, <sup>34,35</sup> and 2) showed that intracellular and extracellular  $\text{H}^{13}\text{CO}_3^-$  could be distinguished in plant cells. <sup>36,37</sup> Our results show that both synthetic **3** and natural product **4**, facilitate  $\text{HCO}_3^-/\text{Cl}^-$  exchange across phospholipid membranes. Figure 3 shows data from the first set of NMR experiments conducted to illustrate transporter-mediated  $\text{HCO}_3^-/\text{Cl}^-$  exchange. This set of NMR experiments was done under similar conditions as described for the  $\text{Cl}^-$  electrode experiments depicted in Figure 1. Thus, EYPC liposomes (5  $\mu\text{m}$ ) filled with 450 mM NaCl were suspended in a sulfate solution and 50 mM  $\text{H}^{13}\text{CO}_3^-$  was added to the NMR sample. A sharp  $^{13}\text{C}$  NMR signal for extravesicular  $\text{H}^{13}\text{CO}_3^-$  was observed at  $\delta$  161 ppm. Upon addition of 0.5 mM  $\text{Mn}^{2+}$ , this signal was broadened into the baseline as the paramagnetic cation interacted with the extravesicular bicarbonate. After addition of transporters (isophthalamide **3** in Figure 3a and prodigiosin **4** in Figure 3b), a sharp  $^{13}\text{C}$  NMR signal for  $\text{H}^{13}\text{CO}_3^-$  ( $\delta$  ~161 ppm) was restored. This renewed  $^{13}\text{C}$  NMR signal must be caused by ligand-mediated transport of  $\text{HCO}_3^-$  into the liposome since the paramagnetic  $\text{Mn}^{2+}$  is impermeable to the phospholipid bilayer. Importantly, the control experiment in which DMSO was added without transporter, did not result in any restoration of  $^{13}\text{C}$  NMR signal (Figure 3c).




**Figure 3.** Representation of titration sequence and  $^{13}\text{C}$  NMR data (a-c) for monitoring transmembrane transport of  $\text{HCO}_3^-$  into  $\text{Cl}^-$ -loaded EYPC liposomes by **3** and **4**: **i**) a  $\text{NaH}^{13}\text{CO}_3$  pulse (50 mM) was added to EYPC vesicles loaded with 450 mM NaCl, 20 mM HEPES (pH 7.3) and dispersed in 150 mM  $\text{Na}_2\text{SO}_4$ , 20 mM HEPES (pH 7.3); **ii**) NMR spectra after addition of 0.5 mM  $\text{Mn}^{2+}$  (1:100  $\text{Mn}^{2+}/\text{H}^{13}\text{CO}_3^-$  ratio); **iii**) NMR spectra after addition of transporter or DMSO (**3** – 1 mol % relative to lipid, **4** – 0.1 mol %, or DMSO – 403 mol%).

Figure 4 shows data from another NMR experiment designed to verify transporter-mediated  $\text{HCO}_3^-/\text{Cl}^-$  exchange. In these experiments we monitored bicarbonate efflux from vesicles loaded with  $\text{H}^{13}\text{CO}_3^-$  upon addition of transporters **3** or **4**. Thus, EYPC vesicles filled with  $\text{H}^{13}\text{CO}_3^-$  and suspended in  $\text{Na}_2\text{SO}_4$  solution were aged overnight at  $4^\circ\text{C}$ . Two  $^{13}\text{C}$  NMR signals separated by 1 ppm ( $\delta \sim 162$  and  $\sim 161$  ppm) were observed, corresponding to separate signals for intravesicular and extravesicular  $\text{H}^{13}\text{CO}_3^-$  (Figure 4a-c). No leakage of  $\text{H}^{13}\text{CO}_3^-$  from these vesicles occurred after addition of 50 mM NaCl. A DMSO solution of the transporters was then added to give ligand-to-lipid ratios of 1 mol % for **3**, or 0.1 mol % for **4**. These transporters promote  $\text{Cl}^-/\text{H}^{13}\text{CO}_3^-$  exchange, as confirmed by observation of only the NMR signal for extravesicular  $\text{H}^{13}\text{CO}_3^-$  (Figure 4a/b). After addition of 0.5 mM  $\text{Mn}^{2+}$  (1:100  $\text{Mn}^{2+}/\text{H}^{13}\text{CO}_3^-$ ), this  $\text{H}^{13}\text{CO}_3^-$  signal was broadened into the baseline, confirming that all of the intravesicular  $\text{H}^{13}\text{CO}_3^-$  ions had been exchanged into the extravesicular milieu (Figure 4a/b). A control experiment confirmed this interpretation (Figure 4c). Thus, after addition of DMSO, the separate signals for intravesicular and extravesicular  $\text{H}^{13}\text{CO}_3^-$  remained unchanged. Addition of  $\text{Mn}^{2+}$  to this control sample erased the



extravesicular  $\text{H}^{13}\text{CO}_3^-$  signal, whereas the intravesicular  $\text{H}^{13}\text{CO}_3^-$  signal remained intact since  $\text{Mn}^{2+}$  cannot cross the lipid membrane.



**Figure 4.** Representation of the titration sequence and NMR stack plots (a-c) for monitoring the transmembrane transport of  $\text{HCO}_3^-$  ions in  $\text{H}^{13}\text{CO}_3^-$ -loaded EYPC liposomes by **3** and **4**. A 50 mM NaCl pulse was added to EYPC vesicles loaded with 100 mM  $\text{NaH}^{13}\text{CO}_3$ , 20 mM HEPES buffer (pH 7.5) and dispersed in 75 mM  $\text{Na}_2\text{SO}_4$ , 20 mM HEPES buffer (pH 7.3), and  $^{13}\text{C}$ -NMR data was acquired before (i) and after (ii) the  $\text{Cl}^-$  pulse. NMR spectra were also collected after the addition of transporter or DMSO (**3** – 1 mol%, **4** – 0.1 mol%, or DMSO – 870 mol% (10  $\mu\text{L}$ ); **iii**), followed by addition of 0.5 mM  $\text{Mn}^{2+}$  (1:100  $\text{Mn}^{2+}/\text{Cl}^-$  ratio; **iv**).

## DISCUSSION

This work identifies “small” molecules with activity as transmembrane bicarbonate carriers. The ion selective electrode assays showed that both the synthetic

isophthalamides **1-3** and the natural product prodigiosin **4** facilitate the release of encapsulated chloride from POPC phospholipid liposomes in the presence of trigonal planar oxoanions such as nitrate and bicarbonate. This efflux is produced *via* an exchange mechanism with external nitrate anions, in agreement with previous studies.<sup>16,29</sup> Replacing nitrate by the more hydrophilic sulfate in the external medium resulted in no anion transport activity as an anion exchange mechanism is not possible due to the higher hydrophilicity of the dianionic sulfate. A subsequent assay in which a second anion (bicarbonate) was added to the external sulfate medium showed that chloride transport by compounds **1-4** was restored, clear evidence that these compounds facilitate  $\text{HCO}_3^-/\text{Cl}^-$  exchange across phospholipid membranes.  $^{13}\text{C}$  NMR assays provided direct evidence for transmembrane bicarbonate transport, as the intra- and extra-vesicular carbon-13 labelled bicarbonate populations could be clearly distinguished. This NMR data, when combined with results from the  $\text{Cl}^-$ -selective electrode experiments, firmly establish that compounds **1-4** enable the transmembrane exchange of  $\text{Cl}^-/\text{HCO}_3^-$  anions.

In conclusion, we have demonstrated for the first time that “small” molecules, including simple synthetic transporters **1-3** and the natural product prodigiosin **4**, are able to mimic the natural chloride/bicarbonate exchange process that is typically mediated by membrane proteins. This is the first report that prodigiosin can facilitate chloride/bicarbonate antiport exchange, an important discovery that may present an alternative mechanism by which such compounds affect biological systems. Synthetic bicarbonate/chloride antiporters may also prove to be useful tools for biomembrane research and membrane based processes. Efforts aimed at producing improved and selective bicarbonate transporters and the investigation of their biological activity are currently underway in our laboratories.

## METHODS

**Preparation of Phospholipid Vesicles.** A chloroform solution of 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) (20 mg/mL) (Genzyme) was evaporated *in vacuo* using a rotary evaporator and the lipid film obtained was dried under high vacuum for at least 2 hours. The lipid film was rehydrated by addition of a sodium chloride solution (488 mM NaCl and 5 mM phosphate buffer, pH 7.2 or 451 mM NaCl and 20 mM phosphate buffer, pH 7.2) and followed by careful vortexing. The lipid suspension was then subjected to nine freeze-thaw cycles and twenty-nine extrusions through a 200 nm polycarbonate Nucleopore membrane using a LiposoFast Basic extruder (Avestin, Inc.) to obtain unilamellar vesicles. The vesicles were dialyzed against a  $\text{NaNO}_3$  solution (488 mM  $\text{NaNO}_3$  and 5 mM phosphate buffer, pH 7.2) or  $\text{Na}_2\text{SO}_4$  solution (150 mM  $\text{Na}_2\text{SO}_4$  and 20 mM phosphate buffer, pH 7.2) to remove unencapsulated NaCl.

**ISE Transport Assays.** Unilamellar vesicles (200 nm mean diameter) composed of POPC containing an encapsulated solution of 488 mM NaCl and 5 mM phosphate buffer pH 7.2 or 451 mM NaCl and 20 mM phosphate buffer pH 7.2, were suspended

in a solution of 488 mM NaNO<sub>3</sub> and 5 mM phosphate buffer pH 7.2, or 150 mM Na<sub>2</sub>SO<sub>4</sub> and 20 mM phosphate buffer pH 7.2, for a final lipid concentration of 1 mM. A DMSO solution of the carrier molecule, typically 10 μL to avoid influence of the solvent molecules in the assay, was added and the chloride release from vesicles was monitored using an Accumet chloride selective electrode for 7 minutes. At a time (t) = 5 min the vesicles were lysed with detergent (polyoxyethylene (8) lauryl ether) to release all chloride ions; the resulting value was considered to represent 100% release and used as such.

For the anion exchange assays in the vesicles suspended in a Na<sub>2</sub>SO<sub>4</sub> solution a solution of NaHCO<sub>3</sub> or NaNO<sub>3</sub> was added for a final concentration of 40 mM at t = 2 min. The chloride efflux was monitored for another 5 minutes and at a time (t) = 7 min the vesicles were lysed with detergent (polyoxyethylene (8) lauryl ether) to release all chloride ions; the resulting value was considered to represent 100% release and used as such.

### <sup>13</sup>C NMR Assays

**Materials.** <sup>13</sup>C NMR spectra were recorded on a Bruker DRX500 instrument operating at 125.77 MHz, with chemical shifts reported in ppm. Egg yolk phosphatidylcholine (EYPC) lipids, nuclepore® polycarbonate membranes and membrane filters were purchased from Avanti Polar Lipids. Deuterated solvents were purchased from Cambridge Isotope Laboratories. Prodigiosin was a gift from the Development Therapeutics Program at the National Cancer Institute, U. S. National Institutes of Health. All other chemicals were purchased from Sigma, Aldrich, Fisher, Fluka, or Acros and used without further purification.

**Liposome Preparation for <sup>13</sup>C NMR Assays.** Giant EYPC liposomes (5 μm) were prepared by evaporating a chloroform solution of EYPC (20 mg/mL) under reduced pressure, resulting in a thin film that was dried *in vacuo* overnight. The liposomes were then formed by rehydrating the lipid film with either 450 mM NaCl in 20 mM HEPES (pH 7.3) buffer (for the liposomes described in Figure 5 of main text, i.e., the Cl<sup>-</sup>-loaded liposomes), or 100 mM NaH<sup>13</sup>CO<sub>3</sub> in 20 mM HEPES (pH 7.5) buffer (for the liposomes described in Figure 4, i.e., the HCO<sub>3</sub><sup>-</sup>-loaded liposomes). Buffer solutions were prepared in a 9:1 H<sub>2</sub>O/D<sub>2</sub>O mixture. After 5 freeze/thaw cycles, the liposomes were extruded through a 5 μm polycarbonate membrane 41 times at room temperature using a high-pressure mini-extruder (Avanti). For the Cl<sup>-</sup>-loaded liposomes, the giant liposome suspension obtained after extrusion was used without further purification in the <sup>13</sup>C NMR transport assays. However, for the HCO<sub>3</sub><sup>-</sup>-loaded liposomes, the resulting giant liposome suspension was separated from extravesicular NaH<sup>13</sup>CO<sub>3</sub> by size exclusion chromatography (SEC) (stationary phase: Sephadex G-25, mobile phase: 9:1 H<sub>2</sub>O/D<sub>2</sub>O, 20 mM HEPES, pH 7.3, 75 mM Na<sub>2</sub>SO<sub>4</sub>). The 30 mL suspension (NaH<sup>13</sup>CO<sub>3</sub> inside, Na<sub>2</sub>SO<sub>4</sub> outside) collected was centrifuged (Eppendorf Centrifuge 5804R) at 10,000 rpm for 30 minutes followed by the removal of the non-liposome containing buffer. The recovered giant liposome suspension was then diluted with the 75 mM Na<sub>2</sub>SO<sub>4</sub> mobile phase buffer and used directly in the <sup>13</sup>C

NMR transport assays. The stock concentrations obtained for the liposomes were 90.3 mM for the Cl<sup>-</sup>-loaded liposomes (assuming 100% lipid retention after extrusion) and 66.6 mM for the HCO<sub>3</sub><sup>-</sup>-loaded liposomes (assuming 80% lipid retention after gel filtration) respectively.

**Bicarbonate Transport in Cl<sup>-</sup>-Loaded Liposomes Monitored by <sup>13</sup>C NMR.** <sup>13</sup>C NMR spectra were recorded using a Bruker DRX500 spectrometer with a 5 mm broad band probe operating at 125.77 MHz, with chemical shifts reported in ppm. The instrument was locked on 9:1 H<sub>2</sub>O/D<sub>2</sub>O. Experimental conditions were: acquisition time, 0.93 s; spectrum width, 35211 Hz; 90° pulse width, 6.70 μs; relaxation delay, 0.2 s; number of scans, 160; temperature, 27 °C. For each experiment, 230 μL of stock (90.3 mM) liposome solution was mixed with 340 μL of 150 mM Na<sub>2</sub>SO<sub>4</sub> in 20 mM HEPES (pH 7.3) buffer in a 5 mm NMR tube to give a liposome suspension containing NaCl inside, and Na<sub>2</sub>SO<sub>4</sub> outside. A NaH<sup>13</sup>CO<sub>3</sub> pulse was then added to the mixture to give 35 mM and 50 mM final concentrations of liposome and H<sup>13</sup>CO<sub>3</sub><sup>-</sup> respectively. The <sup>13</sup>C NMR of the preceding liposome mixture (NaCl inside, Na<sub>2</sub>SO<sub>4</sub> and NaH<sup>13</sup>CO<sub>3</sub> outside) was then taken. After data acquisition, a solution of MnCl<sub>2</sub> was added to give a final Mn<sup>2+</sup> concentration of 0.5 mM (1:100 Mn<sup>2+</sup>/H<sup>13</sup>CO<sub>3</sub><sup>-</sup> ratio), and immediately followed by another set of data acquisition. Finally, a final set of <sup>13</sup>C NMR data was collected after the addition of a solution of the ligand (**1**, **3** or **4**) or DMSO to the mixture. Isophthalamides **1** and **3** were added in a 1 mol% ligand-to-lipid ratio, while prodigiosin **4** was added in a 0.1 mol% ligand-to-lipid ratio. For the DMSO control, 6 μL of the solvent was added corresponding to a 403 mol% DMSO-to-lipid ratio.

**Bicarbonate Transport in HCO<sub>3</sub><sup>-</sup>-Loaded Liposomes Monitored by <sup>13</sup>C NMR.** Instrument details are the same as described above for the Cl<sup>-</sup>-loaded liposomes. The instrument was locked on 9:1 H<sub>2</sub>O/D<sub>2</sub>O. Experimental conditions were: acquisition time, 0.93 s; spectrum width, 35211 Hz; 90° pulse width, 6.70 μs; relaxation delay, 0.2 s; number of scans, 196; temperature, 27 °C. For each experiment, an initial <sup>13</sup>C NMR spectrum of 520 μL of the giant liposome solution was acquired. Then, a NaCl pulse resulting in final extravesicular concentrations of 58 mM and 50 mM for the giant liposomes and Cl<sup>-</sup> respectively was added to the NMR tube. The <sup>13</sup>C NMR of the preceding liposome mixture (NaH<sup>13</sup>CO<sub>3</sub> inside, Na<sub>2</sub>SO<sub>4</sub> and NaCl outside) was taken followed by the addition of a solution of the ligand (**1**, **3** or **4**) or DMSO to the cocktail. Again, isophthalamides **1** and **3** were added in a 1 mol% ligand-to-lipid ratio, while prodigiosin **4** was added in a 0.1 mol% ligand-to-lipid ratio. For the DMSO control, 10 μL of the solvent was added corresponding to an 870 mol% DMSO-to-lipid ratio. A <sup>13</sup>C NMR spectrum of the ligand-containing cocktail was then acquired before and after the addition of a solution of MnCl<sub>2</sub> (0.5 mM final Mn<sup>2+</sup> concentration corresponding to 1:100 Mn<sup>2+</sup>/Cl<sup>-</sup> ratio).

## ACKNOWLEDGMENTS

J. D. and P. A. G. thank the NSF/EPSCRC for funding this research through the "International Collaboration in Chemistry" program under CHE-0822244 and EP/G002576/1. R. Q. thanks the Spanish Ministerio de Educación y Ciencia for support "Jose Castillejo" mobility grant and "Ramon y Cajal" contract. T.T. thanks DGI of Spain (CTQ2006-15456-C04-04BQU) for funding. O.A.O thanks the University of Maryland Graduate School for the Mabel Spencer Fellowship award.

## AUTHOR CONTRIBUTIONS

J.T.D., P.A.G. and R.Q conceived this project, experiments, analyzed data and prepared the manuscript; O.A.O. and R.Q. conducted experiments and analyzed data; P.P. and T.T. contributed reagents, materials, analysis tools and supervised the synthesis; J.C.I.S. and R.Q. synthesized new compounds.

## References

1. Cleland, W. W. *et al.* Mechanism of Rubisco: The Carbamate as General Base *Chem. Rev.* 98, 549-561 (1998).
2. Hempling, H.G. Intracellular Water and the Regulation of Cell Volume and pH in *Principles of Medical Biology* Eds: Bittar E.E. & N. Bittar Elsevier 1996.
3. Lipscomb, W. N. & Strater, N. Recent Advances in Zinc Enzymology. *Chem. Rev.* 96, 2375-2433 (1996).
4. Florman, H. M., Jungnickel & M. K.; Sutton, K. A. What can we learn about fertilization from cystic fibrosis? *Proc. Nat. Acad. Sci.-USA* 104, 11123-11124 (2007).
5. Fujinaga, J., Loisel, F. B. & Casey, J. R. Transport activity of chimaeric AE2-AE3 chloride/bicarbonate anion exchange proteins *Biochem. J.* 371, 687-696 (2003).
6. Choi, J. Y. *et al.* Aberrant CFTR-dependent HCO<sub>3</sub><sup>-</sup> transport in mutations associated with cystic fibrosis *Nature* 410, 94-97 (2001).
7. Casey, J. R. Why Bicarbonate? *Biochem. Cell Biol.* 84, 930-939 (2006)
8. Koropatkin, N. M. *et al.* The Structure of a Cyanobacterial Bicarbonate Transport Protein, CmpA *J. Biol. Chem.* 282, 2606-2614 (2007).
9. Davis, A. P., Shepard, D. N. & Smith, B. D. Development of synthetic membrane transporters for anions *Chem. Soc. Rev.* 36, 348-357 (2007)
10. Okunola, O. A. *et al.* Membrane-active calixarenes: toward 'gating' transmembrane anion transport *Tetrahedron* 63, 10743-10750 (2007).
11. You, L. & Gokel, G. W. Fluorescent, synthetic amphiphilic heptapeptide anion transporters: Evidence for self-assembly and membrane localization in liposomes *Chem. Eur. J.* 14, 5861-5870 (2008).
12. Izzo, I. *et al.* Cationic calix[4]arenes as anion-selective ionophores *Chem. Commun.* 2986-2988 (2008).

13. Whitmarsh, S.D. *et al.* Cationic cyclocholamides; toroidal facial amphiphiles with potential for anion transport *Chem. Commun.* 3669-3671 (2008).
14. Gale, P.A. *et al.* Co-transport of H<sup>+</sup>/Cl<sup>-</sup> by a synthetic prodigiosin mimic, *Chem. Commun.* 3773-3775 (2005).
15. Gale, P. A. *et al.* Conformational control of HCl co-transport: imidazole functionalised isophthalamide vs. 2,6-dicarboxamidopyridine *Chem. Commun.* 1736-1738 (2007).
16. Santacroce, P. V. *et al.* Conformational Control of Transmembrane Cl<sup>-</sup> Transport *J. Am. Chem. Soc.* 129, 1886-1887 (2007).
17. Kavallieratos, K.; Bertao, C.M. & Crabtree R.H. Hydrogen bonding in anion recognition: A family of versatile, non-preorganized neutral and acyclic receptors *J. Org. Chem.* 64, 1675-1683 (1999).
18. Fürstner, A. Chemistry and biology of roseophilin and the prodigiosin alkaloids: A survey of the last 2500 years *Angew. Chem., Int. Ed.*, 42, 3582-3603 (2003).
19. Gerber, N.N. Prodigiosin-like pigments *Crit. Rev. Microbiol.* 3, 469-485 (1975).
20. Bennett, J.W. & Bentley R. Seeing red: The story of prodigiosin *Adv. Appl. Microbiol.* 47, 1-32 (2000).
21. Williamson, N.R. *et al.* Anticancer and immunosuppressive properties of bacterial prodiginines, *Future Microbiol.*, 2, 605–618 (2007).
22. Nguyen, M. *et al.* Small molecule obatoclax (GX15-070) antagonizes MCL-1 and overcomes MCL-1-mediated resistance to apoptosis, *Proc. Natl. Acad. Sci. U. S. A.*, 104, 19512–19517 (2007).
23. Yamamoto, C. *et al.* Cycloprodigiosin hydrochloride, a new H<sup>+</sup>/Cl<sup>-</sup> symporter, induces apoptosis in human and rat hepatocellular cancer cell lines *in vitro* and inhibits the growth of hepatocellular carcinoma xenografts in nude mice *Hepatology* 30, 894-902 (1999).
24. Tanigaki, K. *et al.* BE-18591 as a new H<sup>+</sup>/Cl<sup>-</sup> symport ionophore that inhibits immunoproliferation and gastritis *FEBS Lett.* 524, 37-42 (2002).
25. T. Sato, *et al.*, Prodigiosins as a New Group of H<sup>+</sup>/Cl<sup>-</sup> Symporters That Uncouple Proton Translocators *J. Biol. Chem.*, 273, 21455-21462 (1998).
26. Gottlieb, R.A. *et al.* Apoptosis induced in Jurkat cells by several agents is preceded by intracellular acidification, *Proc. Natl. Acad. Sci. U.S.A.* 93, 654-658 (1996).
27. S. Ohkuma, *et al.* Prodigiosins uncouple vacuolar-type ATPase through H<sup>+</sup>/Cl<sup>-</sup> symport *Biochem. J.* 334, 731-741 (1998).
28. Sessler, J.L. *et al.* Synthesis, anion-binding properties, and *in vitro* anticancer activity of prodigiosin analogues, *Angew. Chem. Int. Ed.* 44, 5989-5992 (2005).
29. Seganish, J.L. & Davis, J.T. Prodigiosin is a chloride carrier that can function as an anion exchanger, *Chem. Commun.*, 5781-5783 (2005).

30. Diaz, R.I.S. *et al.* Chloride anion transport and copper-mediated DNA cleavage by C-ring functionalized prodigiosenes, *Chem. Commun.* 2701-2703 (2007).
31. Koulov, A.V. *et al.* Chloride transport across vesicle and cell membranes by steroid-based receptors, *Angew. Chem. Int. Ed.* 42, 4931-4933 (2003).
32. Hofmeister, F. Zur Lehre von der Wirkung der Salze, Zweite Mittheilung, *Arch. Exp. Pathol. Pharmacol.* 24, 247-260 (1888).
33. Steinle, E.D., Schaller, U. & Meyerhoff, M.E. Response Characteristics of Anion-Selective Polymer membrane Electrodes Based on Gallium(III), Indium(III) and Thallium(III) Porphyrins, *Anal. Sciences*, 14, 79-84 (1998).
34. Riddell, F.G. *et al.* Chloride transport through model biological membranes studied by  $^{35}\text{Cl}$  NMR, *Chem. Commun.* 74-76 (1990).
35. Riddell, F.G. & Zhou, Z.  $\text{Mn}^{2+}$  as a contrast reagent for NMR studies of  $^{35}\text{Cl}^-$  and  $^{81}\text{Br}^-$  transport through model biological membranes, *J. Inorg. Biochem.* 55, 279-293 (1994).
36. Gout, E. *et al.*  $^{13}\text{C}$  nuclear magnetic resonance studies of malate and citrate synthesis and compartmentation in higher plant cells,  $^{81}\text{Br}^-$  transport through model biological membranes, *J. Biol. Chem.* 268, 3986-3992 (1993).
37. Chang, K. & Roberts, J.K.M. Quantitation of rates of transport, metabolic fluxes, and cytoplasmic levels of inorganic carbon in maize root tips during  $\text{K}^+$  ion uptake, *Plant Physiol.* 99, 291-297 (1992).