

Utilization of agro-industrial by-products as substrates for dextranucrase production by *Leuconostoc mesenteroides* T3: process optimization using response surface methodology

Miona G. Miljković¹, Slađana Z. Davidović¹, Aleksandra P. Djukić-Vuković¹, Mila V. Ilić², Milica B. Simović¹, Mirjana D. Rajilić-Stojanović¹ and Suzana I. Dimitrijević-Branković¹

¹University of Belgrade, Faculty of Technology and Metallurgy, Department of Biochemical Engineering and Biotechnology, Karnegijeva 4, Belgrade, Serbia

²University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Njegoševa 12, 11000 Belgrade, Serbia

Abstract

Dextranucrase (DS) is a glucosyltransferase (E. C. 2.4.1.5) that catalyzes the transfer of glucosyl residues from sucrose to dextran polymer and liberates fructose. This enzyme is associated with a wide application range of dextran and oligosaccharides. DS production by *Leuconostoc mesenteroides* T3 was optimized using a Central Composite Design under the Response Surface Methodology. Three variables were chosen for optimization: distillery stillage, sucrose and manganese concentration. The results showed that sucrose and manganese concentrations had a positive linear effect on DS production while all variable interactions (stillage-manganese, stillage-sucrose, and sucrose-manganese) had significant influences on the DS production. The maximal DS yield of 3.391 ± 0.131 U cm⁻³, was obtained in the medium with 64.33 % distillery stillage concentration, 5.30% sucrose concentration and 0.022 % manganese concentration. Our study revealed the potential of distillery stillage combined with sugar beet molasses, supplemented with sucrose and manganese to be employed as a valuable medium growth for lactic acid bacteria and production of DS. Also, taking into consideration the origin of the substrates, utilization of industrial by-products in this way has a great environmental relevance and is in accordance with circular economy.

Keywords: headspace; gas chromatography; alcoholic beverages; cosmetics.

Available on-line at the Journal web address: <http://www.ache.org.rs/HI/>

ORIGINAL SCIENTIFIC PAPER

UDC: 636:087.2:577.151.5:
681.5.015.23

Hem. Ind. 75(3) 135-146 (2021)

1. INTRODUCTION

Dextranucrase (DS) is a glucosyltransferase (E. C. 2.4.1.5) that catalyzes the transfer of glucosyl residues from sucrose to dextran polymer and liberates fructose [1]. This enzyme also catalyzes the so-called acceptor reaction, where in the presence of suitable low molecular weight molecules, for example maltose, the transfer of glucosyl units is redirected from dextran to oligosaccharide synthesis [2]. When water molecule itself is an acceptor, only the reaction of sucrose hydrolysis to glucose and fructose takes place. DS is an extracellular enzyme produced by several species belonging to the genera *Lactobacillus*, *Leuconostoc* and *Streptococcus* [3]. It has been shown that the DS expression is constitutive in *Streptococcus* strains, whereas in *Leuconostoc* spp. DSs can only be produced upon sucrose induction [4,5]. DS is commercially employed in dextran production. Dextran and their derivatives have been profusely used in food, clinical, pharmaceutical, fine chemicals, cosmetics and agricultural industries [6, 7]. Worldwide market for dextran is expected to reach 220 million US\$ in 2024 with the annual rate of 4,2 % [8]. To date, different combinations of cultivation parameters for DS producing strains have been used to obtain the maximal activity of DS.

Corresponding author: Slađana Davidović, University of Belgrade, Faculty of Technology and Metallurgy, Department of Biochemical Engineering and Biotechnology, Karnegijeva 4, 11000 Belgrade, Serbia

E-mail: sdavidovic@tmf.bg.ac.rs

Paper received: 10 July 2020

Paper accepted: 28 April 2021

<https://doi.org/10.2298/HEMIND200710015M>



Various industrial by-products and wastes have a great potential for utilization as raw materials for production of biomolecules, especially enzymes. Taking into consideration that production of enzymes is expensive, mainly due to the high price of the medium and the substrates for microbial fermentations, current challenges in the production of enzymes include testing the possibilities to use wastes or low-cost materials. Several by-products, including sugar beet molasses [9], carob pod extract and cheese whey [10], and cashew apple juice [11] have already been studied as low-cost substrates for the DS production.

Distillery stillage, also termed distillery wastewater, is the major by-product from the distillation of ethanol following fermentation of carbohydrates. The production of bioethanol from biomass, whether from sugar crops, starch crops, dairy products or cellulosic materials, at the same time results in the production of stillage, which exhibits a considerable pollution problem [12]. An average amount of stillage produced in the bioethanol process is approximately 13 L per L of bioethanol [13]. Taking into account a high BOD₅ values and low pH values, its storage represents an important ecological problem in the industrial facilities and the high costs of treatment prior to disposal into watercourses seriously affect the viability and profitability of the process [14]. However, due to its complexity and origin, the stillage from bioethanol production on starch substrates (corn, rye, wheat, potato, etc.) could be a valuable source of nitrogen, vitamins and minerals, which are necessary for the growth of microorganisms [13].

Sugar beet molasses is a by-product of the sugar industry. It is especially attractive, not only because of its low price but also because of the presence of a number of components including minerals, organic compounds and vitamins, which are very useful for the fermentation process [15]. Molasses contains up to 54 % of sugars [16] among which sucrose is the most abundant followed by small quantities of glucose, fructose and raffinose [17]. It is also a valuable source of growth substances such as pantothenic acid, inositol, trace elements, and, to a lesser extent, biotin [18]. Serbia has significant surpluses of sugar beet and in 2011 the export of sugar beet molasses was 59.655 t [19]. Based on its properties, molasses can be considered as a promising starting material or substrate for microbial fermentation [20].

Certain metallic ions like Mg²⁺, Ca²⁺, and especially Mn²⁺ are essential for DS production and bacterial growth according to literature [21,22]. *Leuconostoc* spp. are known to be micro-aerophilic microorganisms [1] and MnSO₄ decreases the oxygen toxicity in the *L. mesenteroides* [23].

The Response Surface Methodology (RSM) represents a useful tool for studying the effects of several factors on one or more responses by varying them simultaneously. In addition, the optimal response is calculated based on the experimental data obtained from a limited number of experiments [24]. RSM has been applied in many areas of biotechnology such as optimization of culture medium, enzyme synthesis, aqueous two-phase separation of proteins, glucan production, *etc.* [25]. Considering possible variations in composition of different agro-industrial wastes, it is necessary to do composition screening (carbon and nitrogen determination) every time when a new batch is obtained. It is also very important to know which raw material is the basis of distillery stillage. For example, bread stillage has significantly more nitrogen than potato stillage [26]. RSM is therefore a useful tool for optimization of a media based on agro-industrial wastes for bacterial growth.

The objective of this study was to test two cheap and abundant by-products from bioethanol production and sugar industry for DS production by *L. mesenteroides* T3. The molasses and distillery stillage as cheap agro-industrial by-products were supplemented with sucrose and manganese to enhance the DS production. The Central Composite Design (CCD) under RSM was used for optimization of fermentation parameters: concentrations of stillage, sucrose and manganese to obtain a maximal DS activity.

2. EXPERIMENTAL

2. 1. Microorganism for dextransucrase production

The microorganism used in this study is *Leuconostoc mesenteroides* T3 natural isolate from water kefir grain, identified as *L. mesenteroides* as described earlier [27].

2. 2. Inoculum and medium preparation for dextransucrase production

For inoculum preparation the organism was grown in slightly modified medium reported by Tsuchiya *et al.* [28] and it contained: sucrose (Lach-Ner, Czech Republic), 40.0 g dm⁻³; yeast extract (Torlak, Serbia), 20.0 g dm⁻³; K₂HPO₄ (Centrohem, Serbia), 20.0 g dm⁻³; MgSO₄·7H₂O (Kemika, Croatia), 0.2 g dm⁻³; MnSO₄·H₂O (Kemika, Croatia), 0.01 g dm⁻³; NaCl (Zorka, Serbia), 0.01 g dm⁻³; CaCl₂ (Lach-Ner, Czech Republic), 0.02 g dm⁻³; FeSO₄·7H₂O (Zorka, Serbia), 0.01 g dm⁻³. All chemicals were of analytical grade. The medium pH was adjusted to 6.9 by using orthophosphoric acid. Erlenmeyer flask cultures were grown for 16h at 23°C under static conditions.

Sugar beet molasses and distillery stillage were obtained from Swan lake d.o.o, Kovin, Serbia and Reahem d.o.o., Srbobran, Serbia, respectively. Fermentation medium was prepared with the fixed concentration of molasses (2.5 wt.%) that was previously defined as optimal for DS production by the same strain [29]. Stillage was added to the molasses medium at different concentrations according to the values given by RSM.

2. 3. Fermentations

All experiments were performed in 100 cm³ Erlenmeyer flasks with a medium volume of 20 cm³. Molasses in 2.5 wt.% concentration was used as a liquid medium for fermentations. In each flask with molasses, stillage was added, separately, in a defined concentration range (15-95 % v/v). Sucrose as a main substrate for DS production was also implemented into the medium in the range from 0-6 % (w/v). Manganese ion was varied in the range from 0-0.024 % (w/v). K₂HPO₄ was added as a buffer substance in a concentration of 20 mM. After sterilization at 121 °C for 20 min, an overnight bacterial culture was inoculated into fresh medium at 23 °C for 12 h under shaking (180 rpm). The culture medium was centrifuged at 6000g for 15 min to remove the cells. The crude cell-free supernatant was analyzed for DS activity.

2. 4. Analytical methods

The dry matter content was determined by a standard drying method in an oven at 105°C to constant mass [24]. The total nitrogen content in molasses was estimated by the Kjeldahl method [24]. The protein content for stillage was estimated by the Kjeldahl method as the total nitrogen and multiplied by the factor of 6.25 [30]. In order to determine the concentration of reducing sugars, the sample solution was hydrolyzed by HCl at 100°C for 10 min and neutralized with NaOH solution. Then, the concentration of reducing sugars was estimated by the 3,5-dinitrosalicylic acid method (DNS) [31]. The metal contents in the liquid distillery stillage and molasses were determined by atomic absorption spectroscopy (Perkin Elmer Analyst 200, Waltham, USA).

The DS activity assay was carried out in a 450 mm³ reaction mixture containing 10 % (w/v) sucrose, 20 mM sodium acetate buffer (pH 5.4) and 50 mm³ cell free supernatant at 30 °C for 15 min. The enzyme activity was determined by measuring the concentration of released reducing sugars by the DNS method, using fructose as a standard [31]. The absorbance was measured at 540 nm by using a spectrophotometer (Ultrospec 3300 pro, Amersham Biosciences). One unit of DS activity was defined as the amount of enzyme releasing 1 μmol of reducing sugars per minute.

2. 5. Experimental design

Based on preliminary "one variable at the time" experiments (data not shown) a CCD was chosen to examine the effects of three independent variables: distillery stillage concentration (A), sucrose concentration (B) and manganese ion concentration (C) within the defined ranges that favored optimal feedback of the DS production response. Each factor in this design was coded in five different levels (Table 1). The data obtained from CCD were analyzed by multiple regressions to fit to a second-order polynomial regression model containing the coefficients of linear, quadratic, and two factor interaction effects.

The model equation of response (Y) on three independent variables (X1, X2 and X3) is given in the following equation:

$$Y = \beta_0 + \beta_1X1 + \beta_2X2 + \beta_3X3 + \beta_{12}X1X2 + \beta_{13}X1X3 + \beta_{23}X2X3 + \beta_{11}X1^2 + \beta_{22}X2^2 + \beta_{33}X3^2 \quad (1)$$

where Y (DS activity, U cm⁻³) is the dependent variable or predicted response associated with each factor level combination; X1 (distillery stillage concentration, % (v/v)), X2 (sucrose concentration, % (w/v)), and X3 (manganese ion



concentration, % (w/v)); β_0 is the intercept term; β_1 , β_2 and β_3 are the linear effects (main effect); β_{11} , β_{22} and β_{33} are the quadratic effects; and β_{12} , β_{13} and β_{23} are the interaction effects.

Table 1. Experimental ranges of the independent variables in the experimental design

Factors	-1	0	1	Axial (- α)	Axial (+ α)
A: Distillery stillage concentration, % (v/v)	35	55	75	15	95
B: Sucrose concentration, % (w/v)	1.5	3	4.5	0	6
C: Mn ²⁺ concentration, % (w/v)	0.006	0.012	0.018	0	0.024

The RSM was applied by using a statistical package, Design-Expert (Version 8, Stat-Ease, Inc., Minneapolis, US).

3. RESULTS AND DISCUSSION

3. 1. Evaluation of substrates for dextransucrase production

The chemical composition and metal ion contents in molasses and distillery stillage used for DS production in this study is presented in Tables 2 and 3, respectively.

Table 2. Chemical compositions of sugar beet molasses and distillery stillage

Parameter	Sugar beet molasses ^a	Distillery stillage ^a
Content of dry matter, wt.%	77.42 ± 0.89	11.55 ± 0.30
Content of total sugars, wt.%	54.80 ± 0.51	9.74 ± 0.04
Content of total nitrogen/ content of total protein, wt.%	1.48 ± 0.16	58.50 ± 0.12

^aValues represent means ± standard deviation calculated from three determinations.

Table 3. Metal ions contents in sugar beet molasses and distillery stillage

Metals	Concentration ± SD, mg dm ⁻³		
	Sugar beet molasses ^a	Distillery stillage ^a	Optimal metal content for LAB [32]
Mg	340.00 ± 0.01	155.00 ± 0.01	480-972
Mn	9.52 ± 0.02	1.34 ± 0.01	≤110
Ca	115000 ^b	210.55 ± 0.03	≤8000
Fe	29.38 ± 0.04	3.02 ± 0.04	≤4
Zn	9.06 ± 0.03	3.78 ± 0.01	<20
Na	135000 ^b	398.02 ± 0.01	≤4000
Cu	0.80 ± 0.01	0.22 ± 0.05	<19

^aAll values represent means ± standard deviation calculated from three determinations, except

^bwhich represents means calculated from three determinations without standard deviation.

Composition of molasses used in this study (Table 2) is in accordance with results reported in the literature [16]. Typically, molasses contains around 50 % of sucrose [17], 30 % non-sugar compounds and around 20 % water. In order to be used in the fermentation, molasses was diluted to the desired sugar content. Dilution of molasses has multiple advantages. Because of the high concentration of sugars, undiluted molasses acts inhibitory on bacterial growth. When molasses is diluted, the concentration of inhibitory components for bacterial growth, fermentation and/or production of DS also decreases. Additionally, dilution also decreases the concentration of salts, normally present in molasses but which can be harmful for microorganisms (Table 3). Among different studied concentrations of molasses (from 1 to 5 %) the best DS production was achieved at the concentration of 2.5 %, which corresponds to the sucrose concentration of 1 % in the fermentation media [29].

Analyses of molasses samples from different sugar factories over several seasons have shown that the total nitrogen content in molasses may vary considerably. According to the literature data, the total nitrogen content in molasses varies in the range of 0.8 to 2.2 %, calculated on the total mass of molasses [33]. This corresponds well with to results (Table 2). However, it has been determined that the betaine content is constant and in the range 33 to 43 % of total nitrogen [17,33]. Since microorganisms cannot use betaine in their metabolic pathways, the content of amino acids is a better criterion for assessing the suitability of molasses for fermentation. These amino acids are easily assimilated by

microorganisms. Diluted molasses was previously used for the growth of *L. mesenteroides*. The content of amino acids in 40-fold diluted molasses was sufficient for the growth of *L. mesenteroides* T3 but still very low and supplementation was needed for enhancement of the DS production [29]. Under these conditions, the contribution of molasses in the total nitrogen content in media is negligible and the origin of the molasses does not play a significant role.

On the other hand, a relatively high amount of proteins in distillery stillage (more than 50 % of dry matter, Table 2) suggests that it could be suitable as a substrate for growth of lactic acid bacteria (LAB). In a combined substrate based on distillery stillage and molasses, distillery stillage primarily acts as an additional source of α -amino nitrogen, which is of great importance since LAB are nutritionally demanding microorganisms, primarily in terms of organic nitrogen sources, such as free amino acids and peptides.

The presence and contents of metals in mixtures of distillery stillage and molasses after appropriate dilution are in correlation with the requirements of LAB [32] and are also below the inhibitory values. Chemical composition of mixtures of molasses and distillery stillage can provide necessary nutrients and fermentable sugars for the growth of *L. mesenteroides* T3 but for the enhancement of DS production it is necessary to add sucrose and Mn^{2+} [34]. In order to obtain the highest possible DS activity, the influence of different concentrations of sucrose and manganese together with the effects of different concentrations of distillery stillage were investigated by RSM.

3. 2. Fitting the process variables

A total of 20 randomized experiments, including six replicates as the center points were carried out according to the experimental design matrix (Table 4) derived from an optimal design for DS production.

Table 4. The design matrix and corresponding responses

Run	Independent variables			Response
	A / %	B / %	C / %	Y / U cm ⁻³
1	55	3.0	0.024	2.322
2	35	4.5	0.018	2.544
3	35	1.5	0.006	1.743
4	55	3.0	0.000	1.862
5	55	3.0	0.012	2.172
6	55	3.0	0.012	2.179
7	95	3.0	0.012	1.485
8	55	3.0	0.012	2.124
9	55	3.0	0.012	2.201
10	75	4.5	0.006	2.054
11	75	1.5	0.018	1.551
12	35	4.5	0.006	2.176
13	55	0.0	0.012	1.062
14	75	1.5	0.006	1.192
15	35	1.5	0.018	1.506
16	15	3.0	0.012	1.597
17	55	3.0	0.012	2.001
18	55	3.0	0.012	1.98
19	75	4.5	0.018	3.077
20	55	6.0	0.012	2.664

A - stillage concentration; B - sucrose concentration; C - manganese concentration; Y - DS activity

For the three examined factors, the CCD model efficiently designed a second order response fit for the surface. The quadratic model was found to be the most suitable model. The statistical significance of the regression model was evaluated by the analysis of variance (ANOVA) (Table 5). For regression analysis, the model was modified by removing the effect of non-significant factors by using backward reduction and the quadratic equation that predicts the maximum yield of DS production:

$$Y = 2.13 - 0.02A + 0.44B + 0.15C + 0.11AB + 0.16AC + 0.16BC - 0.14A^2 - 0.055B^2 \quad (2)$$



where Y (DS activity, U/ml) is the response and A (stillage concentration, %), B (sucrose concentration, %) and C (Mn^{2+} concentration, %) were independent variables while AB , AC and BC present interactions between variables A , B and C .

Table 5. The analysis of variance (ANOVA) for the quadratic model presented by Eq. (2)

	F-value	p-value Prob > F
Model	50.26	< 0.0001 ^a
A	0.57	0.4673 ^b
B	278.22	< 0.0001 ^a
C	32.98	0.0001 ^a
AB	9.37	0.0108 ^a
AC	17.44	0.0015 ^a
BC	17.94	0.0014 ^a
A ²	43.14	< 0.0001 ^a
B ²	7.18	0.0214 ^a
Lack of fit	1.41	0.3619 ^b
R-Squared	0.9734	
Adjusted R-squared	0.9540	
Predicted R-squared	0.8755	
C.V.%	5.36	
Adequate precision	26.981	

^aSignificant coefficient ($P < 0.05$); ^b Non-significant coefficient

As it can be seen in Table 5, the significant factors that influence the response and have a p-value (Prob > F) < 0.05 were B and C , the quadratic coefficients of A and B , as well as the interactions AB , AC and BC . Adequacy of the model for predicting the DS production can be indicated by the non-significant F-value for the lack of fit (1.41) compared to the pure error. The following determination coefficients: R-squared, adjusted R-squared and predicted R-squared were calculated to check the fit of the model. The obtained values of R-squared coefficients were close to 1 which showed a good correlation between the predicted and observed values (Fig. 1A). The actual values were measured response data for a particular run, and the predicted values were evaluated from the model. The adequate precision value of 26.981 was greater than 4, which indicates that the signal was adequate. The value of the coefficient of variation (C.V.) of 5.36 indicated a high degree of precision and reliability of the experimental values, suggesting that the model was reliable and reproducible [35].




Fig. 1. Plots of: (A) the relationship between the experimental and predicted values for the DS production and (B) the perturbation of all the variables

3. 3. Influence of process variables on DS production

The presence of carbon and nitrogen sources is necessary for the bacterial growth and the synthesis of enzymes. In preliminary investigations, medium with molasses and stillage was compared with medium supplemented only with molasses, proving that the combined medium was a better substrate for bacterial growth reaching higher DS activity, and thus, statistical optimization of this medium was performed.

The influence of three process variables on the DS production of *L. mesenteroides* T3 was examined. Our preliminary investigations [34] on the influence of temperature on the enzyme activity showed that the optimal temperature was 23 °C for the maximal enzyme production, and thus this temperature was fixed in further experiments.

The DS activity in fermentation medium obtained under the tested conditions was in the range from 1.062 to 3.077 U cm⁻³. According to the analysis of the experimental data and derived regression model, it can be concluded that two of three linear regression coefficients (β_2 and β_3) are significant (Table 5 and Eq. (2)). On the other hand, two quadratic regression coefficients (β_{11} and β_{22}), are significant and negative, and therefore, the influence of corresponding parameters: stillage and sucrose concentration on the DS production can be described as a quadratic function with a maximum value. Moreover, all interactions between the examined parameters proved to be significant and positive. The significance of each coefficient was determined by p-values which are listed in Table 5. The influence of different variables on the DS production was in the following order: sucrose concentration (B) > Mn²⁺ concentration (C) > stillage concentration (A) (Table 5). Interactions between the stillage concentration and Mn²⁺ concentration, AC, and the sucrose concentration and Mn²⁺ concentration, BC, were of higher significance than the interactions between the stillage concentration and sucrose concentration AB.

The main advantage of the response surface methodology is the possibility to evaluate interactions between tested variables and define the optimum values of the variables such that the response is maximized. The sucrose concentration is the most significant factor which positively affected the DS production. With the increase in sucrose concentration up to 6 %, the DS production increased. In conducted experiments, it was observed that the addition of sucrose at a concentration of 3.0 % to the production medium led to an increase in the DS production by approximately 50 %, when stillage and Mn²⁺ concentrations were maintained constant. The highest enzyme activity of 3.077 U cm⁻³ (according to the CCD model) was achieved in the medium with the addition of 4.5 % of sucrose. But according to the optimal conditions for the DS production, the maximum enzyme activity was achieved when sucrose is added at the concentration of 5.30 % to provide the total sucrose concentration of 7 % in the fermentation medium. Similar sucrose concentration appears to be optimal for DS synthesis by other dextran-producing strains such as *L. mesenteroides* NRRL-B640, as seen in other studies [36]. In order to visualize influence of the independent variables (A, B and C) on the DS production, Eq. (2) was expressed as a response surface plot (Fig. 2). The interaction between the sucrose and stillage concentrations is presented in Fig. 2A. The maximal DS concentration was achieved at the highest concentration of sucrose (6%) and in the range of higher stillage concentrations (55-75 %). Also, it could be noticed that there was the increase in DS production when higher concentrations of sucrose (5-6 %) and Mn²⁺ (0.018-0.024 %) were used (Fig 2C).

According to the Eq. (2) the Mn²⁺ concentration (C), as a single factor, has a positive influence on the DS production and exhibits significant positive interactions: Mn²⁺ - stillage (AC) and Mn²⁺ - sucrose (BC) (Fig. 2 B, C). By comparing the enzymatic activity in the medium with the highest DS activity (Run 19, Table 4), with the medium that contained the same concentrations of stillage and sucrose (Run 10, Table 4) we concluded that the increase in the concentration of Mn²⁺ ions increased the DS activity for 33 %.

The importance of different ions with regard to the enzyme production processes is generally accepted. Purama and Goyal [36] observed a 12 % increase in the enzyme production with the increase in the concentration of MnSO₄ from 0.001 (control) to 0.005 % for *L. mesenteroides* NRRL B-640. The essential role of Mn²⁺ ions for the DS production by *L. mesenteroides* strains was also reported [1]. The addition of amino acids, Mg²⁺ and Mn²⁺ ions stimulated the growth of most *Leuconostoc* strains [23]. It has been also shown that Mn²⁺ suppressed the inhibitory effect of aeration on the growth of *L. mesenteroides* UD-23 [23]. High requirements of Mn²⁺ ions could be explained by its interaction with enzymes and the ability to scavenge toxic oxygen radicals resulting in a protective role. In our previous studies on the DS production, Mn²⁺ ions also showed a positive effect on the activity of partially purified DS obtained from *L. mesenteroides* T3 [34].

In the present experiments, stillage was used as a source of nitrogen. From the perturbation plot (Fig. 1B), the influence of individual factors on the DS production can be seen (Y). A sharp curvature, a function with a maximum value, for stillage concentration (A) (Fig 1.B) shows that the DS production yield is highly sensitive to this parameter and correspondingly the quadratic regression coefficient has a negative value (Eq. 2). With increasing the concentration of stillage, the DS



production yield is increasing, until a maximum is reached after which a further increase in stillage concentration leads to the decrease in DS production. In conducted experiments, the medium with the highest DS activity (Run 19, Table 4) contained approximately 2.5 % of nitrogen, which is similar to the commercial De Man, Rogosa and Sharpe (MRS) medium. Residual yeast from bioethanol production in distillery stillage contributes as a source of assimilative nitrogen and thermo stable vitamins, affecting the efficiency of sugar utilization and promoting the growth of LAB [37].




Fig. 2. Surface plots of interactive effects of: (A) stillage concentration and sucrose concentration (AB), (B) sucrose concentration and Mn^{2+} concentration (BC) and (C) stillage concentration and Mn^{2+} concentration (AC)

There are other studies with waste materials as substrates for DS and dextran productions [10, 40, 41]. According to literature data, molasses was used several times for production of DS and other enzymes. For example, *L. mesenteroides* FT 045B produced DS with the maximum activity (4.03 U cm^{-3}) after 24 h of fermentation while growing on molasses with the addition of corn steep liquor as the nitrogen source [40]. On the other hand, low DS activity $4.3 \text{ DSU cm}^{-3} \text{ h}^{-1}$ (where one DSU was defined as the enzyme quantity that converts 1.0 milligram of sucrose into fructose and dextran in 1.0 h) obtained from *Lactobacillus acidophilus* was reported on molasses as the sole carbon source [41]. In our previous work, we optimized conditions for enhancement of DS production (2.02 U cm^{-3}) on molasses using sugar beet pulp as a support for immobilization of *L. mesenteroides* T3 [29].

In the present study, a 60 % higher DS production ($3.391 \pm 0.131 \text{ U cm}^{-3}$) in comparison to our previous study has been obtained on cheaper and abundant substrate.

3. 4. Validation of the model

The objective of this study was to find the optimal medium composition, using two waste materials, for DS production by *L. mesenteroides* T3. In order to validate the obtained model one point was selected from the numerical optimization results. The experiment was conducted with 64.33 wt % stillage, 5.30 wt % sucrose and 0.022 wt % Mn^{2+} . The predicted value for the outcome DS activity was 3.498 U cm^{-3} with 95 % prediction interval (PI) $3.098 - 3.898$. The

measured value for the parameter fitted within the 95 % PIs, and was very close to the most probable predicated value for the DS activity ($3.391 \pm 0.131 \text{ U cm}^{-3}$), showing that the model is reliable.

The nitrogen: carbon ratio has an important role in optimization of the medium composition for the DS production. After calculation of the total nitrogen and carbon contents in the medium that provided the highest DS activity, the obtained nitrogen: carbon ratio was approximately 0.67:1 (0.85% molasses + 5.30% sucrose + 0.8% stillage or a total of 7% for carbon and 4.7% for nitrogen concentrations). According to literature [4] this is the most suitable nitrogen: carbon ratio for DS production.

4. CONCLUSION

Current trends in the enzyme production include the use of low-cost or waste substrates. Revalorization of agro-industry waste as a substrate for biotechnological production fits within the sustainable development goals [42]. We have demonstrated that two waste substrates, distillery stillage and sugar beet molasses could be combined as cheap and renewable sources of nitrogen, vitamins, minerals and fermentable sugars for the growth of *L. mesenteroides* T3 and for the DS production. The applied optimization process by CCD has shown that 60 % increase in the DS activity ($3.391 \pm 0.131 \text{ U cm}^{-3}$) has been obtained on a cheaper and abundant substrate, as compared to our previous study. Manganese and sucrose are identified as key linear correlating components in media optimization for the DS production. Development of a process for DS production on waste materials with possible reductions of expenses can have both an economic and an ecological significance.

This study proves potentials for using wastes from one industry as the substrates for obtaining valuable biotechnological products in the other industry in accordance with principles of circular economy. It could serve as a basis for the development of a process for DS production with possible reduction of expenses and environmental footprint.

Acknowledgements: This work was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract No. 451-03-9/2021-14/200135).

REFERENCES

- [1] Dols M, Remaud-Simeon M, Willemot RM, Vignon M, Monsan P. Characterization of the different dextransucrase activities excreted in glucose, fructose, or sucrose medium by *Leuconostoc mesenteroides* NRRL B- 1299. *Appl Environ Microbiol* 1998; 64 (4): 1298–1302. <https://doi.org/10.1128/AEM.64.4.1298-1302.1998>
- [2] Paul F, Oriol E, Auriol D, Monsan P. Acceptor reaction of a highly purified dextransucrase with maltose and oligosaccharides. Application to the synthesis of controlled-molecular-weight dextrans. *Carbohydr Res*. 1986; 149: 433–441. [https://doi.org/10.1016/S0008-6215\(00\)90063-6](https://doi.org/10.1016/S0008-6215(00)90063-6)
- [3] Majumder A, Purama RK, Goyal A. An overview of purification methods of glycoside hydrolase family 70 dextransucrase. *Indian J Microbiol* 2007; 47(3): 197–206. <https://doi.org/10.1007/s12088-007-0040-8>
- [4] Lopretti M, Martinez L. Influence of nitrogen/carbon ratio and complementary sugars on dextransucrase production by *Leuconostoc mesenteroides* NRRL B512(f). *Process Biochem*. 2020; 34(9): 879–884. [https://doi.org/10.1016/S0032-9592\(99\)00020-5](https://doi.org/10.1016/S0032-9592(99)00020-5)
- [5] Neubauer H, Bauché A, Mollet B. Molecular characterization and expression analysis of the dextransucrase DsrD of *Leuconostoc mesenteroides* Lcc4 in homologous and heterologous *Lactococcus lactis* cultures. *Microbiology*. 2003; 149(4): 973–982. <https://doi.org/10.1099/mic.0.26029-0>
- [6] Gil EC, Colarte AI, El Ghzaoui A, Durand D, Delarbre JL, Bataille B. A sugar cane native dextran as an innovative functional excipient for the development of pharmaceutical tablets. *Eur J Pharm Biopharm*. 2008; 68 (2): 319–329. <https://doi.org/10.1016/j.ejpb.2007.04.015>
- [7] Patel S, Majumder A, Goyal A. Potentials of Exopolysaccharides from Lactic Acid Bacteria. *Indian J. Microbiol*. 2012; 52 (1): 3-12. <https://doi.org/10.1007/s12088-011-0148-8>
- [8] GLOBAL DEXTRAN MARKET 2019 BY MANUFACTURERS, REGIONS, TYPE AND APPLICATION, FORECAST TO 2024. [Online]. Available: <https://www.360researchreports.com/global-dextran-market-13851154>. [Accessed: 28-Jan-2021].
- [9] Queiroz JHD, Maugerio Filho FG. Optimization of the production of dextransucrase from *Leuconostoc mesenteroides* using molasses as carbon source. *Revista de Microbiologia* (ISSN 0001-3714). 1989; 20 (2): 150–156.
- [10] Santos M, Rodrigues A, Teixeira JA. Production of dextran and fructose from carob pod extract and cheese whey by *Leuconostoc mesenteroides* NRRL B512(f). *Biochem Eng J*. 2005; 25:1-6. <https://doi.org/10.1016/j.bej.2005.01.022>



- [11] Chagas CMA, Honorato TL, Pinto GAS, Maia GA, Rodrigues S. Dextranucrase production using cashew apple juice as substrate: Effect of phosphate and yeast extract addition. *Bioprocess Biosyst Eng.* 2007; 30(3): 207-215. <https://doi.org/doi:10.1007/s00449-007-0117-0>
- [12] Wilkie AC, Riedesel KJ, Owens JM. Stillage characterization and anaerobic treatment of ethanol stillage from conventional and cellulosic feedstocks. *Biomass Bioenerg.* 2000; 19(2): 63–102. [https://doi.org/10.1016/S0961-9534\(00\)00017-9](https://doi.org/10.1016/S0961-9534(00)00017-9)
- [13] Kim J-S, Kim B-G, Lee C-H, Kim S-W, Jee H-S, Koh J-H, Fane AG. Development of clean technology in alcohol fermentation industry. *J Clean Prod.* 1997; 5(4): 263–267. [https://doi.org/10.1016/S0959-6526\(97\)00043-7](https://doi.org/10.1016/S0959-6526(97)00043-7)
- [14] Djukić-Vuković A, Mojović L, Nikolić S, Pejini J, Kocić-Tanackov S, Mihajlovski K. Distillery stillage as a new substrate for lactic acid production in batch and fed-batch fermentation. *Chem Eng Trans* 2013; 34: 97–102. <https://doi.org/10.3303/CET1334017>
- [15] Rodrigues LR, Teixeira JA, Oliveira R. Low-cost fermentative medium for biosurfactant production by probiotic bacteria. *Biochem Eng J.* 2006; 32(3): 135–142. <https://doi.org/10.1016/j.bej.2006.09.012>
- [16] Nigam P, Vogel M. Bioconversion of sugar industry by-products-molasses and sugar beet pulp for single cell protein production by yeasts. *Biomass Bioenerg* 1991; 1(6): 339–345. [https://doi.org/10.1016/0961-9534\(91\)90014-4](https://doi.org/10.1016/0961-9534(91)90014-4)
- [17] Harland JI, Jones CK, Hufford C. Co-products, in *Sugar Beet*, A. Draycott, Ed. Blackwell Publishing Ltd., 2006, pp. 451–455.
- [18] Kalogiannis S, Lakovidou G, Liakopoulou-Kyriakides M, Kyriakidis DA, Skaracis GN. Optimization of xanthan gum production by *Xanthomonas campestris* grown in molasses. *Process Biochem.* 2003; 39(2): 249–256. [https://doi.org/10.1016/S0032-9592\(03\)00067-0](https://doi.org/10.1016/S0032-9592(03)00067-0)
- [19] Mojović L, Nikolić S, Pejini D, Pejini J, Djukić-Vuković A, Kocić-Tanackov S, Semenčenko V. “The potential for sustainable bioethanol production in Serbia: available biomass and new production approaches,” In: Mendez-Vilas A (Ed), *Materials and processes for energy: communicating current research and technological developments*, Formatex Research Center 2013, pp. 380–392.
- [20] Mihajlovski KR, Radovanović NR, Veljović DjN, Šiler-Marinković SS, Dimitrijević-Branković SI. Improved β -amylase production on molasses and sugar beet pulp by a novel strain *Paenibacillus chitinolyticus* CKS1. *Ind Crops Prod.* 2016; 80: 115–122. <https://doi.org/10.1016/j.indcrop.2015.11.025>
- [21] Dols M, Remaud-Simeon M, Monsan PF. Dextranucrase production by *Leuconostoc mesenteroides* NRRL B-1299. Comparison with *L. mesenteroides* NRRL B-512F. *Enzyme Microb Technol.* 1997; 20(7): 523–530. [https://doi.org/10.1016/S0141-0229\(96\)00189-5](https://doi.org/10.1016/S0141-0229(96)00189-5)
- [22] Goyal A, Katiyar S. Effect of certain nutrients on the production of dextranucrase from *Leuconostoc mesenteroides* NRRL B-512F. *J Basic Microbiol.* 1997; 37(3): 197–204. <https://doi.org/10.1002/jobm.3620370308>
- [23] Bellengier P, Richard J, Foucaud C. Nutritional requirements of *Leuconostoc mesenteroides* subsp. *mesenteroides* and subsp. *dextranicum* for growth in milk. *J Dairy Res.* 1997; 64 (1): 95–103. <https://doi.org/10.1017/S0022029996001902>
- [24] Uma Maheswar Rao JM, Satyanarayana T. Improving production of hyperthermostable and high maltose-forming α -amylase by an extreme thermophile *Geobacillus thermoleovorans* using response surface methodology and its applications. *Bioresour Technol.* 2007; 98(2): 345–352. <https://doi.org/10.1016/j.biortech.2005.12.022>
- [25] Majumder A, Bhandari S, Purama RK, Patel S, Goyal A. Enhanced production of a novel dextran from *Leuconostoc mesenteroides* NRRL B-640 by response surface methodology. *Ann Microbiol.* 2009; 59 (2): 309–315. <https://doi.org/10.1007/BF03178333>
- [26] Djukić-Vuković A, Mladenović D, Radosavljević M, Kocić-Tanackov S, Pejini J, Mojović L. Wastes from bioethanol and beer productions as substrates for l(+) lactic acid production - A comparative study. *Waste Manag.* 2016; 48: 478-482. <https://doi.org/10.1016/j.wasman.2015.11.031>
- [27] Davidović S, Miljković MG, Antonović DG, Rajilić-Stojanović MD, Dimitrijević-Branković SI. Water kefir grain as a source of potent dextran producing lactic acid bacteria. *Hem. Ind.* 2015; 69(6): 595-604. <https://doi.org/10.2298/HEMIND140925083D>
- [28] Tsuchiya HM, Koepsell HJ, Corman J, Bryant G, Bogard MO, Feger VH, Jackson RW. The effect of certain cultural factors on production of dextranucrase by *Leuconostoc mesenteroides*. *J Bacteriol.* 1952; 64(4): 521-526. <https://doi.org/10.1128/jb.64.4.521-526.1952>
- [29] Miljković MG, Davidović SZ, Carević MB, Veljović DN, Mladenović DD, Rajilić-Stojanović MD, Dimitrijević-Branković SI. Sugar Beet Pulp as *Leuconostoc mesenteroides* T3 Support for Enhanced Dextranucrase Production on Molasses,” *Appl Biochem Biotechnol.* 2016; 180(5): 1016–1027. <https://doi.org/10.1007/s12010-016-2149-x>
- [30] “Association of Official Analytical Chemists (AOAC),” Off. methods Anal. (17th ed.). Gaithersbg. AOAC Int. (Methods 923.03, 925.09, 930.15, 955.04, 960.39), 2000.
- [31] Miller GL. Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. *Anal. Chem.* 1959; 31(3): 426–428. <https://doi.org/10.1021/ac60147a030>
- [32] Boyaval P. Lactic acid bacteria and metal ions, *Lait.* 1989; 69(2): 87–113.
- [33] Šušić S, Petrov S, Kukić G, Sinobad V, Perunović P, Koronšovac B, Bašić Dj. *Osnovi tehnologije šećera*, Drugo izda. Beograd: Jugošćer, 1995.
- [34] Miljković GM, Davidović SZ, Kralj S, Šiler-Marinković SS, Rajilić-Stojanović MD, Dimitrijević-Branković SI. Characterization of dextranucrase from *Leuconostoc mesenteroides* T3, water kefir grains isolate. *Hem. Ind.* 2017; 71(4): 351-360. <https://doi.org/10.2298/HEMIND160421046M>

- [35] Adinarayana K, Ellaiah P. Response surface optimization of the critical medium components for the production of alkaline protease by a newly isolated *Bacillus* sp. *J Pharm Pharm Sci.* 2002; 5 (3) 272–278. PMID: 12553896.
- [36] Purama RK, Goyal A. Effect of Nutrients by One Variable At A Time (OVAT) Approach on the Dextranucrase Production from *Leuconostoc mesenteroides* NRRL B-640. *Internet J Microbiol.* 2012; 5 (1). <https://doi.org/10.5580/1bc>
- [37] Djukić-Vuković AP, Mojović LV, Vukašinić-Sekulić MS, Nikolić MS, Pejin JD. Integrated production of lactic acid and biomass on distillery stillage. *Bioprocess Biosyst Eng.* 2013; 36 (9): 1157–1164. <https://doi.org/10.1007/s00449-012-0842-x>
- [38] Vedyashkina TA, Revin VV, Gogotov IN. Optimizing the conditions of dextran synthesis by the bacterium *Leuconostoc mesenteroides* grown in a molasses-containing medium. *Appl Biochem Microbiol.* 2005; 41: 361-364. <https://doi.org/10.1007/s10438-005-0061-1>
- [39] Behravan J, Fazly Bazzaz BS, Salimi Z. Optimization of dextran production by *Leuconostoc mesenteroides* NRRL B-512 using cheap and local sources of carbohydrate and nitrogen. *Biotechnol Appl Biochem.* 2003; 38(3): 267-269. <https://doi.org/10.1042/BA20030084>
- [40] Vettori MHPB, de Lima CJB, Blanco KC, Cortezi M, Contiero J. Performance of response surface model for increase of dextranucrase production by leuconostoc mesenteroides FT 045B under different experimental conditions. " *Asian J Biological Life Sci.* 2012; 1(1): 29–35. https://www.ajbls.com/sites/default/files/AsianJBiolLifeSci_1_1_29.pdf
- [41] Abedin RMA, El-Borai AM, Abo Shall M, El-Assar SA. Optimization and statistical evaluation of medium components affecting dextran and dextranucrase production by *Lactobacillus acidophilus* ST76480.01. *Life Sci J.* 2013; 10(1): 1746–1753. <http://www.dx.doi.org/10.7537/marslsj100113.254>
- [42] UN Sustainable Development Goals. <https://sdgs.un.org/goals>. [Accessed: 28-Jan-2021].

SAŽETAK**Iskorišćenje nus-proizvoda agro-industrije za proizvodnju dekstransaharaze pomoću bakterije *Leuconostoc mesenteroides* T3: optimizacija procesa metodom odzivnih površina**

Miona G. Miljković¹, Slađana Z. Davidović^{1*}, Aleksandra P. Djukić-Vuković¹, Mila V. Ilić², Milica B. Simović¹, Mirjana D. Rajilić-Stojanović¹, Suzana I. Dimitrijević-Branković¹

¹Univerzitet u Beogradu, Tehnološko-metalurški fakultet, Katedra za biohemijsko inženjerstvo i biotehnologiju, Karnegijeva 4, Beograd, Srbija

²Univerzitet u Beogradu, Institut za hemiju, tehnologiju i metalurgiju, Njegoševa 12, 11000 Beograd, Srbija

(Naučni rad)

Dekstransaharaza (DS) je glukoziltransferaza (E. C. 2.4.1.5.) koja katalizuje prenos ostataka glukoze iz saharoze u polimer dekstrana, pri čemu se oslobađa fruktoza. Ovaj enzim je povezan sa širokim spektrom primene dekstrana i oligosaharida. Proizvodnja dekstransaharaze pomoću bakterije *Leuconostoc mesenteroides* T3 optimizovana je metodom odzivnih površina korišćenjem centralnog kompozitnog dizajna. Za optimizaciju su izabrane tri promenljive: koncentracija džibre, koncentracija saharoze i koncentracija jona mangana. Rezultati su pokazali da koncentracije saharoze i jona mangana imaju pozitivan linearni efekat na proizvodnju DS dok su sve interakcije (džibra-Mn²⁺, džibra-saharaza i saharoza-Mn²⁺) imale značajan uticaj na proizvodnju DS. Na osnovu eksperimentalnih podataka i numeričke optimizacije, dobijen je maksimalni prinos DS od 3.391 ± 0.131 U cm⁻³ u podlozi sa 64.33 % džibre, 5.30 % saharoze i 0.022 % jona mangana. Naše istraživanje otkrilo je da se džibra u kombinaciji sa melasom šećerne repe kao i saharozom i dodatkom jona mangana može koristiti kao dragocena hranjiva komponenta za rast bakterija mlečne kiseline i proizvodnju DS. Takođe, uzimajući u obzir poreklo supstrata, upotreba industrijskih nusproizvoda na ovaj način ima veliku ekološku važnost.

Ključne reči: bakterije mlečne kiseline; dekstran; proizvodnja enzima; destilerijska džibra; melasa