



Uvaria huana (Annonaceae), a replacement name for an Asian species

RUIJIANG WANG

South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

E-mail: wangrj@scbg.ac.cn

Abstract

Uvaria huana is proposed as a new name for *U. yunnanensis* (Hu) L. L. Zhou, Y. C. F. Su & R. M. K. Saunders (2009), a later homonym of *U. yunnanensis* Li (1978).

Kew words: China, homonyms, taxonomy, *Uvaria yunnanensis*

Introduction

Uvaria L. (1753: 536) is one of the largest genera of Annonaceae, with about 210 woody lianas or scrambling shrubs. The genus is distributed mainly in tropical areas of Africa, Asia, and Australia. It is characterized by a combination of morphological characters, including stellate hairs, valvate sepals, more or less imbricate petals, and many-seeded apocarpous fruits (Meade, 2005). Based on recent molecular phylogenetic analysis (Zhou *et al.*, 2009; Zhou *et al.*, 2010; Attanayake *et al.*, 2011), *Uvaria* is now defined to include *Anomianthus* Zoll. (1858: 324), *Balonga* Le Thomas (1968: 106), *Cyathostemma* Griff. (1854: 707), *Dasoclema* J. Sinclair (1955: 273), *Ellipeia* Hook.f. & Thomson (1855: 104), *Ellipeiopsis* R. E. Fr. (1955: 41), *Rauwenhoffia* Scheff. (1885: 21), and the Australian species of *Melodorum* Lour. (1790: 329, 351).

The ten species of *Cyathostemma* recognized by Utteridge (2000), including the Asian species *C. yunnanensis* Hu, were transferred to *Uvaria* (Zhou *et al.*, 2009). During preparation of an account of the Chinese species of *Uvaria*, it was noticed that Hu's (1940) extant *C. yunnanensis* could not be transferred to *Uvaria* because of the fossil species *U. yunnanensis* Li (Li, 1978), a fact overlooked by Zhou *et al.* A new name, honouring Professor Hsen-Hsu Hu (1894–1968), one of the pioneers of Chinese botanical sciences, is proposed here.

Taxonomic treatment

Uvaria huana R. J. Wang, *nom. nov.*

Replaced name: *Uvaria yunnanensis* (Hu) L. L. Zhou, Y. C. F. Su & R. M. K. Saunders (2009: 255), based on *Cyathostemma yunnanensis* Hu (1940: 121).—Type: CHINA. Yunnan Province: Fo-Hai Hsien, alt. 980m., in mixed forest, woody plants, 3 m high, 12.5 cm in diam., flowers greenish-yellow, Jun 1936, *C. W. Wang* 74547 (holotype: PE! barcode PE00028347; isotype: PE! barcode PE00028346), non Li (1978: 19).

The publication of the fossil-species, *Uvaria yunnanensis* Li, was accompanied by a description in Chinese, along with an illustration and a plate (Li, 1978). This name was validly and effectively published because the protologue satisfied articles 43.1 and 43.2 of ICN (McNeill *et al.*, 2012). It should be noted that the specimen *C. W. Wang* 74879 (PE!) is a paratype of *Cyathostemma yunnanense* and not the holotype as cited by Utteridge (2000).

Distribution:— SW. China (Yunnan province) and N. Vietnam (Yen Bai province).

Acknowledgements

This work was supported by the National Natural Science Foundation of China (grant nos. 30770156, 31070177), National Natural Science Foundation of China (grant no. 31110103911), and Main Direction Program of Knowledge Innovation of Chinese Academy of Sciences (grant no. KSCX2-EW-Z-1).

References

- Attanayake, A.M.A.S., Turner, I.M. & Saunders, R.M.K. (2011) Two new species of *Uvaria* (Annonaceae) from Borneo, with a new nomenclatural combination. *Novon* 21(2): 161–168.
- Fries, R.E. (1955) Verstreute Beobachtungen hinsichtlich der Familie Annonaceae. *Arkiv för Botanik, Andra Serien* 3, 2: 35–42.
- Griffith, W. (1854) *Notulae ad Plantas Asiaticas*, vol. 4. Charles A. Serrao, Calcutta, pp. 706–717.
- Hooker, J.D., & Thomson, T. (1855) Annonaceae. *Flora Indica*, vol. 1. W. Pamplin, London, pp. 86–153.
- Hu, H.H. (1940) Notulae Systematicae ad Floram Sinensem X. *Bulletin of the Fan Memorial Institute of Biology*, Botanical Series 10: 117–172.
- Le Thomas, A. (1968) *Balonga* Le Thomas nouveau genre africain de la famille des Annonacées. *Adansonia*, nova séries, 8: 105–111.
- Li, H.M. (1978) *Uvaria yunnanensis*, sp. nov. In: Editorial Committee of Chinese plants in Cenozoic of The Institute of Botany & Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences (ed.), *Chinese Plant Fossils*, vol. 3: *Chinese plants in Cenozoic*. Science Press, Beijing, p. 19.
- Linnaeus, C. (1753) *Uvaria* L. *Species Plantarum*, vol. 1. Laurentii Salvii, Stockholm, p. 536.
- Loureiro, J. (1790) Genus XXVII. *Melodorum*. *Flora Cochinchinensis*, vol. 1. Typis, et expensis academicis, Lissabon, p. 329, p. 351.
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Mmarhold, K., Prado, J., Reine, W.F.P.V., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012) *International Code of Nomenclature for algae, fungi, and plants (Melbourne Code), adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011*. Regnum Vegetabile 154. Koeltz Scientific Books, 232 pp.
- Meade, C.V. (2005) A new species of *Uvaria* (Annonaceae) from southeast Asia. *Adansonia*, séries 3, 27(1): 17–20.
- Scheffer, R.H.C.C. (1885) Sur Quelques Plantes Nouvelles ou peu Connues de L'Archipel Indien (Annonaceae). *Annales du Jardin Botanique de Buitenzorg* 2: 1–31.
- Sinclair, J. (1955) A revision of the Malayan Annonaceae. *Gardens' Bulletin, Straits Settlements*, series 3, 14(2): 149–516.
- Utteridge, T.M.A. (2000) Revision of the genus *Cyathostemma* (Annonaceae). *Blumea* 45(2): 377–396.
- Zhou, L.L., Su, Y.C.F., Chalermglin, P. & Saunders, R.M.K. (2010) Molecular phylogenetics of *Uvaria* (Annonaceae): relationships with *Balonga*, *Dasoclema* and Australian species of *Melodorum*. *Botanical Journal of the Linnean Society* 163: 33–43.
- Zhou, L.L., Su, Y.C.F. & Saunders, R.M.K. (2009) Molecular phylogenetic support for a broader delimitation of *Uvaria* (Annonaceae), inclusive of *Anomianthus*, *Cyathostemma*, *Ellipeia*, *Ellipeiopsis* and *Rauwenhoffia*. *Systematics and Biodiversity* 7: 249–258.
- Zollinger, H. (1858) Ueber die Anonaceen des ostindischen Archipels. *Linnaea* 29: 297–325.