arXiv:2107.06069v2 [hep-th] 31 Mar 2022

Vacuum Energy and Renormalization of the Field-Independent Term

I. G. Márián,^{1,2} U. D. Jentschura,^{3,1,4} N. Defenu,⁵ A. Trombettoni,^{6,7,8} and I. Nándori^{1,4,2}

¹ MTA-DE Particle Physics Research Group, P.O. Box 51, H-4001 Debrecen, Hungary

²University of Debrecen, P.O.Box 105, H-4010 Debrecen, Hungary

³Department of Physics, Missouri University of Science and Technology, Rolla, Missouri 65409, USA

⁴MTA Atomki, P.O. Box 51, H-4001 Debrecen, Hungary

⁵Institut für Theoretische Physik, ETH Zürich, Wolfgang-Pauli-Str. 27 Zurich, Switzerland

⁶Department of Physics, University of Trieste, Strada Costiera 11, I-34151 Trieste, Italy

⁷CNR-IOM DEMOCRITOS Simulation Center, Via Bonomea 265, I-34136 Trieste, Italy

⁸SISSA and INFN, Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy

Due to its construction, the nonperturbative renormalization group (RG) evolution of the constant, field-independent term (which is constant with respect to field variations but depends on the RG scale k) requires special care within the Functional Renormalization Group (FRG) approach. In several instances, the constant term of the potential has no physical meaning. However, there are special cases where it receives important applications. In low dimensions (d = 1), in a quantum mechanical model, this term is associated with the ground-state energy of the anharmonic oscillator. In higher dimensions (d = 4), it is identical to the Λ term of the Einstein equations and it plays a role in cosmic inflation. Thus, in statistical field theory, in flat space, the constant term could be associated with the free energy, while in curved space, it could be naturally associated with the cosmological constant. It is known that one has to use a subtraction method for the quantum anharmonic oscillator in d = 1 to remove the k^2 term that appears in the RG flow in its high-energy (UV) limit in order to recover the correct results for the ground-state energy. The subtraction is needed because the Gaussian fixed point is missing in the RG flow once the constant term is included. However, if the Gaussian fixed point is there, no further subtraction is required. Here, we propose a subtraction method for k^4 and k^2 terms of the UV scaling of the RG equations for d = 4dimensions if the Gaussian fixed point is missing in the RG flow with the constant term. Finally, comments on the application of our results to cosmological models are provided.

PACS numbers: 98.80.-k, 11.10.Gh, 11.10.Hi

Contents

I.	Introduction	2
II.	Field-Independent Term in Cosmology	4
III.	Field-Independent Term in RG	7
IV.	RG Evolution of the Constant Term	9
v.	Divergences in $d = 1$ Oscillators	9
VI.	Divergences in the Cosmological Model A. General Discussion and Idea B. Double Subtraction C. Comparison with Other Results	10 10 12 13
VII.	Conclusions	14
	Acknowledgments	15
А.	Field–Independent Term and Cosmological Constant	15
	References	15

I. INTRODUCTION

The quantum field theory of gravity, i.e., Quantum Einstein Gravity (QEG) is perturbatively non-renormalizable. It requires infinitely many unknown parameters to be set by experiment. A possible solution to perform renormalization is the use of a nonperturbative treatment. Indeed, nonperturbative renormalizability, which is also referred to as asymptotic safety, provides us with a nontrivial high energy, i.e., ultraviolet (UV) fixed point of the renormalization group (RG) flow. The RG flow leads to a finite number of UV-attractive couplings; so, it is sufficient to perform only a finite number of measurements. In other words, it controls the UV behavior of the dimensionless couplings. They do not need to be small or tend to zero in the UV limit but tend to finite values at the nontrivial UV fixed point. It was shown in [1] that for the simplest truncation of QEG which is the Einstein–Hilbert action, such a nontrivial fixed point is indeed present. Up to the present, many different works have already confirmed that the asymptotic safety scenario is possible. For a recent review, we refer to [2]. For applications to cosmology, one consults Ref. [3, 4], and for applications to black-hole physics and instructive explicit functional renormalization-group computations, we refer to [5–16].

Asymptotic safety is the concept that a theory is UV completed by an interacting theory which, in the RG formalism, is described by a nontrivial (non-Gaussian) UV fixed point and includes the assumption that the RG formalism can be used to solve the so-called cosmological constant problem, too. One can use different methods to test this assumption. The non-perturbative RG is a possible method, but there are also others, including lattice and Monte Carlo methods. In general, the nonperturbative RG method [17–21] has been used successfully in many areas of physics from statistical mechanics to high energy physics [22–30] and in cosmology [31–38] with particular attention to the cosmological constant problem [39–57]. It is obvious that the vacuum energy density induced by the quantum fluctuations would add to the cosmological constant (see Ref. [58, 59]). However, the calculated vacuum energy density is larger by many orders of magnitude than the observed cosmological constant; this is the essence of the cosmological constant problem [60, 61]. Also, it is well known that the bare values of a physical theory are connected to the lowenergy phenomenology by RG transformations [62–67]. So, if one could establish a connection of the bare value of the cosmological constant to its low-energy equivalent by an RG method, then one could possibly solve the problem: The bare value of the cosmological constant carries no physical meaning and is treated as a running parameter. Indeed, in principle, RG running can be used to connect its UV value to the observed low energy, i.e., infrared (IR) one. Thus, in the functional renormalization group (FRG) treatment of gravity, one sees that the cosmological constant corresponds to a relevant direction. It means, there is no cosmological constant problem in the nonperturbative treatment of gravity; it is a free parameter to be fixed by observations. However, let us note that one has to distinguish between the RG scale k and the observational scale, which we denote as p. One should compute the physical limit $k \to 0$ by the RG method first and then compute observables using the full quantum effective action. Unfortunately, this is not yet possible in reasonable approximations, so, one has a qualitative understanding of the implications of asymptotically safe gravity, assuming that the artificial flow with k is a good approximation to the physical flow with respect to the physical momentum p, which one can extract from the effective action in the limit $k \to 0$. The flow generated by the FRG equation can be analysed in regard to the question of whether it allows trajectories that are compatible with the observational constraints on the parameters of the action, since it is related to the k-flow and not to the p-flow. In principle, the effective action, in the limit $k \to 0$, has infinitely many parameters, but only N of these parameters (those associated with the N relevant directions of the non-Gaussian UV fixed point) must be fixed using N experimental inputs.

The RG running could be associated with the temporal evolution of the Universe according to the identification $k \propto 1/H(t)$ where H(t) is the time-dependent Hubble-parameter, according to an idea promoted in Ref. [4]. Thus, one could treat the field independent constant, i.e., the Λ term in Einstein's equations, as a running parameter which varies over the temporal evolution of the Universe. If there is no other scalar field in the theory, the cosmological constant must play an important role in the mechanism of cosmic inflation, too. So, its value must be fixed to be large at a sub-Planckian scales. Thus, renormalization conditions must be chosen very accurately in order to solve the corresponding hierarchy problem. In other words, the running cosmological constant has to be scaled down from high to low energies; this cannot be done straightforwardly in the framework of conventional perturbative renormalization. So, again, a nonperturbative treatment is needed. As an example, a nonperturbative approach has been used in [56] to study the RG evolution in the presence of compact extra dimensions. As a consequence, the studied parameters can be substantially reduced to be comparable with observational values. So, if there is a single scalar in the theory, then one would need to adjust the running of the cosmological constant to accommodate both inflation and late-time dark energy. However, FRG generates infinitely many operators, so that together with the cosmological constant the theory is equipped with an additional scalar degree of freedom (the one of Starobinsky inflation) that can take care of the inflationary era. This has been studied extensively in the literature, see for instance [68, 69].

Therefore, the use of the FRG method and asymptotic safety can solve several important problems of cosmology and quantum gravity. Nevertheless, the FRG approach has its own drawbacks; the evolution of the constant (i.e., field-independent) term could be problematic due to the construction of the method. In particular, divergent terms could appear in the RG flow in its UV limit. These could represent non-physical behaviour. It can be quite a subtle problem to analyze the renormalization of the field-independent term, because of the conditions imposed canonically on the regulator function in the FRG equation (Wetterich equation) [19, 20, 24]. These canonically imposed conditions imply that the solution of the FRG equation reproduces the bare action in the UV only up to a field-independent, but k-dependent term; the latter, in addition, could be infinite. Hence, the analysis becomes subtle. Yet, physical conditions, to be imposed on the RG evolution of the constant term due to consistency considerations, lead to subtracted RG equations whose solution fulfills all physical boundary conditions in IR and UV limits.

Indeed, the idea to remove these UV divergent terms of the RG evolution of the field-independent term in the framework of the FRG method is well known in quantum-mechanical models, and in statistical mechanics. For example, as suggested by quantum mechanical calculations [see Eq. (37) of Ref. [24] and additional discussions in Refs. [25, 26]], the application of a suitable subtraction reduces the quadratic (k^2) divergence of the RG evolution in the UV limit and produces correct results for the free energy in the IR limit. However, it is very important to observe that the non-physical behaviour, i.e., the k^2 divergence of the UV limit is the consequence of the *absence* of the Gaussian fixed point in the RG flow once the constant term is included. Of course, one finds the Gaussian fixed point in the RG flow of the k^2 divergence in the UV limit emerges, and the Gaussian fixed point not only disappears; it becomes unstable against tiny variations of the initial conditions of the RG flow. Conversely, if the β -functions of a certain model allow the existence of the Gaussian fixed point in the presence of the constant coupling, no further subtraction is required.

It is possible to generalise the idea of the subtraction method for higher dimensions. Indeed, a similar procedure has been proposed in Ref. [70] for cosmological applications. In d = 4 dimensions, in the absence of the Gaussian fixed point, k^4 and k^2 divergences appear in the RG flow of the field-independent but k-dependent term in its UV limit which may need special attention.

However, in the framework of the asymptotic-safety (AS) scenario, the situation is more complex. The β -functions of AS gravity (without scalar degrees of freedom) predict the existence of Gaussian and non-Gaussian UV fixed points, see for example [1–3] and the RG flow diagram of Fig.1. At the non-Gaussian UV fixed point, the dimensionless couplings are required to attain a non-zero value. Thus, if we assume that the non-Gaussian UV fixed point is found in the asymptotic region, then at least some of the dimensionful counterparts of the couplings become divergent. For example, if we assume that the dimensionless cosmological constant $\lambda(k) = \Lambda(k) k^{-2}$ tends to a finite value at the non-Gaussian UV fixed point $\lambda(k = k_{\star}) = \lambda_{\star}$, then its dimensionful counterpart $\Lambda(k)$, for large k, can have at most a k^2 divergence. So, one could argue that a subtraction might be required which removes the k^2 divergence and leaves at most a logarithmic divergence, for large k. However, AS gravity (without scalar fields) has the Gaussian UV fixed point, too. The β -functions derived from the FRG equation signal the Gaussian UV fixed point, so the field independent term vanishes if all the other couplings tend to zero in the UV limit. Thus, there is no need of any further subtraction method and the scaling of the potential as k^4 is not a problem in AS gravity since the β -functions predict the existence of the Gaussian UV fixed point, too. However, if one adds scalar degrees of freedom as matter fields to the Einstein-Hilbert action which is the simplest truncation for AS gravity models, the resulting action, i.e., the simplest gravity-scalar system might have problem with the existence of the Gaussian fixed point. Of course, by the application of the subtraction method [70], the Gaussian fixed point can be restored and the method, i.e., the inclusion of subtraction terms does not affect the position of the non-Gaussian UV fixed point.

Therefore, the quantum field theory and the renormalization of gravity and also the consequences of the RG running on cosmology are well defined, well understood due to asymptotic safety (and the presence of the Gaussian fixed point). However, one might ask what would happen if one chooses different approaches to the problem of quantum gravity, for example M-theory or loop quantum gravity. In these cases, one can use quantum field theory maybe together with supersymmetry and RG considerations up to the Planck scale only. In addition, it is maybe required to use an effective quantum field theory beyond the Standard Model. As an example one can mention the effective branon theory in (3+1) dimensions which involves the absolute value of the branon field, leading to a non-differentiable potential and wave function renormalization [71]. For discussions of branons, see, e.g., Refs. [72–78]. This can re-open the discussion on the UV divergent nature of the RG flow of the field-independent term, i.e., the cosmological constant which could play an important role in the mechanism of cosmic inflation, too. For example, it is known that supersymmetric extensions of the Standard Model deal with quadratic divergences. It may resolve major hierarchy problems within the Standard Model, by guaranteeing that quadratic divergences of all orders will cancel out in perturbation theory, however, there is no experimental evidence that a supersymmetric extension to the Standard Model is correct. Due to the negative results from the LHC, it has been a matter of discussion whether the Minimal Supersymmetric Standard Model is no longer able to fully resolve the hierarchy problem. Thus, if not the asymptotic safety scenario is the correct approach to quantum gravity, the problem of UV divergences of the field independent term may require attention.

Indeed, in Ref. [57], the so-called Running Vacuum Model has been studied; it assumes a scale-dependent vacuum,

i.e., a running cosmological constant. The authors of [57] proposed a suitable subtraction of the so-called Minkowski contribution in the framework of the adiabatic regularization method. It results in an RG scaling for the running vacuum described by equation (6.9) of [57], where the running RG scale is denoted by the mass-like term M while its UV value is given by M_0 . The result of their subtraction method is the absence of the M^4 term in the scaling relation for the running vacuum. In the UV limit (when $M \to \infty$), the M^4 is a source of exceedingly large contributions. Due to the subtraction method of Ref. [57], the RG scale-dependence of the running vacuum is characterised with M^2 and $\log(M^2)$ terms only. Our goal here is to show how the subtraction method can be further improved in order to eliminate from the RG scaling of the running vacuum, not just the k^4 (i.e., M^4 of Ref. [57]) but also the k^2 (i.e., M^2 of Ref. [57]) terms which results in a purely logarithmic dependence on the scale.

II. FIELD-INDEPENDENT TERM IN COSMOLOGY

While, in ordinary particle physics, a constant, field-independent term of the potential carries no physical meaning, it has great importance in the case of gravity. For example, in order to describe the observed accelerated expansion of the Universe at present [81, 82] a possible solution is the inclusion of a constant term into Einstein's equation which reads (in the absence of matter)

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = -\Lambda g_{\mu\nu}$$
(1)

where R is the scalar curvature and Λ is the cosmological constant. The latter is assumed to be related to dark energy [83–87] and is expected to cause the accelerated expansion of the universe observed today. Indeed, by using the Friedmann–Lemaître–Robertson–Walker (FLRW, see Ref. [88–91]) metric (in our units, the speed of light is $c \equiv 1$ and $\hbar \equiv 1$), $g_{\mu\nu} = \text{diag}(-1, a^2, a^2, a^2)$ the scale factor a(t) of the expanding homogeneous and isotropic Universe can be calculated which results in an exponentially fast expansion, $a(t) \sim \exp(\sqrt{\Lambda/3} t)$.

There is, however, a serious problem with this scenario, namely the discrepancy between the theoretical prediction for the cosmological constant by quantum field theory (vacuum energy) and the energy density needed to explain the accelerated expansion of the present Universe. In other words, the estimated value is 120 orders of magnitude greater than the energy density of all the other matter. Thus, the cosmological constant which correctly describes the acceleration rate is very small compared to the Planck scale. This is the so-called cosmological constant problem.

One possible solution to this problem could be the existence of a hypothetical scalar field, referred to as quintessence [92–101], which is minimally coupled to gravity. Compared to other scalar-field models such as k-essence [102–104], quintessence is the simplest scenario where the slowly varying field along a potential results in a negative pressure and accelerated expansion. This mechanism is very similar to the particle physics model for cosmic inflation in the early Universe, but the difference is that non-relativistic matter cannot be ignored and the quintessence potential is much smaller than that of the inflaton potential. Let us first review the inflationary mechanism of the early Universe.

The key observation is that scalar fields can mimic the equation of state for negative pressure. Thus they represent an excellent model for inflation,

$$S = \int d^4x \sqrt{-g} \left[\frac{m_p^2}{2} R + \mathcal{L}_\phi \right], \qquad \mathcal{L}_\phi = -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \, \partial_\nu \phi - V(\phi) \,, \tag{2}$$

where $\sqrt{-g} = \sqrt{-\det(g_{\mu\nu})} = a^3$ with the (reduced) Planck mass $m_p^2 = 1/(8\pi G)$, and G is Newton's constant. The Einstein equation has to be written in the presence of matter fields,

$$G_{\mu\nu} = 8\pi G T_{\mu\nu}, \quad T^{\mu}_{\nu} = \text{diag}(-\rho, p, p, p),$$
(3)

where the stress-energy tensor of the scalar field has the following form,

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}\mathcal{L}_{\phi})}{\delta g^{\mu\nu}} = \partial_{\mu}\phi \,\partial_{\nu}\phi + g_{\mu\nu}\,\mathcal{L}_{\phi}.$$
(4)

Since over the inflation the field can be considered to be homogeneous $(\nabla \phi/a = 0)$, the relation between the density and pressure reads as

$$\omega = \frac{p}{\rho} = \frac{\frac{1}{2}\dot{\phi}^2 - V}{\frac{1}{2}\dot{\phi}^2 + V} \quad \text{if} \quad \frac{1}{2}\dot{\phi}^2 \ll V \implies \omega = -1,$$
(5)

$$\epsilon \equiv \hat{V}^{2}/(2\hat{V}^{2}) \ll 1, \qquad \eta \equiv \hat{V}^{\prime\prime}/\hat{V} \ll 1,$$
(6)

which have to be fulfilled by a suitable potential for a prolonged exponential inflation with slow roll down. The e-fold number $N \equiv -\int_{\phi_i}^{\phi_f} d\phi \frac{V}{V^{\tau}}$ should be in the range 50 < N < 60 where ϕ_i and ϕ_f are the initial and final configurations of the field, respectively. The power spectra of scalar (\mathcal{P}_S) and tensor (\mathcal{P}_{τ}) fluctuations can be characterized by their scale dependence, i.e., $\mathcal{P}_S \sim k^{n_s-1}$, where k is the comoving wave number. Then, slow-roll parameters are encoded in expressions for the scalar tilt $n_s - 1 \approx 2\eta - 6\epsilon$ and for the tensor-to-scalar ratio $r = \mathcal{P}_{\tau}/\mathcal{P}_S \approx 16\epsilon$, which can be directly compared to CMBR data [105–107].

The potential is determined by the slow-roll conditions (6) up to an overall multiplicative factor, but this factor is fixed by the absolute normalisation. According to Eq. (23) of Ref. [108] and Eq. (218) of Ref. [109], the normalisation condition is

$$V(\phi_i) \equiv \frac{r}{0.01} (10^{16} \,\text{GeV})^4 \,. \tag{7}$$

The tensor-to-scalar ratio r is given by the slow-roll parameters which are fixed at the scale of inflation (k_i) , according to remarks preceding Eq. (218) of Ref. [109]. Thus, the scale of inflation is given by the following relation

$$V(\phi_i) \equiv k_i^4, \qquad k_i = \left(\frac{r}{0.01}\right)^{\frac{1}{4}} 10^{16} \text{ GeV},$$
(8)

which entirely fixes the inflationary potential including the constant term. Therefore, the field-independent term is fixed at the scale of inflation, too.

Let us now come back to the idea of quintessence. The cosmological constant is static which means once it is fixed (e.g., to describe the rate of accelerating expansion of the Universe today) its value remains constant over the history of the Universe. Thus, extrapolating back in time to the early Universe, it has a very small value compared to the Planck mass. It would be more natural for the dark energy to start with an energy density similar to the density of matter and radiation in the early Universe. The concept of quintessence was introduced to overcome this problem which assumes a scalar field similar to inflaton with negative pressure but with a very large wavelength. The equation of state of the quintessence is dynamic, time-evolving and given by (5) which has been used for the inflaton case, too. Therefore, it does not matter whether one relies on the idea of quintessence or on the cosmological constant. The field-independent term of the action is involved in the theoretical model, and its value is determined.

Thus, the Standard Model of Cosmology requires two periods of accelerated expansion: in the early Universe, when the Universe doubles in size in every 10^{-35} s, and today, when the doubling time is 50 orders of magnitude greater. The field-independent term of the action has to be fine-tuned in each period of acceleration, thus a reliable theory should take into account the change of this term over the time-evolution of the Universe. If the time-evolution of the Universe and the momentum scale of renormalization is related to each other (inversely), then the required variation of the cosmological constant in time can be produced by RG methods.

Indeed, the essential idea of the nonperturbative RG analysis is to describe the evolution of the self-interaction potential from the UV (from the Planck scale or from over the Planck scale) to the scale of inflation and towards to the low-energy (IR) limit, and of course, it is an important question how to describe the RG running of the constant term. The asymptotic safety scenario of quantum gravity is designed for that purpose, and its essence is the existence of a non-Gaussian UV fixed point. To summarise its cornerstones, one can start from the simplest realization of Quantum Einstein Gravity (QEG) which is the Einstein-Hilbert truncation of the effective average action

$$\Gamma_k = \frac{1}{16\pi G_k} \int d^4x \sqrt{-g} \left(R - 2\Lambda_k\right),\tag{9}$$

where g is the determinant of the metric tensor, R is the Ricci scalar and the scale-dependent parameters are the cosmological constant Λ_k and the Newton coupling G_k . The field independent term $V(\phi = 0)$ of the scalar potential (2) is related to the cosmological constant, i.e., $V(\phi = 0) = m_p^2 \Lambda$. The scale-dependence of Λ_k implies the scale-dependence of $V_k(\phi = 0)$, and it is analyzed in terms of dimensionless couplings, $\lambda_k \equiv \Lambda_k k^{-2}$, $g_k \equiv G_k k^2$ with the help of the β -functions, see for example [3]

$$k\partial_k g_k = \beta_g , \qquad k\partial_k \lambda_k = \beta_\lambda \tag{10}$$

which are calculated by the Litim regulator [111]

$$\beta_g = (2 + \eta_N)g_k, \qquad \beta_\lambda = (\eta_N - 2)\lambda_k + \frac{g_k}{12\pi} \left[\frac{30}{1 - 2\lambda_k} - 24 - \frac{5}{1 - 2\lambda_k} \eta_N \right], \tag{11}$$

where the anomalous dimension of Newton's constant $\eta_N = G_k^{-1} k \partial_k G_k$ is given by

$$\eta_N = \frac{g_k B_1}{1 - g_k B_2} \,, \tag{12}$$

where

$$B_1 = \frac{1}{3\pi} \left[\frac{5}{1 - 2\lambda_k} - \frac{9}{(1 - 2\lambda_k)^2} - 7 \right], \qquad B_2 = -\frac{1}{12\pi} \left[\frac{5}{1 - 2\lambda_k} - \frac{6}{(1 - 2\lambda_k)^2} \right]. \tag{13}$$

The RG flow diagram based on the above β -functions is plotted on Fig. 1.




FIG. 1: We present the RG flow diagram of QEG based on the Einstein-Hilbert trunction given in Eq. (9).

The β -functions contain the information on fixed points g_{\star} and λ_{\star} of the RG flow where the beta functions vanish simultaneously. They give rise to two fixed points: the Gaussian (G) UV fixed point situated at $(g_{\star}, \lambda_{\star}) = (0,0)$ and the non-Gaussian (NG) UV fixed point located at $(g_{\star}, \lambda_{\star}) = (0.707, 0.193)$. As we argued in the introduction, the existence of the non-Gaussian UV fixed point can solve important problems of quantum gravity. In order to find cosmological applications, the running RG cutoff k is identified with a typical length scale of the system [5]. There are several types of cutoff identifications [5]; among these, one finds $k \sim t^{-1}$, where t is the cosmic time, or $k \sim H(t)$ where H(t) is the Hubble parameter, or $k \sim T$, where T is the temperature of the cosmic plasma. The idea is to use RG running to connect the physics of various energy scales. For example, one should find the non-Gaussian fixed point above the Planck scale $k \gg m_p = 2.4 \times 10^{27}$ eV. By contrast, cosmic inflation takes place below the Planck scale $k = k_{inf} = 10^{22}$ eV, and the well-known value of Newton's constant is fixed by laboratory experiments $G_k = G = 6.67 \times 10^{-57} \text{ eV}^{-1}$ at low-energies $k = k_{lab} = 10^{-5} \text{ eV}$. Finally, one should mention the accelerated expansion of the Universe at present which requires $\Lambda_k = \Lambda = 4 \times 10^{-66} \text{ eV}^2$ at the scale $k = k_{\text{Hub}} = 10^{-33} \text{ eV}$. The nonperturbative RG (using various extension of the Einstein-Hilbert truncation, see for example [6]) is capable to build up connection between these scales and cover many orders of magnitude in change of couplings, like the Newton and the cosmological constants.

Our goal here is to consider the RG flow of the field independent term in the presence of scalar fields, i.e. when the coupling constants of the action (2) are considered as running parameters similarly to (9),

$$\Gamma_k[\phi] = \int d^4x \sqrt{-g} \left[\frac{1}{16\pi G_k} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \,\partial_\nu \phi - V_k(\phi) \right], \tag{14}$$

where the scalar potential is usually expanded in terms of the field. If this expansion is terminated at the quartic order, it has the following form

$$V_k(\phi) = V_k(0) + \frac{1}{2}m_k^2\phi^2 - \frac{1}{4!}\lambda_{4,k}\phi^4, \qquad V_k(0) \equiv \frac{2\Lambda_k}{16\pi G_k},$$
(15)

which is one of the simplest scenarios when a single real scalar field is coupled to gravity. For a detailed study of scalar fields coupled to asymptotically safe quantum gravity see for example [79] where the existence of the non-Gaussian fixed point was shown for the simple case of (15) with non-minimal coupling to gravity. The fixed points of the RG flow for a scalar field in curved space with non-minimal coupling is discussed in [80], too where the RG equation for the scalar potential in the so-called Local Potential Approximation of the Wetterich equation with the Litim cutoff reads as

$$k\partial_k V_k(\phi) = \mu_d k^d \frac{k^2}{k^2 + \partial_\phi^2 V_k(\phi)},$$
(16)

with $\mu_d = 1/[(4\pi)^{d/2}\Gamma(d/2+1)]$. The existence of the Gaussian (G) and non-Gaussian (NG) fixed points and the RG flow of the full (dimensionless) potential is discussed, for example, in Refs. [79] and [80] but no β function is given for the field-independent term. In this work, we are interested in the RG flow of the field-independent term, $V_k(0)$, with a special attention on its UV limit.

Let us discuss the RG flow of the cosmological constant in the absence of quantum gravity effects, but take into account the RG equation (16) for the scalar potential (15). We use the relation $\tilde{V}_k(0) \equiv \frac{2\lambda_k}{16\pi g_k}$ which connects dimensionless couplings and gives

$$k\partial_k\lambda_k = 8\pi \left(g_k k\partial_k \tilde{V}_k(0) + \tilde{V}_k(0) k\partial_k g_k\right)$$
(17)

In the absence of quantum gravity effects, i.e., assuming a scale-independent dimensionful Newton's constant, $G_k = G$, the anomalous dimension vanishes because $\eta_N = G^{-1} k \partial_k G = 0$, and one finds $k \partial_k g_k = 2g_k$ which results in a trivial RG scaling $g_k \sim k^2$. The RG flow equation for the dimensionless field-independent term obtained from (16) reads as

$$k\partial_k \tilde{V}_k(0) = \frac{1}{32\pi^2} \left(\frac{1}{1+\tilde{m}_k^2}\right) - 4\tilde{V}_k(0)$$
(18)

which can be used to obtain the RG flow equation for the dimensionless cosmological constant,

$$k\partial_k \lambda_k = \frac{1}{4\pi} g_k \left(\frac{1}{1+\tilde{m}_k^2}\right) - 2\lambda_k.$$
⁽¹⁹⁾

In this approximation, g_k has a trivial scaling, $g_k \sim k^2$, it tends to infinity in the UV limit. The UV scaling of the dimensionless mass term is $\tilde{m}_k^2 \sim k^{-2}$, so, it tends to zero in the UV limit. Although the UV Gaussian fixed point formally exists, it cannot be reached because the corresponding β -function diverges in the UV limit, $g_k \left(\frac{1}{1+\tilde{m}_k^2}\right) \to \infty$ if $k \to \infty$. Our goal in this work is to show that one has to apply additional subtraction terms in order to restore the Gaussian fixed point in (19).

We will discuss that, even though the Gaussian fixed point is known to exist in many quantum field theories, there may be questions regarding the proper definition and retention of the Gaussian fixed point due to presence of the divergent constant term. A point emerging in our discussion is that with the use of subtractions, one can restore the Gaussian fixed point of the pure scalar theory in the RG flow based on the β -functions even if the field-independent term is included. We will show that the inclusion of these subtraction terms do not influence the non-Gaussian fixed point of the gravity-scalar models, but could modify their IR behavior.

In summary, the constant term, i.e. the cosmological constant plays an important role (i) in the physics above the Planck scale, (iii) in the mechanism of inflation in the early Universe, (iii) in the accelerating expansion at present. This justifies our main interest in the RG evolution of the constant term in general, which we extensively discuss in the next section.

III. FIELD-INDEPENDENT TERM IN RG

The modern formulation of nonperturbative RG usually referred as the Wetterich RG equation [19, 20] has the following form for the one-component scalar field theory:

$$k \partial_k \Gamma_k[\phi] = \frac{1}{2} \int \frac{d^d p}{(2\pi)^d} \frac{k \partial_k R_k(p)}{R_k(p) + \Gamma_k^{(2)}[\phi]},\tag{20}$$

8

where k is the RG scale, $\Gamma_k[\phi]$ is the running effective action with its Hessian $\Gamma_k^{(2)}[\phi]$, and $R_k(p)$ is the so-called regulator function. It is illustrative to discuss its connection to the effective action, which has the following form at the one-loop level:

$$\Gamma_{\text{eff}}[\phi] = S_{\Lambda}[\phi] + \frac{1}{2} \int \frac{d^d p}{(2\pi)^d} \ln\left[S_{\Lambda}^{(2)}[\phi]\right] + \mathcal{O}(\hbar^2), \qquad (21)$$

where S_{Λ} is the classical (bare) action. A Pauli-Villars approach is used to regularise the momentum integral which can be divergent at its upper (UV) and lower (IR) bounds. This can be achieved by adding a momentum dependent mass term $\frac{1}{2} \int R_k(p) \phi^2$ to the bare action, and introduce a scale-dependent action

$$\Gamma_k[\phi] \equiv S_{\Lambda}[\phi] + \frac{1}{2} \int \frac{d^d p}{(2\pi)^d} \ln\left[R_k(p) + S_{\Lambda}^{(2)}[\phi]\right], \qquad (22)$$

which recovers the effective action (at one-loop) in the IR limit if the regulator function $R_k(p)$ fulfils the requirements, $R_{k\to 0}(p) = 0$, $R_k(p \to 0) > 0$ [see Eqs. (13)—(15) of Ref. [24]]. The latter condition is important to avoid IR divergences. However, one canonically also imposes the condition

$$R_{k \to \Lambda}(p) = \infty \tag{23}$$

(see Ref. [24]), and thus, in the UV limit, the scale-dependent action reproduces the classical (bare) action only up to a field-independent, constant term. If one can differentiate Eq. (22) with respect to the running scale k (and multiplies both sides by k), then one finds

$$k\partial_k\Gamma_k[\phi] = \frac{1}{2} \int \frac{d^d p}{(2\pi)^d} \frac{k\partial_k R_k(p)}{R_k(p) + S_\Lambda^{(2)}[\phi]},$$
(24)

which recovers the "exact" Wetterich RG equation (20) up to the replacement $S_{\Lambda}^{(2)} \to \Gamma_k^{(2)}$.

Let us come back to various limits of the scale-dependent action Eq. (22). It recovers the effective action in the limit $k \to 0$ and the bare action for $k \to \Lambda$, up to a field-independent but k-dependent term, which we will denote as $V_k(0)$ for reasons which will become obvious immediately,

$$\Gamma_{k\to\Lambda}[\phi] = \Gamma_{\Lambda}[\phi] = S_{\Lambda}[\phi] + \text{const.} = S_{\Lambda}[\phi] + \int d^d x \, V_{k\to\Lambda}(0) \,. \tag{25}$$

This clearly signals that the formulation of the RG evolution of the constant, field-independent term $V_k(0)$ requires special care within the nonperturbative approach implied by the Wetterich equation (see also Sec. 2.3 of Ref. [24]). Moreover, if we implement the condition $R_{k\to\Lambda}(p) = \infty$ on the regulator, then it turns out that in many cases, the "constant term" $V_k(0)$ in Eq. (25), actually is given by a divergent integral.

Therefore, the constant term $V_k(0)$ needs a special treatment in the framework of the nonperturbative RG method. In the following, we will consider cases where $V_k(0)$ can naturally be identified with the zeroth-order term (in ϕ) obtained from the scale-dependent potential $V_k(\phi)$. One might argue that, for many purposes, the precise form of the function $V_k(0)$ is physically irrelevant as it constitutes a field-independent constant. However, there are special cases where the RG evolution of a constant (field-independent) part of the potential has physical meaning. For example, if one aims at a determination of the free energy in a flat background or of the cosmological constant in a general non-flat background, then the problem of unambiguously determining $V_k(0)$ has to be seriously considered.

The explicit form of the nonperturbative RG equation which is suitable for application can be obtained by from (20) using various approximations. Derivative expansion is one of the widely used approximation and its leading order is the local potential approximation (LPA) [110]. Within the LPA, the RG equation (20), using the so-called Litim cutoff [111], reads as

$$k\partial_k V_k = \frac{2\alpha_d}{d} \frac{k^{d+2}}{k^2 + \partial_\phi^2 V_k},$$
(26)

where V_k is the dimensionful scaling potential in d dimensions, k is the running momentum, and $\alpha_d = \Omega_d/(2(2\pi)^d)$ is related to the d-dimensional solid angle $\Omega_d = 2\pi^{d/2}/\Gamma(d/2)$. Eq. (26) has been obtained projecting the exact functional RG equation [17–21] [see Eq. (20)] on a functional ansatz for the scalar field effective action, which contains only the bare kinetic term plus a scale dependent effective potential V_k . Then, Eq. (26) represents the non-perturbative β -function of the effective potential in absence of any renormalization for the non-local operators in the action [22– 24, 27–30].

In the following, we will investigate the role of the RG evolution of the field-independent terms in the framework of the nonperturbative RG approach. We would like to use the RG treatment of the constant term developed for low-dimensional quantum mechanical systems [24] and extend this method to higher dimensions.

IV. RG EVOLUTION OF THE CONSTANT TERM

We now come to the most important point to be discussed in the context of the current work. Namely, from Eq. (26), one could in principle derive an RG equation for the constant term $V_k(\phi = 0)$. Written for the dimensionful potential, one would naïvely obtain the following RG equation for $V_k(0)$ from Eq. (26) in d = 4 dimensions:

$$k\partial_k V_k(0) \stackrel{?}{=} \frac{k^4}{32\pi^2} \frac{k^2}{k^2 + \partial_{\phi}^2 V_k(\phi)|_{\phi=0}} \,. \tag{27}$$

It has already been mentioned above that, within the nonperturbative RG equations used by us, extra care is needed in the analysis of the field-independent terms.

The essence of the problem of the RG scaling of the constant term is the (possible) absence of the Gaussian fixed point which, otherwise, is present, if the constant term is not considered. In other words, the β -function of the constant term should vanish if all couplings are set to zero. If the Gaussian fixed point is retained when the RG flow of the constant term is considered, which is the case in QEG, then one finds no problem with the RG scaling of the constant term. However, if the Gaussian fixed point is missing once the field-independent coupling is included, then one finds problematic UV divergences which requires a subtraction method. It was shown that such divergences occur for the quantum anharmonic oscillator and if one considers it in higher dimensions one has to generalise the subtraction method for d = 4 dimensions. Indeed, in the limit $k^2 \gg \partial_{\phi}^2 V_k(0)$, the solution to Eq. (27) is

$$V_k(0) \stackrel{?}{=} V_{\Lambda}(0) + \frac{k^4 - \Lambda^4}{128\pi^2}, \qquad (28)$$

which would otherwise indicate a rampant quartic divergence of $V_k(0)$ in the limit of large Λ and lead to a considerable change in $V_k(0)$ between the Planck and the GUT scales, possibly requiring some fine-tuning of our model in the UV, i.e., at the Planck scale. In view of the quartic divergence of $V_k(0)$ in the UV, the fine-tuning problem would be of quite an extreme nature and conceivably render the model rather questionable.

In order to address the problem, it is necessary to include a longer discussion. The problem is exacerbated by the fact that $V_k(0)$ could in principle be associated with a cosmological constant term. As shown in the discussion surrounding Eq. (A2)—(A5), for an FLRW metric, one can bring the action into a form resembling a flat-space theory, but upon going back to the original FLRW coordinates, one would realize that $V_k(0)$ indeed can take the role of a cosmological constant [see also Eq. (A6)].

One should observe that the rampant k^4 behavior in Eq. (28) persists, even if we set, e.g., all coupling terms of the model to zero. In principle, one might think that it would be difficult to argue that a quartic divergence could be obtained for the RG running of the constant term $V_k(0)$ of a potential that completely vanishes in the IR. The suspicion arises that the behavior implied by Eq. (28) cannot be physical and must be spurious.

V. DIVERGENCES IN d = 1 OSCILLATORS

We start this section by observing that questions related to the RG running of field-independent constant terms belong to the more subtle questions connected with nonperturbative RG equations. In Sec. 2.3 of Ref. [24], a (0 + 1)dimensional model problem is studied which illustrates the spurious nature of these terms, namely, an anharmonic quantum mechanical oscillator. For a potential of the form

$$V_k(x) = E_{0k} + \frac{1}{2}\omega_k^2 x^2 + \frac{1}{24}\lambda_k^2 x^4 + \dots = \sum_{n=0}^{N_{\text{cut}}} \frac{g_{2n,k}}{(2n)!} x^{2n},$$
(29)

where $E_{0k} \equiv g_{0,k}$, $\omega_k^2 \equiv g_{2,k}$ and $\lambda_k^2 \equiv g_{4,k}$.

One may investigate what happens if one drops the anharmonic term $(\lambda = 0)$, which implies that $\omega_k = \omega_{k=\Lambda} \equiv \omega$. In this limit, the following RG equation is obtained for the constant term E_{0k} [see Eq. (40) of Ref. [24]],

$$\frac{d}{dk}E_{0k} \stackrel{?}{=} \frac{1}{\pi} \frac{k^2}{k^2 + \omega^2} \,. \tag{30}$$

When integrating this equation, one would obtain a spurious linearly divergent ground-state energy of the anharmonic oscillator in the harmonic limit ($\lambda_k \to 0$, i.e., when the anharmonicity vanishes).

This behavior cannot be physical and cannot be trusted. The answer is connected with the observation surrounding Eq. (25), which implies the necessity of subtracting a spurious field-independent constant term if one would like to

10

recover the bare action (in the UV) from the nonperturbative RG equation. In the case of the (0 + 1)-dimensional field theory, the solution is given by subtracting the spurious term from the right-hand side RG equation (30), and to solve instead

$$\frac{d}{dk}E_{0k} \stackrel{!?}{=} \frac{1}{\pi} \left(\frac{k^2}{k^2 + \omega^2} - 1\right).$$
(31)

Integrating this equation, one obtains the correct ground-state energy $\frac{1}{2}\omega$ [see Eq. (40) of Ref. [24]].

Let us consider what happens if one keeps the anharmonic term $(\lambda \neq 0)$. In order to study the dependence of the results on the truncation N_{cut} we use the general form for the potential where the couplings are denoted by g_{2n} . One can derive the flow equations for the running couplings $g_{2n,k}$ by using the subtraction method explained above. For example by using the Litim regulator [111], and the truncation $N_{\text{cut}} = 2$ the flow equations become

$$k\partial_k g_{0,k} = \frac{k}{\pi} \left(\frac{k^2}{k^2 + g_{2,k}} - 1 \right), \tag{32}$$

$$k\partial_k g_{2,k} = -\frac{1}{\pi} \frac{k^3 g_{4,k}}{(k^2 + g_{2,k})^2},\tag{33}$$

$$k\partial_k g_{4,k} = \frac{6}{\pi} \frac{k^3 g_{4,k}^2}{(k^2 + g_{2,k})^3}.$$
(34)

By solving these differential equations we computed the ground state energy, $E_0 = g_{0,k=0}$, with different initial conditions and got consistent results with the solutions of the Schrödinger equation as shown in Fig. 2 and Fig. 3.




FIG. 2: The dependence of the ground state energy of the quantum anharmonic oscillator on the $g_{4,\Lambda}$ initial condition, obtained for a single-well potential ($g_{2,\Lambda} = 4$, $g_{0,\Lambda} = 0$), using the truncation $N_{\text{cut}} = 2$. The solid curve is from the solutions of the Schrödinger equations, while the dashed curve is computed by FRG.

The unphysical behaviour of the ground state energy for the double-well potential in low $g_{4,\Lambda}$ is just the result of our approximations. In fact better approximations improve these results, for example improving the truncation $(N_{\text{cut}} = 3)$ gives an improved curve, see Fig. 4.

Therefore, the use of the subtraction method for the set of RG flow equations recovers the correct results for the quantum anharmonic oscillator in d = 1.

VI. DIVERGENCES IN THE COSMOLOGICAL MODEL

A. General Discussion and Idea

If the need for such subtractions arises in the context of (0 + 1)-dimensional field theories, then we can expect similar or even aggravated problems in our (3 + 1)-dimensional case. We should clarify that the subtraction leading from Eq. (30) to Eq. (31) is not rigorously derived in Ref. [24], but constitutes more than an *ad hoc* subtraction. Namely, as pointed out in the text preceding Eq. (37) of Ref. [24], the physical requirement is that the constant term in the potential must remain zero under the RG in the limit of vanishing parameters $\omega \to 0$, and $\lambda \to 0$. As pointed out in between Eqs. (36) and (37) of Ref. [24], this requirement fixes the subtraction term. The subtraction leading to Eq. (31) is justified by the fact that it reproduces the known ground-state energy of the harmonic oscillator.




FIG. 3: The dependence of the ground state energy of the quantum anharmonic oscillator on the $g_{4,\Lambda}$ initial condition, obtained for a double-well potential $(g_{2,\Lambda} = -4, g_{0,\Lambda} = \frac{3g_{2,\Lambda}^2}{2g_{4,\Lambda}})$, using the truncation $N_{\text{cut}} = 2$. The solid curve is from the solutions of the Schrödinger equations, while the dashed curve is computed by FRG.




FIG. 4: The dependence of the ground state energy of the quantum anharmonic oscillator on the $g_{4,\Lambda}$ initial condition, obtained for a double-well potential $(g_{2,\Lambda} = -4, g_{0,\Lambda} = \frac{3g_{2,\Lambda}^2}{2g_{4,\Lambda}})$, using the truncation $N_{\text{cut}} = 3$. The solid curve is from the solutions of the Schrödinger equations, while the dashed curve is computed by FRG.

Based on the analogy with Ref. [24], one might investigate if a valid subtraction scheme for the field-independent term in a nonperturbative RG could be obtained by subtracting from the naive RG equation (27) the spurious asymptotic (UV) terms which cause the unphysical divergences. One should take into account that one single subtraction, as in Eq. (31), may not be enough. In a different context, namely, in the case of perturbatively renormalizable theories such as quantum electrodynamics (QED), one sometimes has to introduce more than one subtraction term. E.g., the regularization of the vacuum polarization integral, discussed in Eq. (7.3) ff. of Ref. [112] necessitates the introduction of more than one "heavy fermion" in order to eliminate a spurious quadratic divergence, for large loop momenta. In the case of the nonperturbative RG method used here, one can show that, in general, the RG equation for $V_k(0)$ implies a dependence with the functional form k^d in the UV (when no subtractions are applied), consistent with the cases d = 1 discussed in Eq. (30) and the case d = 4 discussed in Eq. (28). This observation might suggest that more than one subtraction could be required for higher dimensions.

Let us try to formulate possible, but not unique, further requirements, in terms of conjectures, that could be imposed on the subtracted, physical, RG evolution of the constant term. These constitute mild generalizations of the considerations reported in Ref. [24].

- (i) The RG evolution of $V_k(0)$ must vanish in the UV, in the limit of vanishing expansion coefficients of the model.
- (ii) The subtraction terms, originating from the UV, should have a polynomial functional form (in k), as they are obtained from an asymptotic expansion of integrals obtained in the limit $k \to \Lambda$, which typically gives rise to an asymptotic expansion with polynomial terms.
- (*iii*) The subtractions cannot possibly influence the IR behavior of the RG, as they originate purely from the UV behavior of the regulator given in Eq. (23).

Condition *(iii)* implies that the subtractions cannot induce IR divergences. This means that one cannot subtract "too many" terms; otherwise one incurs infrared problems.

B. Double Subtraction

As suggested by the above general considerations, we observe that a single subtraction, based on the replacement

$$k\partial_k V_k(0) = \frac{k^4}{32\pi^2} \frac{k^2}{k^2 + \partial_\phi^2 V_k(0)} \to \frac{k^4}{32\pi^2} \left[\frac{k^2}{k^2 + \partial_\phi^2 V_k(0)} - 1 \right] \to -\frac{k^2}{32\pi^2} \partial_\phi^2 V_k(0) , \qquad k \to \infty , \tag{35}$$

in Eq. (27), leaves a quadratic term (in k) on the right-hand side of Eq. (27) in the UV limit. This subtraction term modifies the RG evolution of the cosmological constant. Thus, Eq. (19) is changed as

$$k\partial_k\lambda_k = \frac{1}{4\pi}g_k\left(\frac{1}{1+\tilde{m}_k^2}-1\right) - 2\lambda_k,\tag{36}$$

from which one observes that the corresponding β -function does not diverge in the UV limit but tends to a non-vanishing finite value, i.e., the expression $g_k \left(\frac{1}{1+\tilde{m}_k^2}-1\right)$ approaches a negative constant, where we keep in mind that $g_k \sim k^2$ and $\tilde{m}_k^2 \sim k^{-2}$ in the UV limit.

It is well known that comparable subtractions produce the correct result for the free energy of certain interacting quantum-statistical mechanics models (see Chap. 8 of Ref. [113]). Further remarks on related issues can also be found in Ref. [114, 115]. One might argue that the negative value of $V_k(0)$ in the UV, implied if we assume Eq. (35) to be valid in the UV, is akin to the negative mass square acquired by the Higgs particle upon considering tadpole diagrams involving fermionic loops (Yukawa coupling), which also diverges quadratically in Λ . Furthermore, one might argue that this problem would be on the same level as the so-called hierarchy problem of the Standard Model. Conversely, if a subtraction is carried out in the RG evolution of the field-independent term [see Eq. (35)], the hierarchy problem for the field-independent term in our model for the inflaton, is reduced to the same level at which the Higgs particle mass of the Standard Model itself suffers from a comparable RG running and concomitant hierarchy problem. (The lack of experimental evidence for supersymmetric particles, whose presence could potentially alleviate the hierarchy problem of the Standard Model, does not need to be stressed in the current context.)

The leading term on the right-hand side of Eq. (35) has a peculiar property: It vanishes for vanishing couplings of the model but diverges for $\Lambda \to \infty$. The physical condition of a vanishing RG evolution for the field-independent term in the limit of vanishing couplings is implemented in Eq. (35), but the limit is not approached uniformly in the sense that, colloquially speaking, if Λ goes to infinity faster than the couplings go to zero, there is a remaining term. If, in addition to conjectures (i), (ii), and (iii), we also conjecture that the limit of vanishing parameters (expansion coefficients) be approached smoothly, then the second subtraction would be required. Of course, this additional subtraction eventually has to be justified on a calculation based on first principles. It is permissible, though, to consider a more favourable scenario where the use of an optimized regulator, which enables us to recover the bare action from the solution of the RG equation in the UV, will translate into a modified RG evolution for the constant term, which involves more than one subtraction in the UV, for our four-dimensional case. Let us therefore investigate the doubly-subtracted RG evolution

$$k\partial_k V_k(0) = \frac{k^4}{32\pi^2} \left(\frac{k^2}{k^2 + \partial_\phi^2 V_k(0)} - 1 + \frac{\partial_\phi^2 V_k(0)}{k^2} \right), \qquad k \to \infty,$$
(37)

which is still infrared safe because of the k^4 prefactor. The subtraction of even more asymptotic terms would contradict conjecture *(iii)*. In the UV limit, one then obtains

$$k\partial_k V_k(0) \sim \frac{[\partial_\phi^2 V_k(0)]^2}{32\pi^2}, \qquad k \to \infty.$$
(38)

If the double subtraction implied by Eq. (37) holds for a modified RG evolution which avoids the unphysical divergences, then this would lead to a cosmological constant that diverges in the UV, but only logarithmically.

Indeed, the RG flow equation (19) of the dimensionless cosmological constant is modified as

$$k\partial_k\lambda_k = \frac{1}{4\pi}g_k\left(\frac{1}{1+\tilde{m}_k^2} - 1 + \tilde{m}_k^2\right) - 2\lambda_k,\tag{39}$$

from which one observes that the β -function tends to zero in the UV limit, since we have the asymptotic behavior that $g_k \left(\frac{1}{1+\tilde{m}_k^2}-1+\tilde{m}_k^2\right) \to 0$ if $k \to \infty$, where we keep in mind that $g_k \sim k^2$ and $\tilde{m}_k^2 \sim k^{-2}$ in the UV limit.

In Ref. [57] the authors show that no M^4 term is present (due to the subtraction) and the RG scale-dependence is characterised by M^2 and $\log(M^2)$ terms, where M denotes the running RG scale. Here we use a double subtraction in (38), thus, there is no k^4 term in the UV limit of the RG equation. If one neglects the scaling of the mass term $m_k \to m$, the solution of the flow equation can be obtained and reads

$$V_k(0) \sim \frac{1}{32\pi^2} m^4 \log(k), \qquad k \to \infty.$$

$$\tag{40}$$

Relating the RG scales $k \sim M$ in (40) and by setting $\xi = 0$ in equation (6.9) of [57], the two runnings can be compared, as is done in Sec. VIC.

Let us explore a (possibly curious) analogy to the emergence of other logarithmic divergences in quantum field theory, which seem to have a long history, ever since Bethe, in his first calculation of the Lamb shift [116], obtained a logarithmically divergent result (expressed in terms of a UV cutoff parameter), in addition to the logarithmic sum over hydrogenic excited state which bears his name. The logarithmic divergence was classified as nonproblematic in Ref. [116], as it was clear that a natural cutoff (in the case of Ref. [116], the electron mass scale) exists. (In our case, of course, a natural UV cutoff is found at $k \sim \Lambda$, the Planck scale.) The problem was later analyzed in greater detail by French and Weisskopf [117], and Kroll and Lamb [118], as well as Feynman [119], who clarified the matching of UV and IR divergences (see p. 777 of Ref. [119]). Without drawing any further analogies here, we note that the original subtraction introduced by Bethe in Ref. [116] was completely *ad hoc* at the time; the full physical picture was clarified later. May it be permissible to mention that corresponding subtractions, to obtain physically acceptable results in problems of advanced classical electrodynamics, have recently been discussed in Chap. 8 of Ref. [120].

The subtractions introduced in Eqs. (35) or (37) are not unique, and we do not have a rigorous derivation at present beyond the considerations described above. However, the physical requirements that should to be fulfilled by the subtracted RG evolution of the constant term, as formulated in the conjectures (i)-(iii) above, are in agreement with the physical requirements expressed between Eqs. (36) and (37) of Ref. [24]. The conjecture and the subtraction terms enter as an additional input into our calculations. In general, it might be possible to avoid the unphysical divergences of the constant, field-independent terms via a suitable modification of the condition (23) imposed on the regulator function.

C. Comparison with Other Results

In the previous subsection we proposed a new subtraction in order to get rid of the spurious divergent terms k^4 and k^2 in the RG flow equation of the constant (field-indepedent) term of the action, which is related to the cosmological constant. In the literature, the RG running of the cosmological constant has been already suggested using subtraction ideas in order to solve the cosmological constant problem. Thus, let us compare them to our double-subtracted RG equation (38) and its solution.

In [121], slightly generalized DeWitt-Schwinger adiabatic renormalization subtractions are proposed in curved space to include an arbitrary renormalization mass scale μ . The running of the cosmological constant is obtained in equation (21) of Ref. [121]. A similar running is given by Eq. (14) of Ref. [122], which reads as

$$\Lambda(\mu) = \Lambda_0 - \frac{1}{128\pi^2} \left(-(\mu^4 - \mu_0^4) + 2m^2(\mu^2 - \mu_0^2) - 2m^4 \log\left(\frac{m^2 + \mu^2}{m^2 + \mu^4}\right) \right), \tag{41}$$

and it only differs from equation (21) of [121] in the prefactor of the second term of the right hand side by a multiplicative factor of 2. The prefactor is, however, scheme dependent and it also depends on the physical content of the studied Lagrangian. Thus, it is not surprising that one finds different values in the literature [see Eq. (33) of [122]]. In Eq. (41), the running RG scale is denoted by μ with the UV value μ_0 . Regarding the RG running, three distinct scale dependencies are present; a term that is proportional to μ^4 , one with μ^2 and lastly a logarithmic term log μ^2 . The RG running of the first two, especially of the μ^4 term, can cause a rampant divergent behavior.

Now, let us take a look at the leading terms of Eq. (27) that do not yet contain subtractions. Insted of μ we denoted the RG running scale by k. In the limit $k^2 \gg \partial_{\phi}^2 V_k(0)$, the solution to Eq. (27) is Eq. (28),

$$V_k(0) \sim +\frac{1}{128\pi^2}k^4, \qquad k \to \infty,$$
 (42)

which indicates the same rampant quartic divergence of $V_k(0)$. The prefactor $1/(128\pi^2)$ is also identical to the one in Eq. (41). If one uses a single subtraction (35) also proposed in [24], then the leading term in the UV yields the solution

$$V_k(0) \sim -\frac{1}{64\pi^2} m^2 k^2, \qquad k \to \infty.$$
 (43)

This still contains a k^2 divergence in the UV limit, so, we propose a double subtraction, see equation (38), which has only a mild logarithmic divergence in the UV limit. For constant mass, its solution can be written as (40)

$$V_k(0) \sim +\frac{1}{64\pi^2} m^4 \log(k^2), \qquad k \to \infty.$$
 (44)

Again, the prefactors $1/(64\pi^2)$ also match exactly the prefactors of equation (41).

In Ref. [57] the so-called Running Vacuum Model has been studied which assumes a scale-dependent vacuum, i.e, a running cosmological constant. The authors of [57] propose a suitable subtraction of the so-called Minkowski contribution in the framework of the adiabatic regularization method which results in an RG scaling for the running vacuum described by equation (6.9) of [57]

$$\rho_{\rm vac}(M) = \rho_{\rm vac}(M_0) + \frac{3}{16\pi^2} \left(\xi - \frac{1}{6}\right) H^2 \left[M^2 - M_0^2 - m^2 \ln \frac{M^2}{M_0^2}\right] - \frac{9}{16\pi^2} \left(\xi - \frac{1}{6}\right)^2 \left(\dot{H}^2 - 2H\ddot{H} - 6H^2\dot{H}\right) \ln \frac{M^2}{M_0^2}, \qquad (45)$$

where the running RG scale is denoted by the mass-like term M while its UV value is given by M_0 . The result of the author's subtraction method is the absence of the M^4 term in the scaling relation for the running vacuum. This has great importance; in the UV limit (when $M \to \infty$), the M^4 is a well-known source of exceedingly large contributions. The term ξ stands for the non-minimal coupling to gravity.

One can relate the RG scale of Ref. [57] [denoted as M^2 in Eq. (45)] and the one used by us (which we denote as k); the result is the proportionality $k \propto M$. Therefore, one can find very interesting similarities between our approaches. In our case, we have minimal coupling, i.e., we do not include an extra $\xi R \phi^2$ coupling between gravity and the matter content, and thus $\xi = 0$. Thus, for minimal coupling, i.e., for $\xi = 0$, the numerical prefactors of the M^2 and $\log(M^2)$ terms are $-1/(32\pi^2)$ and $1/(32\pi^2)$. Despite the factor of 2 difference compared to our results, we can say that the functional forms encountered in the RG running in Ref. [57] and here are compatible. In Ref. [57] the authors show that no M^4 term is present (due to their subtraction of the Minkowskian contribution), and the RG scale-dependence is characterised by the M^2 and $\log(M^2)$ terms only. Arguments for the absence of the quartic M^4 term are also presented in Refs. [123, 124], where similar flow equations are obtained. If a double-subtraction method is used (40), only a logarithmic scale dependence remains.

VII. CONCLUSIONS

In this work, we have discussed the role of the constant term in the non-perturbative RG evolution. In particular, we investigated whether the rampant divergent terms k^2 and k^4 which naturally appear in the RG equation for d = 4 dimensions for the scalar inflaton field (in the absence of the Gaussian fixed point) can be removed by a suitable subtraction method. These divergent terms are the consequence of the construction of the functional RG method and considered as unphysical. They make the application of the functional RG method on the proper treatment of the cosmological constant very questionable if the Gaussian fixed point is absent.

Renormalisation and RG scaling could be a possible tool to handle the cosmological problem and they are under intense debate, see for example the recent works, [56, 57]. In Ref. [57] the Running Vacuum Model has been studied which assumes a running cosmological constant where the running vacuum described by equation (6.9) of [57] and the running RG scale is denoted by the mass-like term M. Based on the subtraction method of [57] it is possible to remove the divergent M^4 term and RG scale-dependence of the running vacuum is characterised with M^2 and $\log(M^2)$ terms only.

Here we showed how the subtraction method can be improved further in order to eliminate from the RG scaling of the running vacuum, not just the k^4 (i.e., M^4 of Ref. [57]) but also the k^2 (i.e., M^2 of Ref. [57]) terms which results in a purely logarithmic dependence on the scale, thus, the unphysical rampant behaviour of the naive approach is handled. As a future improvement on our subtraction method we think it would be interesting to include the contributions of a theory that is non-minimally coupled to gravity ($\xi \neq 0$).

Acknowledgments

This work was supported by the ÚNKP-20-4-I New National Excellence Program of the Ministry for Innovation and Technology from the source of the National Research, Development and Innovation Fund and by the National Science Foundation (Grant PHY–2110294). Useful discussions with Z. Trocsanyi and G. Somogyi are gratefully acknowledged. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC2181/1-390900948 (the Heidelberg STRUCTURES Excellence Cluster). The CNR/MTA Italy-Hungary 2019-2021 Joint Project "Strongly interacting systems in confined geometries" also is gratefully acknowledged.

Appendix A: Field-Independent Term and Cosmological Constant

The standard model of cosmology implies the requirement of an exponentially fast expansion [125–129] of the early Universe which is usually achieved by assuming a hypothetic inflaton field which slowly rolls down from a potential hill towards its minimum. Particle physics provides us with possible candidates for the inflaton field. A first guess involves a scalar field, ϕ , as a candidate for the inflaton field, with a (Euclidean) action that includes the scalar curvature R with minimal coupling to gravity,

$$S[\phi] = \int d^4x \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi + V(\phi) + \frac{m_p^2}{2} R \right], \qquad (A1)$$

where the Planck mass $m_p^2 \equiv 1/(8\pi G)$ has been used and $\sqrt{-g} = \sqrt{-\det g_{\mu\nu}} = a^3$ with $g_{\mu\nu} = \operatorname{diag}(-1, a^2, a^2, a^2)$. Here, the scale factor a = a(t) describes the cosmological scaling in the Friedmann–Lemaître–Robertson–Walker (FLRW, see Ref. [88–91]) metric (in our units, the speed of light is $c \equiv 1$ and $\hbar \equiv 1$).

The FLRW metric is given by

$$g_{\mu\nu} = \text{diag}(-1, a(t)^2, a(t)^2, a(t)^2), \qquad (A2)$$

with the line element

$$ds^2 = -dt^2 + a(t)\,d\vec{r}^2\,,\tag{A3}$$

where $\vec{r} = (x, y, z)$. For scalar fields, the partial and covariant derivatives coincide, and hence, we can replace, in Eq. (A1),

$$g^{\mu\nu} \nabla_{\mu} \phi \nabla_{\nu} \phi \to -\phi \nabla^{\mu} \nabla_{\mu} \phi = -\phi \left(-\frac{\partial^2}{\partial t^2} + \frac{1}{a(t)^2} \frac{\partial^2}{\partial \vec{r}^2} \right) \phi.$$
(A4)

One may stretch the spatial coordinates, in a "flattening" transformation, according to

$$\vec{r}' = a(t')\vec{r}, \qquad \phi(t,\vec{r}) = \phi'(t',\vec{r}'), \qquad t' = t.$$
 (A5)

After this transformation, the action is brought into the form

$$S[\phi] = \int dt' \int d^3r' \left[-\frac{1}{2}\phi' \left(-\frac{\partial^2}{\partial t'^2} + \frac{\partial^2}{\partial \bar{r}'^2} \right) \phi' + V(\phi') + \frac{m_p^2}{2}R \right], \tag{A6}$$

If we now re-identify

$$t' \to i t$$
 (Wick rotation), $\vec{r}' \to \vec{r}$, $\phi'(t', \vec{r}') = \phi(t, \vec{r})$, $V(\phi') \to V(\phi)$, (A7)

and ignore the curvature term which is the cosmological constant term, then we obtain

$$S[\phi] = \int d^4x \left[\frac{1}{2} (\partial_\mu \phi)^2 + V(\phi) \right], \qquad (A8)$$

with a fully local (Euclidean) action of our theory, formulated in Euclidean four-dimensional space. In comparison to Eqs. (A1), Eq. (A8) has the same structure as would otherwise be expected for flat space, and makes the model amenable to a nonperturbative RG analysis using established techniques [111].

^[1] M. Reuter, Nonperturbative evolution equation for quantum gravity, Phys. Rev. D. 57, 971–285 (1998).

- [2] M. Reuter, F. Saueressig, Quantum Gravity and the Functional Renormalization Group: The Road towards Asymptotic Safety (Cambridge Monographs on Mathematical Physics, Cambridge University, 2019).
- [3] A. Bonanno, F. Saueressig, Asymptotically safe cosmology a status report, C. R. Phys. 18, 254–264 (2017).
- [4] A. Platania, From renormalization group flows to cosmology, Front. Phys. 8, 188 (2020).
- G. Gubitosia, R. Ooijera, C. Ripkena, F. Saueressig, Consistent early and late time cosmology from the RG flow of gravity, J. Cosmol. Astropart. Phys. 1805, 004 (2018).
- [6] A. Bonanno, A. Platania, Asymptotically safe inflation from quadratic gravity, Phys. Lett. B 750, 638-642 (2015).
- [7] J. Biemans, A. Platania, F. Saueressig, Renormalization group fixed points of foliated gravity-matter systems, J. High Energy Phys. 1705, 1–41 (2017).
- [8] J. Biemans, A. Platania, F. Saueressig, Quantum gravity on foliated spacetimes: Asymptotically safe and sound, Phys. Rev. D 95, 086013 (2017).
- [9] A. Bonanno, A. Platania, F. Saueressig, Cosmological bounds on the field content of asymptotically safe gravity-matter models, Phys. Lett. B 784, 229–236 (2018).
- [10] A. Bonanno, B. Koch, A. Platania, Cosmic censorship in quantum Einstein gravity, Class. Quant. Grav. 34, 095012 (2017).
- [11] A. Bonanno, S. J. G. Gionti, A. Platania, Bouncing and emergent cosmologies from ADM RG flows, Class. Quant. Grav. 35, 065004 (2018).
- [12] A. Eichhorn, A. Platania, M. Schiffer, Lorentz invariance violations in the interplay of quantum gravity with matter, Phys. Rev. D 102, 026007 (2020).
- [13] A. Bonanno, B. Koch, A. Platania, Gravitational collapse in quantum Einstein gravity, Found. Phys. 48, 1393–1406 (2018).
- [14] A. Platania, F. Saueressig, Functional Renormalization Group Flows on Friedman-Lemaitre-Robertson-Walker backgrounds, Found. Phys. 48, 1291–1304 (2018).
- [15] S. Nagy, J. Krizsan, K. Sailer, Infrared fixed point in quantum Einstein gravity, J. High Energy Phys. 1207, 1–18 (2012).
- [16] P. Dona, A. Eichhorn, R. Percacci, Matter matters in asymptotically safe quantum gravity, Phys. Rev. D 89, 084035 (2014).
- [17] F. J. Wegner, A. Houghton, Renormalization group equation for critical phenomena, Phys. Rev. A 8, 401–412 (1973).
- [18] J. Polchinski, *Renormalization and effective Lagrangians*, Nucl. Phys. B **231**, 269–295 (1984).
- [19] C. Wetterich, Exact evolution equation for the effective potential, Phys. Lett. B 301, 90–94 (1993).
- [20] T. R. Morris, The Exact renormalization group and approximate solutions, Int. J. Mod. Phys. A 9, 2411–2449 (1994).
- [21] J. Alexandre, J. Polonyi, Functional Callan-Symanzik equation, Ann. Phys. (N. Y.) 288, 37–51 (2001).
- [22] J. Berges, N. Tetradis, C. Wetterich, Non-perturbative renormalization flow in quantum field theory and statistical physics, Phys. Rep. 363, 223–386 (2002).
- [23] J. Polonyi, Lectures on the functional renormalization group method, Cent. Eur. J. Phys. 1, 1–71 (2003).
- [24] H. Gies, Introduction to the Functional RG and Applications to Gauge Theories, in Springer Notes in Physics vol. 852,
 J. Polonyi, A. Schwenk (Eds.), pp. 287–348 Springer (Heidelberg), 2012. See also e-print hep-ph/0611146.
- [25] G. P. Vacca, L. Zambelli, Functional renormalization group flow equation: Regularization and coarse-graining in phase space, Phys. Rev. D 83, 125024 (2011).
- [26] J. Kovacs, S. Nagy, K. Sailer, Optimized regulator for the quantized anharmonic oscillator, Int. J. Mod. Phys. A 30, 1550058 (2015).
- [27] J. M. Pawlowski, Aspects of the Functional Renormalisation Group, Ann. Phys. (N. Y.) 322, 2831–2915 (2007).
- [28] B. Delamotte, An Introduction to the Nonperturbative Renormalization Group, in Springer Notes in Physics vol. 852, J. Polonyi, A. Schwenk (Eds.), pp. 49–85, Springer (Heidelberg), 2012. See also e-print cond-mat/0702365.
- [29] O. J. Rosten, Fundamentals of the exact renormalization group, Phys. Rep. 511, 177–272 (2012).
- [30] S. Nagy, Lectures on renormalization and asymptotic safety, Ann. Phys. (N. Y.) **350**, 310–346 (2014).
- [31] A. Bonanno, M. Reuter, Cosmology of the Planck era from a renormalization group for quantum gravity, Phys. Rev. D 65, 043508 (2002).
- [32] A. Bonanno, M. Reuter, Cosmology with self-adjusting vacuum energy density from a renormalization group fixed point, Phys. Lett. B 527, 9–17 (2002).
- [33] I. L. Shapiro, J. Sola, H. Stefancic, Running G and Λ at low energies from physics at M_X : possible cosmological and astrophysical implications, J. Cosmol. Astropart. Phys. **2005**, 012 (2005).
- [34] J. Serreau, Renormalization group flow and symmetry restoration in de Sitter space, Phys. Lett. B 730, 271–274 (2014).
- [35] M. Reuter, H. Weyer, Renormalization group improved gravitational actions: A Brans-Dicke approach, Phys. Rev. D 69, 104022 (2004).
- [36] A. Babic, B. Guberina, R. Horvat, H. Stefancic, Renormalization-group running cosmologies: A scale-setting procedure, Phys. Rev. D 71, 124041 (2005).
- [37] M. Hindmarsh, I. D. Saltas, f(R) gravity from the renormalization group, Phys. Rev. D 86, 064029 (2012).
- [38] E. J. Copeland, C. Rahmede, I. D. Saltas, Asymptotically safe Starobinsky inflation, Phys. Rev. D 91, 103530 (2015).
- [39] I. L. Shapiro, J. Sola, Scaling behavior of the cosmological constant and the possible existence of new forces and new light degrees of freedom, Phys. Lett. B 475, 236–246 (2000).
- [40] A. Babic, B. Guberina, R. Horvat, H. Stefancic, Renormalization group running of the cosmological constant and its implication for the Higgs particle boson mass in the standard model, Phys. Rev. D 65, 085002 (2002).
- [41] I. L. Shapiro, J. Sola, The scaling evolution of cosmological constant, J. High Energy Phys. 2002, 006 (2002).
- [42] I. L. Shapiro, J. Sola, C. Espana-Bonet, P. Ruiz-Lapuente, Variable Cosmological Constant as a Planck Scale Effect,

Phys. Lett. B 574, 149–155 (2003).

- [43] I. L. Shapiro, J. Sola, On the possible running of the cosmological constant, Phys. Lett. B 682, 105–113 (2009).
- [44] J. Sola, A. Gomez-Valent, J. de Cruz Perez, First Evidence of Running Cosmic Vacuum: Challenging the Concordance Model, Astrophys. J. 836, 43 (2017).
- [45] S. Basilakos, D. Polarski, J. Sola, Generalizing the running vacuum energy model and comparing with the entropic-force models, Phys. Rev. D 86, 043010 (2012).
- [46] V. Salvatelli, N. Said, M. Bruni, A. Melchiorri, D. Wands, Indications of a Late-Time Interaction in the Dark Sector, Phys. Rev. Lett. 113, 181301 (2014).
- [47] A. Gomez-Valent, J. Sola, S. Basilakos, Dynamical vacuum energy in the expanding Universe confronted with observations: a dedicated study, J. Cosmol. Astropart. Phys. 1501, 004 (2015).
- [48] E. D. Valentino, A. Melchiorri, J. Silk, Reconciling Planck with the local value of H_0 in extended parameter space, Phys. Lett. B **761**, 242–246 (2016).
- [49] Y. H. Li, J. F. Zhang, X. Zhang, Testing models of vacuum energy interacting with cold dark matter, Phys. Rev. D 93, 023002 (2016).
- [50] Y. H. Li, J. F. Zhang, X. Zhang, Exploring the full parameter space for an interacting dark energy model with recent observations including redshift-space distortions: Application of the parametrized post-Friedmann approach, Phys. Rev. D 90, 123007 (2014).
- [51] J. Sola, A. Gomez-Valent, J. de Cruz Perez, The H_0 tension in light of vacuum dynamics in the universe, Phys. Lett. B 774, 317–324 (2017).
- [52] E. D. Valentino, A. Melchiorri, O. Mena, Can interacting dark energy solve the H_0 tension?, Phys. Rev. D **96**, 043503 (2017).
- [53] A. A. Costa, X. D. Xu, B. Wang, E. Abdalla, Constraints on interacting dark energy models from Planck 2015 and redshift-space distortion data, J. Cosmol. Astropart. Phys. 1701, 028 (2017).
- [54] C.-G. Park, B. Ratra, Observational Constraints on the Tilted Spatially Flat and the Untilted Nonflat Φ CDM Dynamical Dark Energy Inflation Models, Astrophys. J. 868, 83 (2018).
- [55] E. D. Valentino, A. Melchiorri, O. Mena, S. Vagnozzi, Nonminimal dark sector physics and cosmological tensions, Phys. Rev. D 101, 063502 (2020).
- [56] S. G. Rubin, How to make the physical parameters small, Adv. High Energy Phys. 2020, 1048585 (2020).
- [57] C. Moreno-Pulido, J. Sola Peracaula, Running vacuum in quantum field theory in curved spacetime: renormalizing $\rho_{vac}(H)$ without m^4 terms, Eur. Phys. J. C 80, 1–23 (2020).
- [58] Y. B. Zeldovich, Cosmological Constant and Elementary Particles, JETP Lett. 6, 316–317 (1967) [ZhETF Pis'ma 6, 883–884 (1967)].
- [59] Y. B. Zeldovich, The Cosmological Constant and the Theory of Elementary Particles, Soviet Physics Uspekhi 11, 381–393 (1968).
- [60] S. Weinberg, The cosmological constant problem, Rev. Mod. Phys. 61, 1–23 (1989).
- [61] E. Witten, The Cosmological Constant From The Viewpoint Of String Theory, Contribution to the 4th International Symposium on Sources and Detection of Dark Matter in the Universe (DM 2000), 27–36, hep-ph/0002297.
- [62] K. G. Wilson, Renormalization Group and Strong Interactions, Phys. Rev. D 3, 1818–1846 (1971).
- [63] K. G. Wilson, J. Kogut, The renormalization group and the ϵ expansion, Phys. Rep. 12, 75–200 (1974).
- [64] K. G. Wilson, The renormalization group: Critical phenomena and the Kondo problem, Rev. Mod. Phys. 47, 773–840 (1975).
- [65] J. Zinn-Justin, Quantum field theory and critical phenomena, Oxford University Press (Oxford), [Int.Ser.Monogr.Phys. 113, (2002)].
- [66] J. C. Le Guillou, J. Zinn-Justin, Critical Exponents for the n-Vector Model in Three Dimensions from Field Theory, Phys. Rev. Lett. 39, 95–98 (1977).
- [67] J. C. Le Guillou, J. Zinn-Justin, Critical exponents from field theory, Phys. Rev. B 21, 3976–3998 (1980).
- [68] A. Bonanno, A. Platania, Asymptotically safe inflation from quadratic gravity, Phys. Lett. B 750, 638–642 (2015).
- [69] G. Gubitosi, R. Ooijer, C. Ripken, F. Saueressig, Consistent early and late time cosmology from the RG flow of gravity, J. Cosmol. Astropart. Phys. 12, 004 (2018).
- [70] I. G. Márián, N. Defenu, U. D. Jentschura, A. Trombettoni, I. Nándori, *Renormalization-Group Running Induced Cosmic Inflation*, J. Cosmol. Astropart. Phys. **2020**, 028 (2020).
- [71] J. Alexandre, N. Defenu, G. Grigolia, I. G. Marian, D. Mdinaradze, A. Trombettoni, J. Turoci, I. Nandori, Quantization and renormalization of non-differentiable potentials, arXiv:2112.14696 [hep-th].
- [72] T. Kugo, K. Yoshioka, Probing extra dimensions using Nambu-Goldstone bosons, Nucl. Phys. B 594, 301–328 (2001).
- [73] M. Bando, T. Kugo, T. Noguchi, K. Yoshioka, Brane fluctuations and suppression of Kaluza-Klein mode couplings, Phys. Rev. Lett. 83, 3601–3604 (1999).
- [74] A. Dobado, A. L. Maroto, The dynamics of the Goldstone bosons on the brane, Nucl. Phys. B 592, 203–218 (2001).
- [75] J. A. R. Cembranos, A. Dobado, A. L. Maroto, Brane-Skyrmions and wrapped states, Phys. Rev. D 65, 026005 (2001)
- [76] J. Alexandre, D. Yawitch, Branon stabilization from fermion-induced radiative corrections, Phys. Lett. B 676, 184–187 (2009).
- [77] Y. Burnier, K. Zuleta, Effective action of a five-dimensional domain wall, J. High Energy Phys. 0905, 065 (2009).
- [78] J. Alexandre, D. Yawitch, Comments on branon dressing and the standard model, New J. Phys. 12, 043027 (2010).
- [79] A. Eichhorn, M. Pauly, Constraining power of asymptotic safety for scalar fields, Phys. Rev. D 103, 026006 (2021).
- [80] B. S. Merzlikin, I. L. Shapiro, A. Wipf, O. Zanusso, Renormalization group flows and fixed points for a scalar field in

curved space with nonminimal $F(\phi)R$ coupling, Phys. Rev. D 96, 125007 (2017).

- [81] A. G. Riess et al., Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant, Astron. J. 116, 1009–1038 (1998).
- [82] S. Perlmutter et al., Measurements of Ω and Λ from 42 High-Redshift Supernovae Astrophys. J. 517, 565–586 (1999).
- [83] V. Sahni, A. A. Starobinsky, The case for a positive cosmological Λ-term, Int. J. Mod. Phys. D 9, 373–443 (2000).
- [84] S. M. Carroll, The Cosmological Constant, Living Rev. Relativ. 4, 1–56 (2001).
- [85] P. J. E. Peebles, B. Ratra, The cosmological constant and dark energy, Rev. Mod. Phys. 75, 559–606 (2003).
- [86] T. Padmanabhan, Cosmological constant-the weight of the vacuum, Phys. Rep. 380, 235–320 (2003).
- [87] E. J. Copeland, M. Sami, S. Tsujikawa, Dynamics of dark energy, Int. J. Mod. Phys. D 15, 1753–1935 (2006).
- [88] A. Friedmann, Über die Krümmung des Raumes, Z. Phys. A 10, 377–386 (1922).
- [89] G. Lemaître, A Homogeneous Universe of Constant Mass and Increasing Radius accounting for the Radial Velocity of Extra-galactic Nebulae, Mon. Not. R. Astron. Soc. 91, 483–490 (1931).
- [90] H. P. Robertson, Kinematics and World-Structure, Astrophys. J. 82, 284–301 (1935).
- [91] A. G. Walker, On Milne's Theory of World-Structure, Proc. London Math. Soc. 42, 90–127 (1937).
- [92] Y. Fujii, Origin of the gravitational constant and particle masses in a scale-invariant scalar-tensor theory, Phys. Rev. D 26, 2580–2588 (1982).
- [93] L. H. Ford, Cosmological-constant damping by unstable scalar fields, Phys. Rev. D 35, 2339–2344 (1987).
- [94] C. Wetterich, Cosmology and the fate of dilatation symmetry, Nucl. Phys. B 302, 668–696 (1988).
- [95] B. Ratra, P. J. E. Peebles, Cosmological consequences of a rolling homogeneous scalar field, Phys. Rev. D 37, 3406–3427 (1988).
- [96] T. Chiba, N. Sugiyama, T. Nakamura, Cosmology with x-matter, Mon. Not. R. Astron. Soc. 289, L5–L9 (1997).
- [97] P. G. Ferreira, M. Joyce, Structure Formation with a Self-Tuning Scalar Field, Phys. Rev. Lett. 79, 4740–4743 (1997).
- [98] P. G. Ferreira, M. Joyce, Cosmology with a primordial scaling field, Phys. Rev. D 58, 023503 (1998).
- [99] E. J. Copeland, A. R. Liddle, D. Wands, Exponential potentials and cosmological scaling solutions, Phys. Rev. D 57, 4686–4690 (1998).
- [100] R. R. Caldwell, R. Dave, P. J. Steinhardt, Cosmological Imprint of an Energy Component with General Equation of State, Phys. Rev. Lett. 80, 1582–1585 (1998).
- [101] I. Zlatev, L. M. Wang, P. J. Steinhardt, Quintessence, Cosmic Coincidence, and the Cosmological Constant, Phys. Rev. Lett. 82, 896–899 (1999).
- [102] T. Chiba, T. Okabe, M. Yamaguchi, Kinetically driven quintessence, Phys. Rev. D 62, 023511 (2000).
- [103] C. Armendariz-Picon, V. F. Mukhanov, P. J. Steinhardt, Dynamical Solution to the Problem of a Small Cosmological Constant and Late-Time Cosmic Acceleration, Phys. Rev. Lett. 85, 4438–4441 (2000).
- [104] C. Armendariz-Picon, V. F. Mukhanov, P. J. Steinhardt, Essentials of k-essence, Phys. Rev. D 63, 103510 (2001).
- [105] P. A. R. Ade et al. [Planck collaboration], Planck 2015 results. XIII. Cosmological parameters, Astron. Astrophys. 594, A13 (2016).
- [106] P. A. R. Ade et al. [Planck collaboration], Planck 2015 results. XX. Constraints on inflation, Astron. Astrophys. 594, A20 (2016).
- [107] P. A. R. Ade et al. [BICEP2 and Keck Array collaboration], Improved Constraints on Cosmology and Foregrounds from BICEP2 and Keck Array Cosmic Microwave Background Data with Inclusion of 95 GHz Band, Phys. Rev. Lett. 116, 031302 (2016).
- [108] D. H. Lyth, Particle Physics Models of Inflation, Lect. Notes Phys. 738 81–118 (2008).
- [109] D. Baumann, TASI Lectures on Inflation, available as arXiv:0907.5424 [hep-th].
- [110] N. Dupuis, L. Canet, A. Eichhorn, W. Metzner, J. M. Pawlowski, M. Tissier, N. Wschebor, The nonperturbative functional renormalization group and its applications, Phys. Rep. 910, 1–114 (2021).
- [111] D. F. Litim, Optimisation of the exact renormalisation group, Phys. Lett. B 486, 92–99 (2000).
- [112] C. Itzykson, J. B. Zuber, *Quantum Field Theory*, (McGraw-Hill, New York, 1980).
- [113] C. J. Pethick, H. Smith, Bose-Einstein condensation in dilute gases, (Cambridge University Press, Cambridge, 2008).
- [114] P. Kopietz, L. Bartosch, F. Schutz, Introduction to the functional renormalization group, (Springer, Berlin, 2010).
- [115] E. H. Lieb, R. Seiringer, J. P. Solovej, J. Yngvason, The mathematics of the Bose gas and its condensation, (Birkhäuser, Basel, 2005).
- [116] H. A. Bethe, The Electromagnetic Shift of Energy Levels, Phys. Rev. 72, 339–341 (1947).
- [117] J. B. French, V. F. Weisskopf, The Electromagnetic Shift of Energy Levels, Phys. Rev. 75, 1240–1248 (1949).
- [118] N. M. Kroll, W. E. Lamb, On the Self-Energy of a Bound Electron, Phys. Rev. 75, 388–398 (1949).
- [119] R. P. Feynman, Space-Time Approach to Quantum Electrodynamics, Phys. Rev. 76, 769–789 (1949).
- [120] U. D. Jentschura, Advanced Classical Electrodynamics, (World Scientific, Singapore, 2017).
- [121] A. Ferreiro, J. Navarro-Salas, Running gravitational couplings, decoupling, and curved spacetime renormalization, Phys. Rev. D 102, 045021 (2020).
- [122] A. Ferreiro, S. Nadal-Gisbert, J. Navarro-Salas, Renormalization, running couplings and decoupling for the Yukawa model in curved spacetime, Phys. Rev. D 104, 025003 (2021).
- [123] E. Kh. Akhmedov, Vacuum energy and relativistic invariance, arXiv:0204048 [hep-th]
- [124] G. Ossola, A. Sirlin, Considerations concerning the contributions of fundamental particles to the vacuum energy density, Eur. Phys. J. C 31, 165–175 (2003).
- [125] A. H. Guth, Inflationary universe: A possible solution to the horizon and flatness problems, Phys. Rev. D 23, 347–356

(1981).

- [126] A. A. Starobinsky, A new type of isotropic cosmological models without singularity, Phys. Lett. B 91, 99–102 (1980).
- [127] V. F. Mukhanov, G. V. Chibisov, Quantum fluctuations and a nonsingular universe, JETP Lett. 33, 532–535 (1981)
 [Pisma Zh. Eksp. Teor. Fiz. 33, 549–553 (1981)].
- [128] A. D. Linde, A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems, Phys. Lett. B 108, 389–393 (1982).
- [129] A. Albrecht, P. J. Steinhardt, Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking, Phys. Rev. Lett. 48, 1220–1223 (1982).