

VALUATION OF EQ-5D-3L HEALTH STATES IN SLOVENIA: VAS BASED AND TTO BASED VALUE SETS

VREDNOTENJE ZDRAVSTVENIH STANJ EQ-5D-3L V SLOVENIJI: VREDNOSTI ZDRAVSTVENIH STANJ, PRIDOBLENE Z METODAMA VAS IN TTO

Valentina PREVOLNIK RUPEL^{1*}, Andrej SRAKAR¹, Kim RAND²

¹Institute for Economic Research, Kardeljeva ploščad 17, 1000 Ljubljana, Slovenia

²Health Services Research Unit, Akershus University Hospital, Lørenskog, Norway

Received: May 21, 2019

Accepted: Sep 26, 2019

Original scientific article

ABSTRACT

Introduction: The two primary objectives of this paper were (a) to develop first logically consistent TTO based EQ-5D-3L value sets for Slovenia and (b) to revisit earlier developed VAS based EQ-5D-3L value sets.

Keywords:

EQ-5D-3L, Slovenia, quality-adjusted life-years, social value set, utility

Methods: Between September 2005 and April 2006, face-to-face interviews with 225 individuals in Slovenia were conducted. Protocols from the Measurement and Value of Health study were followed closely. Each respondent valued 15 health states out of a total of 23. Model selection was informed by the criteria monotonicity/logical consistency. Predictive accuracy was assessed in terms of mean square difference between out-of-sample predictions and corresponding observed means, as well as Lin's Concordance Correlation Coefficient.

Results: Modelling was based on 2,717 VAS and 2,831 TTO values elicited from 225 respondents. A 6-parameter constrained regression model with a supplementary power term was selected for VAS and TTO value sets, as it produces monotonic values, and proved superior in terms of out-of-sample predictive accuracy over the tested alternatives.

Conclusion: This is the first EQ-5D-3L TTO based value set in Slovenia and the second in Central and Eastern Europe (besides Poland). It is also the first monotonic and logically consistent VAS value set in Central and Eastern Europe. Comparisons with Polish and UK TTO values show considerable differences, mostly due to mobility with having a substantially greater weight in Slovenia. The UK value set generally produces lower values and the Polish value set higher values for mild states.

IZVLEČEK

Uvod: Dva osnovna cilja raziskave sta (a) prikazati prvi logično konsistentni vrednostni set EQ-5D-3L za Slovenijo, ki temelji na metodi časovne izmenjave, (b) izboljšati prejšnji vrednostni set EQ-5D-3L za Slovenijo, ki temelji na vrednostni lestvici (VAS-metodi).

Ključne besede:

EQ-5D-3L, Slovenija, kakovostno prilagojena leta življenja, vrednostni set VAS, koristnost

Metode: Od septembra 2005 do aprila 2006 je bilo opravljenih 225 osebnih intervjujev s posamezniki iz 40 slovenskih občin. Študija je natančno sledila protokolu študije MVH o merjenju in vrednotenju zdravja, ki je bila izvedena v Združenem kraljestvu. Vsak anketiranec je ocenil 15 od skupno 23 zdravstvenih stanj. Izbira modela za izračun vrednosti zdravstvenih stanj je temeljila na dveh osnovnih merilih: monotonosti in logični doslednosti vrednosti. Napovedno moč smo vrednotili s povprečno kvadrirano razliko med napovedmi izven vzorca in pripadajočimi ocenjenimi povprečji ter s pomočjo Linovega konkordančnega korelacijskega koeficienta.

Rezultati: Izbrana modela temeljita na vrednostih zdravstvenih stanj 2,717 VAS in 2,831 TTO, ki smo jih pridobili v 225 osebnih intervjujih. Za oceno vrednosti VAS in TTO smo izbrali šestparametrski regresijski model z omejitvami in dodanim potenčnim faktorjem, saj se je izkazalo, da so ocenjene vrednosti na temelju tega modela monotone in imajo boljšo napovedno moč ocen izven vzorca kot vsi drugi ocenjevani modeli.

Zaključek: V študiji smo prikazali prvi slovenski vrednostni set EQ-5D, ki temelji na metodi TTO, hkrati pa je to drugi set, izračunan v srednji in vzhodni Evropi (poleg Poljske). Gre tudi za prvi monotoni in logično dosledni vrednostni VAS-set tako v Sloveniji kot srednji in vzhodni Evropi. Primerjave z vrednostmi poljskega in britanskega TTO kažejo precejšnje razlike med vrednostmi posameznih zdravstvenih stanj, predvsem zaradi dimenzije pokretnosti, ki ima bistveno večjo težo v Sloveniji. Vrednosti TTO v Združenem Kraljestvu so na splošno nižje za manj težavna zdravstvena stanja, poljske vrednosti zdravstvenih stanj pa so na splošno višje.

*Corresponding author: Tel. + 386 1 478 6870; E-mail: katkarupel@yahoo.com; katka.rupel@gmail.com

1 INTRODUCTION

Slovenia passed the regulation that required economic evaluation to inform drug and health technology reimbursement decision-making in the 1990s. Health technologies are assessed by various bodies (1, 2). The latest evaluation guidelines by the Health Insurance Institute of Slovenia recommend that the benefits of the treatment are expressed as quality-adjusted life years (QALYs).

QALY is a measure that encapsulates a treatment's impact on a patient's life length and also on their health-related quality of life (HRQOL), which is recognized as a key indicator of treatment outcomes (3). The QALY requires data that expresses health-related quality of life (HRQOL) in the form of a single value, sometimes known as a health state utility value, which is scored on a scale that assigns a value of 1 to a state equivalent to full health and 0 to a state equivalent to death (4). Weinstein and Stason (1977) connected QALYs with utilities, specifically expected utility, rather than the "weights" of the earlier literature; and this connection has remained although, not everybody agrees with the concession of the term "quality" to refer only to expected utility-based measures (5). Anyhow, in health economics, utilities (values) are typically combined with survival estimates and aggregated across individuals to generate quality-adjusted life years (QALYs) for use in cost-utility analyses of healthcare interventions (6).

There are many methods available regarding how the health states can be valued and grouped into two broad categories of measures: direct and indirect methods of measurement. The direct valuation methods include standard gamble (SG), time trade-off (TTO), DCE (discrete choice experiment), rating scales, equivalence technique, ratio scaling and person trade-off. The SG approach is the classic method of measuring preferences in economics under conditions of uncertainty, and is based on von Neumann Morgenstern utility theory (7). The theoretical underpinnings of all other methods are less clear. TTO valuation methodology does not conform to utility-under-uncertainty requirements under expected utility theory, but is still a dominant method in the valuation sets across countries (8). Regarding VAS values, there are a lot of criticisms and opposing views on their suitability for use in cost utility analysis. Mostly, criticisms consist of VAS values not being choice based and their lack of theoretical foundation (5). Due to these issues, most health economists would recommend a choice-based value set, derived from TTO or DCE data, especially for economic studies where cost-utility analysis is anticipated. If a choice-based value set is not available for the country/region, a choice-based value set can be selected from a country/region that most closely approximates the country where the study is being conducted. Alternatively, a VAS-based value set can be

used if that is available for the country/region (9). Due to these issues, most health economists would recommend a choice-based value set, derived from TTO or discrete choice experiment (DCE) data to be used in studies that estimate the value of health states of any population. If a choice-based value set is not available for the country/region, a choice-based value set can be selected from a country/region that most closely approximates the country where the study is being conducted. Alternatively, a VAS-based value set can be used if that is available for the country/region (9).

Utilities (values representing preferences) for healthcare priority setting are typically obtained indirectly by asking the general population (or patients) to fill in a questionnaire and attach value to hypothetical health state, later on converting the results to a value set for all health states, using population (or patient) values. There are at least two advantages that contributed to the popularity of the indirect methods: the pool of health states is already defined and so are their values (value set). When a patient defines his own health state in subsequent studies, a value can thus be attached to his/her health state from the value set.

Some of the established questionnaires are the Health Utility Index, the Short Form 6D, 15D instrument, Assessment of Quality of Life (AQOL) and the EuroQol 5D (EQ-5D). The EQ-5D is a prominent example of preference-based measures developed by the EuroQol Group (9). It has been suggested that these are the most widely used preference-based measures in the world (10). To improve the instrument's sensitivity and to reduce ceiling effect, EuroQol Group developed a new version in 2009, called EQ-5D-5L. The new version kept its original 5 dimensions, but expanded the response options from 3 to 5 levels. As there are a lot of existing 3L value sets in many countries, for comparison reasons a non-parametric model was set up to transform any EQ-5D-3L values into EQ-5D-5L values. In this way, 5L values can also be used in cases when 5L preferences directly elicited from representative general population samples are not yet available (11).

The EQ-5D-3L descriptive system has been formally translated and validated into the Slovenian language in 1999/2000 (12). The two primary objectives of this paper were (a) to develop first logically consistent EQ-5D-3L TTO-based value sets for Slovenia and (b) to revisit earlier developed VAS-based EQ-5D-3L value sets for Slovenia (13). Some issues that went undetected with the previous VAS value set have been identified, and methodological advances seem to make it possible to improve on earlier modelling.

2 METHODS

2.1 Study Overview

The study was a multicentre, population-based study, using face-to-face interviews. The sample was prepared by the Statistical Office of Slovenia using the Central Population Register. In the sample, 1,000 individuals aged 18+ from 40 Slovenian municipalities were included. At the first level, 40 municipalities were randomly selected and later on 25 individuals were selected from each municipality. Each person carried a name, last name, address, house number, postcode, municipality, age and gender. The investigators started the interviews in September 2005 and finished in April 2006. Participant recruitment was conducted primarily through landline telephone numbers for each participant in the sample. 225 participants agreed to participate in the survey. Interviews were conducted by three interviewers, who underwent one-day training on the health state valuations, the purpose of the interviews and TTO procedures. To facilitate the training, the interview book prepared by Gudex (14) was translated into Slovenian language and used for training of the interviewers and in the pilot training. Each investigator conducted 5 test interviews.

2.2 EQ-5D

EQ-5D consists of a descriptive system and EQ visual analogue scale (EQ-VAS). The EQ-5D descriptive system consists of 5 dimensions: mobility (MO), self-care (SC), usual activities (UA), pain/discomfort (PD), and anxiety/depression (AD). Each dimension has 3 levels: no problems, some problems, and extreme problems (9). The respondent is asked to indicate his or her health state by ticking the box that marks the most appropriate level of problems in each dimension. A unique health state can be described by using a 5-digit vector formed according to the responses to the 5 questions. For example, no problems in MO and SC, some problems in UA and PD and extreme problems in AD can be referred to as "11223." Health states defined in this way may be converted into a single summary index by applying a formula that attaches values to each of the levels in each dimension. A total of 243 possible health states can be defined.

2.3 Health State Selection

In the valuation task, each investigator had 3 sets of health states, and investigators decided randomly which set to use with each respondent. The sets were named A, B and C. Each set contained 15 health states. Some health states were included in all three sets, but some were not. Health states in each set represent the complete scale of health states, from worst to best health states. Sets B and C were developed in 2000 (16). The number of all various directly valued health states in all three sets is 23 plus unconscious and dead. These states are 11211,

11111, 21111, 12111, 11112, 11121, 11122, 11113, 11131, 11133, 11312, 13311, 32211, 22222, 21232, 22323, 22233, 32223, 32313, 23232, 33321, 33323, 33333, unconscious and dead. Health states can also be divided into mild, moderate and severe states (17) in such a way that all the categories were represented in all three sets.

2.4 Interview Process

The questionnaire consists of four parts. In the first part, the respondent indicated his/her own health state on the day of the interview using an EQ-5D descriptive system. Furthermore, the respondent marks how good or bad his/her health state is on a visual analogue scale (VAS) from 0 to 100 (where 0 represents the worst health state imaginable and 100 represents the best health state imaginable).

The second part of the questionnaire is a valuation of the 15 selected health states. Once the respondent had familiarised himself/herself with the health states by reading them, he/she ranked the selected states from worst to best. After ranking he/she attached the value from 0 to 100 to all 15 health states.

The third part of the interview is the valuation of the same selected 15 health states using time trade off (TTO) method. The interviewers follow the adapted Measurement and Value of Health study (MVH) protocol (14). The MVH study was a large exercise, in which 3,395 respondents valued 13 different health states. Because of the limited budget, we included 23 health states altogether. Out of 23 health states, we made three different sets of 15 health states (sets A, B and C) as described in Chapter 2.3.

The objective of the TTO is to determine the length of lifetime the respondent would be willing to forego to live in a better health state (typically 'full health') and to avoid living in a bad health state. This is achieved by presenting respondents with a series of choice tasks, each involving two alternative hypothetical lives. The two lives are presented so that the respondent is forced to choose between a longer life in the health state of interest and a shorter life in better health (15).

The last part of the interview collects social demographic data: gender, age, education, work experience, smoking habits, experience with illness and postcode.

2.5 Preference Elicitation Techniques

In the TTO procedure, the interviewers used a TTO board and a set of health state cards. A TTO board was made of three layers of thick cardboard and incorporated a sliding scale from 0 to 10 years. Both sides of the board were used, one for states better than dead and the other for health states worse than dead. The respondent was taken through each of 15 health states to be valued,

one at a time, with the interviewer moving the scale as appropriate. The respondent needed to make a series of choices between two hypothetical lives: one involving x years of healthy life, followed by death (Life A) and the other involving t years in a worse health state (where $x < t$), followed by death (Life B). Time t was fixed, whereas time x was varied until the respondent reached their point of indifference. This iterative procedure continued until the respondent was unable to choose between the two lives. In our study, the respondent started with a choice between living Life A (health state 11111) for 10 years followed by death and living Life B (worse health state) for 10 years followed by death. Life A was chosen - the next choice was between Immediate Death ($x=0$ for Life A) and 10 years of Life B, followed by death. In the next choice, x was set at 5 years; in case Life A was chosen, x was decreased and the opposite until the point of indifference was found. The value of Life B was calculated according to how much healthy time the respondent was willing to forgo at this point of indifference - the utility value of the health state was at this point calculated as $x/10$. In case of states worse than dead, the respondent was again given two alternatives, but this time Life A was a combination of health state l for y number of years followed by full health for x number of years ($x+y=10$), followed by death. Life B was a certain outcome of immediate death. Time x was again varied until the respondent was indifferent between both alternatives. At this point, the utility value for health state l was calculated as $-x/(10-x)$. Respondents were allowed to trade time in months and weeks.

In VAS procedure, the respondents ranked the health states and, in the second phase, attached them a value from 0 to 100. VAS values were later rescaled using the mean observed values for state 11111 and death. Health state “Unconscious” was not used.

2.6 Statistical Analyses

Historically, values for EQ-5D-3L have been modelled by use of ordinary least squares (OLS) regression, using dummy variables representing the presence or absence of different levels of problems on each of the five dimensions. Built on this framework, different interaction terms have been added in different national valuation studies. More recently, the introduction of the EQ-5D-5L has resulted in a range of innovations in terms of modelling, including hybrid models combining TTO and DCE data (24), random intercept models, censored/interval regression to account for censoring at -1, and use of constrained, non-linear regression models (18, 22, 23).

For EQ-5D-3L, the standard, additive 10-parameter model, hereafter referred to as ADD10, has parameters representing levels 2 and “3” for each dimension. Let y represent the observed disutility of a health state, represent x_{dl} the dummy variable indicating the presence

of problems on dimension d at level l and β_{dl} the coefficient representing the estimated disutility of having problems on dimension d at level l (e.g. β_{MO3} representing the disutility of having moderate problems on mobility). The mathematical function of ADD10 is as follows:

$$y = \sum_l \sum_d \beta_{dl} x_{dl} + e$$

$$= \beta_{MO2} x_{MO2} + \beta_{SC2} x_{SC2} + \beta_{UA2} x_{UA2} + \beta_{PD2} x_{PD2} + \beta_{AD2} x_{AD2} + \beta_{MO3} x_{MO3} + \beta_{SC3} x_{SC3} + \beta_{UA3} x_{UA3} + \beta_{PD3} x_{PD3} + \beta_{AD3} x_{AD3} + e \quad (1)$$

An EQ-5D-3L variant of the constrained model approach used in the Chinese and Malaysian EQ-5D-5L valuation studies employs six primary parameters: one for each dimension, representing the disutility of having problems at level 3 ($\beta_{MO}, \beta_{SC}, \beta_{UA}, \beta_{PD}, \beta_{AD}$), which for level 2 are multiplied by parameters for level 2 (L_2). Thus, the disutility of having moderate problems on mobility is $\beta_{MO} \times L_2$. The mathematical function of this model, hereafter MULT6, is as follows (note that x_{dl} still represents the dummy variable representing the presence of problems on dimension d at level l):

$$y = \sum_l \sum_d \beta_{dl} x_{dl} L_l + e$$

$$= (\beta_{MO2} x_{MO2} + \beta_{SC2} x_{SC2} + \beta_{UA2} x_{UA2} + \beta_{PD2} x_{PD2} + \beta_{AD2} x_{AD2}) L_2 + \beta_{MO3} x_{MO3} + \beta_{SC3} x_{SC3} + \beta_{UA3} x_{UA3} + \beta_{PD3} x_{PD3} + \beta_{AD3} x_{AD3} + e \quad (2)$$

This constrained model assumes that the relative severity of level 2, “moderate problems”, is similar across dimensions. This assumption reduces the number of parameters to be fitted, and thereby provides more robust results than unconstrained models, particularly with smaller samples of data.

We tested the ADD10 and MULT6 models, with and without the inclusion of a constant term (intercept) representing any deviation from full health. The model variants including intercept are denoted with an “i”, e.g. ADD10i. We also tested an extension of MULT6 in which the full expression is exponentiated by a separately fitted parameter P :

$$y = ((\beta_{MO2} x_{MO2} + \beta_{SC2} x_{SC2} + \beta_{UA2} x_{UA2} + \beta_{PD2} x_{PD2} + \beta_{AD2} x_{AD2}) L_2 + \beta_{MO3} x_{MO3} + \beta_{SC3} x_{SC3} + \beta_{UA3} x_{UA3} + \beta_{PD3} x_{PD3} + \beta_{AD3} x_{AD3})^P + e \quad (3)$$

This model, hereafter MULT6P, was included under the assumption that respondents may display diminishing sensitivity to health problems when combined, so that the perceived disutility of problems on two separate dimensions at the same time may be smaller than the sum of the disutility of each problem in isolation.

Standard error estimation is non-trivial in regression models involving multiplication of two or more (presumably normally distributed) parameters. Consequently, standard errors for model parameters and modelled values were estimated for MULT6 and MULT6P using bootstrapping

(22). Briefly, 10,000 bootstrap samples were drawn (with replacement) at the level of individual study participants, each subsample of the same size as the observed data. The regression models were fitted to each bootstrap sample, and standard errors were calculated by taking the standard deviation for the resulting coefficients and the predicted health state values.

Given the limited number of valued health states, and the relatively small sample size used in this study, we were concerned that regular regression models might produce results that were highly susceptible to random error. We, therefore, tested the included model variants using penalised regression, including Lasso (20), Ridge regression (17-19), and Elastic net (21).

Model selection was informed by two primary criteria, being monotonicity/logical consistency. Modelled state values should reflect the hierarchical structure of the EQ-5D descriptive system, so that further problems on any dimension should always result in worse (lower) values. Monotonic models were compared in terms of out-of-sample predictive accuracy for observed means. This was compared using leave-out-by-state cross-validation (18, 22). Predictive accuracy was assessed in terms of mean square difference between out-of-sample predictions and corresponding observed means, as well as Lin's Concordance Correlation Coefficient.

The final Slovenian TTO model was compared visually to the Polish EQ-5D-3L value set (25) and the UK MVH value set (26), and the final VAS model was compared visually to the EU VAS value set (27).

All statistical analyses were performed in the R statistical package, version 3.3.2, in the RStudio environment, using ggplot for graphical output (28-30). Regression models were run in the xreg package (31).

3 RESULTS

In total, 225 respondents completed the interview, of which 126 (56%) were female. Distribution of the respondents according to social and demographic variables is shown in Table 1. The sample was well representative of the Slovenian population in terms of age, educational level and activity with students being slightly underrepresented and unemployed being slightly overrepresented. Regarding gender distribution, women were overrepresented in the sample. The majority of problems reported in the EQ-5D descriptive system were pain/discomfort, followed by problems with usual activities and mobility. A really small share of the sample had problems with self-care (9.3%). The mean health state recorded on the EQ-VAS was 72.15 (SD 20.2) and the mean estimated interview difficulty was 2.87 (1 is very easy and 5 is very difficult).

Table 1. Study sample characteristics compared with the Slovenian general population data 2005.

Group	Mean pre-test scores (SD)	Mean post-test scores (SD)
Age		
18-24	27 (12%)	190,239 (11.5%)
25-34	48 (21.3%)	300,793 (18.2%)
35-44	43 (19.1%)	304,490 (18.5%)
45-54	39 (17.3%)	310,757 (18.9%)
55-64	28 (12.5%)	229,580 (13.9%)
65+	40 (17.8%)	312,874 (19%)
Mean age (SD)	45.3 (17.4)	n/a
Gender		
Male	99 (44%)	981,465 (49%)
Female	126 (56%)	1,021,893 (51%)
Educational level		
Primary	53 (23.6%)	494 (28.8%)
Secondary	147 (65.3%)	952 (55.5%)
High	25 (11.1%)	267 (15.6%)
Work		
Employed	111 (49.3%)	813,100 (47.3%)
Retired	62 (27.5%)	529,622 (30.8%)
Housewife	8 (3.6%)	n/a
Student	20 (8.9%)	112,228 (6.5%)
Unemployed	18 (8%)	92,575 (5.4%)
Other	6 (2.7%)	n/a
EQ-5D dimension problems (%)		n/a
Mobility	68 (30.2%)	
Self-care	21 (9.3%)	
Usual activities	69 (30.7%)	
Pain/discomfort	101 (44.9%)	
Anxiety/Depression	64 (28.4%)	
EQ VAS own health (SD)	72.15 (20.2)	n/a

Cross-validation fit statistics can be found in Table 2. The fitted parameters of ADD10 were not monotonic. MULT6 and MULT6P with no intercept were monotonic for both VAS and TTO, while the version with an intercept was monotonic for TTO only. MULT6 displayed poor fit, both in direct estimation and in cross-validation. Ridge regression improved out-of-sample predictive accuracy for ADD10 and MULT6, but not for MULT6P. By comparison, MULT6P had substantially improved fit, outperforming all other tested variants in terms of out of sample predictive accuracy, both for VAS and TTO data. MULT6P with an intercept did not improve predictions for TTO, and did not converge for VAS. MULT6P was selected for generating VAS and TTO value sets. Coefficients and bootstrap-based SE estimates for the two models can be found in Table 3.

Table 2. Cross-Validation fit statistics.

TTO	ADD10	ADD10i	MULT6	MULT6i	MULT6P	MULT6iP
Monotonicity	-	-	✓	✓	✓	✓
R	0.920	0.941	0.934	0.930	0.966	0.966
CCC	0.894	0.938	0.893	0.929	0.966	0.966
MSE	0.046	0.022	0.048	0.024	0.012	0.012
MAE	0.181	0.114	0.182	0.126	0.087	0.087

VAS	ADD10	ADD10i	MULT6	MULT6i	MULT6P	MULT6iP
Monotonicity	-	-	✓	✓	✓	-
R	0.926	0.919	0.891	0.923	0.971	-
CCC	0.879	0.897	0.886	0.883	0.941	-
MSE	0.02	0.015	0.015	0.021	0.01	-
MAE	0.123	0.094	0.096	0.102	0.082	-

R - Pearson's correlation coefficient, CCC - concordance correlation coefficient, MSE - mean squared error, MAE - mean absolute error

Table 3. Coefficients and bootstrap-based SE estimates.

	TTO		VAS	
	Coefficient	SE	Coefficient	SE
MO	0.943	0.126	0.424	0.070
SC	0.243	0.052	0.105	0.029
UA	0.202	0.039	0.103	0.028
PD	0.448	0.049	0.180	0.012
AD	0.239	0.037	0.137	0.021
L2	0.125	0.043	0.176	0.025
P	0.551	0.044	0.423	0.020

4 DISCUSSION

The Slovenian VAS and TTO based value sets are presented in Annex 1 and 2. The first VAS value set for Slovenia was calculated back in year 2000, however, the values of the health state were not monotonic: some of the logically superior health states displayed lower values (12). In 2012, a new improved set was published (13), however, again due to some methodological issues discovered later, it cannot be recommended for Slovenia's priority setting. With the advanced methodology, for the first time in Slovenia it was possible to obtain a logically consistent and monotonic VAS based value set as well as a 3L TTO based value set, which is also the second 3L TTO based value set in Central and Eastern Europe.

Figure 1 displays the TTO value set compared to observed mean values, along with TTO-based values from a UK MVH study and the Polish TTO-based EQ-5D-3L valuation study. Figure 2 presents the Slovenian VAS value set, observed mean values, and the EU VAS value set.




Figure 1. Graphical comparison of Slovenian EQ-5D-3L TTO value set versus (a) UK TTO and (b) Polish TTO value sets.




Figure 2. Graphical comparison of Slovenian VAS value set versus EU VAS value set.

For the TTO value set, there are two main drivers of differences between the national value sets: first, mobility has a substantially greater weight in the Slovenian value set. Second, the UK value set produces lower values for mild states, while the Polish value set produces higher values. The VAS value set is more in line with the EU VAS value set, but generally produces somewhat higher values. Due to considerable differences between TTO value sets in Slovenia in comparison to the UK and Poland, its use should be recommended for Slovenian studies.

After testing various modelling approaches, the Slovenian TTO and VAS value sets were fitted using a 6-parameter constrained regression model with a supplementary power term, which produces monotonic values and was superior in terms of out-of-sample predictive accuracy over the tested alternatives.

The Slovenian TTO-based value set, being a choice-based method, is recommended for use in all studies, including economic analysis. Systematic pairwise comparison across all conditions and value sets in previous studies (32) revealed the greatest differences between the TTO and VAS-based value sets, as well as the varying sensitivity of the disease burden evaluations of chronic disease conditions to the choice of value sets. Therefore, using a VAS value set in the presence of newly developed TTO value set in Slovenia would unnecessarily produce incomparable results. However, in order to allow for comparisons with previous studies where VAS values were used due to the absence of a TTO-based value set, it is suggested to present VAS-based results in parallel. Another option is also the presentation of the results in parallel with the UK TTO value set, given that it has been the most used value set in the region (33).

Further analysis of the differences between the first two TTO based value sets in Central and Eastern Europe (CEE) is recommended - it has always been claimed that CEE countries display more similar values of health states, which differ from value sets in Western European countries, however, the first glance at both value sets does not confirm such speculations.

The main limitation of the study is the year of the data collection: the completion of the valuation study has been substantially delayed (from the data collection in 2005), as earlier modelling attempts produced non-monotonic values. Attempts at ameliorating these modelling issues through the application of exclusion criteria failed, indicating that the observed non-monotonicities were not reflective of a small subgroup of respondents displaying conflicting preferences. The improved fit of the chosen model, which included a power term below 1, indicates that respondents display substantially diminishing sensitivity to increasing health problems. Whether or not this diminishing sensitivity is unique to this study population may warrant further investigation.

Besides the modelling issues, sample size and the low number of health states valued were additional reasons why it was difficult to obtain the monotonic value set. Currently, the EuroQol Group Association recommends the sample size of 1,000 units to complete the valuation study with sufficient statistical power. Back in 2005, such recommendations were not in place and our data collection was limited in financial terms as well as timewise.

5 CONCLUSION

The article presents the first TTO-based EQ-5D value set for Slovenia. There have been two previous attempts to present an EQ-5D VAS based value set in Slovenia, once in 2000 (12) and once in 2012 (13), however, those value sets either lacked logical consistency or consistent modelling techniques. The use of a constrained ordinary least squares approach built upon experiences from EQ-5D-5L valuation studies in China, but extended to handle diminishing sensitivity to increasing health problems, allowed us to generate logically consistent value sets for VAS and TTO. The two value sets presented in this paper are recommended for use in EQ-5D-3L studies in Slovenia.

CONFLICT OF INTEREST

VPR and KR are members of the EuroQol Group, a not-for-profit organisation that develops and distributes instruments for measuring and valuing health.

FUNDING

The study received funding from the EuroQol Research Foundation.

ETHICAL APPROVAL

Not required as the data in the study is not personal, but values of hypothetical health states.

REFERENCES

- Health Insurance Institute of Slovenia. Rules of the Commission on enlisting drugs for public financing [Slovenian]. Ljubljana: Health Insurance Institute of Slovenia, 2014. Accessed March 26th, 2019 at: <http://www.pisrs.si/Pis.web/pregledPredpisa?id=PRAV11493>.
- Ministry of Health of Republic of Slovenia. Procedures on handling the applications for new health care programmes [Slovenian]. Ljubljana: Ministry of Health, 2015. Accessed April 20th, 2019 at: http://www.mz.gov.si/si/o_ministrstvu/zdravstveni_svet_in_ostala_posvetovalna_telesa/zdravstveni_svet/postopek_za_vloge/.
- Whitehead SJ, Ali S. Health outcomes in economic evaluation: the QALY and utilities. *Br Med Bull.* 2010;96:5-21. doi: 10.1093/bmb/ldq033.
- Torrance GW. Measurement of health state utilities for economic appraisal. *J Health Econ.* 1986;5:1-30.
- Parkin D, Devlin N. Is there a case for using Visual Analogue Scale valuations in Cost-Utility Analysis? Report No. 04/03. London: Department of Economics, City University London, 2004. Accessed March 20th, 2019 at: http://openaccess.city.ac.uk/1428/1/0403_parkin-devlin.pdf.
- Tolley K. What are health utilities? NPR09/1103. Accessed February 17th, 2019 at: http://www.bandolier.org.uk/painres/download/What%20is%202009/What_are_health_util.pdf.
- von Neumann J, Morgenstern O. *Theory of games and economic behavior*. Princeton: University Press, 1944.
- Buckingham K, Devlin N, Tabberer M. A theoretical framework for TTO valuations and a taxonomy of TTO approaches: results from a pilot study. Department of Economics Discussion Paper Series 04/07. London: City University London, Department of Economics, 2007.
- EQ-5D, an instrument to describe and value health. Accessed April 29th, 2019 at: <https://euroqol.org/euroqol/>.
- Brauer CA, Rosen AB, Greenberg D, Neumann PJ. Trends in the measurement of health utilities in published cost-utility analyses. *Value Health.* 2006;9:213-8. doi: 10.1111/j.1524-4733.2006.00116.x.
- van Hout B, Janssen MF, Feng YS, Kohlmann T, Busschbach J, Golicki D, et al. Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Health.* 2012;15:708-15. doi: 10.1016/j.jval.2012.02.008.
- Prevolnik Rupel V, Rebolj M. The Slovenian VAS tariff based on valuations of EQ-5D health states from the general population. In: Cabases JM, Gaminde I. 17th Plenary Meeting of the EuroQol Group, Discussion Papers. Pamplona: EuroQol Group, 2000.
- Prevolnik Rupel V, Ogorevc M. The EQ-5D health states value set for Slovenia. *Zdr Varst* 2012;51:128-40. doi: 10.2478/v10152-012-0015-y.
- Gudex C. *Time trade-off user manual: props and self-completion methods*. York: The University of York, Centre for Health Economics, 1994.
- Oppe M, Rand-Hendriksen K, Shah K, Ramos-Goñi JM, Luo N. EuroQol protocols for time trade-off valuation of health outcomes. *Pharmacoeconomics.* 2016;34:993-1004. doi: 10.1007/s40273-016-0404-1.
- Devlin NJ, Hansen P, Kind P, William AH. The health states preferences and logical inconsistencies of New Zealanders: a tale of two tariffs. In: Cabases JM, Gaminde I. Pamplona: 17th Plenary Meeting of the EuroQol Group, Discussion Papers, 2000.
- Kind P. Improving the MVH protocol: learning from the past. Work in progress.
- Rand-Hendriksen K., Ramos-Goñi JM, Augestad LA, Luo N. Less is more: cross-validation testing of simplified nonlinear regression model specifications for EQ-5D-5L health state values. *Value Health.* 2017;20:945-52. doi: 10.1016/j.jval.2017.03.013.
- Hoert AE, Kennard RW. Ridge regression: biased estimation for nonorthogonal problems. *Technometrics.* 1970;12:55-67. doi: 10.2307/1271436.
- Tibshirani R. Regression shrinkage and selection via the lasso. *J. R. Statist. Soc. B.* 1996;58:267-88.
- Zou H, Hastie T. Regularization and variable selection via the elastic net. *J. R. Statist. Soc. B.* 2005;67:301-20.
- Nan L, Liu G, Li M, Guan H, Jin X, Rand-Hendriksen K. Estimating an EQ-5D-5L value set for China. *Value Health.* 2017;20:662-9. doi: 10.1016/j.jval.2016.11.016.
- Shafie AA, Vasanthakumar A, Lim CJ, Luo N, Rand-Hendriksen K, Md Yusof FA. EQ-5D-5L valuation for the Malaysian population. *Pharmacoeconomics.* 2019;37:715-25. doi: 10.1007/s40273-018-0758-7.
- Ramos-Goñi JM, Pinto-Prades JL, Oppe M, Cabases JM, Serrano-Aguilar P, Rivero-Arias O. Valuation and modeling of EQ-5D-5L health states using a hybrid approach. *Med Care.* 2017;55:e51-8. doi: 10.1097/MLR.0000000000000283.
- Golicki D, Jakubczyk M, Niewada M, Wrona W, van Busschbach JJ. Valuation of EQ-5D health states in Poland: first TTO-based social value set in Central and Eastern Europe. *Value Health.* 2010;13:289-97. doi: 10.1111/j.1524-4733.2009.00596.x.
- Dolan P. Modeling valuations for EuroQol health states. *Med Care.* 1997;35:1095-108.
- Greiner W, Weijnen T, Nieuwenhuizen M, et al. A single European currency for EQ-5D health states. *Eur J Health Econ.* 2003;4:222-31. doi: 10.1007/s10198-003-0182-5.
- R Core Team. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing, 2016. Accessed February 17th, 2019 at: <https://www.R-project.org/>.
- RStudio Team. RStudio: integrated development for R. Accessed May 20th, 2019 at: www.rstudio.com.
- Wickham H. *Ggplot2 - Elegant Graphics for Data Analysis*. 2nd ed. New York: Springer-Verlag, 2016.
- Rand K. xreg: flexible multi-frame non-linear regression model. Accessed May 17th, 2019 at: <https://www.github.com/intelligentaccident/xreg>.
- Zrubka Z, Beretzky Z, Hermann Z, Brodsky V, Gulácsi L, Rencz F, et al. A comparison of European, Polish, Slovenian and British EQ-5D-3L value sets using a Hungarian sample of 18 chronic diseases. *Eur J Health Econ.* 2019;20(Suppl 1):119-32. doi: 10.1007/s10198-019-01069-8.
- Rencz F, Gulácsi L, Drummond M, Golicki D, Prevolnik Rupel V, Simon J, et al. EQ-5D in Central and Eastern Europe: 2000-2015. *Qual Life Res.* 2016;25:2693-710. doi: 10.1007/s10198-019-01069-8.

Annex 1. Slovenian EQ-5D-3L VAS value set.

State	Utility	State	Utility	State	Utility	State	Utility
11111	1	23131	0.351316	32222	0.243814	12313	0.435993
21111	0.6664	33131	0.135372	13222	0.51722	22313	0.371997
31111	0.304328	11231	0.496121	23222	0.440402	32313	0.149199
12111	0.81524	21231	0.423003	33222	0.19261	13313	0.362674
22111	0.63367	31231	0.181925	11322	0.54139	23313	0.307537
32111	0.291652	12231	0.476714	21322	0.45996	33313	0.105132
13111	0.61471	22231	0.406756	31322	0.20431	11123	0.528806
23111	0.51633	32231	0.171722	12322	0.519463	21123	0.44983
33111	0.236074	13231	0.396696	22322	0.442235	31123	0.198291
11211	0.817021	23231	0.337714	32322	0.19372	12123	0.507633
21211	0.634367	33231	0.126114	13322	0.431312	22123	0.432532
31211	0.291936	11331	0.414277	23322	0.367949	32123	0.187808
12211	0.753447	21331	0.353134	33322	0.146516	13123	0.421857
22211	0.605136	31331	0.136599	11132	0.489607	23123	0.359742
32211	0.279555	12331	0.398349	21132	0.417575	33123	0.141042
13211	0.587925	22331	0.339169	31132	0.17854	11223	0.508098
23211	0.496312	32331	0.12711	12132	0.470522	21223	0.432916
33211	0.22514	13331	0.330462	22132	0.401527	31223	0.188043
11311	0.618425	23331	0.2786	32132	0.168393	12223	0.488074
21311	0.519047	33331	0.084473	13132	0.391584	22223	0.416293
31311	0.237527	11112	0.793055	23132	0.333207	32223	0.177737
12311	0.590712	21112	0.624379	33132	0.123018	13223	0.406013
22311	0.498429	31112	0.287819	11232	0.470944	23223	0.345901
32311	0.226314	12112	0.73681	21232	0.401884	33223	0.131702
13311	0.485851	22112	0.596115	31232	0.168621	11323	0.423985
23311	0.414433	32112	0.275534	12232	0.452735	21323	0.361592
33311	0.17657	13112	0.579401	22232	0.38639	31323	0.142281
11121	0.767966	23112	0.489795	32232	0.158637	12323	0.407701
21121	0.612567	33112	0.221496	13232	0.376774	22323	0.347381
31121	0.282798	11212	0.737905	23232	0.320094	32323	0.132707
12121	0.718127	21212	0.596725	33232	0.113935	13323	0.338527
22121	0.58538	31212	0.275809	11332	0.393568	23323	0.285876
32121	0.270628	12212	0.69438	21332	0.334958	33323	0.089715
13121	0.569227	22212	0.570879	31332	0.124222	11133	0.384893
23121	0.481929	32212	0.263795	12332	0.378355	21133	0.327293
33121	0.217045	13212	0.555435	22332	0.321498	31133	0.118936
11221	0.719125	23212	0.471123	32332	0.114913	12133	0.369963
21221	0.585969	33212	0.210836	13332	0.313094	22133	0.314037
31221	0.2709	11312	0.582693	23332	0.262868	32133	0.109702
12221	0.678933	21312	0.49232	33332	0.073032	13133	0.305756
22221	0.560972	31312	0.222913	11113	0.568446	23133	0.256196
32221	0.258995	12312	0.557946	21113	0.481322	33133	0.068137
13221	0.545982	22312	0.473103	31113	0.216699	11233	0.370296
23221	0.463627	32312	0.211981	12113	0.544737	21233	0.314334
33221	0.206466	13312	0.46132	22113	0.462634	31233	0.109909
11321	0.572413	23312	0.393717	32113	0.205884	12233	0.355824
21321	0.484402	33312	0.163382	13113	0.451156	22233	0.301411
31321	0.218451	11122	0.705034	23113	0.385039	32233	0.100801
12321	0.548422	21122	0.577498	33113	0.157758	13233	0.29333
22321	0.465569	31122	0.266943	11213	0.545255	23233	0.244864
32321	0.207603	12122	0.667085	21213	0.463048	33233	0.059767
13321	0.454006	22122	0.553138	31213	0.206126	11333	0.307413
23321	0.387478	32122	0.255124	12213	0.523088	21333	0.257703
33321	0.159344	13122	0.538492	22213	0.445189	31333	0.069245
11131	0.516128	23122	0.457638	32213	0.195504	12333	0.294662
21131	0.439509	33122	0.202938	13213	0.434188	22333	0.24608
31131	0.192067	11222	0.66788	23213	0.370437	32333	0.060669
12131	0.49567	21222	0.55367	33213	0.148167	13333	0.238788
22131	0.422629	31222	0.255388	11313	0.453441	23333	0.194682
32131	0.181692	12222	0.634971	21313	0.386994	33333	0.021893
13131	0.412197	22222	0.530963	31313	0.15903		

Annex 2. Slovenian EQ-5D-3L TTO value set.

State	Utility	State	Utility	State	Utility	State	Utility
11111	1	23131	0.128439	32222	-0.03062	12313	0.34613
21111	0.661462	33131	-0.25294	13222	0.430732	22313	0.207237
31111	0.130169	11231	0.389626	23222	0.282156	32313	-0.18975
12111	0.853651	21231	0.245965	33222	-0.13124	13313	0.218732
22111	0.606623	31231	-0.1593	11322	0.46964	23313	0.091724
32111	0.097211	12231	0.349529	21322	0.31599	33313	-0.28289
13111	0.623976	22231	0.210278	31322	-0.10538	11123	0.49996
23111	0.444812	32231	-0.18735	12322	0.426358	21123	0.342034
33111	-0.01073	13231	0.221799	22322	0.278326	31123	-0.08574
11211	0.850867	23231	0.094535	32322	-0.13419	12123	0.455282
21211	0.60504	33231	-0.28058	13322	0.290467	22123	0.303556
31211	0.096229	11331	0.254042	23322	0.157111	32123	-0.11484
12211	0.770598	21331	0.124011	33322	-0.22977	13123	0.315952
22211	0.554403	31331	-0.25653	11132	0.385375	23123	0.180134
32211	0.063968	12331	0.218141	21132	0.242199	33123	-0.21131
13211	0.570517	22331	0.091182	31132	-0.16224	11223	0.453969
23211	0.401345	32331	-0.28333	12132	0.34543	21223	0.302415
33211	-0.04194	13331	0.1018	22132	0.20661	31223	-0.11571
11311	0.616823	23331	-0.01638	32132	-0.19025	12223	0.411364
21311	0.439082	33331	-0.37271	13132	0.2181	22223	0.265157
31311	-0.0148	11112	0.841454	23132	0.091144	32223	-0.14437
12311	0.565379	21112	0.59959	33132	-0.28336	13223	0.277172
22311	0.397094	31112	0.092834	11232	0.344249	23223	0.145055
32311	-0.04503	12112	0.763187	21232	0.205552	33223	-0.23949
13311	0.410581	22112	0.549314	31232	-0.19109	11323	0.310866
23311	0.264468	32112	0.060641	12232	0.305699	21323	0.175548
33311	-0.1449	13112	0.565321	22232	0.170885	31323	-0.21497
11121	0.779161	23112	0.397046	32232	-0.21871	12323	0.273356
21121	0.560205	33112	-0.04506	13232	0.182084	22323	0.141589
31121	0.067744	11212	0.76107	23232	0.058026	32323	-0.24229
12121	0.711564	21212	0.54785	33232	-0.31063	13323	0.152565
22121	0.512356	31212	0.059681	11332	0.213388	23323	0.030755
32121	0.03604	12212	0.69593	21332	0.086822	33323	-0.33326
13121	0.527634	22212	0.5007	31332	-0.28691	11133	0.237367
23121	0.365528	32212	0.028129	12332	0.17853	21133	0.108786
33121	-0.06823	13212	0.515767	22332	0.054748	31133	-0.26893
11221	0.709673	23212	0.355487	32332	-0.31335	12133	0.201903
21221	0.510957	33212	-0.07569	13332	0.065126	22133	0.076272
31221	0.035094	11312	0.558733	23332	-0.05061	32133	-0.29558
12221	0.65042	21312	0.391579	33332	-0.4016	13133	0.08679
22221	0.465742	31312	-0.04905	11113	0.592636	23133	-0.03037
32221	0.003995	12312	0.510968	21113	0.419503	33133	-0.38449
13221	0.48022	22312	0.351413	31113	-0.0288	11233	0.200849
23221	0.325111	32312	-0.07872	12113	0.542813	21233	0.075303
33221	-0.09848	13312	0.364335	22113	0.378293	31233	-0.29638
11321	0.521351	23312	0.223497	32113	-0.0588	12233	0.166296
21321	0.360218	33312	-0.17692	13113	0.391539	22233	0.043453
31321	-0.07217	11122	0.703183	23113	0.247659	32233	-0.32271
12321	0.475612	21122	0.506132	33113	-0.15798	13233	0.05376
22321	0.321143	31122	0.031824	11213	0.541361	23233	-0.06124
32321	-0.10148	12122	0.644578	21213	0.377075	33233	-0.41062
13321	0.333726	22122	0.461154	31213	-0.0597	11333	0.082479
23321	0.196118	32122	0.000785	12213	0.494567	21333	-0.0344
33321	-0.19857	13122	0.47556	22213	0.337423	31333	-0.38789
11131	0.432582	23122	0.321098	32213	-0.0892	12333	0.05049
21131	0.283774	33122	-0.10151	13213	0.350186	22333	-0.0643
31131	-0.13	11222	0.642901	23213	0.210866	32333	-0.41322
12131	0.390857	21222	0.459833	33213	-0.18688	13333	-0.05462
22131	0.247056	31222	-0.00014	11313	0.386101	23333	-0.16321
32131	-0.15845	12222	0.589543	21313	0.242843	33333	-0.498
13131	0.258901	22222	0.416978	31313	-0.16174		