

## Erratum to: Variables associated with physical health-related quality of life in Parkinson's disease patients presenting for deep brain stimulation

Alexandre Paim Diaz<sup>1,2,3</sup> · Fernando Cini Freitas<sup>2,3</sup> · Maria Emília de Oliveira Thais<sup>2</sup> ·  
Fernando Zanela da Silva Areas<sup>2</sup> · Marcelo Liborio Schwarzbold<sup>2,4</sup> ·  
Rodrigo Debona<sup>2</sup> · Jean Costa Nunes<sup>2,5</sup> · Ricardo Guarnieri<sup>2</sup> ·  
Daniel Martinez-Ramirez<sup>6</sup> · Rui Daniel Prediger<sup>7</sup> · Aparna Wagle Shukla<sup>6</sup> ·  
Marcelo Neves Linhares<sup>2,3</sup> · Roger Walz<sup>2,4</sup>

Published online: 25 August 2016  
© Springer-Verlag Italia 2016

### Erratum to: Neurol Sci DOI 10.1007/s10072-016-2681-z

The authors would like to correct the following errors in the publication of the original article:

- In Table 1, below “MDD”, the words “sim” and “nao” should be read as “yes” and “no”, respectively. The correct table is given below:

The online version of the original article can be found under doi:[10.1007/s10072-016-2681-z](https://doi.org/10.1007/s10072-016-2681-z).

✉ Alexandre Paim Diaz  
alexandrepaimdiaz@gmail.com

- <sup>1</sup> Department of Psychiatry, New York State Psychiatric Institute, Columbia University College of Physicians and Surgeons, 1051 Riverside Drive Unit 24, New York, NY 10032, USA
- <sup>2</sup> Centro de Ciências da Saúde (CCS), Centro de Neurociências Aplicadas (CeNAp), Hospital Universitário (HU), Universidade Federal de Santa Catarina (UFSC), Florianópolis, SC, Brazil
- <sup>3</sup> Unidade de Distúrbios do Movimento, Hospital Governador Celso Ramos (HGCR), Florianópolis, SC, Brazil
- <sup>4</sup> Departamento de Clínica Médica, CCS, HU, UFSC, Florianópolis, SC, Brazil
- <sup>5</sup> Laboratório de Neuropatologia, Serviço de Patologia, HU, UFSC, Florianópolis, Brazil
- <sup>6</sup> Department of Neurology, Center for Movement Disorders and Neurorestoration, University of Florida, 3450 Hull Road, Gainesville, FL 32607, USA
- <sup>7</sup> Departamento de Farmacologia, Centro de Ciências Biológicas (CCB), UFSC, Florianópolis, SC, Brazil

**Table 1** Sample characteristics of PD patients selected for DBS surgery

Variables	n (%) mean (±SD)
Sex	
Female	23 (39)
Male	36 (61)
Age (years)	58 (8)
Years of education, median (IQ range)	5 (4/11)
Married/unmarried	
Married	44 (75)
Unmarried	15 (25)
Disease duration (years)	13 (5)
UPDRS MDS III	27 (17)
Hoehn and Yahr <sup>a</sup>	2 (2/3)
MDD <sup>b</sup>	
Yes	11 (19)
No	48 (81)
HADS scores	12 (8)
Non-psychiatric diseases <sup>c</sup>	
Yes	25 (43)
No	33 (57)
PDQ 39 mobility domain	61 (26)
PDQ 39 activities of daily living domain	64 (29)
PDQ 39 bodily discomfort domain	56 (29)
Use of levodopa	57 (97)
Use of dopamine agonist	33 (56)
Use of antidepressant	42 (71)

*HADS* Hospital anxiety and depression scale, *PDQ* Parkinson's disease questionnaire

<sup>a</sup> Median and interquartile range

<sup>b</sup> MDD: major depressive disorder at moment of evaluation

<sup>c</sup> One missing case

- 
- In the section “Results *Bodily discomfort domain scores*”, in the last phrase of the first paragraph, the text “mobility domain scores” should be read as “bodily discomfort domain scores”. Thus, the right text is “The model explained 24 % of the variability of the bodily discomfort domain scores (Table 5)”.