Variational Approach to Quantum State Tomography based on Maximal Entropy Formalism

Rishabh Gupta,[†] Manas Sajjan,[†] Raphael D. Levine,^{‡,¶} and Sabre Kais*,§

†Department of Chemistry, Purdue University, West Lafayette, IN, USA

‡The Fritz Haber Center for Molecular Dynamics and Institute of Chemistry, The Hebrew

University of Jerusalem, Jerusalem 91904, Israel

¶Department of Chemistry and Biochemistry and Department of Molecular and Medical Pharmacology, David Geffen School of Medicine, University of California, Los Angeles, CA 90095, USA

§Department of Chemistry, Department of Physics and Astronomy, and Purdue Quantum Science and Engineering Institute, Purdue University, West Lafayette, IN, USA

E-mail: kais@purdue.edu

Abstract

Quantum state tomography is an integral part of quantum computation and offers the starting point for the validation of various quantum devices. One of the central tasks in the field of state tomography is to reconstruct with high fidelity, the quantum states of a quantum system. From an experiment on a real quantum device, one can obtain the mean measurement values of different operators. With such a data as input, in this report we employ the maximal entropy formalism to construct the least biased mixed quantum state that is consistent with the given set of expectation values. Even though in principle, the reported formalism is quite general and should work for

an arbitrary set of observables, in practice we shall demonstrate the efficacy of the algorithm on an informationally complete (IC) set of Hermitian operators. Such a set possesses the advantage of uniquely specifying a single quantum state from which the experimental measurements have been sampled and hence renders the rare opportunity to not only construct a least-biased quantum state but even replicate the exact state prepared experimentally within a preset tolerance. The primary workhorse of the algorithm is re-constructing an energy function which we designate as the effective Hamiltonian of the system, and parameterizing it with Lagrange multipliers, according to the formalism of maximal entropy. These parameters are thereafter optimized variationally so that the reconstructed quantum state of the system converges to the true quantum state within an error threshold. To this end, we employ a parameterized quantum circuit and a hybrid quantum-classical variational algorithm to obtain such a target state making our recipe easily implementable on a near-term quantum device.

1 Introduction

The method of uniquely characterizing the quantum mechanical state of a quantum system based on a series of measurements of an informationally complete (IC) set of Hermitian operators is called quantum state tomography (QST) [1,2,3,4,5] and forms an important basis for testing and validating quantum devices. However, the traditional approaches to QST are being exhausted to its limits [6] because of certain limitations that accompanies those approaches. Some of these limitations correspond to exponential scaling of the traditional QST techniques with system size, which in turn require exponential amounts of storage and processing power to carry out computations. Along with this, since we are in the era of noisy intermediate-scale quantum (NISQ) [7] devices, the fidelity of measurements is also a limiting factor for performing state tomography efficiently as noisy measurements can lead to low fidelity of the reconstructed quantum state [8]. Another challenging task within the domain of QST is to reconstruct high fidelity quantum states [9] that can be used as a

starting point while addressing problems in the field of condensed-matter physics and also in the validation of quantum technologies [10]. Several research approaches have already been proposed that attempts to address one or the other limitations and it has paved the way for further advancements in this field. Some of these tomographic techniques include maximum likelihood estimation (MLE) [11,12], Bayesian mean estimation (BME) [13,14], quantum overlap tomography [15], shadow tomography [16,17], neural network tomography [18,19,20,21], and others [22,23,24,25]. In our previous work we proposed a method of QST based on the formalism of maximal entropy from an incomplete set of measurements [26,27]. With that motivation, this research is an attempt to address the challenge of quantum state preparation in the field of QST based on a variational approach that can be easily implemented on a near-term quantum device.

The maximal entropy formalism [23,28,29] provides the most unbiased probability distribution, based on the maximization of the von Neumann entropy of the system, subject to the constraints of the problem [22,30,31,32]. As a natural consequence, when combined with the method of Lagrange multipliers it leads to an expression of density operator given by Eq. (1), that can serve as an optimal candidate for variational Gibbs sampling [9]. Inspired by this, the current work focuses on reconstructing the quantum state of a system, represented by the quantum Gibbs state and based on the formalism of maximal entropy, from mean measurement values of IC set of Hermitian operators. Sampling from a probability distribution corresponding to quantum Gibbs state plays an important role in a variety of diverse fields within and not limited to many-body physics [33,10], quantum simulations [34], quantum optimization [35], and quantum machine learning [36,37]. However, preparing Gibbs state of a given Hamiltonian at arbitrary low temperature is not an easy task [38] and various approaches have been proposed, both classical and quantum [39,40,41,42], to prepare Gibbs state under certain specified conditions. Some of these techniques include algorithms based on quantum rejection sampling [43], dynamics simulation [44,45], dimension reduction [46] but the overhead quantum resource cost of implementing these approaches is very high and

not suitable for execution on near-term quantum devices. In order to find applications of quantum algorithms on NISQ devices the underlying quantum circuit should be shallow with low circuit depth and low number of qubits. Variational quantum algorithm (VQA) [47] is one such class of hybrid quantum-classical algorithm that follows a heuristic approach based on the variational principle and has been quite popular in the recent years [48,49,50,51,52,53] owing to their implementation on NISQ devices with shallow quantum circuits.

To prepare a quantum Gibbs state on a NISQ device using VQAs, several methods have been proposed [54,55,56,57,58,59]. In this work, we have employed the approach by Wang et al. [38] wherein the loss function for preparing the Gibbs state on the quantum circuit involves the truncation of the Taylor series for the entropy and is shown to prepare Gibbs state for a given Hamiltonian with fidelity of over 99%. The physical Hamiltonian of the system is unknown and is in fact unnecessary in this protocol. One only has access to the expectation values of arbitrary set of Hermitian operators. In principle, using the formalism one can generate a least-biased quantum state consistent with such an arbitrary and even incomplete set of mean measurements, yet in this report we use an IC set for testing and validation with the hope of affording a near-exact re-construction of the unknown pure quantum state used for sampling. This is attained by constructing a Hermitian matrix H, parameterized by Lagrange multipliers. The latter serves as a proxy Hamiltonian for the construction of the Gibbs state that represents the tomographic reconstruction of the state of the quantum system. The methodology is elaborately discussed in Section 2. To validate the proposed approach, the formalism is implemented in IBM Qiskit [60] and the results corresponding to the fidelity and trace distance between the reconstructed quantum state and the true state are shown in Section 3.

2 Methodology

The reconstruction of an unknown quantum state requires the information of a complete set of observables that are obtained through experimental measurements of Hermitian operators usually defined as positive-operator-valued measures (POVMs). The formalism of maximal entropy provides a unique characterization of the quantum state subject to the expectation values of a given set of operators that serve as the constraints of the problem. It also ensures that the Von Neumann entropy of the proposed distribution is maximum under the given constraints. The maximal entropy formalism when combined with the method of Lagrange multipliers $\lambda_k \in \mathbb{C}^2$ [23,29], yields the following expression for the density operator of the unknown quantum state [22,32]:

$$\hat{\rho} = \frac{1}{Z(\lambda_1, \dots, \lambda_k)} \exp\{-\sum_k \lambda_k \hat{f}_k\}$$
(1)

where \hat{f}_k corresponds to the operators whose expectation values are known and $Z(\lambda_1, \dots, \lambda_k) = Tr(\exp\{-\sum_k \lambda_k \hat{f}_k\})$ insures normalization as $Tr(\hat{\rho}) = 1$. The formalism of maximal entropy in general outputs a mixed state that is parameterized by the Lagrange multipliers as shown in Eq. (1). In our approach since the target state is pure, these Lagrange multipliers are optimized such that the initial mixed state from the recipe gradually approaches idempotency during the training process and hence converges within an ϵ -neighborhood of the pure target state with ϵ being the error tolerance specified by the user. For example, consider the case a 2-qubit quantum system that can be uniquely described by the informationally complete (IC) set of Hermitian operators given by:

$$\{|1\rangle\langle 1|, |2\rangle\langle 2|, |3\rangle\langle 3|, |4\rangle\langle 4|, (|1\rangle\langle 2| \pm |2\rangle\langle 1|), (|1\rangle\langle 3| \pm |3\rangle\langle 1|), (|1\rangle\langle 4| \pm |4\rangle\langle 1|), (|2\rangle\langle 3| \pm |3\rangle\langle 2|), (|2\rangle\langle 4| \pm |4\rangle\langle 2|), (|3\rangle\langle 4| \pm |4\rangle\langle 3|)\}$$

$$(2)$$

The expectation values of the initial four of these operators correspond to the probabilities and the rest are the coherences of the 2-qubit quantum system [31]. The IC set of these operators can be obtained using the linear combinations of Pauli string operators (σ_x , σ_y , σ_z , and σ_i) [61,62,26]:

$$x_{11} = \langle |1\rangle \langle 1| \rangle = \frac{1}{4} (\sigma_z^2 \sigma_z^1 + \sigma_z^2 \sigma_i^1 + \sigma_i^2 \sigma_z^1 + \sigma_i^2 \sigma_i^1)$$

$$x_{12} = \langle |1\rangle \langle 2| + |2\rangle \langle 1| \rangle = \frac{1}{2} (\sigma_z^2 \sigma_x^1 + \sigma_i^2 \sigma_x^1)$$

$$x_{22} = \langle |2\rangle \langle 2| \rangle = \frac{1}{4} (\sigma_z^2 \sigma_z^1 - \sigma_z^2 \sigma_i^1 + \sigma_i^2 \sigma_z^1 - \sigma_i^2 \sigma_i^1)$$

and so on.

Analogous to the maximal entropy formalism, parameterized by a single parameter $\beta = 1/k_BT$ where k_B is the Boltzmann's constant and T is the temperature, the quantum Gibbs state for a given Hamiltonian H is defined as:

$$\hat{\rho} = \frac{\exp\{-\beta H\}}{tr(\exp\{-\beta H\})} \tag{3}$$

Constructing the Gibbs state of a given Hamiltonian on a parameterized quantum circuit requires minimization of the Helmholtz free energy described by the function:

$$F(\rho) = tr(\rho H) - \beta^{-1} S(\rho) \tag{4}$$

where $S(\rho) = -\text{tr}(\rho \ln \rho)$ corresponds to the von Neumann entropy of ρ . However, the most challenging part of constructing the loss function that minimizes the free energy of the Hamiltonian is estimating the entropy of the parameterized quantum state [63]. In this work, to address the problem we adopted the method proposed by Wang *et al* [38] wherein they used the Taylor series of entropy and truncate it at order K and therefore, the truncated free energy is set as the loss function of the variational quantum algorithm. This method is practical in its implementation on a near-term quantum device as essentially the

loss function involves estimating higher-order state overlaps, $tr(\rho^k)$, that corresponds to the truncated von Neumann entropy and can be carried out on a quantum device using swap tests [64,65,54,66,67].

The core idea of the current research work stems from the combination of Eq. (1) and (3) as in Eq. (1) we are interested in maximizing the entropy by optimizing the unknown Lagrange multipliers λ_k according to the constraints of the expectation values of the set of Hermitian operators and in Eq. (3) we want to optimize the parameters of the quantum circuit in order to minimize the free energy that yields the quantum Gibbs state for a particular Hamiltonian. In our methodology, the exponent term in Eq. (1) is a Hermitian matrix H that constitutes the Hamiltonian for which the Gibbs state given by Eq. (3) is constructed on a quantum circuit using a variational algorithm. Thus, the hybrid variational quantum algorithm that is employed basically involves two levels of optimization and is termed as inner and outer optimization levels as shown in Figure 1. For a fixed set of Lagrange multipliers λ_k , the Hamiltonian **H** is constructed and passed onto the inner optimization level where the circuit parameters are variationally optimized using the truncated free energy as the loss function to yield a quantum Gibbs state corresponding to **H**. The constructed Gibbs state is then sent to the outer optimization level where the expectation values of the set of Hermitian operators are computed using the generated Gibbs state and then the Lagrange multipliers λ_k are updated so as to minimize the mean square error between the generated and true expectation values of the POVMs. The updated Lagrange multipliers yield the new proxy Hamiltonian H that is again sent back to the inner optimization level and this process continues until convergence of the generated expectation values to the true values. This variational approach to QST based on maximal entropy is theoretically generalizable to any number of qubits.

The variational quantum circuit comprises of n-qubits that corresponds to the size of the quantum system and also an additional ancilla qubit. The circuit incorporates a series of parameterized single qubit rotational gates on every qubit and each qubit is entangled to the




Figure 1: The hybrid quantum-classical variational algorithm for quantum state tomography based on the formalism of maximal entropy.

next qubit using the controlled-NOT (CNOT) gates. This sequence of rotation and CNOT gates is repeated depending on the expressivity that is required to obtain high fidelity of the prepared quantum states. The scaling of the algorithm in terms of quantum resource allocation is strictly polynomial as to re-construct a generic n-qubit pure quantum state, we require n+1 qubits and $\mathcal{O}(Dn)$ quantum gates wherein D is the depth (number of repeating layers) of the circuit ansatz used.

Using the aforesaid procedure, in this work, we were able to obtain a high fidelity of 0.99 by setting D=2 for two qubits system and D=6 for the six qubits quantum system. The choice of the single qubit rotational gates depends on whether the reconstructed state needs to be real or complex. In case of real quantum states, parameterized R_y gates can be used else for generalized complex states one can choose R_x gates in the sequence.

3 Results and discussion

In this current work we propose a variational approach based on maximal entropy formalism to perform quantum state tomography on a near-term quantum device. This procedure outcomes a reconstructed quantum state that is prepared on a parameterized quantum circuit by using the expectation values of an IC set of Hermitian operators as input. The approach is tested and validated through numerical simulations conducted on IBM's Qiskit [60] using its prototype quantum simulators for quantum systems consisting of up to 6 qubits. There are various backends available in Qiskit and we used the noise-free *statevector_simulator* backend to corroborate the theory.

Different quantum circuits ranging from 2-6 qubits and consisting of one and two qubit




Figure 2: Sample 4-qubit and 6-qubit quantum circuits for which the quantum states are reconstructed using the proposed approach based on maximal entropy formalism.

quantum gates such as rotational, Hadamard, CNOT gates, etc. are used to prepare the sample states whose measurement statistics are reproduced using the reconstructed quantum state from the proposed maximal entropy based variational approach. Sample 4-qubit and 6-qubit quantum circuits are shown in Figure 2. As discussed in Section 2, at the end of full execution of inner optimization level a quantum Gibbs state is generated that is used to calculate the expectation values of the considered Hermitian operators. The mean square error (MSE) between the generated and the true expectation values serve as the loss function for updating the Lagrange multipliers in the outer optimization level. The MSE

loss is plotted as a function of the epochs in Figure 3. As can be seen in Figure 3, the MSE loss converges to zero faster for smaller systems and the convergence becomes more erratic as the size of the quantum system increases.

To perform quantum state tomography it is imperative that the reconstructed quantum




Figure 3: For the considered quantum systems with qubits ranging from 2-6, this plot shows the mean square error (MSE) loss as a function of the number of epochs between the true expectation values and the generated expectation values, of the IC set of Hermitian operators, obtained using the reconstructed quantum state upon each step of optimization.

state should be in close agreement with the true state. To demonstrate this a plot of fidelity between the reconstructed quantum state and the true state as a function of the number of epochs is shown in Figure 4 for the different quantum systems with varying number of qubits. The figure shows the convergence of the reconstructed state to the true state as the Lagrange multipliers are updated with each step and the fidelity approaches 1 near the end of the optimization cycle. Another measure to test the performance of our approach is the trace distance between the true state (σ) and the reconstructed quantum state (ρ) . Trace distance given by $T(\rho,\sigma) = \frac{1}{2}||\rho-\sigma||_1$ is a measure of the closeness between two states. Figure 5 shows the trace distance between the reconstructed quantum state, obtained from the converged set of λ parameters, and the true state and as can be seen the trace distance for all the quantum systems considered is below 0.05 that also validates the successful reconstruction of the quantum state.




Figure 4: As a measure of convergence between the true and the reconstructed state, the fidelity between the two states is plotted as a function of the number of epochs. The fidelity close to 1 at the end of the optimization cycle shows that the proposed method based on maximal entropy formalism is able to successfully reconstruct the quantum state using the experimental mean measurement values of IC set of operators.




Figure 5: Trace distance between the true state and the final reconstructed quantum state obtained using the converged Lagrange multipliers at the end of the optimization process.

4 Concluding remarks

In this research we have shown the successful reconstruction of the quantum state based on a variational approach to QST by utilizing the formalism of maximal entropy and can be easily implemented on a near-term quantum device. The reconstructed state can reproduce the measurement statistics of the IC set of Hermitian operators that are considered while formulating the cost function for updating the variational parameters of the Hamiltonian arising from maximal entropy. The fidelity between the converged state and the true state is over 0.99 for all the considered quantum systems and the trace distance is below 0.05 that depicts the validation of the proposed approach for QST. This proposed variational approach can be applicable to a variety of different directions where QST is essential. For example, we intend to further study this approach for analyzing, characterizing, and mitigating single and double qubit quantum gates' errors while running computations on prototype quantum devices. This method can also be used as an efficient approach for sampling Gibbs state and therefore, can serve as a starting point for preparing targeted quantum state for quantum simulations.

Acknowledgement

We acknowledge the financial support from the National Science Foundation under Award No. 1955907. We also like to acknowledge the partial financial support by the U.S. Department of Energy (Office of Basic Energy Sciences) under Award No. DE-SC0019215. We also acknowledge use of the IBM Q for this work. The views expressed here are those of the authors and do not reflect the official policy or position of IBM or the IBM Q team.

References

- (1) Nielsen, M. A.; Chuang, I. L. Quantum Computation and Quantum Information: 10th Anniversary Edition, 10th ed.; Cambridge University Press: USA, 2011.
- (2) Kais, S. Quantum Information and Computation for Chemistry; Wiley and Sons: Hoboken: NJ, 2014; Vol. 154.
- (3) Altepeter, J.; Jeffrey, E.; Kwiat, P. In *Photonic State Tomography*; Berman, P., Lin, C., Eds.; Advances In Atomic, Molecular, and Optical Physics; Academic Press, 2005; Vol. 52; pp 105 159.

- (4) James, D. F. V.; Kwiat, P. G.; Munro, W. J.; White, A. G. Measurement of qubits. Phys. Rev. A 2001, 64, 052312.
- (5) Banaszek, K.; Cramer, M.; Gross, D. Focus on quantum tomography. New J. Phys. **2013**, 15, 125020.
- (6) Song, C. et al. 10-Qubit Entanglement and Parallel Logic Operations with a Superconducting Circuit. *Phys. Rev. Lett.* **2017**, *119*, 180511.
- (7) Preskill, J. Quantum Computing in the NISQ era and beyond. Quantum 2018, 2, 79.
- (8) Chen, Y.; Farahzad, M.; Yoo, S.; Wei, T.-C. Detector tomography on IBM quantum computers and mitigation of an imperfect measurement. Phys. Rev. A 2019, 100, 052315.
- (9) Aharonov, D.; Arad, I.; Vidick, T. Guest column: the quantum PCP conjecture. *Acm sigact news* **2013**, *44*, 47–79.
- (10) Anshu, A.; Arunachalam, S.; Kuwahara, T.; Soleimanifar, M. Sample-efficient learning of interacting quantum systems. *Nature Physics* **2021**, *17*, 931–935.
- (11) Hradil, Z. Quantum-state estimation. *Phys. Rev. A* **1997**, *55*, R1561–R1564.
- (12) Baumgratz, T.; Nüßeler, A.; Cramer, M.; Plenio, M. B. A scalable maximum likelihood method for quantum state tomography. *New Journal of Physics* **2013**, *15*, 125004.
- (13) Lukens, J. M.; Law, K. J.; Jasra, A.; Lougovski, P. A practical and efficient approach for Bayesian quantum state estimation. *New Journal of Physics* **2020**, *22*, 063038.
- (14) Lukens, J. M.; Law, K. J.; Bennink, R. S. A Bayesian analysis of classical shadows. arXiv preprint arXiv:2012.08997 2020,
- (15) Cotler, J.; Wilczek, F. Quantum Overlapping Tomography. *Phys. Rev. Lett.* **2020**, *124*, 100401.

- (16) Aaronson, S. Shadow Tomography of Quantum States. Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing. New York, NY, USA, 2018; p 325–338.
- (17) Huang, H.-Y.; Kueng, R.; Preskill, J. Predicting many properties of a quantum system from very few measurements. *Nature Physics* **2020**, *16*, 1050–1057.
- (18) Torlai, G.; Mazzola, G.; Carrasquilla, J.; Troyer, M.; Melko, R.; Carleo, G. Neuralnetwork quantum state tomography. *Nature Physics* **2018**, *14*, 447–450.
- (19) Carrasquilla, J.; Torlai, G.; Melko, R. G.; Aolita, L. Reconstructing quantum states with generative models. *Nature Machine Intelligence* **2019**, *1*, 155–161.
- (20) Palmieri, A. M.; Kovlakov, E.; Bianchi, F.; Yudin, D.; Straupe, S.; Biamonte, J. D.; Kulik, S. Experimental neural network enhanced quantum tomography. npj Quantum Information 2020, 6, 20.
- (21) Xin, T.; Lu, S.; Cao, N.; Anikeeva, G.; Lu, D.; Li, J.; Long, G.; Zeng, B. Local-measurement-based quantum state tomography via neural networks. *npj Quantum Information* **2019**, *5*, 1–8.
- (22) Wichmann, E. H. Density Matrices Arising from Incomplete Measurements. *J. Math. Phys.* **1963**, *4*, 884–896.
- (23) Jaynes, E. T. Information Theory and Statistical Mechanics. II. *Phys. Rev.* **1957**, *108*, 171–190.
- (24) Katz, A. Principles of Statistical Mechanics: The Information Theory Approach; W. H. Freeman, 1967.
- (25) Sajjan, M.; Li, J.; Selvarajan, R.; Sureshbabu, S. H.; Kale, S. S.; Gupta, R.; Kais, S. Quantum computing enhanced machine learning for physico-chemical applications. arXiv preprint arXiv:2111.00851 2021,

- (26) Gupta, R.; Xia, R.; Levine, R. D.; Kais, S. Maximal Entropy Approach for Quantum State Tomography. *PRX Quantum* **2021**, 2, 010318.
- (27) Gupta, R.; Levine, R. D.; Kais, S. Convergence of a Reconstructed Density Matrix to a Pure State Using the Maximal Entropy Approach. The Journal of Physical Chemistry A 2021, 125, 7588–7595.
- (28) Levine, R. D.; Tribus, M. *Maximum Entropy Formalism*; Cambridge, Mass. : MIT Press: USA, 1979.
- (29) Agmon, N.; Alhassid, Y.; Levine, R. D. An Algorithm for Finding the Distribution of Maximal Entropy. J. Comput. Phys. 1979, 30, 250–258.
- (30) Bužek, V.; Drobný, G.; Adam, G.; Derka, R.; Knight, P. L. Reconstruction of quantum states of spin systems via the Jaynes principle of maximum entropy. *Journal of Modern Optics* 1997, 44, 2607–2627.
- (31) Alhassid, Y.; Levine, R. D. Connection between the maximal entropy and the scattering theoretic analyses of collision processes. *Phys. Rev. A* **1978**, *18*, 89–116.
- (32) Dagan, S.; Dothan, Y. Evaluation of an incompletely measured spin density matrix.

 Phys. Rev. D 1982, 26, 248–260.
- (33) Lewin, M.; Nam, P. T.; Rougerie, N. Classical field theory limit of many-body quantum Gibbs states in 2D and 3D. *Inventiones mathematicae* **2021**, *224*, 315–444.
- (34) Childs, A. M.; Maslov, D.; Nam, Y.; Ross, N. J.; Su, Y. Toward the first quantum simulation with quantum speedup. *Proceedings of the National Academy of Sciences* **2018**, *115*, 9456–9461.
- (35) Somma, R. D.; Boixo, S.; Barnum, H.; Knill, E. Quantum simulations of classical annealing processes. *Physical review letters* **2008**, *101*, 130504.

- (36) Kieferová, M.; Wiebe, N. Tomography and generative training with quantum Boltzmann machines. *Physical Review A* **2017**, *96*, 062327.
- (37) Biamonte, J.; Wittek, P.; Pancotti, N.; Rebentrost, P.; Wiebe, N.; Lloyd, S. Quantum machine learning. *Nature* **2017**, *549*, 195–202.
- (38) Wang, Y.; Li, G.; Wang, X. Variational quantum Gibbs state preparation with a truncated Taylor series. *Physical Review Applied* **2021**, *16*, 054035.
- (39) Poulin, D.; Wocjan, P. Sampling from the thermal quantum Gibbs state and evaluating partition functions with a quantum computer. *Physical review letters* **2009**, *103*, 220502.
- (40) Temme, K.; Osborne, T. J.; Vollbrecht, K. G.; Poulin, D.; Verstraete, F. Quantum metropolis sampling. *Nature* **2011**, *471*, 87–90.
- (41) Kastoryano, M. J.; Brandao, F. G. Quantum Gibbs samplers: The commuting case.

 Communications in Mathematical Physics 2016, 344, 915–957.
- (42) Brandão, F. G.; Kastoryano, M. J. Finite correlation length implies efficient preparation of quantum thermal states. *Communications in Mathematical Physics* **2019**, *365*, 1–16.
- (43) Yung, M.-H.; Aspuru-Guzik, A. A quantum-quantum Metropolis algorithm. *Proceedings of the National Academy of Sciences* **2012**, *109*, 754–759.
- (44) Kaplan, D. B.; Klco, N.; Roggero, A. Ground states via spectral combing on a quantum computer. arXiv preprint arXiv:1709.08250 2017,
- (45) Riera, A.; Gogolin, C.; Eisert, J. Thermalization in nature and on a quantum computer. Physical review letters **2012**, 108, 080402.
- (46) Bilgin, E.; Boixo, S. Preparing thermal states of quantum systems by dimension reduction. *Physical review letters* **2010**, *105*, 170405.

- (47) McClean, J. R.; Romero, J.; Babbush, R.; Aspuru-Guzik, A. The theory of variational hybrid quantum-classical algorithms. *New Journal of Physics* **2016**, *18*, 023023.
- (48) Bravo-Prieto, C.; LaRose, R.; Cerezo, M.; Subasi, Y.; Cincio, L.; Coles, P. J. Variational quantum linear solver. arXiv preprint arXiv:1909.05820 2019,
- (49) Huang, H.-Y.; Bharti, K.; Rebentrost, P. Near-term quantum algorithms for linear systems of equations. arXiv preprint arXiv:1909.07344 2019,
- (50) LaRose, R.; Tikku, A.; O'Neel-Judy, É.; Cincio, L.; Coles, P. J. Variational quantum state diagonalization. *npj Quantum Information* **2019**, *5*, 1–10.
- (51) Cerezo, M.; Sharma, K.; Arrasmith, A.; Coles, P. J. Variational quantum state eigensolver. arXiv preprint arXiv:2004.01372 2020,
- (52) Peruzzo, A.; McClean, J.; Shadbolt, P.; Yung, M.-H.; Zhou, X.-Q.; Love, P. J.; Aspuru-Guzik, A.; O'brien, J. L. A variational eigenvalue solver on a photonic quantum processor. *Nature communications* **2014**, *5*, 1–7.
- (53) Chen, R.; Song, Z.; Zhao, X.; Wang, X. Variational quantum algorithms for trace distance and fidelity estimation. *Quantum Science and Technology* **2021**, *7*, 015019.
- (54) Islam, R.; Ma, R.; Preiss, P. M.; Eric Tai, M.; Lukin, A.; Rispoli, M.; Greiner, M. Measuring entanglement entropy in a quantum many-body system. *Nature* 2015, 528, 77–83.
- (55) Verdon, G.; Marks, J.; Nanda, S.; Leichenauer, S.; Hidary, J. Quantum Hamiltonian-based models and the variational quantum thermalizer algorithm. arXiv preprint arXiv:1910.02071 2019,
- (56) Wu, J.; Hsieh, T. H. Variational thermal quantum simulation via thermofield double states. *Physical review letters* **2019**, *123*, 220502.

- (57) McArdle, S.; Jones, T.; Endo, S.; Li, Y.; Benjamin, S. C.; Yuan, X. Variational ansatz-based quantum simulation of imaginary time evolution. *npj Quantum Information* **2019**, *5*, 1–6.
- (58) Yuan, X.; Endo, S.; Zhao, Q.; Li, Y.; Benjamin, S. C. Theory of variational quantum simulation. *Quantum* **2019**, *3*, 191.
- (59) Chowdhury, A. N.; Low, G. H.; Wiebe, N. A variational quantum algorithm for preparing quantum Gibbs states. arXiv preprint arXiv:2002.00055 2020,
- (60) Abraham, H. et al. Qiskit: An Open-source Framework for Quantum Computing. 2019.
- (61) Kandala, A.; Mezzacapo, A.; Temme, K.; Takita, M.; Brink, M.; Chow, J. M.; Gambetta, J. M. Hardware-efficient variational quantum eigensolver for small molecules and quantum magnets. *Nature* 2017, 549, 242–246.
- (62) Bian, T.; Murphy, D.; Xia, R.; Daskin, A.; Kais, S. Quantum computing methods for electronic states of the water molecule. *Molecular Physics* **2019**, *117*, 2069–2082.
- (63) Gheorghiu, A.; Hoban, M. J. Estimating the entropy of shallow circuit outputs is hard. arXiv preprint arXiv:2002.12814 2020,
- (64) Buhrman, H.; Cleve, R.; Watrous, J.; De Wolf, R. Quantum fingerprinting. *Physical Review Letters* **2001**, *87*, 167902.
- (65) Gottesman, D.; Chuang, I. Quantum digital signatures. arXiv preprint quant-ph/0105032 **2001**,
- (66) Patel, R. B.; Ho, J.; Ferreyrol, F.; Ralph, T. C.; Pryde, G. J. A quantum Fredkin gate. Science advances 2016, 2, e1501531.
- (67) Linke, N. M.; Johri, S.; Figgatt, C.; Landsman, K. A.; Matsuura, A. Y.; Monroe, C. Measuring the Rényi entropy of a two-site Fermi-Hubbard model on a trapped ion quantum computer. *Physical Review A* 2018, 98, 052334.