

Vascular Function and Inflammation in Rheumatoid Arthritis: the Role of Physical Activity

George S. Metsios^{1,2,*}, Antonios Stavropoulos-Kalinoglou^{1,2}, Aamer Sandoo², Jet J.C.S. Veldhuijzen van Zanten², Tracey E. Toms², Holly John² and George D. Kitas²

¹*School of Sport Performing Arts and Leisure, University of Wolverhampton, Walsall, West Midlands, United Kingdom*

²*Department of Rheumatology, Dudley Group of Hospitals NHS, Foundation Trust, Russell's Hall Hospital, Dudley, West Midlands, United Kingdom*

Abstract: Inflammation disturbs biochemical pathways involved in homeostasis of the endothelium. Research has established clear links between inflammatory mediators, particularly C-reactive protein and tumour necrosis factor alpha, endothelial dysfunction, and atherosclerosis. Endothelial dysfunction and atherosclerosis may be subclinical at early stages, and thus the ability to detect them with non-invasive techniques is crucially important, particularly in populations at increased risk for cardiovascular disease, such as those with rheumatoid arthritis. This may allow the identification of interventions that may reverse these processes early on. One of the best non-pharmacological interventions that may achieve this is physical activity. This review explores the associations between inflammation, endothelial dysfunction, and atherosclerosis and discusses the role of exercise in blocking specific pathways in the inflammation, endothelial dysfunction - atherosclerosis network.

Keywords: Inflammation, Atherosclerosis, Endothelial function, Cytokines, Vascular function, Exercise, Physical activity, Cardiovascular disease.

INTRODUCTION

The endothelium is now regarded as a dynamic organ lining the entire vascular system. It is responsible for many atheroprotective functions *via* its regulation of the immune response, fibrinolysis, coagulation, as well as multiple other functions related to vascular growth, vasoprotection, and vasoregulation. Disruption of normal endothelial function impedes its protective effects on vascular homeostasis, an early event characterizing the development of cardiovascular disease (CVD) [1-3]. Most importantly, endothelial dysfunction and the early stages of CVD may occur in the absence of any clinical symptoms. Hence, finding methods for assessing vascular dysfunction during the early stages of the disease is important, particularly in patient groups at high CVD risk [4].

Rheumatoid arthritis (RA) is the most common chronic inflammatory arthritis characterized by symptoms and signs of joint and systemic inflammation, joint destruction, body composition alterations, and physical disability. It also associates with high CVD morbidity and mortality, which are not fully explained by the presence of traditional CVD risk factors [5-8]. In RA, increased local (synovial) and systemic expression of specific inflammatory mediators may have detrimental effects on the endothelium by enhancing biochemical processes leading to its dysfunction and eventually

the development of atherosclerosis [2, 9]. Indeed, the inflammatory pathways involved in the pathogenesis of RA and atherosclerosis share many characteristics [10]. Inflammation in RA is often evaluated by measuring the acute phase reactant C-reactive protein (CRP) and/or the erythrocyte sedimentation rate (ESR), which, in combination with other clinical markers (e.g. disease activity or X-ray scores), are used to assess the disease state and response to treatment [11]. The acute phase response is attributed to increased levels of pro-inflammatory cytokines, such as tumour necrosis factor alpha (TNF-alpha) and interleukin-6 (IL-6) [10], which are individually or synergistically involved in many other disease processes, including joint destruction [12] altered body composition [13], and changes in vascular homeostasis [14, 15].

INFLAMMATION AND ENDOTHELIAL DYSFUNCTION

The endothelium controls vascular function by releasing vasoactive factors such as nitric oxide (NO), prostacyclin (PGI₂), and endothelin-1 (ET-1). Endothelial dysfunction occurs when the balance between these vasoactive factors is disturbed. The down-regulation of endothelial NO synthase (eNOS) plays an important role in this process [16]. Altered endothelial homeostasis due to the disturbed NO production is characterized by diminished endothelium-dependent dilation and increased expression of adhesion molecules, as well as changes in endothelial cell phenotype and increased endothelial permeability [17]. Such changes in the structure and function of the endothelium are thought to be important as

*Address correspondence to this author at the School of Sport Performing Arts and Leisure, University of Wolverhampton, Wolverhampton, West Midlands, United Kingdom; Tel: +44-0121-4561302; E-mail: G.Metsios@wlv.ac.uk

initial stages in the pathogenesis of atherosclerosis and CVD [18].

Several studies have established a strong link between inflammation and endothelial dysfunction. *In vitro*, CRP modulates eNOS activity of human endothelial cells [19, 20] *via* uncoupling eNOS and subsequently increasing superoxide anions (oxidative stress) from NADPH oxidase [21]. As a result, phosphorylation of eNOS is decreased, and this subsequently reduces bioavailability of NO [22]. Apart from NO, CRP upregulates the expression of ET-1 which is a vasoconstrictor [23]. Studies also reveal a direct effect of TNF-alpha on downregulating the expression of eNOS and diminishing endothelial NO [15]. TNF-alpha downregulates at the same time enzymatic activities of eNOS and argininosuccinate synthetase (regulator of citrulline/NO cycle) [24] and induces an increase in mRNA degradation of both eNOS and neuronal NO synthase promoting vasoconstriction [25]. In addition, by activating NADPH oxidase, TNF-alpha stimulates the generation of reactive oxygen species within the endothelial environment further impairing NO-mediated vasodilation [26]. Apart from inhibition of NO production, TNF-alpha may also be responsible for the reduction in the bioavailability of NO [27]. This is perhaps the main reason why TNF-alpha inhibition is directly linked to improved endothelial function in both RA [28] and non-RA populations [29].

Endothelial function and morphology are reported to be significantly worse in RA compared to healthy controls and this is attributed in large part to the high-grade inflammatory state of RA [30, 31]; in fact, acute inflammation (e.g. in response to immunization) and low-grade chronic systemic inflammation (detected using high sensitivity CRP assays) in healthy individuals has been directly linked with arterial stiffness and vascular dysfunction [32, 33]. However, despite this biologically plausible link between inflammation and vascular dysfunction, there are still some controversies and inconsistent results in RA [34, 35], where some studies report a lack of association between systemic inflammatory load and endothelial function [36-38]. This may be attributed to different methodological approaches (e.g. using small, diverse patient populations cross-sectionally, or even smaller populations longitudinally but with short time courses), or lack of sufficient statistical power and no correction for multiple potential confounders (e.g. age and many CVD risk factors). The link between inflammation and vascular function appears to be more consistently demonstrated in homogeneous RA samples, such as young patients, patients with new onset RA and without established CVD [31, 39]. Vascular function may progressively deteriorate during the course of RA [40]. Hence, RA patients with long disease duration may already have significantly impaired endothelial function at the time of assessment, irrespective of disease "current" activity. Given the fluctuations in inflammation characteristic of RA, it might be that longstanding, though intermittent, inflammatory insult to the endothelium is of more importance to vascular function in RA than the current level of inflammation [41]. Studies that took into account long-term inflammatory burden have been able to demonstrate associations between this and vascular function [36, 39].

INFLAMMATION AND ATHEROSCLEROSIS

Endothelial cell dysfunction, evident as NO deficiency, appears to be amongst the earliest processes involved in atherosclerosis. Reduction of NO bioavailability disrupts the balance of the vasoactive factors, which allow ET-1 levels to increase and cause vasoconstriction [16]. Along with the biochemical mediators of endothelial function, mechanical forces and in particular, shear stress are also responsible for altering the phenotype of endothelial cells. Shear stress is the tangential stress that is applied to the artery wall and induces distension of the arteries; it is long-known that atherosclerotic lesions originate mainly in areas of low shear stress, where flow is disturbed [42, 43]. In these areas, endothelial cells promote a pro-atherogenic phenotype [44], enhancing both the local selectivity of plaque formation, as well as vessel wall remodeling, which, in turn, affects plaque vulnerability [45].

At the initial stages of atherosclerosis, leukocytes, particularly monocytes, progressively transmigrate through the endothelium with the help of monocyte chemoattractant protein 1 (MCP-1) and IL-8 [14]. Atherosclerosis progresses *via* the differentiation of monocytes into macrophages that retain low-density lipoproteins and become foam cells. Increased permeability of RA endothelial cells further enhances the entry of low-density lipoproteins and further promotes this cascade of events [46]. It has now been well-established that inflammation is involved in all phases of the atherothrombotic processes [2]. Whereas MCP-1 and IL-8 regulate monocyte transmigration, IL-8 can be upregulated in endothelial cells by CRP *via* the NF-kB pathway [47]. CRP can also increase the expression of macrophage colony stimulating factor (M-CSF), responsible for the differentiation of monocytes into macrophages. Subsequently, macrophage proliferation increases, thereby enhancing the progression of atherosclerosis *via* the promotion of the formation of foam cells [48]. An alternative effect of CRP is to destabilize the atheromatous plaque at later stages of atherosclerosis, by upregulation of matrix metalloproteinases [49] which causes weakening of the plaque's, fibrous cap. Apart from the above effects of CRP on these individual steps of atherosclerosis, it also enhances fibrinolytic procedures within the endothelial environment. CRP increases plasminogen activator inhibitor-1 mRNA through induction of the NF-kB signaling pathway [50] and downregulates endothelial tissue plasminogen activator, a serine protease that is responsible for fibrinolysis. This biochemical process is orchestrated by TNF-alpha and IL-1b [51]. Other important atherogenic properties of CRP include CRP-induced oxidative stress *via* overproduction of reactive oxygen species [52]. Reactive oxygen species in the endothelium enhance degradation of eNOS *via* eNOS uncoupling and subsequent increase in superoxide [53]. In general, superoxide reacts rapidly with NO, forming peroxynitrite which significantly impairs NO bioactivity [54] and deteriorates endothelial function.

TNF-alpha has direct effects on endothelial permeability by altering the distribution of endothelial cadherin-catenin and inhibiting the re-structure of F-actin fibers [55]. This procedure may enable monocyte transmigration into the endothelium and amplify the cascade of procedures that favor the formation of foam cells from macrophages [54]. Fur-

thermore, the proliferation of macrophages and increased formation of foam cells enhances secretion of TNF-alpha in the endothelium, perpetuating this vicious atherogenic cycle [3]. TNF-alpha also upregulates membrane expression of vascular cell adhesion molecule-1 [56], which plays a well established role in atherosclerosis [57] and it precedes the development of thickened intima with foam cell lesions [58]. Apart from these *in vitro* observations, increased expression of serum TNF-alpha has been found in patients with atherosclerotic complications in population studies [59, 60]. In RA patients, TNF-alpha enhances prothrombotic states such as dyslipidemia [61], but, most importantly, pooled evidence reveals that anti-TNF-alpha treatment may be in part responsible for reducing the risk for CVD events in RA [62]. Recent studies also reveal that TNF-alpha is involved in the more advanced processes of atherosclerosis, specific to formation of an advanced lesion *via* the inhibition of endothelial progenitor cells [63]. In RA patients, endothelial progenitor cell numbers are decreased compared to the healthy population [63], a phenomenon which reverses *via* blockade of TNF-alpha [64].

Inflammation in RA associates with dramatic increases of IL-6, IL-1 and TNF-alpha levels, which can trigger the acute-phase response resulting in high CRP, and have direct effects on the endothelium that may enhance atherosclerotic processes. As such, uncontrolled inflammation is thought to be amongst the prime factors involved in the accelerated atherosclerosis in RA [1, 2].

THE EFFECTS OF EXERCISE ON VASCULAR BIOLOGY

Increasing physical activity has been identified as one of the most important non-pharmacological interventions in both preventing and rehabilitating patients with non-

communicable chronic diseases such as CVD. This is because exercise reverses endothelial dysfunction and has important anti-atherogenic and anti-inflammatory effects (Fig. 1). Most importantly, even compared to standardized interventional strategies, exercise may improve survival at lower treatment costs [65].

Endothelial Function and Exercise

Exercise exerts significant effects on the endothelial system, both acutely and in the long term. These stimuli intervene and beneficially modify endothelial cell phenotype [17]. The acute effects of exercise are characterized by significant increase in blood flow pertaining to the increased metabolic demands of the exercising muscles and are closely related to the muscle fibre type recruitment (slow or fast twitch) as well as the duration and intensity of exercise [66]. The major mechanism responsible for exercise-induced hyperaemia is endothelial NO [67] and occurs due to the increased endothelial eNOS protein expression [68] as well as eNOS activation *via* Akt phosphorylation [69]. Exercise acutely increases local expression of endothelial progenitor cells (endothelial repair) as well as cultured/circulating angiogenic cells which promote angiogenesis and endothelial growth [70]. Shear stress which is the result of exercise-induced blood flow enhances intracellular mechanisms which increase eNOS transcription and upregulate anti-oxidative mechanisms. Specifically, tyrosine kinase c-Src acts to increase eNOS and extracellular superoxide dismutase expression, both of which relate to better endothelial function [71]. In addition, it has been well-established that shear stress increases the expression of PG I₂, a vasodilator and inhibitor of platelet aggregation [72]. Furthermore, vascular endothelial growth factor, which regulates capillary supply and exercise-induced angiogenesis, is significantly increased in response to exercise in both untrained and




Fig. (1). The effects of physical activity on endothelial function, atherosclerosis and inflammation.

trained individuals [73]. In the long term, the above mentioned physiological responses promote vascular remodelling, a necessary adaptation for improved oxygen exchange and blood flow delivery [74]. These long-term improvements are characterized by improved NO bioavailability and/or endothelium-dependent vasodilation that may even reverse age-related vascular deterioration [75, 76]. Improved endothelium-dependent vasodilation in response to long-term exercise is a consistent finding in both healthy [77] and disease populations [78].

Anti-Atherogenic Effects of Exercise

Acutely, due to exercise-induced shear stress, eNOS mRNA expression increases while ET-1 decreases [79, 80], while concomitant changes involve down-regulation of vascular cell adhesion molecule 1 [81]. Oxidative stress accelerates atherogenesis *via* oxidation of retained low-density lipoprotein, enhancing formation of foam cells [3]. An important long-term adaptation to exercise is the improvement in the anti-oxidative capacity of the human body, i.e. resistance to oxidative stress [82]. In particular, the enzymes catalase and superoxide dismutase which decompose reactive oxygen species are significantly increased as a result of habitual physical activity [83]. Increased eNOS in the endothelium increases anti-oxidant molecules such as superoxide dismutases 1 and 3 and angiotensin receptor type 2 and decreases oxidative molecules such as NADPH oxidase and angiotensin receptor type 1 [84]. Endothelial expression of adhesion molecules, selectins and MCP-1, all of which promote atherosclerosis, also decreases [85].

Anti-Inflammatory Effects of Exercise

Physical activity and/or moderate intensity exercise have profound anti-inflammatory effects in the healthy population as well as patients with chronic diseases [86]. In population-based studies, regular physical activity is consistently associated with a reduction in CRP levels [87, 88] which is also the case in patients with RA [89, 90]. The exact mechanisms whereby regular exercise reduces CRP have not yet been fully elucidated, but it seems that this association may exist *via* exercise-induced reduction in hypertension, triglycerides, and apolipoproteins, factors which are directly related to CRP concentration [91]. Most importantly, exercise reduces adipose tissue which is largely responsible for secretion of IL-6, a major trigger for hepatic production of CRP [92, 93]. However, these are physiological responses of moderate physical activity/exercise, whereas strenuous exercise may acutely promote an inflammatory response. CRP may increase immediately after prolonged strenuous exercise but returns to basal levels after approximately 48 hours, while this proportionate rise in CRP depends on the intensity of the exercise and the subsequent exercise-induced muscle damage [94, 95].

Recent advances in muscle physiology reveal that muscle tissue is responsible for the production of myokines (muscle cytokines) suggesting that muscle is also an endocrine organ [96]. Muscle contractions regulate expression of specific cytokines such as IL-6, -8, -10, and -15, as well as IL-1 receptor antagonist and TNF-alpha [96-98]. It is thought that, although IL-6 is predominantly a pro-inflammatory cytokine,

it may also have under certain circumstances anti-inflammatory properties [99]. In that light, and based on the fact that IL-6 increases exponentially with acute exercise up to 100 times compared to resting levels and returns to resting levels post-exercise [100], it has been suggested that IL-6 is the most important cytokine that induces the anti-inflammatory effects of exercise [96]. In contrast, pro-inflammatory cytokines IL-1b and TNF-alpha do not generally increase in response to moderate intensity exercise [101], whereas anti-inflammatory cytokines such as IL-1ra and IL-10, significantly increase [99, 102]. Interestingly, in response to regular exercise, IL-6 exerts its anti-inflammatory effects by primarily inhibiting the expression of TNF-alpha [99]. Data have demonstrated that IL-6 inhibits lipopolysaccharide-induced TNF-alpha in human blood mononuclear cells [103], whereas TNF-alpha levels are overexpressed in IL-6 deficient mice [104]. Recently, the transcriptional coactivator PGC1a has also attracted attention as it suppresses various inflammatory responses and regulates the effects of exercise [105] by coactivating transcription factors involved in biogenesis of mitochondria, oxidative phosphorylation and fatty acid oxidation [106, 107]. All these biological changes in cytokine expression in response to exercise, lead to beneficial long-term effects. Indeed, a consistent finding in the literature is that regular exercise relates to decreased systemic inflammation in both healthy and diseased populations [88, 89, 108, 109].

EFFECTS OF EXERCISE ON CVD IN RA

The extensive damage of the joint structures in RA has initially led to the notion that RA patients should rest, as exercise may enhance joint damage [110]. However, during the last decade substantial evidence deriving from randomised trials reveals that exercise inhibits the progression of the disease and improves both wellbeing and functional ability of RA patients [90]. This is due to normal physiological processes which develop as a result of exercise training such as the improved muscle co-ordination and hypertrophy, reduced fat mass and better immune function. Patients that indeed have to refrain from specific exercises are those with extensive structural damage; in this occasion load of the damaged joints has to be avoided and can be replaced with alternative types of exercise. The beneficial effects of different exercise training types on disease activity and severity have been confirmed several times by various studies [110, 111] and hence, based on these evidence, exercise has now been incorporated in the management of RA [112]. It is, therefore imperative that RA patients embark on exercise training programmes. Although exercise is beneficial, increasing physical activity in this population may prove difficult [113] perhaps due to the frequent advice from rheumatology specialists to RA patients, that exercise may exacerbate disease symptoms [114]. However, data from published studies [110] of excellent quality as well as anecdotal evidence, reveal that RA patients should and can exercise at high intensities, and this is beneficial for various outcomes of this chronic disease. Moreover, studies have collectively shown that overall the adherence rates in exercise programmes can be high, presumably due to the fact that the significant improvements achieved *via* exercise promote self motivation [90].

In a systematic review on exercise, CVD and RA, we found that, surprisingly, no studies have investigated the effects of exercise on CVD outcomes in this population [90]. However, given the ample evidence of the beneficial effects of physical activity/exercise on inhibiting disease progression and improving disease outcomes, we have proposed a model in which exercise should be incorporated in the management of the disease in order to prevent CVD, which is highly prevalent in this population. In this model, we suggest that effective treatment and control of inflammation should precede involvement in exercise in order to prohibit further damage and discomfort for the RA patient. Various drugs are available for patients with RA, including biological agents such as TNF-alpha, which target different biological pathways for inhibiting disease progression. Overall, medication strategies are effective in ameliorating disease activity (i.e. reduced expression of pro-inflammatory cytokines and CRP), a phenomenon that is also apparent as a response to exercise training. Nevertheless, given the lack of evidence regarding the effects of exercise on RA, potential changes in medication should always be based on robust clinical outcomes and patient feedback.

Exercise involvement and/or increased physical activity are included in the management of many chronic diseases. In RA, where patients experience increased prevalence of CVD, exercise may be even more important due to its anti-atherogenic and anti-inflammatory effects [17, 99]. The majority of studies on RA and exercise investigated the effects of physical activity/exercise regimens on improving RA-related disease outcomes. We have recently shown that physically inactive RA patients have worse CVD risk profile compared to physically active patients. In this cross-sectional study, we have investigated classical and novel risk factors, the prevalence of established CVD as well as the risk assessment of developing CVDs in physically active as well as inactive RA patients. Regarding the novel CVD risk factors, our results revealed that parameters associated with vascular function (e.g. von Willebrand factor), fibrinolysis atherogenesis (lipoproteins), pro-inflammatory cytokines, and CRP were markedly improved in patients who demonstrated increased levels of physical activity [89]. However, more research is required in order to investigate the potential associations of exercise on vascular function and CVD risk in this population as well as the mechanisms that underlie these associations.

CONCLUSION

Evidence from basic research supports a clear association between inflammation, vascular dysfunction and atherosclerosis. In RA, although evidence exists to support this association for atherosclerosis, the results are equivocal at least for endothelial dysfunction, potentially due to the differences in the methodological designs that do not take into account important factors that could influence endothelial function and atherosclerosis in RA. In addition, exercise has a protective effect *via* inhibiting the expression of inflammatory markers and their effects on both endothelial dysfunction and promotion of atherosclerosis. Future studies in RA should specifically investigate the effects of exercise regimes on

endothelial function and atherosclerosis both in population studies and at a molecular level.

REFERENCES

- Libby P. Role of inflammation in atherosclerosis associated with rheumatoid arthritis. *Am J Med* 2008; 121: S21-31.
- Sattar N, McCarey DW, Capell H, McInnes IB. Explaining how "high-grade" systemic inflammation accelerates vascular risk in rheumatoid arthritis. *Circulation* 2003; 108: 2957-63.
- Lamon BD, Hajjar DP. Inflammation at the molecular interface of atherogenesis: an anthropological journey. *Am J Pathol* 2008; 173: 1253-64.
- Gonzalez MA, Selwyn AP. Endothelial function, inflammation, and prognosis in cardiovascular disease. *Am J Med* 2003; 115 (Suppl 8A): 99S-106S.
- Kitas GD and Erb N. Tackling ischaemic heart disease in rheumatoid arthritis. *Rheumatology (Oxford)* 2003; 42: 607-13.
- Douglas KM, Pace AV, Trehan GJ, *et al.* Excess recurrent cardiac events in rheumatoid arthritis patients with acute coronary syndrome. *Ann Rheum Dis* 2006; 65: 348-53.
- Panoulas VF, Douglas KM, Milionis HJ, *et al.* Prevalence and associations of hypertension and its control in patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2007; 46: 1477-82.
- Panoulas VF, Metsios GS, Pace AV, *et al.* Hypertension in rheumatoid arthritis. *Rheumatology (Oxford)* 2008; 47: 1286-98.
- Stevens RJ, Douglas KM, Saratzis AN, Kitas GD. Inflammation and atherosclerosis in rheumatoid arthritis. *Expert Rev Mol Med* 2005; 7: 1-24.
- Brennan FM, McInnes IB. Evidence that cytokines play a role in rheumatoid arthritis. *J Clin Invest* 2008; 118: 3537-45.
- McInnes IB. Rheumatoid arthritis. From bench to bedside. *Rheum Dis Clin North Am* 2001; 27: 373-87.
- Choy EH, Panayi GS. Cytokine pathways and joint inflammation in rheumatoid arthritis. *N Engl J Med* 2001; 344: 907-16.
- Metsios GS, Stavropoulos-Kalinoglou A, Panoulas VF, *et al.* New resting energy expenditure prediction equations for patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2008; 47: 500-6.
- Devaraj S, Singh U, Jialal I. The evolving role of C-reactive protein in atherothrombosis. *Clin Chem* 2009; 55: 229-38.
- Zhang H, Park Y, Wu J, *et al.* Role of TNF-alpha in vascular dysfunction. *Clin Sci (Lond)* 2009; 116: 219-30.
- Vanhoutte PM. Say NO to ET. *J Auton Nerv Syst* 2000; 81: 271-7.
- Laughlin MH, Roseguini B. Mechanisms for exercise training-induced increases in skeletal muscle blood flow capacity: differences with interval sprint training versus aerobic endurance training. *J Physiol Pharmacol* 2008; 59 (Suppl 7): 71-88.
- Lerman A, Zeiher AM. Endothelial function: cardiac events. *Circulation* 2005; 111: 363-8.
- Venugopal SK, Devaraj S, Yuhanna I, Shaul P, Jialal I. Demonstration that C-reactive protein decreases eNOS expression and bioactivity in human aortic endothelial cells. *Circulation* 2002; 106: 1439-41.
- Nagaoka T, Kuo L, Ren Y, Yoshida A, Hein TW. C-reactive protein inhibits endothelium-dependent nitric oxide-mediated dilation of retinal arterioles *via* enhanced superoxide production. *Invest Ophthalmol Vis Sci* 2008; 49: 2053-60.
- Qamirani E, Ren Y, Kuo L, Hein TW. C-reactive protein inhibits endothelium-dependent NO-mediated dilation in coronary arterioles by activating p38 kinase and NAD(P)H oxidase. *Arterioscler Thromb Vasc Biol* 2005; 25: 995-1001.
- Singh U, Devaraj S, Vasquez-Vivar J, Jialal I. C-reactive protein decreases endothelial nitric oxide synthase activity *via* uncoupling. *J Mol Cell Cardiol* 2007; 43: 780-91.
- Verma S, Li SH, Badiwala MV, *et al.* Endothelin antagonism and interleukin-6 inhibition attenuate the proatherogenic effects of C-reactive protein. *Circulation* 2002; 105: 1890-6.
- Goodwin BL, Pendleton LC, Levy MM, Solomonson LP, Eichler DC. Tumor necrosis factor-alpha reduces argininosuccinate

- synthase expression and nitric oxide production in aortic endothelial cells. *Am J Physiol Heart Circ Physiol* 2007; 293: H1115-21.
- [25] De Palma C, Meacci E, Perrotta C, Bruni P, Clementi E. Endothelial nitric oxide synthase activation by tumor necrosis factor alpha through neutral sphingomyelinase 2, sphingosine kinase 1, and sphingosine 1 phosphate receptors: a novel pathway relevant to the pathophysiology of endothelium. *Arterioscler Thromb Vasc Biol* 2006; 26: 99-105.
- [26] Sorescu D, Griendling KK. Reactive oxygen species, mitochondria, and NAD(P)H oxidases in the development and progression of heart failure. *Congest Heart Fail* 2002; 8: 132-40.
- [27] Gao X, Belmadani S, Picchi A, *et al.* Tumor necrosis factor-alpha induces endothelial dysfunction in Lepr(db) mice. *Circulation* 2007; 115: 245-54.
- [28] Hurlimann D, Forster A, Noll G, *et al.* Anti-tumor necrosis factor-alpha treatment improves endothelial function in patients with rheumatoid arthritis. *Circulation* 2002; 106: 2184-7.
- [29] Fichtlscherer S, Rossig L, Breuer S, Vasa M, Dimmeler S, Zeiher AM. Tumor necrosis factor antagonism with etanercept improves systemic endothelial vasoreactivity in patients with advanced heart failure. *Circulation* 2001; 104: 3023-5.
- [30] Roman MJ, Moeller E, Davis A, *et al.* Preclinical carotid atherosclerosis in patients with rheumatoid arthritis. *Ann Intern Med* 2006; 144: 249-56.
- [31] Hannawi S, Marwick TH, Thomas R. Inflammation predicts accelerated brachial arterial wall changes in patients with recent-onset rheumatoid arthritis. *Arthritis Res Ther* 2009; 11: R51.
- [32] Vlachopoulos C, Dima I, Aznaouridis K, *et al.* Acute systemic inflammation increases arterial stiffness and decreases wave reflections in healthy individuals. *Circulation* 2005; 112: 2193-200.
- [33] Hingorani AD, Cross J, Kharbanda RK, *et al.* Acute systemic inflammation impairs endothelium-dependent dilatation in humans. *Circulation* 2000; 102: 994-9.
- [34] Wallberg-Jonsson S, Caidahl K, Klintland N, Nyberg G, Rantapaa-Dahlqvist S. Increased arterial stiffness and indication of endothelial dysfunction in long-standing rheumatoid arthritis. *Scand J Rheumatol* 2008; 37: 1-5.
- [35] Dessein PH, Joffe BI, Singh S. Biomarkers of endothelial dysfunction, cardiovascular risk factors and atherosclerosis in rheumatoid arthritis. *Arthritis Res Ther* 2005; 7: R634-43.
- [36] Gonzalez-Gay MA, Gonzalez-Juanatey C, Pineiro A, Garcia-Porrúa C, Testa A, Llorca J. High-grade C-reactive protein elevation correlates with accelerated atherogenesis in patients with rheumatoid arthritis. *J Rheumatol* 2005; 32: 1219-23.
- [37] Klocke R, Cockcroft JR, Taylor GJ, Hall IR, Blake DR. Arterial stiffness and central blood pressure, as determined by pulse wave analysis, in rheumatoid arthritis. *Ann Rheum Dis* 2003; 62: 414-8.
- [38] Dessein PH, Joffe BI, Veller MG, *et al.* Traditional and nontraditional cardiovascular risk factors are associated with atherosclerosis in rheumatoid arthritis. *J Rheumatol* 2005; 32: 435-42.
- [39] Vaudo G, Marchesi S, Gerli R, *et al.* Endothelial dysfunction in young patients with rheumatoid arthritis and low disease activity. *Ann Rheum Dis* 2004; 63: 31-5.
- [40] Del Rincon I, O'Leary DH, Freeman GL, Escalante A. Acceleration of atherosclerosis during the course of rheumatoid arthritis. *Atherosclerosis* 2007; 195: 354-60.
- [41] Veldhuijzen van Zanten JJCS, Kitas GD. Inflammation, carotid intima-media thickness and atherosclerosis in rheumatoid arthritis. *Arthritis Res Ther* 2008; 10: 102.
- [42] Friedman MH, Deters OJ, Barger CB, Hutchins GM, Mark FF. Shear-dependent thickening of the human arterial intima. *Atherosclerosis* 1986; 60: 161-71.
- [43] Ku DN, Giddens DP, Phillips DJ, Strandness DE, Jr. Hemodynamics of the normal human carotid bifurcation: *in vitro* and *in vivo* studies. *Ultrasound Med Biol* 1985; 11: 13-26.
- [44] Malek AM, Alper SL, Izumo S. Hemodynamic shear stress and its role in atherosclerosis. *JAMA* 1999; 282: 2035-42.
- [45] Cunningham KS, Gotlieb AI. The role of shear stress in the pathogenesis of atherosclerosis. *Lab Invest* 2005; 85: 9-23.
- [46] Hahn BH, Grossman J, Chen W, McMahon M. The pathogenesis of atherosclerosis in autoimmune rheumatic diseases: roles of inflammation and dyslipidemia. *J Autoimmun* 2007; 28: 69-75.
- [47] Wang Q, Zhu X, Xu Q, Ding X, Chen YE, Song Q. Effect of C-reactive protein on gene expression in vascular endothelial cells. *Am J Physiol Heart Circ Physiol* 2005; 288: H1539-45.
- [48] Devaraj S, Yun JM, Duncan-Staley C, Jialal I. C-reactive protein induces M-CSF release and macrophage proliferation. *J Leukoc Biol* 2009; 85: 262-7.
- [49] Williams TN, Zhang CX, Game BA, He L, Huang Y. C-reactive protein stimulates MMP-1 expression in U937 histiocytes through Fe[gamma]RII and extracellular signal-regulated kinase pathway: an implication of CRP involvement in plaque destabilization. *Arterioscler Thromb Vasc Biol* 2004; 24: 61-6.
- [50] Nakakuki T, Ito M, Iwasaki H, *et al.* Rho/Rho-kinase pathway contributes to C-reactive protein-induced plasminogen activator inhibitor-1 expression in endothelial cells. *Arterioscler Thromb Vasc Biol* 2005; 25: 2088-93.
- [51] Singh U, Devaraj S, Jialal I. C-reactive protein decreases tissue plasminogen activator activity in human aortic endothelial cells: evidence that C-reactive protein is a procoagulant. *Arterioscler Thromb Vasc Biol* 2005; 25: 2216-21.
- [52] Prasad K. C-reactive protein increases oxygen radical generation by neutrophils. *J Cardiovasc Pharmacol Ther* 2004; 9: 203-9.
- [53] Landmesser U, Dikalov S, Price SR, *et al.* Oxidation of tetrahydrobiopterin leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J Clin Invest* 2003; 111: 1201-9.
- [54] Landmesser U, Hornig B, Drexler H. Endothelial function: a critical determinant in atherosclerosis? *Circulation* 2004; 109: I127-33.
- [55] Wojciak-Stothard B, Entwistle A, Garg R, Ridley AJ. Regulation of TNF-alpha-induced reorganization of the actin cytoskeleton and cell-cell junctions by Rho, Rac, and Cdc42 in human endothelial cells. *J Cell Physiol* 1998; 176: 150-65.
- [56] Iademaro MF, Barks JL, Dean DC. Regulation of vascular cell adhesion molecule-1 expression by IL-4 and TNF-alpha in cultured endothelial cells. *J Clin Invest* 1995; 95: 264-71.
- [57] Cybulsky MI, Gimbrone MA, Jr. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Science* 1991; 251: 788-91.
- [58] Li H, Cybulsky MI, Gimbrone MA, Jr., Libby P. An atherogenic diet rapidly induces VCAM-1, a cytokine-regulatable mononuclear leukocyte adhesion molecule, in rabbit aortic endothelium. *Arterioscler Thromb* 1993; 13: 197-204.
- [59] Ozeren A, Aydin M, Tokac M, *et al.* Levels of serum IL-1beta, IL-2, IL-8 and tumor necrosis factor-alpha in patients with unstable angina pectoris. *Mediators Inflamm* 2003; 12: 361-5.
- [60] Naya M, Tsukamoto T, Morita K, *et al.* Plasma interleukin-6 and tumor necrosis factor-alpha can predict coronary endothelial dysfunction in hypertensive patients. *Hypertens Res* 2007; 30: 541-8.
- [61] Dixon WG, Symmons DP. What effects might anti-TNFalpha treatment be expected to have on cardiovascular morbidity and mortality in rheumatoid arthritis? A review of the role of TNFalpha in cardiovascular pathophysiology. *Ann Rheum Dis* 2007; 66: 1132-6.
- [62] Avouac J, Allanore Y. Cardiovascular risk in rheumatoid arthritis: effects of anti-TNF drugs. *Expert Opin Pharmacother* 2008; 9: 1121-8.
- [63] Grisar J, Aletaha D, Steiner CW, *et al.* Depletion of endothelial progenitor cells in the peripheral blood of patients with rheumatoid arthritis. *Circulation* 2005; 111: 204-11.
- [64] Grisar J, Aletaha D, Steiner CW, *et al.* Endothelial progenitor cells in active rheumatoid arthritis: effects of tumour necrosis factor and glucocorticoid therapy. *Ann Rheum Dis* 2007; 66: 1284-8.
- [65] Hambrecht R, Walther C, Mobius-Winkler S, *et al.* Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease: a randomized trial. *Circulation* 2004; 109: 1371-8.
- [66] Laughlin MH, Armstrong RB. Muscle blood flow during locomotory exercise. *Exerc Sport Sci Rev* 1985; 13: 95-136.

- [67] Tschakovsky ME, Joyner MJ. Nitric oxide and muscle blood flow in exercise. *Appl Physiol Nutr Metab* 2008; 33: 151-61.
- [68] McAllister RM, Jaspere JL, Laughlin MH. Nonuniform effects of endurance exercise training on vasodilation in rat skeletal muscle. *J Appl Physiol* 2005; 98: 753-61.
- [69] Hambrecht R, Adams V, Erbs S, *et al.* Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation* 2003; 107: 3152-8.
- [70] Rehman J, Li J, Parvathaneni L, *et al.* Exercise acutely increases circulating endothelial progenitor cells and monocyte/macrophage-derived angiogenic cells. *J Am Coll Cardiol* 2004; 43: 2314-8.
- [71] Davis ME, Cai H, McCann L, Fukai T, Harrison DG. Role of c-Src in regulation of endothelial nitric oxide synthase expression during exercise training. *Am J Physiol Heart Circ Physiol* 2003; 284: H1449-53.
- [72] Frangos JA, Eskin SG, McIntire LV, Ives CL. Flow effects on prostacyclin production by cultured human endothelial cells. *Science* 1985; 227: 1477-9.
- [73] Kraus RM, Stallings HW, 3rd, Yeager RC, Gavin TP. Circulating plasma VEGF response to exercise in sedentary and endurance-trained men. *J Appl Physiol* 2004; 96: 1445-50.
- [74] Yang HT, Prior BM, Lloyd PG, *et al.* Training-induced vascular adaptations to ischemic muscle. *J Physiol Pharmacol* 2008; 59 (Suppl 7): 57-70.
- [75] Hambrecht R, Niebauer J, Marburger C, *et al.* Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *J Am Coll Cardiol* 1993; 22: 468-77.
- [76] DeSouza CA, Shapiro LF, Clevenger CM, *et al.* Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation* 2000; 102: 1351-7.
- [77] Clarkson P, Montgomery HE, Mullen MJ, *et al.* Exercise training enhances endothelial function in young men. *J Am Coll Cardiol* 1999; 33: 1379-85.
- [78] Higashi Y, Sasaki S, Sasaki N, *et al.* Daily aerobic exercise improves reactive hyperemia in patients with essential hypertension. *Hypertension* 1999; 33: 591-7.
- [79] Tuttle JL, Nachreiner RD, Bhuller AS, *et al.* Shear level influences resistance artery remodeling: wall dimensions, cell density, and eNOS expression. *Am J Physiol Heart Circ Physiol* 2001; 281: H1380-9.
- [80] Malek A, Izumo S. Physiological fluid shear stress causes down-regulation of endothelin-1 mRNA in bovine aortic endothelium. *Am J Physiol* 1992; 263: C389-96.
- [81] Himburg HA, Dowd SE, Friedman MH. Frequency-dependent response of the vascular endothelium to pulsatile shear stress. *Am J Physiol Heart Circ Physiol* 2007; 293: H645-53.
- [82] Radak Z, Taylor AW, Ohno H, Goto S. Adaptation to exercise-induced oxidative stress: from muscle to brain. *Exerc Immunol Rev* 2001; 7: 90-107.
- [83] Seals DR, Desouza CA, Donato AJ, Tanaka H. Habitual exercise and arterial aging. *J Appl Physiol* 2008; 105: 1323-32.
- [84] Adams V, Linke A, Krankel N, *et al.* Impact of regular physical activity on the NAD(P)H oxidase and angiotensin receptor system in patients with coronary artery disease. *Circulation* 2005; 111: 555-62.
- [85] Chien S. Mechanotransduction and endothelial cell homeostasis: the wisdom of the cell. *Am J Physiol Heart Circ Physiol* 2007; 292: H1209-24.
- [86] Kaspis C, Thompson PD. The effects of physical activity on serum C-reactive protein and inflammatory markers: a systematic review. *J Am Coll Cardiol* 2005; 45: 1563-9.
- [87] Albert MA, Glynn RJ, Ridker PM. Effect of physical activity on serum C-reactive protein. *Am J Cardiol* 2004; 93: 221-5.
- [88] Geffken DF, Cushman M, Burke GL, Polak JF, Sakkinen PA, Tracy RP. Association between physical activity and markers of inflammation in a healthy elderly population. *Am J Epidemiol* 2001; 153: 242-50.
- [89] Metsios GS, Stavropoulos-Kalinoglou A, Panoulas VF, *et al.* Association of physical inactivity with increased cardiovascular risk in patients with rheumatoid arthritis. *Eur J Cardiovasc Prev Rehabil* 2009; 16: 188-94.
- [90] Metsios GS, Stavropoulos-Kalinoglou A, Veldhuijzen van Zanten JJCS, *et al.* Rheumatoid arthritis, cardiovascular disease and physical exercise: a systematic review. *Rheumatology (Oxford)* 2008; 47: 239-48.
- [91] Ford ES. Does exercise reduce inflammation? Physical activity and C-reactive protein among U.S. adults. *Epidemiology* 2002; 13: 561-8.
- [92] McLaughlin T, Abbasi F, Lamendola C, *et al.* Differentiation between obesity and insulin resistance in the association with C-reactive protein. *Circulation* 2002; 106: 2908-12.
- [93] Abbasi F, Brown BW, Jr., Lamendola C, McLaughlin T, Reaven GM. Relationship between obesity, insulin resistance, and coronary heart disease risk. *J Am Coll Cardiol* 2002; 40: 937-43.
- [94] Taylor C, Rogers G, Goodman C, *et al.* Hematologic, iron-related, and acute-phase protein responses to sustained strenuous exercise. *J Appl Physiol* 1987; 62: 464-9.
- [95] Siegel AJ, Stec JJ, Lipinska I, *et al.* Effect of marathon running on inflammatory and hemostatic markers. *Am J Cardiol* 2001; 88: 918-20, A9.
- [96] Pedersen BK, Akerstrom TC, Nielsen AR, Fischer CP. Role of myokines in exercise and metabolism. *J Appl Physiol* 2007; 103: 1093-8.
- [97] Akerstrom T, Steensberg A, Keller P, Keller C, Penkowa M, Pedersen BK. Exercise induces interleukin-8 expression in human skeletal muscle. *J Physiol* 2005; 563: 507-16.
- [98] Febbraio MA, Pedersen BK. Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. *FASEB J* 2002; 16: 1335-47.
- [99] Petersen AM and Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol* 2005; 98: 1154-62.
- [100] Steensberg A, van Hall G, Osada T, Sacchetti M, Saltin B, Klarlund PB. Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *J Physiol* 2000; 529 Pt 1: 237-42.
- [101] Mathur N, Pedersen BK. Exercise as a mean to control low-grade systemic inflammation. *Mediators Inflamm* 2008; 2008: 109502.
- [102] Febbraio MA, Pedersen BK. Contraction-induced myokine production and release: is skeletal muscle an endocrine organ? *Exerc Sport Sci Rev* 2005; 33: 114-9.
- [103] Schindler R, Mancilla J, Endres S, Ghorbani R, Clark SC, Dinarello CA. Correlations and interactions in the production of interleukin-6 (IL-6), IL-1, and tumor necrosis factor (TNF) in human blood mononuclear cells: IL-6 suppresses IL-1 and TNF. *Blood* 1990; 75: 40-7.
- [104] Mizuhara H, O'Neill E, Seki N, *et al.* T cell activation-associated hepatic injury: mediation by tumor necrosis factors and protection by interleukin 6. *J Exp Med* 1994; 179: 1529-37.
- [105] Handschin C, Spiegelman BM. The role of exercise and PGC1alpha in inflammation and chronic disease. *Nature* 2008; 454: 463-9.
- [106] Mootha VK, Lindgren CM, Eriksson KF, *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 2003; 34: 267-73.
- [107] Huss JM, Kopp RP, Kelly DP. Peroxisome proliferator-activated receptor coactivator-1alpha (PGC-1alpha) coactivates the cardiac-enriched nuclear receptors estrogen-related receptor-alpha and -gamma. Identification of novel leucine-rich interaction motif within PGC-1alpha. *J Biol Chem* 2002; 277: 40265-74.
- [108] Fischer CP, Berntsen A, Perstrup LB, Eskildsen P, Pedersen BK. Plasma levels of interleukin-6 and C-reactive protein are associated with physical inactivity independent of obesity. *Scand J Med Sci Sports* 2007; 17: 580-7.
- [109] Tisi PV, Hulse M, Chulakadabba A, Gosling P, Shearman CP. Exercise training for intermittent claudication: does it adversely affect biochemical markers of the exercise-induced inflammatory response? *Eur J Vasc Endovasc Surg* 1997; 14: 344-50.

- [110] de Jong Z, Munneke M, Zwinderman AH, *et al.* Is a long-term high-intensity exercise program effective and safe in patients with rheumatoid arthritis? Results of a randomized controlled trial. *Arthritis Rheum* 2003; 48: 2415-24.
- [111] Noreau L, Martineau H, Roy L, Belzile M. Effects of a modified dance-based exercise on cardiorespiratory fitness, psychological state and health status of persons with rheumatoid arthritis. *Am J Phys Med Rehabil* 1995; 74: 19-27.
- [112] ARC. Guidelines for the management of rheumatoid arthritis: 2002 Update. *Arthritis Rheum* 2002; 46: 328-46.
- [113] Gordon MM, Thomson EA, Madhok R, Capell HA. Can intervention modify adverse lifestyle variables in a rheumatoid population? Results of a pilot study. *Ann Rheum Dis* 2002; 61: 66-9.
- [114] Munneke M, de Jong Z, Zwinderman AH, *et al.* High intensity exercise or conventional exercise for patients with rheumatoid arthritis? Outcome expectations of patients, rheumatologists, and physiotherapists. *Ann Rheum Dis* 2004; 63: 804-8.

Received: November 20, 2009

Revised: November 30, 2009

Accepted: December 14, 2009

© Metsios *et al.*; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.