

*Pacific
Journal of
Mathematics*

**VECTOR BUNDLE VALUED DIFFERENTIAL FORMS ON
 $\mathbb{N}Q$ -MANIFOLDS**

LUCA VITAGLIANO

VECTOR BUNDLE VALUED DIFFERENTIAL FORMS ON $\mathbb{N}Q$ -MANIFOLDS

LUCA VITAGLIANO

Geometric structures on $\mathbb{N}Q$ -manifolds, i.e., nonnegatively graded manifolds with a homological vector field, encode nongraded geometric data on Lie algebroids and their higher analogues. A particularly relevant class of structures consists of vector bundle valued differential forms. Symplectic forms, contact structures and, more generally, distributions are in this class. We describe vector bundle valued differential forms on nonnegatively graded manifolds in terms of nongraded geometric data. Moreover, we use this description to present, in a unified way, novel proofs of known results, and new results about degree-one $\mathbb{N}Q$ -manifolds equipped with certain geometric structures, namely symplectic structures, contact structures, involutive distributions (already present in literature), locally conformal symplectic structures, and generic vector bundle valued higher order forms, in particular presymplectic and multisymplectic structures (not yet present in literature).

1. Introduction	449
2. Vector valued forms on graded manifolds	452
3. Vector valued 1-forms on $\mathbb{N}Q$ -manifolds	461
4. Vector Valued 2-forms on $\mathbb{N}Q$ -manifolds	468
5. Higher-degree forms on degree-one $\mathbb{N}Q$ -manifolds	474
Appendix A. Locally conformal symplectic manifolds revisited	477
Appendix B. Lie algebroids and their representations	479
References	481

1. Introduction

Graded geometry encodes (nongraded) geometric structures in an efficient way. For instance, a vector bundle is the same as a degree-one nonnegatively graded manifold. In this respect $\mathbb{N}Q$ -manifolds, i.e., nonnegatively graded manifolds equipped with a homological vector field, are of a special interest. Namely, they

MSC2010: 53D17, 58A50.

Keywords: graded manifolds, $\mathbb{N}Q$ -manifolds, vector bundle valued differential forms, Lie algebroids, Spencer operators.

encode Lie algebroids in degree one, and higher versions of Lie algebroids (including homotopy Lie algebroids) in higher degrees [Voronov 2010] (see also [Bruce 2011; Kjeseth 2001b; Sati et al. 2012; Sheng and Zhu 2011; Bonavolontà and Poncin 2013; Vitagliano 2015b], and [Kjeseth 2001a; Vitagliano 2014] for applications of homotopy Lie algebroids). Accordingly, geometric data on Lie algebroids, or higher versions of them, that are compatible with the algebroid structure, can be encoded by suitable geometric structures on an $\mathbb{N}Q$ -manifold that are preserved by the homological vector field. This is a general rule with various examples scattered in the literature. For instance, degree-one *symplectic $\mathbb{N}Q$ -manifolds* are equivalent to Poisson manifolds (which can be understood as Lie algebroids of a special kind) [Roytenberg 2002]. Similarly, degree-one *contact $\mathbb{N}Q$ -manifolds* [Grabowski 2013; Mehta 2013] are equivalent to Jacobi manifolds, and degree-one $\mathbb{N}Q$ -manifolds equipped with a compatible involutive distribution are equivalent to Lie algebroids equipped with an *IM foliation* (see [Jotz and Ortiz 2011] for a definition) [Zambon and Zhu 2012]. More examples can be presented in degree two. For instance, degree-two symplectic $\mathbb{N}Q$ -manifolds are equivalent to Courant algebroids [Roytenberg 2002], and degree-two contact $\mathbb{N}Q$ -manifolds encode a contact version of Courant algebroids: *Grabowski's contact-Courant algebroids* [Grabowski 2013].

In all examples above the geometric structure on the $\mathbb{N}Q$ -manifold is, or can be understood as, a *differential form with values in a vector bundle*. This motivates the study of vector bundle valued differential forms (vector valued forms, in the following) on graded manifolds, and, in particular, $\mathbb{N}Q$ -manifolds. In this paper, we describe vector valued forms on nonnegatively graded manifolds in terms of nongraded geometric data (Theorem 10). Later we apply this description to the study of degree-one $\mathbb{N}Q$ -manifolds equipped with a compatible vector valued form. In this way, we get a unified formalism for the description of degree-one contact $\mathbb{N}Q$ -manifolds, symplectic $\mathbb{N}Q$ -manifolds, and $\mathbb{N}Q$ -manifolds equipped with a compatible involutive distribution. In particular, we manage to present alternative proofs of results (in degree one) of Roytenberg [2002], Grabowski [2013], Mehta [2013], and Zambon and Zhu [2012]. We also discuss three new examples. Namely, we show that degree-one presymplectic $\mathbb{N}Q$ -manifolds (with an additional nondegeneracy condition) are basically equivalent to Dirac manifolds (Corollary 31). We also show that degree-one locally conformal symplectic $\mathbb{N}Q$ -manifolds are equivalent to locally conformal Poisson manifolds (Theorem 35), and, more generally, degree-one $\mathbb{N}Q$ -manifolds equipped with a compatible, higher degree, vector valued form are equivalent to Lie algebroids equipped with *Spencer operators* (Theorem 36). The latter have been recently introduced in [Crainic et al. 2015] (see also [Salazar 2013]) as infinitesimal counterparts of multiplicative vector valued forms on Lie groupoids. In particular, degree-one multisymplectic $\mathbb{N}Q$ -manifolds are equivalent to Lie algebroids equipped with an *IM multisymplectic*

structure [Bursztyn et al. 2015] (Theorem 39). We stress that we do only consider differential forms with values in vector bundles generated in one single degree (which, up to a shift, are actually generated in degree zero). This hypothesis simplifies the discussion a lot. We hope to discuss the general case, as well as higher-degree cases, elsewhere.

The paper is divided into three main sections and two appendixes. In Section 2, after a short review of vector valued Cartan calculus on graded manifolds, we present our description of vector valued forms on \mathbb{N} -manifolds in terms of nongraded geometric data (Theorem 10). As already remarked, this description allows one to present in a unified way various results scattered in the literature about the correspondence between geometric structures on degree-one $\mathbb{N}Q$ -manifolds and (nongraded) geometric structures on Lie algebroids. In Section 3, we discuss 1-forms on degree-one $\mathbb{N}Q$ -manifolds. Surjective 1-forms are the same as distributions and we discuss in some detail the contact and involutive cases. The results of this section (Theorem 23 and Theorem 25) are already present in literature, but they are presented here in a new and unified way that allows a straightforward generalization to (possibly degenerate) differential forms of higher order. In Section 4, we discuss 2-forms (on degree-one $\mathbb{N}Q$ -manifolds). In particular, we present a novel proof of the remark of Roytenberg [2002] that degree-one symplectic $\mathbb{N}Q$ -manifolds are equivalent to Poisson manifolds (Theorem 27). We also generalize Roytenberg result in two different directions, namely to presymplectic forms on one side (Theorem 30 and Corollary 31) and to locally conformal symplectic structures on the other side (Theorem 35). In Section 5 we discuss the general case of a differential form of arbitrarily high order. In particular, we relate compatible vector valued forms on $\mathbb{N}Q$ -manifold and the Spencer operators of Crainic–Salazar–Struchiner [Crainic et al. 2015; Salazar 2013] (Theorem 36). Finally, we discuss degree-one multisymplectic $\mathbb{N}Q$ -manifolds (Theorem 39). The paper is complemented by two appendixes. In Appendix A, we revisit briefly the concept of locally conformal symplectic manifolds [Vaisman 1985], and give a slightly more intrinsic definition of them. We also briefly review the relation between locally conformal symplectic manifolds and locally conformal Poisson manifolds [Vaisman 2007]. In Appendix B, we review the definition of Lie algebroids and their representations. As already remarked they play a key role in the paper.

1.1. Notation and conventions. Let $V = \bigoplus_i V_i$ be a graded vector space. We denote by $|v|$ the degree of a homogeneous element, i.e., $|v| = i$ whenever $v \in V_i$, unless otherwise stated.

Let \mathcal{M} be a (graded) manifold, and $\mathcal{E} \rightarrow \mathcal{M}$ a (graded) vector bundle on it. We denote by M the support of \mathcal{M} . In the case when \mathcal{M} is nonnegatively graded, M is also the degree-zero shadow of \mathcal{M} . Moreover, we denote by $C_i^\infty(\mathcal{M})$, (resp.

$\mathfrak{X}_i(\mathcal{M})$, $\Gamma_i(\mathcal{E})$) the vector space of degree- i smooth functions on \mathcal{M} (resp. vector fields on \mathcal{M} , sections of \mathcal{E}). We also denote by $\mathfrak{X}_-(\mathcal{M})$ the graded vector space of negatively graded vector fields on \mathcal{M} . Sometimes, if there is no risk of confusion, we denote by \mathcal{E} the (graded) $C^\infty(\mathcal{M})$ -module of sections of \mathcal{E} . Similarly, we often identify (graded) vector bundle morphisms and (graded) homomorphisms between modules of sections.

We adopt the Einstein summation convention.

2. Vector valued forms on graded manifolds

2.1. $\mathbb{N}Q$ -manifolds and vector $\mathbb{N}Q$ -bundles. We refer to [Roytenberg 2002; Mehta 2006; Cattaneo and Schätz 2011] for details about graded manifolds, and, in particular, \mathbb{N} -manifolds. In the following, we just recall some basic facts which will be often used below. We will work with the simplest possible notion of a *graded manifold*. Namely, any graded manifold \mathcal{M} in this paper is equipped with one single \mathbb{Z} -grading in its algebra $C^\infty(\mathcal{M})$ of smooth functions (unless otherwise stated). Moreover, $C^\infty(\mathcal{M})$ is graded commutative with respect to the grading. We will call *degree* the grading. We will focus on \mathbb{N} -manifolds, i.e., *nonnegatively graded* manifolds. Recall that the *degree of an \mathbb{N} -manifold* is the highest degree of its coordinates. Similarly, the degree of a vector \mathbb{N} -bundle, i.e., a nonnegatively graded vector bundle over an \mathbb{N} -manifold, is the highest degree of its fiber coordinates.

Example 1. Every degree-one \mathbb{N} -manifold \mathcal{M} is of the form $A[1]$ for some non-graded vector bundle $A \rightarrow M$, and one has $C^\infty(\mathcal{M}) = \Gamma(\wedge^* A^*)$. In particular, degree-zero functions on \mathcal{M} identify with functions on M , and degree-one functions on \mathcal{M} identify with sections of A^* . Accordingly, vector fields of degree -1 on \mathcal{M} identify with sections of A . In the following, we will tacitly understand the identifications $C_0^\infty(\mathcal{M}) \simeq C^\infty(M)$, $C_1^\infty(\mathcal{M}) \simeq \Gamma(A^*)$, and $\mathfrak{X}_{-1}(\mathcal{M}) \simeq \Gamma(A)$. The action of a vector field $X \in \Gamma(A)$ of degree -1 on a degree-one function $f \in \Gamma(A^*)$ is given by the *duality pairing*: $X(f) = \langle X, f \rangle$.

Example 2. Recall that every \mathbb{N} -manifold \mathcal{M} is fibered over its degree-zero shadow M . Every degree-zero vector \mathbb{N} -bundle \mathcal{E} over \mathcal{M} is of the form $\mathcal{M} \times_M E$ for some nongraded vector bundle $E \rightarrow M$, and one has $\Gamma(\mathcal{E}) = C^\infty(\mathcal{M}) \otimes \Gamma(E)$ (where the tensor product is over $C^\infty(M)$). In particular, degree-zero sections of \mathcal{E} identify with sections of E . In the following, we will tacitly understand the identification $\Gamma_0(\mathcal{E}) \simeq \Gamma(E)$.

A Q -manifold is a graded manifold \mathcal{M} equipped with a homological vector field Q , i.e., a degree-one vector field Q such that $[Q, Q] = 0$. An $\mathbb{N}Q$ -manifold is a nonnegatively graded Q -manifold.

Example 3. Every degree-one $\mathbb{N}Q$ -manifold (\mathcal{M}, Q) is of the form $(A[1], d_A)$ for some nongraded Lie algebroid $A \rightarrow M$ (see Appendix B for a definition of Lie

algebroid). Here d_A is the homological derivation induced in $\Gamma(\wedge^* A^*) = C^\infty(\mathcal{M})$. The Lie bracket $[[-, -]]$ in $\Gamma(A)$ and the anchor $\rho : \Gamma(A) \rightarrow \mathfrak{X}(\mathcal{M})$ can be recovered from Q via formulas

$$[[X, Y]] = [[Q, X], Y], \quad \rho(X)(f) = [Q, X](f),$$

where $X, Y \in \Gamma(A)$ are also interpreted as vector fields of degree -1 on \mathcal{M} (so that $[[Q, X], Y]$ also has degree -1), and $f \in C^\infty(\mathcal{M})$.

Similarly, we call a Q -vector bundle (resp. $\mathbb{N}\mathbb{Q}$ -vector bundle) a graded vector bundle $\mathcal{E} \rightarrow \mathcal{M}$ (resp. a vector \mathbb{N} -bundle) equipped with a homological derivation. In this respect, recall that a (graded) derivation of \mathcal{E} is a graded, \mathbb{R} -linear map $\mathbb{X} : \Gamma(\mathcal{E}) \rightarrow \Gamma(\mathcal{E})$ such that

$$\mathbb{X}(fe) = X(f)e + (-)^{|f||\mathbb{X}|} f\mathbb{X}(e), \quad f \in C^\infty(\mathcal{M}), \quad e \in \Gamma(\mathcal{E}),$$

for a (necessarily unique) vector field $X \in \mathfrak{X}(\mathcal{M})$ called the *symbol of \mathbb{X}* . Clearly, a derivation of \mathcal{E} is completely determined by its symbol and its action on generators of $\Gamma(\mathcal{E})$.

Example 4. Denote by Δ the grading vector field on \mathcal{M} , i.e., $\Delta(f) = |f|f$, for all homogeneous functions f on \mathcal{M} . The grading $\Delta_{\mathcal{E}} : \Gamma(\mathcal{E}) \rightarrow \Gamma(\mathcal{E})$, $e \mapsto |e|e$, is a distinguished degree-zero derivation. Obviously, the symbol of $\Delta_{\mathcal{E}}$ is Δ .

Example 5. Let \mathcal{M} be an \mathbb{N} -manifold, and let $\mathcal{E} = \mathcal{M} \times_M E$ be a degree-zero vector \mathbb{N} -bundle on it. Since $\Gamma(\mathcal{E})$ is generated in degree zero, then a negatively graded derivation \mathbb{X} of \mathcal{E} is completely determined by its symbol X and, therefore, it is the same as a negatively graded vector field on \mathcal{M} . Specifically, for a section of \mathcal{E} of the form $f \otimes e$, $f \in C^\infty(\mathcal{M})$, $e \in \Gamma(E)$, one has $\mathbb{X}(f \otimes e) = X(f) \otimes e$. In the following, we will tacitly identify negatively graded derivations of \mathcal{E} and negatively graded vector fields on \mathcal{M} .

Derivations of $\Gamma(\mathcal{E})$ are sections of a (graded) Lie algebroid $D\mathcal{E}$ over \mathcal{M} with bracket given by the (graded) commutator, and anchor given by the symbol. A *homological derivation of \mathcal{E}* is a degree-one derivation \mathbb{Q} , with symbol Q , such that $[\mathbb{Q}, \mathbb{Q}] = 0$ (in particular, Q is a homological vector field).

Example 6. Any degree-zero $\mathbb{N}\mathbb{Q}$ -vector bundle $(\mathcal{E}, \mathbb{Q})$ over a degree-one \mathbb{N} -manifold \mathcal{M} is of the form $(A[1] \times_M E, d_E)$ for some nongraded Lie algebroid $A \rightarrow M$ equipped with a representation $E \rightarrow M$. Here d_E is the homological derivation induced on $\Gamma(\wedge^* A^* \otimes E) = \Gamma(\mathcal{E})$. The algebroid structure on A corresponds to the symbol Q of \mathbb{Q} , while the (flat) A -connection ∇^E in E can be recovered from \mathbb{Q} via formula

$$\nabla_X^E e = [\mathbb{Q}, X](e) = \mathbb{Q}(X(e)),$$

where $X \in \Gamma(A)$ is also interpreted as a derivation of \mathcal{E} of degree -1 (see [Example 5](#)), and $e \in \Gamma(E)$.

2.2. Vector valued Cartan calculus on graded manifolds. Let \mathcal{M} be a graded manifold and let \mathcal{E} be a graded vector bundle over \mathcal{M} . Differential forms on \mathcal{M} are functions on $T[1]\mathcal{M}$ which are polynomial on fibers of $T[1]\mathcal{M} \rightarrow \mathcal{M}$. In particular, the algebra $\Omega(\mathcal{M})$ of differential forms on \mathcal{M} is equipped with two gradings: the “form” degree and the “internal, manifold” degree, which is usually referred simply as the *degree* (or, sometimes, the *weight*). The “total” degree is the sum of the form degree and the degree. Notice that the algebra $\Omega(\mathcal{M})$ is graded commutative with respect to the total degree. Similarly, \mathcal{E} -valued differential forms on \mathcal{M} are sections of the vector bundle $T[1]\mathcal{M} \times_{\mathcal{M}} \mathcal{E} \rightarrow T[1]\mathcal{M}$ which are polynomial on fibers of $T[1]\mathcal{M} \rightarrow \mathcal{M}$. The $\Omega(\mathcal{M})$ -module $\Omega(\mathcal{M}, \mathcal{E}) \simeq \Omega(\mathcal{M}) \otimes \Gamma(\mathcal{E})$ of \mathcal{E} -valued forms is equipped with two gradings. The “internal” degree will be referred to simply as the *degree*. We will denote by $|\omega|$ the degree of a homogeneous (with respect to the internal degree) \mathcal{E} -valued form ω .

Now, we briefly review the \mathcal{E} -valued version of Cartan calculus. Let \mathbb{X} be a derivation of \mathcal{E} . There are unique derivations $i_{\mathbb{X}}$ and $L_{\mathbb{X}}$ of the vector bundle $T[1]\mathcal{M} \times_{\mathcal{M}} \mathcal{E} \rightarrow T[1]\mathcal{M}$ such that

- (1) the symbol of $i_{\mathbb{X}}$ is the insertion i_X of the symbol X of \mathbb{X} ,
- (2) $i_{\mathbb{X}}$ vanishes on $\Gamma(\mathcal{E})$,
- (3) the symbol of $L_{\mathbb{X}}$ is the Lie derivative L_X along the symbol X of \mathbb{X} ,
- (4) $L_{\mathbb{X}}$ agrees with \mathbb{X} on $\Gamma(\mathcal{E})$.

Notice that, actually, $i_{\mathbb{X}}$ does only depend on the symbol of \mathbb{X} . For this reason, we will sometimes write i_X for $i_{\mathbb{X}}$.

Example 7. For any homogenous \mathcal{E} -valued form ω , $L_{\Delta_{\mathcal{E}}}\omega = |\omega|\omega$.

The following \mathcal{E} -valued Cartan identities hold:

$$(1) \quad [i_{\mathbb{X}}, i_{\mathbb{X}'}] = 0, \quad [L_{\mathbb{X}}, i_{\mathbb{X}'}] = i_{[\mathbb{X}, \mathbb{X}']}, \quad [L_{\mathbb{X}}, L_{\mathbb{X}'}] = L_{[\mathbb{X}, \mathbb{X}']},$$

for all $\mathbb{X}, \mathbb{X}' \in \Gamma(D\mathcal{E})$, where the bracket $[-, -]$ denotes the graded commutator. Moreover,

$$(2) \quad i_{f\mathbb{X}} = fi_{\mathbb{X}}, \quad L_{f\mathbb{X}} = fL_{\mathbb{X}} + (-)^{|f|+|\mathbb{X}|}df i_{\mathbb{X}},$$

for all $f \in C^\infty(\mathcal{M})$.

Now suppose that \mathcal{E} is equipped with a flat connection ∇ . Recall that a *connection* in \mathcal{E} is a graded, homogeneous, $C^\infty(\mathcal{M})$ -linear map $\nabla : \mathfrak{X}(\mathcal{M}) \rightarrow \Gamma(D\mathcal{E})$, denoted $X \mapsto \nabla_X$, such that the symbol of ∇_X is precisely X . In particular $|\nabla| = 0$. Derivation ∇_X is called the *covariant derivative along X*. A connection ∇ is *flat* if it is a

morphism of (graded) Lie algebras, i.e., $[\nabla_X, \nabla_Y] = \nabla_{[X, Y]}$, for all $X, Y \in \mathfrak{X}(\mathcal{M})$. A connection ∇ in \mathcal{E} determines a unique degree-one derivation d_∇ of the vector bundle $T[1]\mathcal{M} \times_{\mathcal{M}} \mathcal{E} \rightarrow T[1]\mathcal{M}$ such that

- (1) the symbol of d_∇ is the de Rham differential $d \in \mathfrak{X}(T[1]\mathcal{M})$,
- (2) $i_X d_\nabla e = \nabla_X e$ for all $e \in \Gamma(\mathcal{E})$ and $X \in \mathfrak{X}(\mathcal{M})$.

The derivation d_∇ is the *de Rham differential* of ∇ . It is a homological derivation if and only if ∇ is flat.

Let ∇ be a flat connection in \mathcal{E} . The following identities hold:

$$(3) \quad [i_{\nabla_X}, d_\nabla] = L_{\nabla_X}, \quad [L_{\nabla_X}, d_\nabla] = 0, \quad [d_\nabla, d_\nabla] = 0,$$

for all $X \in \mathfrak{X}(\mathcal{M})$.

Remark 8. Specialize to the case when \mathcal{M} is an \mathbb{N} -manifold and \mathcal{E} is of degree zero. Then $\mathcal{E} = \mathcal{M} \times_M E$ for some vector bundle E over the degree-zero shadow M of \mathcal{M} . A connection ∇^0 in E induces a unique connection ∇ in \mathcal{E} such that

$$\nabla_X e = \nabla_{\underline{X}}^0 e,$$

for all $e \in \Gamma(E)$ and $X \in \mathfrak{X}_0(\mathcal{M})$, where $\underline{X} \in \mathfrak{X}(M)$ is the projection of X onto M . The connection ∇ is flat if and only if ∇^0 is flat. Moreover, every connection in \mathcal{E} is of this kind. Notice that if ∇ is flat, whatever it is, the covariant derivative along the grading vector field Δ coincides with the grading derivation $\Delta_\mathcal{E}$. To see this it is enough to compare the action of ∇_Δ and $\Delta_\mathcal{E}$ on generators. Locally, $\Gamma(\mathcal{E})$ is generated by ∇^0 -flat sections of E . Thus, let $e \in \Gamma(E)$ be ∇^0 -flat. Then $\nabla_\Delta e = 0 = \Delta_\mathcal{E} e$. As an immediate consequence, every d_∇ -closed E -valued differential form ω on \mathcal{M} of positive degree n is also d_∇ -exact, i.e., $\omega = d_\nabla \vartheta$, for a suitable ϑ . One can choose, for instance, $\vartheta = n^{-1} i_{\Delta_\mathcal{E}} \omega$. Indeed,

$$d_\nabla \left(\frac{1}{n} i_{\Delta_\mathcal{E}} \omega \right) = \frac{1}{n} [d_\nabla, i_{\Delta_\mathcal{E}}] \omega = \frac{1}{n} L_{\Delta_\mathcal{E}} \omega = \omega.$$

2.3. An alternative description of vector valued forms on \mathbb{N} -manifolds. In the following, we will only consider the case when \mathcal{M} is an \mathbb{N} -manifold and \mathcal{E} is generated in one single degree. Let M be the degree-zero shadow of \mathcal{M} . Then, up to an irrelevant shift, \mathcal{E} is isomorphic to a pull-back $\mathcal{M} \times_M E$, where E is a nongraded vector bundle over M (see [Example 2](#)). Accordingly, we will often write $C^\infty(\mathcal{M}, E)$ for $\Gamma(\mathcal{E})$ and $\Omega(\mathcal{M}, E)$ for $\Omega(\mathcal{M}, \mathcal{E})$.

Remark 9. Despite the huge simplifications inherent to the hypothesis $\mathcal{E} \simeq \mathcal{M} \times_M E$, this case still captures many interesting situations. For instance, a degree- n symplectic \mathbb{N} -manifold [[Roytenberg 2002](#)] or contact \mathbb{N} -manifold [[Grabowski 2013](#); [Mehta 2013](#)] can each be understood as an \mathbb{N} -manifold \mathcal{M} equipped with a degree- n differential form with values in a vector bundle concentrated in just one degree

(the trivial bundle $\mathcal{M} \times \mathbb{R}$ in the symplectic case, and a generically nontrivial line bundle concentrated in degree n in the contact case). We hope to discuss the case of a general vector bundle \mathcal{E} elsewhere.

Theorem 10. *Let n be a positive integer. A degree- n differential k -form on \mathcal{M} with values in E is equivalent to the following data:*

- a degree- n (first order) differential operator $D : \mathfrak{X}_-(\mathcal{M}) \rightarrow \Omega^k(\mathcal{M}, E)$, and
- a degree- n $C^\infty(M)$ -linear map $\ell : \mathfrak{X}_-(\mathcal{M}) \rightarrow \Omega^{k-1}(\mathcal{M}, E)$,

such that

$$(4) \quad D(fX) = fD(X) + (-)^{|X|}df \ell(X),$$

and, moreover,

$$(5) \quad L_X D(Y) - (-)^{|X||Y|} L_Y D(X) = D([X, Y]),$$

$$(6) \quad L_X \ell(Y) - (-)^{|X|(|Y|-1)} i_Y D(X) = \ell([X, Y]),$$

$$(7) \quad i_X \ell(Y) - (-)^{(|X|-1)(|Y|-1)} i_Y \ell(X) = 0.$$

for all $X, Y \in \mathfrak{X}_-(\mathcal{M})$, and $f \in C^\infty(M)$.

Remark 11. By induction on n , **Theorem 10** provides a description of E -valued differential forms in terms of nongraded data. Indeed, D and ℓ take values in lower-degree forms and one can use degree-zero forms, namely E -valued forms on M as base of induction.

Proof. Let ω be a degree- n , E -valued differential k -form on \mathcal{M} . Define $D : \mathfrak{X}_-(\mathcal{M}) \rightarrow \Omega^k(\mathcal{M}, E)$ and $\ell : \mathfrak{X}_-(\mathcal{M}) \rightarrow \Omega^{k-1}(\mathcal{M}, E)$ by putting

$$(8) \quad D(X) := L_X \omega, \quad \text{and} \quad \ell(X) := i_X \omega, \quad X \in \mathfrak{X}_-(\mathcal{M}).$$

Properties (4), (5), (6), and (7) immediately follow from identities (1), and (2).

Conversely, let D and ℓ be as in the statement of the theorem, and prove that there exists a unique degree- n differential form $\omega \in \Omega^k(\mathcal{M}, E)$ fulfilling (8). We propose a local proof. One can pass to the global setting by partition of unity arguments. Let \dots, z^a, \dots be positively graded coordinates on \mathcal{M} and $\partial_a := \partial/\partial z^a$. In particular the grading derivation $\Delta_{\mathcal{E}}$ is locally given by

$$\Delta_{\mathcal{E}} = |z^a| z^a \partial_a.$$

Moreover, $\mathfrak{X}_-(\mathcal{M})$ is locally generated, as a $C^\infty(M)$ -module, by vector fields

$$z^{b_1} \dots z^{b_k} \partial_b, \quad |z^{b_1}| + \dots + |z^{b_k}| - |z^b| < 0.$$

Put

$$\omega := \frac{|z^a|}{n} (z^a D(\partial_a) + dz^a \ell(\partial_a))$$

and prove (8). First of all, for $X = X^a \partial_a$,

$$\begin{aligned} L_X \omega &= \frac{|z^a|}{n} L_X (z^a D(\partial_a) + dz^a \ell(\partial_a)) \\ &= \frac{|z^a|}{n} (X^a D(\partial_a) + (-)^{|z^a||X|} L_X D(\partial_a) + (-)^{|X|} dX^a \ell(\partial_a) + (-)^{|X|(|z^a|+1)} dz^a L_X \ell(\partial_a)). \end{aligned}$$

In view of (5) and (6),

$$(-)^{|z^a||X|} L_X D(\partial_a) = L_{\partial_a} D(X) - D([\partial_a, X])$$

and

$$(-)^{|X|(|z^a|+1)} L_X \ell(\partial_a) = i_{\partial_a} D(X) - (-)^{|X|} \ell([\partial_a, X]),$$

so that

$$\begin{aligned} (9) \quad L_X \omega &= \frac{1}{n} L_{\Delta_\varepsilon} D(X) + \frac{|z^a|}{n} (X^a D(\partial_a) + (-)^{|X|} dX^a \ell(\partial_a) \\ &\quad - z^a D([\partial_a, X]) - (-)^{|X|} dz^a \ell([\partial_a, X])) \\ &= \frac{n+|X|}{n} D(X) + \frac{|z^a|}{n} (X^a D(\partial_a) + (-)^{|X|} dX^a \ell(\partial_a) \\ &\quad - z^a D([\partial_a, X]) - (-)^{|X|} dz^a \ell([\partial_a, X])). \end{aligned}$$

Similarly

$$(10) \quad i_X \omega = \frac{n+|X|}{n} \ell(X) + \frac{|z^a|}{n} (X^a \ell(\partial_a) - z^a \ell([\partial_a, X])).$$

Using $X = \partial_b$ in (9) and (10) gives

$$(11) \quad L_{\partial_b} \omega = D(\partial_b), \quad \text{and} \quad i_{\partial_b} \omega = \ell(\partial_b).$$

In view of identity (4), in order to prove (8), it is enough to restrict to vector fields X of the form $z^{b_1} \cdots z^{b_k} \partial_b$. This case can be treated by induction on k , using (11) as the base. Namely, use $X = z^{b_1} \cdots z^{b_k} \partial_b$ in (9), with $k > 0$. Since

$$[\partial_a, X] = \sum_i (-)^{(|z^{b_1}|+\cdots+|z^{b_i-1}|)|z^a|} \delta_a^{b_i} z^{b_1} \cdots \widehat{z^{b_i}} \cdots z^{b_k} \partial_b,$$

where a hat “ $\widehat{}$ ” denotes omission, by the induction hypothesis we have

$$D([\partial_a, X]) = L_{[\partial_a, X]} \omega, \quad \text{and} \quad \ell([\partial_a, X]) = i_{[\partial_a, X]} \omega.$$

A direct computation shows that the second summand in the right-hand side of (9) is equal to $-(|X|/n)L_X \omega$. Similarly, the second summand in the right-hand side of (10) is equal to $-(|X|/n)i_X \omega$. Notice that, since $k > 0$, we have $|X| > -n$ and can conclude that $L_X \omega = D(X)$, and similarly, $i_X \omega = \ell(X)$.

To prove uniqueness, it is enough to show that a degree- n differential form ω with values in E is completely determined by contraction with, and Lie derivative

along, negatively graded derivations. Thus,

$$n\omega = L_{\Delta_\varepsilon}\omega = |z^a|(z^a L_{\partial_a}\omega + dz^a i_{\partial_a}\omega).$$

In particular ω is completely determined by $L_{\partial_a}\omega$ and $i_{\partial_a}\omega$. □

We will refer to the data (D, ℓ) corresponding to a vector valued form ω as the *Spencer data* of ω . Indeed, as we will show in [Section 5](#), they are a vast generalization of the *Spencer operators* considered in [\[Crainic et al. 2015; Salazar 2013\]](#).

Example 12. Let $E \rightarrow M$ be a nongraded vector bundle equipped with a flat connection ∇ , and let \mathcal{M} be an \mathbb{N} -manifold. As discussed in [Section 2.2](#), ∇ induces a flat connection in the graded vector bundle $\mathcal{M} \times_M E \rightarrow \mathcal{M}$ which we denote again by ∇ . In its turn, the induced connection determines a homological derivation d_∇ of the vector bundle $T[1]\mathcal{M} \times_M E \rightarrow T[1]\mathcal{M}$ of E -valued forms on \mathcal{M} . Notice that d_∇ maps k -forms to $(k + 1)$ -forms. Now, let $\omega \in \Omega^k(\mathcal{M}, E)$ and let (D, ℓ) be the corresponding Spencer data. We want to describe the Spencer data (D', ℓ') of $d_\nabla\omega$. To do this, we first observe that a discussion similar to that in [Remark 8](#) shows that, whatever ∇ , the covariant derivative along a negatively graded vector field $X \in \mathfrak{X}_-(\mathcal{M})$ satisfies $\nabla_X = X$. Hence, from [\(3\)](#)

$$D'(X) = L_X d_\nabla\omega = L_{\nabla_X} d_\nabla\omega = d_\nabla L_X\omega = d_\nabla D(X)$$

and

$$\ell'(X) = i_X d_\nabla\omega = i_{\nabla_X} d_\nabla\omega = L_X\omega - (-)^{|X|} d_\nabla i_X\omega = D(X) - (-)^{|X|} d_\nabla\ell(X),$$

which completely describe (D', ℓ') in terms of (D, ℓ) and d_∇ .

Example 13. Let $E \rightarrow M$ be a nongraded vector bundle. The first jet bundle $J^1 E \rightarrow M$ fits in an exact sequence

$$(12) \quad 0 \longrightarrow \Omega^1(M, E) \longrightarrow \Gamma(J^1 E) \xrightarrow{p} \Gamma(E) \longrightarrow 0$$

of $C^\infty(M)$ -linear maps, where p is the canonical projection. Sequence [\(12\)](#) splits (beware, *over* \mathbb{R} not over $C^\infty(M)$) via the universal first order differential operator $j^1 : \Gamma(E) \rightarrow \Gamma(J^1 E)$. Accordingly, there is a first order differential operator $S : \Gamma(J^1 E) \rightarrow \Omega^1(M, E)$ sometimes called the *Spencer operator*. The degree- n \mathbb{N} -manifold $\mathcal{M} = J^1 E[n]$ comes equipped with an E -valued, degree- n *Cartan 1-form* θ . In order to define θ , recall that negatively graded vector fields on \mathcal{M} are concentrated in degree $-n$, and $\mathfrak{X}_{-n}(\mathcal{M})$ identifies with $\Gamma(J^1 E)$ as a $C^\infty(M)$ -module. Now, θ is uniquely defined by the properties

$$(13) \quad i_{j^1 e}\theta = e \quad \text{and} \quad L_{j^1 e}\theta = 0,$$

for all $e \in \Gamma(E)$. It immediately follows from (13) that the Spencer data (D, ℓ) of θ identify with $(-)^n$ times the Spencer operator $S : \Gamma(J^1 E) \rightarrow \Omega^1(M, E)$ and the projection $\Gamma(J^1 E) \rightarrow \Gamma(E)$ respectively.

Example 14. Let M be a nongraded manifold. The degree- n \mathbb{N} -manifold $\mathcal{M} = T^*[n]M$ comes equipped with the obvious tautological, degree- n 1-form ϑ . Consider the degree- n 2-form $\omega = d\vartheta$. Negatively graded vector fields on \mathcal{M} are concentrated in degree $-n$, and $\mathfrak{X}_{-n}(\mathcal{M})$ is naturally isomorphic to $\Omega^1(M)$ as a $C^\infty(M)$ -module. It is easy to see that ω is uniquely defined by the properties

$$(14) \quad i_{df}\omega = df \quad \text{and} \quad L_{df}\omega = 0,$$

for all $f \in C^\infty(M)$. It immediately follows from (14) that the Spencer data (D, ℓ) of ω identify with $(-)^n$ times the exterior differential $d : \Omega^1(M) \rightarrow \Omega^2(M)$ and the identity $\text{id} : \Omega^1(M) \rightarrow \Omega^1(M)$ respectively.

Example 15. Let $E \rightarrow M$ be a nongraded vector bundle equipped with a flat connection ∇ . The degree- n \mathbb{N} -manifold $\mathcal{M} = T^*[n]M \otimes E$ is equipped with a tautological, degree- n E -valued 1-form ϑ . The flat connection ∇ induces a flat connection in the graded vector bundle $\mathcal{M} \times_M E \rightarrow \mathcal{M}$ which we denote again by ∇ . Consider the homological derivation d_∇ as in Example 12. Notice that d_∇ agrees with the de Rham differential of ∇ on degree-zero forms, i.e., elements of $\Omega(M, E)$. Consider the degree- n 2-form $\omega = d_\nabla\vartheta$ with values in E . Negatively graded vector fields on \mathcal{M} are concentrated in degree $-n$, and $\mathfrak{X}_{-n}(\mathcal{M})$ is isomorphic to $\Omega^1(M, E)$ as a $C^\infty(M)$ -module. It is easy to see that ω is uniquely defined by the properties

$$(15) \quad i_{d_\nabla e}\omega = d_\nabla e \quad \text{and} \quad L_{d_\nabla e}\omega = 0,$$

for all $e \in \Gamma(E)$. It immediately follows from (15) that the Spencer data (D, ℓ) of ω identify with $(-)^n$ times the de Rham differential $d_\nabla : \Omega^1(M, E) \rightarrow \Omega^2(M, E)$ and the identity $\text{id} : \Omega^1(M, E) \rightarrow \Omega^1(M, E)$ respectively.

In the three remaining sections we use Theorem 10 (and Proposition 17 below) to describe degree-one $\mathbb{N}Q$ -manifolds equipped with a compatible vector valued differential form (see below) in terms of nongraded data. In particular, we manage to give alternative proofs of known results about compatible contact structures [Grabowski 2013; Mehta 2013], involutive distributions [Zambon and Zhu 2012], and symplectic forms [Roytenberg 2002] on degree-one $\mathbb{N}Q$ -manifolds. We also manage to find new results about compatible, presymplectic and locally conformal symplectic structures, and, more generally, higher order vector valued forms on degree-one $\mathbb{N}Q$ -manifolds. It turns out (Theorem 36) that a compatible degree-one differential k -form on a degree-one $\mathbb{N}Q$ -manifold (\mathcal{M}, Q) is equivalent to a Lie algebroid equipped with a structure recently identified in [Crainic et al. 2015] as the

infinitesimal counterpart of a multiplicative vector valued form on a Lie groupoid (see also [Salazar 2013]), namely, a k -th order Spencer operator.

Let \mathcal{M} be an \mathbb{N} -manifold, with degree-zero shadow M , and let $(\mathcal{E}, \mathbb{Q})$ be an $\mathbb{N}\mathbb{Q}$ -vector bundle over it. We denote by Q the symbol of \mathbb{Q} .

Definition 16. An \mathcal{E} -valued differential form on \mathcal{M} , ω , is *compatible with \mathbb{Q}* if $L_{\mathbb{Q}}\omega = 0$.

Suppose \mathcal{E} is of degree zero. Then $\mathcal{E} = \mathcal{M} \times_M E$ for a nongraded vector bundle $E \rightarrow M$. For later use, we conclude this section expressing the compatibility of an E -valued form ω on \mathcal{M} with \mathbb{Q} in terms of Spencer data.

Proposition 17. Let $\omega \in \Omega^k(\mathcal{M}, E)$ be an E -valued k -form on \mathcal{M} of degree $n > 0$, and let (D, ℓ) be its Spencer data. Then ω is compatible with \mathbb{Q} , i.e., $L_{\mathbb{Q}}\omega = 0$, if and only if

$$(16) \quad A(X, Y) := D([[Q, X], Y]) - L_{[Q, X]}D(Y) - (-)^{|X||Y|}(L_{[Q, Y]}D(X) - L_{\mathbb{Q}}L_YD(X)) = 0,$$

$$(17) \quad B(X, Y) := \ell([[Q, X], Y]) - (-)^{|Y|}i_{[Q, X]}D(Y) - (-)^{|X||Y|}(L_{[Q, Y]}\ell(X) - L_{\mathbb{Q}}L_Y\ell(X)) = 0,$$

$$(18) \quad C(X, Y) := i_{[Q, X]}\ell(Y) + (-)^{(|X|-1)(|Y|-1)}(i_{[Q, Y]}\ell(X) - L_{\mathbb{Q}}i_Y\ell(X)) = 0.$$

Remark 18. By induction on n , Proposition 17 provides a description of the compatibility condition between ω and \mathbb{Q} in terms of nongraded data (see also Remark 19 below). Notice that, when $|\omega| = 1$, the last summand in (16), (17) and (18) vanishes by degree reasons.

Proof. First of all, notice that $[[Q, X], Y]$ is negatively graded for all X, Y . Hence it identifies with $[[Q, X], Y]$. In particular, the left-hand side of (16), (17) and (18) are well-defined. Now, for any ω as in the statement, $L_{\mathbb{Q}}\omega$ is a form of degree $n + 1$. Since every form of positive degree on \mathcal{M} is completely determined by its Spencer data, $L_{\mathbb{Q}}\omega$ vanishes if and only if

$$L_YL_XL_{\mathbb{Q}}\omega = i_YL_XL_{\mathbb{Q}}\omega = L_Yi_XL_{\mathbb{Q}}\omega = i_Yi_XL_{\mathbb{Q}}\omega = 0,$$

for all $X, Y \in \mathfrak{X}_-(\mathcal{M})$. It immediately follows from the second Cartan identity (1) that condition $i_YL_XL_{\mathbb{Q}}\omega = 0$ is actually redundant. It remains to compute $L_YL_XL_{\mathbb{Q}}\omega$, $L_Yi_XL_{\mathbb{Q}}\omega$, and $i_Yi_XL_{\mathbb{Q}}\omega$. So

$$\begin{aligned} L_YL_XL_{\mathbb{Q}}\omega &= L_YL_{[X, \mathbb{Q}]} \omega - (-)^{|Y|}L_YL_{\mathbb{Q}}L_X\omega \\ &= (-)^{|X|+|Y|(|X|+1)}(L_{[[Q, X], Y]}\omega - L_{[Q, X]}L_Y\omega) \\ &\quad - (-)^{|X|+|Y|}(L_{[Q, Y]}L_X\omega - L_{\mathbb{Q}}L_YL_X\omega), \end{aligned}$$

which differs from $A(X, Y)$ in (16) by an overall sign $(-)^{|X|+|Y|(|X|+1)}$. Similarly,

$$\begin{aligned} L_Y i_X L_{\mathbb{Q}} \omega &= (-)^{|X|} L_Y i_{[\mathbb{Q}, X]} \omega - (-)^{|X|} L_Y L_{\mathbb{Q}} i_X \omega \\ &= (-)^{(|X|+1)(|Y|+1)} (i_{[[\mathbb{Q}, X], Y]} \omega - (-)^{|Y|} i_{[\mathbb{Q}, X]} L_Y \omega) \\ &\quad + (-)^{|X|+|Y|} (L_{[\mathbb{Q}, Y]} i_X \omega - L_{\mathbb{Q}} L_Y i_X \omega), \end{aligned}$$

which differs from $B(X, Y)$ in (17) by an overall sign $(-)^{(|X|+1)(|Y|+1)}$. Finally,

$$\begin{aligned} i_Y i_X L_{\mathbb{Q}} \omega &= (-)^{|X|} i_Y i_{[\mathbb{Q}, X]} \omega - (-)^{|X|} i_Y L_{\mathbb{Q}} i_X \omega \\ &= (-)^{|X||Y|} i_{[\mathbb{Q}, X]} i_Y \omega - (-)^{|X|+|Y|} (i_{[\mathbb{Q}, Y]} i_X \omega - L_{\mathbb{Q}} i_Y i_X \omega), \end{aligned}$$

which differs from $C(X, Y)$ in (18) by an overall sign $(-)^{|X||Y|}$. □

Remark 19. When \mathcal{M} is the total space of a negatively graded vector bundle $V \rightarrow M$ (which is always the case up to a noncanonical isomorphism), a homological vector field on \mathcal{M} is the same as an L_{∞} -algebroid structure on $\Gamma(V^*)$ (see, e.g., [Bonavolontà and Poncin 2013; Bruce 2011; Sati et al. 2012; Vitagliano 2015b]). We conjecture the existence of formulas expressing the compatibility between ω and \mathbb{Q} in terms of the higher brackets (and the anchor) of this L_{∞} -algebroid, and the Spencer data of ω . Similarly, when no isomorphism $\mathcal{M} \simeq V$ is assigned, there should be formulas involving *Getzler higher derived brackets* on $\mathfrak{X}_-(\mathcal{M})$ [Getzler 2010]. Finding these formulas goes beyond the scopes of this paper and we postpone this task to a subsequent publication.

3. Vector valued 1-forms on $\mathbb{N}\mathcal{Q}$ -manifolds

3.1. Vector valued 1-forms and distributions. Let \mathcal{M} be a degree- n \mathbb{N} -manifold, $n > 0$, and let $(\mathcal{E} = \mathcal{M} \times_M E, \mathbb{Q})$ be a degree-zero $\mathbb{N}\mathcal{Q}$ -vector bundle over it. According to Definition 16, a degree- n 1-form θ with values in E is *compatible* with \mathbb{Q} if, by definition, $L_{\mathbb{Q}} \theta = 0$. Several interesting geometric structures are described by compatible 1-forms. For instance, compatible distributions on an $\mathbb{N}\mathcal{Q}$ -manifold are equivalent to surjective compatible 1-forms. Namely, Let $(\mathcal{M}, \mathcal{Q})$ be a degree- n $\mathbb{N}\mathcal{Q}$ -manifold, and let $\mathcal{D} \subset T\mathcal{M}$ be a distribution on \mathcal{M} . Consider the normal bundle $T\mathcal{M}/\mathcal{D}$. Projection $T\mathcal{M} \rightarrow T\mathcal{M}/\mathcal{D}$ can be interpreted as a degree-zero surjective 1-form with values in $T\mathcal{M}/\mathcal{D}$. We say that \mathcal{D} is *cogenerated* in degree k if $T\mathcal{M}/\mathcal{D}$ is generated in degree $-k$. In this case, $T\mathcal{M}/\mathcal{D} = \mathcal{M} \times_M E[k]$ for a suitable nongraded vector bundle $E \rightarrow M$, and the projection $\theta_{\mathcal{D}} : T\mathcal{M} \rightarrow \mathcal{M} \times_M E$ can be interpreted as a degree- k , surjective, E -valued 1-form such that $\ker \theta_{\mathcal{D}} = \mathcal{D}$. Conversely, if $E \rightarrow M$ is a nongraded vector bundle and θ is a degree- k , surjective 1-form with values in E , then $\mathcal{D} := \ker \theta$ is a distribution such that $\theta_{\mathcal{D}} = \theta$.

Definition 20. A distribution \mathcal{D} on \mathcal{M} , cogenerated in degree n , is *compatible* with \mathcal{Q} if $[\mathcal{Q}, \Gamma(\mathcal{D})] \subset \Gamma(\mathcal{D})$.

Now, let \mathcal{D} be a distribution cogenerated in degree- n and E be such that $T\mathcal{M}/\mathcal{D} = \mathcal{M} \times_M E[-n]$. If \mathcal{D} is compatible with Q , then the commutator with Q restricts to a homological derivation of \mathcal{D} , hence it also descends to a homological derivation of the vector bundle $\mathcal{M} \times_M E$ which we denote by \mathbb{Q} . Moreover, $L_{\mathbb{Q}}\theta_{\mathcal{D}} = 0$. Indeed, for every vector field $X \in \mathfrak{X}(\mathcal{M})$,

$$(19) \quad i_X L_{\mathbb{Q}}\theta_{\mathcal{D}} = (-)^{|X|} (i_{[Q, X]}\theta_{\mathcal{D}} - \mathbb{Q}(i_X\theta_{\mathcal{D}})) = 0.$$

Conversely, if $(\mathcal{E}, \mathbb{Q})$ is a degree-zero $\mathbb{N}Q$ -bundle and θ is a degree- n surjective 1-form with values in E , then it follows from (19) that $L_{\mathbb{Q}}\theta = 0$ if and only if 1) $\ker \theta$ is a distribution compatible with the symbol Q of \mathbb{Q} , and 2) \mathbb{Q} is induced on $\Gamma(\mathcal{E})$ by the adjoint operator $[Q, -]$ on $\mathfrak{X}(\mathcal{M})$. One concludes that compatible distributions are the same as compatible surjective 1-forms.

Remark 21. The above discussion is actually independent of the degree of Q . Hence, it shows that an infinitesimal symmetry of \mathcal{D} , i.e., any vector field X such that $[X, \Gamma(\mathcal{D})] \subset \Gamma(\mathcal{D})$, determines a derivation \mathbb{X} of $T\mathcal{M}/\mathcal{D}$ via

$$\mathbb{X}(Y \bmod \mathcal{D}) := [X, Y] \bmod \mathcal{D}.$$

The symbol of \mathbb{X} is precisely X . Moreover, one can compute the Lie derivative $L_{\mathbb{X}}\theta_{\mathcal{D}}$, and find $L_{\mathbb{X}}\theta_{\mathcal{D}} = 0$.

Now recall that a distribution \mathcal{D} on a (graded) manifold \mathcal{M} comes equipped with a *curvature form*

$$\omega_{\mathcal{D}} : \Gamma(\mathcal{D}) \times \Gamma(\mathcal{D}) \rightarrow \Gamma(T\mathcal{M}/\mathcal{D}), \quad (X, Y) \mapsto [X, Y] \bmod \Gamma(\mathcal{D}).$$

The curvature form $\omega_{\mathcal{D}}$ measures how far is \mathcal{D} from being involutive. The two limit cases $\omega_{\mathcal{D}}$ nondegenerate and $\omega_{\mathcal{D}} = 0$ are of special interest. The first one corresponds to *maximally nonintegrable distributions*, the second one to *involutive distributions*.

3.2. Degree one contact $\mathbb{N}Q$ -manifolds. Let \mathcal{M} be an \mathbb{N} -manifold and let C be a hyperplane distribution on it. Since $\mathcal{L} := T\mathcal{M}/C$ is a line bundle, it is generated in one single degree. A *degree- n contact \mathbb{N} -manifold* is an \mathbb{N} -manifold \mathcal{M} equipped with a *degree- n contact structure*, i.e., a hyperplane distribution C , such that the line bundle $\mathcal{L} := T\mathcal{M}/C$ is generated in degree n , and the curvature form ω_C is nondegenerate (see [Grabowski 2013] for an alternative definition exploiting the “symplectization trick”).

Example 22. Let $L \rightarrow M$ be a nongraded line bundle. The kernel of the Cartan form θ on $J^1L[n]$ (see Example 13) is a degree- n contact structure.

It follows from the definition that, if (\mathcal{M}, C) is a degree- n contact \mathbb{N} -manifold, then the degree of \mathcal{M} is at most n . When \mathcal{L} is a trivial line bundle, C is the

kernel of a (nowhere vanishing) 1-form α which can be used to simplify the theory significantly (see [Mehta 2013]). In this case the contact structure is said to be *coorientable* and a choice of α provides a coorientation (i.e., an orientation of \mathcal{L}). In the general case, $\mathcal{L} := \mathcal{M} \times_M L[n]$ for a nongraded line bundle $L \rightarrow M$, and C is the kernel of a (degree- n) 1-form θ_C with values in a (generically nontrivial) line bundle L .

A degree- n contact structure on \mathcal{M} determines a nondegenerate Jacobi bracket $\{-, -\}$ of degree $-n$ on $\Gamma(\mathcal{L})$, i.e., a Lie bracket of degree $-n$ which is a graded first order differential operator in each entry and such that the associated morphism $J^1\mathcal{L} \otimes J^1\mathcal{L} \rightarrow \mathcal{L}$ is nondegenerate (see also Appendix A). For the details about how to define the Jacobi bracket $\{-, -\}$ from C in the nongraded case see, for instance, [Crainic and Salazar 2015]. The generalization to the graded case can be carried out straightforwardly and the obvious details are left to the reader. A *degree- n contact $\mathbb{N}Q$ -manifold* is a degree- n contact manifold (\mathcal{M}, C) equipped with a homological vector field Q such that $[Q, \Gamma(C)] \subset \Gamma(C)$, in other words it is an $\mathbb{N}Q$ -manifold equipped with a compatible degree- n contact structure. If (\mathcal{M}, C, Q) is a contact $\mathbb{N}Q$ -manifold, the homological vector field Q induces a homological derivation \mathbb{Q} of \mathcal{L} as discussed above. Thus, equivalently, a degree- n contact $\mathbb{N}Q$ -manifold is a degree- n contact manifold (\mathcal{M}, C) equipped with a homological derivation \mathbb{Q} of \mathcal{L} such that $L_{\mathbb{Q}}\theta_C = 0$.

Theorem 23 [Mehta 2013; Grabowski 2013]. *Every degree-one contact \mathbb{N} -manifold (\mathcal{M}, C) is of the kind $(J^1L[1], \ker \theta)$, up to contactomorphisms, where $L \rightarrow M$ is a (nongraded) line bundle, and θ is the Cartan form on $J^1L[1]$. Moreover, there is a one-to-one correspondence between degree-one contact $\mathbb{N}Q$ -manifolds and (nongraded) manifolds equipped with an abstract Jacobi structure (see the appendixes).*

Notice that Mehta does only discuss the case when C is coorientable, i.e., $T\mathcal{M}/C$ is globally trivial. Moreover, he selects a contact form, which amounts to selecting a global trivialization $T\mathcal{M}/C \simeq \mathcal{M} \times \mathbb{R}[1]$ (see [Mehta 2013] for details). On the other hand, in independent work Grabowski discusses the general case (he actually treats the degree-two case as well). His proof relies on the “symplectization trick” which consists in understanding a contact manifold as a homogeneous symplectic manifold (see [Grabowski 2013]) and then using already known results in the symplectic case. We propose an alternative proof avoiding the “symplectization trick” and focusing on the Spencer data of the structure 1-form of C . We refer to [Crainic and Salazar 2015] for details on abstract Jacobi structures.

Proof. Let $(\mathcal{M} = A[1], C)$ be a degree-one contact \mathbb{N} -manifold, and let $\mathcal{L} = T\mathcal{M}/C$ be the associated degree-one line-bundle. Then $\mathcal{L} = \mathcal{M} \times_M L[1]$ for a nongraded line bundle $L \rightarrow M$, and θ_C is a degree-one L -valued 1-form on \mathcal{M} . Denote by (D, ℓ)

the Spencer data of θ_C . The Jacobi bracket $\{-, -\}$ determines a isomorphism of graded vector bundles of degree -1 between $J^1\mathcal{L}$ and $D\mathcal{L}$. Since negatively graded derivations are completely determined by their symbol, this gives an isomorphism $\Gamma(J^1L) \simeq \mathfrak{X}_{-1}(\mathcal{M}) \simeq \Gamma(A)$, hence a diffeomorphism $\mathcal{M} \simeq J^1L[1]$. The diagram

$$\begin{CD} \Gamma(L) @<\ell<< \Gamma(A) @>D>> \Omega^1(M, L) \\ @| @VV\wr V @| \\ \Gamma(L) @<p<< \Gamma(J^1L) @>-S>> \Omega^1(M, L) \end{CD}$$

commutes, as can be easily seen. This shows that the diffeomorphism $\mathcal{M} \simeq J^1L[1]$ identifies θ_C with the Cartan form θ (see [Example 13](#)), thus proving the first part of the statement. In the following we identify \mathcal{M} and $J^1L[1]$. For the second part of the statement, let \mathbb{Q} be a homological derivation of \mathcal{L} and let Q be its symbol. Moreover, let $(J^1L, \rho, \llbracket -, - \rrbracket)$ and (L, ∇^L) be the Lie algebroid and the Lie algebroid representation associated to \mathbb{Q} . We use [Proposition 17](#) to see when is $(\mathcal{M}, C, \mathbb{Q})$ a contact $\mathbb{N}\mathbb{Q}$ -manifold. Since θ is a 1-form, [Equation \(18\)](#) is automatically satisfied, and θ is compatible with \mathbb{Q} if and only if $A(X, Y) = B(X, Y) = 0$, with $\omega = \theta$ and $X, Y \in \mathfrak{X}_{-1}(\mathcal{M}) \simeq \Gamma(J^1L)$. In fact, one can even restrict to X, Y in the form $j^1\lambda, j^1\mu$, with $\lambda, \mu \in \Gamma(L)$. In this case, one gets

$$A(X, Y) = D(\llbracket Q, j^1\lambda \rrbracket, j^1\mu) = -S\llbracket j^1\lambda, j^1\mu \rrbracket,$$

and

$$B(X, Y) = \ell(\llbracket Q, j^1\lambda \rrbracket, j^1\mu) + L_{[Q, j^1\mu]}\lambda = p\llbracket j^1\lambda, j^1\mu \rrbracket + \nabla_{j^1\mu}^L \lambda,$$

where we used that $\ell(j^1\lambda) = i_{j^1\lambda}\theta = \lambda$, and $D(j^1\lambda) = L_{j^1\lambda}\theta = 0$ (see [Example 13](#)). Concluding, $(\mathcal{M}, C, \mathbb{Q})$ is a contact $\mathbb{N}\mathbb{Q}$ -manifold if and only if $p\llbracket j^1\lambda, j^1\mu \rrbracket = -\nabla_{j^1\mu}^L \lambda = \nabla_{j^1\lambda}^L \mu$ and $S\llbracket j^1\lambda, j^1\mu \rrbracket = 0$, i.e., if and only if $(J^1L, \llbracket -, - \rrbracket, \rho)$ is the Lie algebroid associated to a Jacobi structure on $L \rightarrow M$, and ∇^L is its natural representation (see [Appendix B](#)). □

3.3. Involutive distributions on degree-one $\mathbb{N}\mathbb{Q}$ -manifolds. Compatible involutive distributions (cogenerated in degree one) on a degree-one $\mathbb{N}\mathbb{Q}$ -manifold are equivalent to infinitesimally multiplicative (IM) foliations of a special kind. Let $(A, \llbracket -, - \rrbracket, \rho)$ be a Lie algebroid over a manifold M , and let $F \subset TM$ be an involutive distribution. An *IM foliation of A over F* [[Jotz and Ortiz 2011](#)] is a triple consisting of

- an involutive distribution F ,
- a Lie subalgebroid $B \subset A$,
- a flat F -connection ∇ in the quotient bundle A/B ,

such that

- (1) sections X of A such that $X \bmod B$ is ∇ -flat form a Lie subalgebra in $\Gamma(A)$ with sections of B as a Lie ideal,
- (2) ρ takes values in the stabilizer of F ,
- (3) $\rho|_B$ takes values in F .

As the terminology suggests, IM foliations are infinitesimal counterparts of involutive multiplicative distributions on Lie groupoids [Jotz and Ortiz 2011]. Zambon and Zhu [2012] proved that IM foliations can be also understood as degree-one $\mathbb{N}\mathcal{Q}$ -manifolds equipped with an involutive distribution preserved by the homological vector field. In the following, we restrict to distributions cogenerated in degree one. In this particularly simple situation, we can provide an alternative proof of Zambon–Zhu result exploiting the description of vector valued forms in terms of their Spencer data.

Lemma 24. *Let $(A, \llbracket -, - \rrbracket, \rho)$ be a Lie algebroid over a manifold M . If (TM, B, ∇) is an IM foliation of A over TM , then there is a flat A -connection $\nabla^{A/B}$ in A/B such that*

$$(20) \quad \nabla_X^{A/B}(Y \bmod B) = \nabla_{\rho(Y)}(X \bmod B) - \llbracket Y, X \rrbracket \bmod B,$$

and, moreover,

$$(21) \quad d_\nabla(\llbracket X, Y \rrbracket \bmod B) = L_{\nabla_X^{A/B}} d_\nabla(Y \bmod B) - L_{\nabla_Y^{A/B}} d_\nabla(X \bmod B),$$

for all $X, Y \in \Gamma(A)$. Conversely, if $B \subset A$ is a vector subbundle, ∇ is a flat connection in A/B , and $\nabla^{A/B}$ is a flat A -connection in A/B satisfying (20) and (21), then (TM, B, ∇) is an IM foliation of A over TM .

Proof. For the first part of the statement, let (TM, B, ∇) be an IM foliation as in the statement. Denote by Γ_∇ the sheaf on M consisting of sections X of A such that $X \bmod B$ is ∇ -flat. Since $\Gamma(A/B)$ is locally generated by flat sections, $\Gamma(A)$ is locally generated by Γ_∇ . Now, the left-hand side of (20) is clearly $C^\infty(M)$ -linear in X . Moreover, it vanishes whenever $Y \in \Gamma(B)$. To see this, it is enough to compute on local generators $X \in \Gamma_\nabla$. In this case, the left-hand side of (20) reduces to $-\llbracket Y, X \rrbracket \bmod B$ which vanishes by property (1) of IM foliations whenever $Y \in \Gamma(B)$. One concludes that (20) defines a differential operator $\nabla_X^{A/B}$ in $\Gamma(A/B)$ for all $X \in \Gamma(A)$. It is easy to see that, besides being $C^\infty(M)$ -linear in X , $\nabla_X^{A/B}$ is actually a derivation with symbol $\rho(X)$. Thus $\nabla^{A/B}$ is a well-defined A -connection in A/B . To see that it is flat, check that the curvature

$$R(X, Y)(Z \bmod B) := (\llbracket \nabla_X^{A/B}, \nabla_Y^{A/B} \rrbracket - \nabla_{\llbracket X, Y \rrbracket}^{A/B})(Z \bmod B)$$

vanishes on all X, Y, Z . Since R is linear in the first two arguments, it is enough to check that it vanishes on $X, Y \in \Gamma_\nabla$. In this case $\llbracket X, Y \rrbracket \in \Gamma_\nabla$ as well and

$$R(X, Y)(Z \bmod B) = \llbracket \llbracket Z, Y \rrbracket, X \rrbracket - \llbracket \llbracket Z, X \rrbracket, Y \rrbracket + \llbracket Z, \llbracket X, Y \rrbracket \rrbracket = 0$$

by the Jacobi identity.

Finally, notice that Equation (21) is equivalent to

$$(22) \quad \nabla_Z(\llbracket X, Y \rrbracket \bmod B) = (\nabla_X^{A/B} \nabla_Z - \nabla_{[\rho(X), Z]}) (Y \bmod B) - (\nabla_Y^{A/B} \nabla_Z - \nabla_{[\rho(Y), Z]}) (X \bmod B),$$

$X, Y \in \Gamma(A)$, and $Z \in \mathfrak{X}(M)$. Actually, (22) can be easily obtained from (21), by inserting Z in both sides, and using

$$[i_Z, L_{\nabla_X^{A/B}}] = i_{[Z, \rho(X)]}.$$

Thus it is enough to check that the expression

$$S(X, Y; Z) := \nabla_Z(\llbracket X, Y \rrbracket \bmod B) - (\nabla_X^{A/B} \nabla_Z - \nabla_{[\rho(X), Z]}) (Y \bmod B) + (\nabla_Y^{A/B} \nabla_Z - \nabla_{[\rho(Y), Z]}) (X \bmod B)$$

vanishes for all X, Y, Z . A direct computation shows that S is $C^\infty(M)$ -linear in its first two arguments. Therefore, it is enough to compute $S(X, Y; Z)$ for $X, Y \in \Gamma_\nabla$. In this case $\llbracket X, Y \rrbracket \in \Gamma_\nabla$ as well and $S(X, Y; Z)$ vanishes.

The second part of the statement immediately follows from (20) and (21). \square

Theorem 25 [Zambon and Zhu 2012]. *There is a one-to-one correspondence between degree-one $\mathbb{N}Q$ -manifolds equipped with a compatible involutive distribution, cogenerated in degree one, and Lie algebroids $A \rightarrow M$ equipped with an IM foliation over TM .*

Proof. Let $\mathcal{M} = A[1]$ be a degree-one \mathbb{N} -manifold, and let \mathcal{D} be an involutive distribution on it, cogenerated in degree one. Denote by $\pi : \mathcal{M} \rightarrow M$ the projection of \mathcal{M} onto its zero dimensional shadow. The quotient bundle $T\mathcal{M}/\mathcal{D}$ identifies with $\mathcal{M} \times_M E[1]$ for a nongraded vector bundle $E \rightarrow M$, and $\theta_{\mathcal{D}}$ is a degree-one E -valued 1-form on \mathcal{M} . Moreover, \mathcal{D} projects surjectively onto TM , i.e., $\pi_* \mathcal{D} = TM$. In particular, for any vector field Z on M there is a (degree-zero) vector field $\tilde{Z} \in \Gamma(\mathcal{D})$ that is π -related to Z .

Denote by (D, ℓ) the Spencer data of $\theta_{\mathcal{D}}$. In particular, $\ell : \Gamma(A) \rightarrow \Gamma(E)$ is surjective. Let $B = \ker \ell$ so that E identifies with A/B and ℓ identifies with the projection $\Gamma(A) \rightarrow \Gamma(A/B)$. In the following we will understand this isomorphism. There is a unique first order differential operator $\delta : \Gamma(A/B) \rightarrow \Omega^1(M, A/B)$ such

that diagram

$$\begin{array}{ccc} \Gamma(A) & \xrightarrow{D} & \Omega^1(M, A/B) \\ \downarrow & \nearrow \delta & \\ \Gamma(A/B) & & \end{array}$$

commutes. To see this it is enough to show that $\Gamma(B) \subset \ker D$. Since $\Gamma(B) = \Gamma_{-1}(\mathcal{D})$, and sections of \mathcal{D} are infinitesimal symmetries by involutivity, $D(X) = L_X\theta_{\mathcal{D}} = 0$ for all $X \in \Gamma(B)$ (see [Remark 21](#)). It follows from (4) that $\delta(f\alpha) = f\delta\alpha - df \otimes \alpha$ for all $f \in C^\infty(M)$, and $\alpha \in \Gamma(A/B)$. Therefore, δ is minus the (first) de Rham differential of a unique connection ∇ in A/B . We claim that ∇ is a flat connection. Indeed, first of all, notice that for all $Z \in \mathfrak{X}(M)$ and $X \in \Gamma(A)$,

$$\nabla_Z(X \bmod B) = i_Z d_{\nabla}(X \bmod B) = -i_Z D(X) = -i_{\tilde{Z}} L_X \theta_{\mathcal{D}},$$

where \tilde{Z} is any degree-zero vector field on \mathcal{M} that is π -related to Z . We can choose $\tilde{Z} \in \Gamma(\mathcal{D})$ so that

$$\nabla_Z(X \bmod B) = -i_{\tilde{Z}} L_X \theta_{\mathcal{D}} = -i_{[\tilde{Z}, X]} \theta_{\mathcal{D}} = [\tilde{Z}, X] \bmod B.$$

Now, let Y, Z be vector fields on M , and let \tilde{Y}, \tilde{Z} be vector fields in \mathcal{D} that are π -related to them. Then, by involutivity, $[\tilde{Y}, \tilde{Z}]$ is in \mathcal{D} and it is π -related to $[Y, Z]$. Thus

$$\begin{aligned} \nabla_{[Y, Z]}(X \bmod B) &= [[\tilde{Y}, \tilde{Z}], X] \bmod B \\ &= ([\tilde{Y}, [\tilde{Z}, X]] - [\tilde{Z}, [\tilde{Y}, X]]) \bmod B \\ &= [\nabla_Y, \nabla_Z](X \bmod B). \end{aligned}$$

Conversely, let $B \subset A$ be a vector subbundle and let ∇ be a flat connection in A/B . Denote by $\ell : A \rightarrow A/B$ the projection. Then $(-d_{\nabla} \circ \ell, \ell)$ are Spencer data for an A/B -valued 1-form θ on \mathcal{M} . Put $\mathcal{D} = \ker \theta$. To see that \mathcal{D} is involutive, notice that $\Gamma_{-1}(\mathcal{D}) = \ker \ell = \Gamma(B)$. Moreover, \mathcal{D} projects surjectively on TM , therefore $\Gamma(\mathcal{D})$ is generated by 1) sections of B and 2) degree-zero vector fields in \mathcal{D} that are projectable onto M . Commuting the latter with the former, one gets sections of B which are again in \mathcal{D} . It remains to show that the commutator of two projectable vector fields \tilde{Z}, \tilde{Y} in \mathcal{D} is again in \mathcal{D} , i.e., $i_{[\tilde{Y}, \tilde{Z}]} \theta = 0$. Now $i_{[\tilde{Y}, \tilde{Z}]} \theta = 0$ if and only if $L_X i_{[\tilde{Y}, \tilde{Z}]} \theta = 0$ for all $X \in \mathfrak{X}_-(\mathcal{M}) = \Gamma(A)$. The same computation as above shows that

$$L_X i_{[\tilde{Y}, \tilde{Z}]} \theta = ([\nabla_Y, \nabla_Z] - \nabla_{[Y, Z]})(X \bmod B) = 0,$$

where $Y = \pi_* \tilde{Y}$, and $Z = \pi_* \tilde{Z}$. We conclude that involutive distributions \mathcal{D} on \mathcal{M} cogenerated in degree one are equivalent to the following (nongraded) data: a vector subbundle $B \subset A$ and a flat connection in A/B .

Finally, let \mathcal{D} be an involutive distribution on \mathcal{M} cogenerated in degree one and let (B, ∇) be the corresponding nongraded data. Moreover, let \mathbb{Q} be a homological derivation of $T\mathcal{M}/\mathcal{D} = \mathcal{M} \times_M A/B$, let Q be its symbol, and let $(A, \llbracket -, - \rrbracket, \rho)$ and $(A/B, \nabla^{A/B})$ be the Lie algebroid and the Lie algebroid representation corresponding to \mathbb{Q} . The distribution \mathcal{D} is compatible with Q , and \mathbb{Q} is induced by $[Q, -]$, if and only if $\theta_{\mathcal{D}}$ is compatible with \mathbb{Q} . To see when this is the case, we use again [Proposition 17](#). Identity (18) is automatically satisfied by $\omega = \theta_{\mathcal{D}}$. Additionally, for $\omega = \theta_{\mathcal{D}}$ and $X, Y \in \mathfrak{X}_-(\mathcal{M}) = \Gamma(A)$, one gets

$$\begin{aligned} A(X, Y) &= D(\llbracket [Q, X], Y \rrbracket) - L_{[Q, X]}D(Y) + L_{[Q, Y]}D(X) \\ &= -d_{\nabla}(\llbracket X, Y \rrbracket \bmod B) + L_{\nabla_X^{A/B}}d_{\nabla}(Y \bmod B) - L_{\nabla_Y^{A/B}}d_{\nabla}(X \bmod B), \end{aligned}$$

and

$$\begin{aligned} B(X, Y) &= \ell(\llbracket [Q, X], Y \rrbracket) + i_{[Q, X]}D(Y) + L_{[Q, Y]}\ell(X) \\ &= \llbracket X, Y \rrbracket \bmod B - \nabla_{\rho(X)}(Y \bmod B) + \nabla_Y^{A/B}(X \bmod B), \end{aligned}$$

where we used that $D = -d_{\nabla} \circ \ell$. [Proposition 17](#) and [Lemma 24](#) then show that \mathcal{D} is compatible with Q , and \mathbb{Q} is induced by $[Q, -]$, if and only if (B, ∇, TM) is an IM foliation of A over TM , and $\nabla^{A/B}$ is the A -connection in the statement of the lemma. □

4. Vector Valued 2-forms on $\mathbb{N}Q$ -manifolds

4.1. Degree one symplectic $\mathbb{N}Q$ -manifolds. Recall that a degree- n symplectic \mathbb{N} -manifold is an \mathbb{N} -manifold \mathcal{M} equipped with a degree- n symplectic structure, i.e., a degree- n nondegenerate, closed 2-form ω .

It immediately follows from the definition that, if (\mathcal{M}, ω) is a degree- n symplectic \mathbb{N} -manifold, then the degree of \mathcal{M} is at most n . If $n > 0$, then $\omega = d\vartheta$, with $\vartheta = n^{-1}i_{\Delta}\omega$.

Example 26. The degree- n 2-form ω on $T^*[n]M$ (see [Example 14](#)) is a degree- n symplectic structure.

A degree- n symplectic $\mathbb{N}Q$ -manifold is a degree- n symplectic manifold (\mathcal{M}, ω) equipped with a homological vector field Q such that $L_Q\omega = 0$.

Theorem 27 [Roytenberg 2002]. *Every degree-one symplectic \mathbb{N} -manifold (\mathcal{M}, ω) is of the kind $(T^*[1]M, d\vartheta)$, up to symplectomorphisms, where ϑ is the tautological degree-one 1-form on $T^*[1]M$ (see [Example 14](#)). Moreover, there is a one-to-one correspondence between degree-one symplectic $\mathbb{N}Q$ -manifolds and (nongraded) Poisson manifolds.*

Roytenberg’s proof exploits explicitly the Poisson bracket. We propose an alternative proof focusing on Spencer data. The advantage is that we can apply the

same strategy to degenerate ([Theorem 30](#) and [Corollary 31](#)) or higher-degree forms ([Theorems 36](#) and [39](#)) in a straightforward way.

Proof. Let (\mathcal{M}, ω) be a degree-one symplectic \mathbb{N} -manifold, and let (D, ℓ) be the Spencer data of ω . In particular, $\mathcal{M} = A[1]$ for some vector bundle $A \rightarrow M$. By nondegeneracy $\ell : \mathfrak{X}_{-1}(\mathcal{M}) \rightarrow \Omega^1(\mathcal{M})$ is an isomorphism $\Gamma(A) \simeq \Omega^1(M)$, i.e., $\mathcal{M} = A[1] \simeq T^*[1]M$. Moreover, since ω is closed, the diagram

$$(23) \quad \begin{array}{ccc} \Gamma(A) & \xrightarrow{D} & \Omega^2(M) \\ \ell \downarrow & \nearrow -d & \\ \Omega^1(M) & & \end{array}$$

commutes. This shows that diffeomorphism $\mathcal{M} \simeq T^*[1]M$ identifies ω with the canonical symplectic structure on $T^*[1]M$ (see [Example 14](#)), thus proving the first part of the statement. In the following we identify \mathcal{M} and $T^*[1]M$. For the second part of the statement, let Q be a homological vector field on \mathcal{M} and let $(T^*M, \rho, \llbracket -, - \rrbracket)$ be the corresponding Lie algebroid. Similarly to the previous section, (\mathcal{M}, ω, Q) is a symplectic $\mathbb{N}Q$ -manifold if and only if it satisfies [\(17\)](#), and [\(18\)](#), for all $X, Y \in \mathfrak{X}_{-1}(\mathcal{M}) \simeq \Omega^1(M)$. Indeed, since ω is closed, condition [\(16\)](#) is actually a consequence of [\(17\)](#), and [\(18\)](#). It is easy to see that one can even restrict to X, Y in the form df, dg , with $f, g \in C^\infty(M)$. In this case, one gets

$$\begin{aligned} B(X, Y) &= \ell(\llbracket Q, df \rrbracket, dg) + L_{[Q, df]}\ell(dg) = \llbracket df, dg \rrbracket + L_{\rho(df)}dg \\ &= \llbracket df, dg \rrbracket + d\rho(df)(g), \end{aligned}$$

and

$$C(X, Y) = i_{[Q, df]}\ell(dg) + i_{[Q, dg]}\ell(df) = \rho(df)(g) + \rho(dg)(f),$$

where we used that $\ell(df) = i_{df}\omega = df$, and $D(df) = L_{df}\omega = 0$ (see [Example 14](#)). Concluding, (\mathcal{M}, ω, Q) is a symplectic $\mathbb{N}Q$ -manifold if and only if

$$\rho(df)(g) + \rho(dg)(f) = 0 \quad \text{and} \quad \llbracket dg, df \rrbracket = -d\rho(df)(g) = d\rho(dg)(f),$$

i.e., if and only if $(T^*M, \rho, \llbracket -, - \rrbracket)$ is the Lie algebroid associated to a Poisson structure on M (see [Appendix B](#)). \square

4.2. Degree one presymplectic $\mathbb{N}Q$ -manifolds. In this subsection we relax the hypothesis about nondegeneracy of the 2-form in the previous subsection.

Definition 28. A *degree- n presymplectic \mathbb{N} -manifold* is a degree- n \mathbb{N} -manifold \mathcal{M} equipped with a degree- n presymplectic structure, i.e., a degree- n (possibly degenerate) closed 2-form ω . A *degree- n presymplectic $\mathbb{N}Q$ -manifold* is a degree- n presymplectic manifold (\mathcal{M}, ω) equipped with a homological vector field Q such that $L_Q\omega = 0$.

Remark 29. Unlike the symplectic case, the existence of a presymplectic form on an \mathbb{N} -manifold \mathcal{M} doesn't bound the degree of \mathcal{M} . This is the reason why we added a condition on the degree of \mathcal{M} in the definition of a degree- n presymplectic \mathbb{N} -manifold above.

In what follows we show that degree-one presymplectic $\mathbb{N}Q$ -manifolds (with an additional nondegeneracy condition) are basically equivalent to *Dirac manifolds*. Recall that a Dirac manifold is a manifold M equipped with a *Dirac structure*, i.e., a subbundle $\mathcal{D} \subset TM \oplus T^*M$ such that 1) \mathcal{D} is maximally isotropic with respect to the canonical, split signature, symmetric form on $TM \oplus T^*M$

$$(24) \quad \langle\langle (X, \sigma), (X', \sigma') \rangle\rangle = i_X \sigma' + i_{X'} \sigma,$$

and 2) sections of \mathcal{D} are preserved by the Dorfman (equivalently, Courant) bracket

$$(25) \quad [(X, \sigma), (X', \sigma')]_{\mathcal{D}} := ([X, X'], L_X \sigma' - i_{X'} d\sigma),$$

$X, X' \in \mathfrak{X}(M)$, $\sigma, \sigma' \in \Omega^1(M)$. Any Dirac structure \mathcal{D} is a Lie algebroid, with anchor given by projection $TM \oplus T^*M \rightarrow TM$ and bracket given by the Dorfman bracket (25). Dirac manifolds encompass presymplectic and Poisson manifolds (see [Courant 1990; Bursztyn 2013] for more details). They are sometimes regarded as Lagrangian submanifolds in certain degree-two symplectic $\mathbb{N}Q$ -manifolds. Corollary 31 below shows that they can be also regarded as suitable degree-one presymplectic $\mathbb{N}Q$ -manifolds.

Let M be a manifold, denote by $\text{pr}_T : TM \oplus T^*M \rightarrow TM$, and $\text{pr}_{T^*} : TM \oplus T^*M \rightarrow T^*M$ the canonical projections.

Theorem 30. *There is a one-to-one correspondence between degree-one presymplectic $\mathbb{N}Q$ -manifolds and (nongraded) Lie algebroids $A \rightarrow M$ equipped with a vector bundle morphism $\Phi : A \rightarrow TM \oplus T^*M$ such that*

- (1) *the anchor of A equals the composition $\text{pr}_T \circ \Phi$,*
- (2) *the image of Φ is an isotropic subbundle with respect to (24), and*
- (3) *Φ intertwines the Lie bracket $\llbracket -, - \rrbracket$ on $\Gamma(A)$ and the Dorfman bracket (25) on $\Gamma(TM \oplus T^*M)$, i.e., $\Phi \llbracket X, Y \rrbracket = [\Phi(X), \Phi(Y)]_{\mathcal{D}}$ for all $X, Y \in \Gamma(A)$.*

Proof. Let (\mathcal{M}, ω) be a degree-one presymplectic \mathbb{N} -manifold, and let (D, ℓ) be the corresponding Spencer data. Moreover, let Q be a homological vector field on \mathcal{M} , and let $(A, \rho, \llbracket -, - \rrbracket)$ be the corresponding Lie algebroid. Since ω is closed, diagram (23) commutes and ω is completely determined by ℓ . Now, combine ℓ and $\rho : A \rightarrow TM$ in a vector bundle morphism $\Phi := (\rho, \ell) : A \rightarrow TM \oplus T^*M$. In particular, $\rho = \text{pr}_T \circ \Phi$, i.e., Φ satisfies property (1) in the statement. Similarly as in the proof of Theorem 27, (\mathcal{M}, ω, Q) is a presymplectic $\mathbb{N}Q$ -manifold if and only

if (17) and (18) are satisfied for all $X, Y \in \mathfrak{X}_{-1}(\mathcal{M}) \simeq \Omega^1(M)$. One gets

$$\begin{aligned} B(X, Y) &= \ell(\llbracket [Q, X], Y \rrbracket) + i_{[Q, X]}D(Y) + L_{[Q, Y]}\ell(X) \\ &= \ell(\llbracket X, Y \rrbracket) - i_{\rho(X)}d\ell(Y) + L_{\rho(X)}\ell(X) \\ &= \text{pr}_{T^*}(\Phi\llbracket X, Y \rrbracket - [\Phi(X), \Phi(Y)]_{\mathbb{D}}), \end{aligned}$$

and

$$C(X, Y) = i_{[Q, X]}\ell(Y) + i_{[Q, Y]}\ell(X) = i_{\rho(X)}\ell(Y) - i_{\rho(Y)}\ell(X) = \langle\langle \Phi(X), \Phi(Y) \rangle\rangle.$$

Since $\text{pr}_T([\Phi(X), \Phi(Y)]_{\mathbb{D}}) = [\rho(X), \rho(Y)] = \rho[X, Y] = \text{pr}_T \Phi\llbracket X, Y \rrbracket$, one concludes that (\mathcal{M}, ω, Q) is a presymplectic $\mathbb{N}Q$ -manifold if and only if Φ , besides satisfying property (1) in the statement, does also satisfy properties (2) and (3).

Conversely, Let $\Phi : A \rightarrow TM \oplus T^*M$ be a vector bundle morphism. Put $\ell := \text{pr}_{T^*} \circ \Phi$. It is easy to see that $(\ell, -d \circ \ell)$ is a pair of Spencer data corresponding to a degree-one presymplectic form ω on \mathcal{M} . If, additionally, Φ satisfies properties (1), (2), and (3) in the statement, then the same computations as above show that (\mathcal{M}, ω, Q) is a presymplectic $\mathbb{N}Q$ -manifold. \square

Corollary 31. *There is a one-to-one correspondence between degree-one presymplectic $\mathbb{N}Q$ -manifolds (\mathcal{M}, ω, Q) such that*

- (1) $\text{rank } A = \dim M$, and
- (2) $\ker \ell \cap \ker \rho = 0$,

where $(A \rightarrow M, \rho, \llbracket -, - \rrbracket)$ is the Lie algebroid corresponding to (\mathcal{M}, Q) , and (ℓ, D) are Spencer data corresponding to ω , and (nongraded) Lie algebroids $A \rightarrow M$ equipped with a Lie algebroid isomorphism $\Phi : A \simeq \mathfrak{D}$ with values in a Dirac structure $\mathfrak{D} \subset TM \oplus T^*M$ over M .

Proof. Let (\mathcal{M}, ω, Q) be a degree-one presymplectic $\mathbb{N}Q$ -manifold and let (A, Φ) be the corresponding nongraded data as in [Theorem 30](#). The vector bundle morphism Φ is injective if and only if condition (2) in the statement is satisfied. In this case, Φ is an isomorphism onto its image \mathfrak{D} . Additionally, \mathfrak{D} is maximally isotropic in $TM \oplus T^*M$, hence a Dirac structure, if and only if $\text{rank } \mathfrak{D} = \text{rank } A$ is precisely $\dim M$, i.e., condition (1) in the statement is satisfied. \square

4.3. Degree one locally conformal symplectic $\mathbb{N}Q$ -manifolds. The original definition of a locally conformal symplectic (lcs) structure is (equivalent to) the following [[Vaisman 1985](#)]: an *lcs structure* on a manifold M is a pair (ϕ, ω) , where ϕ is a closed 1-form and ω is a nondegenerate 2-form on M such that $d\omega = \phi \wedge \omega$. Ordinary symplectic manifolds and lcs manifolds share some properties, but the latter are manifestly more general. Moreover, they are examples of Jacobi manifolds. In this paper we adopt an approach to lcs manifolds which is slightly more intrinsic

than the traditional one (see [Appendix A](#)) in the same spirit as the *intrinsic approach* to contact and Jacobi geometry of [[Crainic and Salazar 2015](#)].

Definition 32. A degree- n abstract lcs \mathbb{N} -manifold is an \mathbb{N} -manifold \mathcal{M} equipped with a degree- n abstract lcs structure, i.e., a triple $(\mathcal{L}, \nabla, \omega)$ where $\mathcal{L} \rightarrow \mathcal{M}$ is a line \mathbb{N} -bundle, ∇ is a flat connection in \mathcal{L} , and ω is a degree- n , nondegenerate, d_∇ -closed, \mathcal{L} -valued 2-form ω .

First of all, notice that \mathcal{L} , being a line bundle, is actually generated in one single degree $-k$. Up to a shift in degree in the above definition we may (and we actually will) assume $k = 0$. In particular, $\mathcal{L} = \mathcal{M} \times_M L$, for some (nongraded) vector bundle L on the degree-zero shadow M of \mathcal{M} , and ∇ is actually induced from a flat connection on L . Exactly as in the symplectic case [[Roytenberg 2002](#)] one shows that, if \mathcal{M} possesses a degree- n abstract lcs structure, then, by nondegeneracy, the degree of \mathcal{M} is at most n . If $n > 0$, then $\omega = d_\nabla \vartheta$, with $\vartheta = n^{-1} i_{\Delta_\mathcal{L}} \omega$.

Example 33. Consider the degree- n 2-form ω of [Example 15](#). If $E = L$ is a line bundle then the triple $(T[1]\mathcal{M} \times_M L, \nabla, \omega)$ is a degree- n abstract lcs structure.

A degree- n abstract lcs symplectic $\mathbb{N}\mathcal{Q}$ -manifold is a degree- n abstract lcs manifold $(\mathcal{M}, \mathcal{L}, \nabla, \omega)$ equipped with a homological derivation \mathbb{Q} of \mathcal{L} such that $L_{\mathbb{Q}}\omega = 0$. The proposition below shows that, actually, \mathbb{Q} is completely determined by its symbol.

Proposition 34. *Let $(\mathcal{M}, \mathcal{L}, \nabla, \omega)$ be an abstract lcs \mathbb{N} -manifold with homological derivation \mathbb{Q} , and let Q be the symbol of \mathbb{Q} . Then \mathbb{Q} is the covariant derivative along Q .*

Proof. The derivations \mathbb{Q} and ∇_Q share the same symbol Q and, therefore, their difference $\mathbb{Q} - \nabla_Q$ is a degree-one endomorphism of \mathcal{L} which can only consist in multiplying sections by a degree-one function f on \mathcal{M} . Thus,

$$0 = L_{\mathbb{Q}}\omega = L_{\nabla_Q}\omega + f\omega = -d_\nabla i_{\mathbb{Q}}\omega + f\omega$$

So that $f\omega = d_\nabla i_{\mathbb{Q}}\omega$. It follows that

$$0 = d_\nabla(f\omega) = df \cdot \omega.$$

Hence, by nondegeneracy, $df = 0$. Since f is a function of positive degree, one concludes that $f = 0$. \square

Theorem 35. *Every degree-one abstract lcs \mathbb{N} -manifold $(\mathcal{M}, \mathcal{L}, \nabla, \omega)$ is of the form $(T^*[1]M \otimes L, (T^*[1]M \otimes L) \times_M L, \nabla, d_\nabla \vartheta)$, up to isomorphisms (and a shift in the degree of \mathcal{L}), where ϑ is the tautological degree-one L -valued 1-form on $T^*[1]M \otimes L$ and ∇ is a flat connection in the line bundle $L \rightarrow M$ (see [Example 15](#)). Moreover, there is a one-to-one correspondence between degree-one abstract lcs*

$\mathbb{N}Q$ -manifolds and (nongraded) abstract locally conformal Poisson manifolds (see Appendix A for a definition).

Proof. The proof is a suitable adaptation of both the proofs of Theorem 23 and Theorem 27. Let $(\mathcal{M}, \mathcal{L}, \nabla, \omega)$ be a degree-one abstract lcs \mathbb{N} -manifold, and let (D, ℓ) be the Spencer data of ω . In particular, $\mathcal{M} = A[1]$ for some vector bundle $A \rightarrow M$, and $\mathcal{L} = A[1] \times_M L$ for some line bundle $L \rightarrow M$ (up to a shift). Moreover ∇ is induced in \mathcal{L} by a flat connection in L which, abusing the notation, we denote by ∇ again.

By nondegeneracy $\ell : \mathfrak{X}_{-1}(\mathcal{M}) \rightarrow \Omega^1(\mathcal{M}, L)$ is an isomorphism $\Gamma(A) \simeq \Omega^1(M, L)$, i.e., $\mathcal{M} = A[1] \simeq T^*[1]M \otimes L$. Moreover, since ω is d_∇ -closed, the diagram

$$\begin{array}{ccc} \Gamma(A) & \xrightarrow{D} & \Omega^2(M, L) \\ \ell \downarrow & \nearrow -d_\nabla & \\ \Omega^1(M, L) & & \end{array}$$

commutes. This shows that the diffeomorphism $\mathcal{M} \simeq T^*[1]M \otimes L$ identifies ω with the canonical L -valued 2-form on $T^*[1]M \otimes L$ (see Example 15), thus proving the first part of the statement. In the following we identify \mathcal{M} and $T^*[1]M \otimes L$. For the second part of the statement, let \mathbb{Q} be a homological derivation of $\mathcal{L} = \mathcal{M} \times_M L$. The derivation \mathbb{Q} is equivalent to a Lie algebroid $(T^*M \otimes L, \rho, \llbracket -, - \rrbracket)$ equipped with a representation (L, ∇^L) (beware not to confuse the Lie algebroid connection ∇^L and the standard connection ∇). As above $(\mathcal{M}, \mathcal{L}, \nabla, \omega)$ is an abstract lcs $\mathbb{N}Q$ -manifold with homological derivation \mathbb{Q} if and only if (17), and (18) are satisfied for all $X, Y \in \mathfrak{X}_{-1}(\mathcal{M}) \simeq \Omega^1(M, L)$. Similarly to the proof of Theorem 27, one can even restrict to X, Y in the form $d_\nabla \lambda, d_\nabla \mu$, with $\lambda, \mu \in \Gamma(L)$. In this case one gets

$$\begin{aligned} B(X, Y) &= \ell(\llbracket \mathbb{Q}, d_\nabla \lambda \rrbracket, d_\nabla \mu) + L_{[\mathbb{Q}, d_\nabla \lambda]} \ell(d_\nabla \mu) \\ &= \llbracket d_\nabla \lambda, d_\nabla \mu \rrbracket + \nabla_{\rho(d_\nabla \lambda)}^L d_\nabla \mu \\ &= \llbracket d_\nabla \lambda, d_\nabla \mu \rrbracket + d_\nabla \nabla_{\rho(d_\nabla \lambda)}^L \mu, \end{aligned}$$

and

$$C(X, Y) = i_{[\mathbb{Q}, d_\nabla \lambda]} \ell(d_\nabla \mu) + i_{[\mathbb{Q}, d_\nabla \mu]} d_\nabla \lambda = \nabla_{\rho(d_\nabla \lambda)}^L \mu + \nabla_{\rho(d_\nabla \mu)}^L \lambda,$$

where we used that $\ell(d_\nabla \lambda) = i_{d_\nabla \lambda} \omega = d_\nabla \lambda$, and $D(d_\nabla \lambda) = L_{d_\nabla \lambda} \omega = 0$ (see Example 14). Concluding, $(\mathcal{M}, \mathcal{L}, \nabla, \omega)$ is an abstract lcs $\mathbb{N}Q$ -manifold with homological derivation \mathbb{Q} if and only if $\nabla_{\rho(d_\nabla \lambda)}^L \mu + \nabla_{\rho(d_\nabla \mu)}^L \lambda = 0$ and $\llbracket d_\nabla \lambda, d_\nabla \mu \rrbracket = -d_\nabla \nabla_{\rho(d_\nabla \lambda)}^L \mu = d_\nabla \nabla_{\rho(d_\nabla \mu)}^L \lambda$, i.e., if and only if $(T^*M \otimes L, \rho, \llbracket -, - \rrbracket)$ is the Lie algebroid associated to a locally conformal Poisson structure (L, ∇, P) on M , and (L, ∇^L) is its canonical representation. \square

5. Higher-degree forms on degree-one $\mathbb{N}Q$ -manifolds

5.1. Vector valued forms on degree-one $\mathbb{N}Q$ -manifolds and Spencer operators.

In this section, we discuss general degree-one compatible vector valued forms on degree-one $\mathbb{N}Q$ -manifolds. It turns out that they are equivalent to the recently introduced *Spencer operators on Lie algebroids* [Crainic et al. 2015]. Let $(A, \llbracket -, - \rrbracket, \rho)$ be a Lie algebroid over a manifold M , (E, ∇^E) a representation of A , and let k be a nonnegative integer. An E -valued k -Spencer operator [Crainic et al. 2015] is a pair consisting of

- a (first order) differential operator $D : \Gamma(A) \rightarrow \Omega^k(M, E)$, and
- a $C^\infty(M)$ -linear map $\ell : \Gamma(A) \rightarrow \Omega^{k-1}(M, E)$,

such that

$$D(fX) = fD(X) - df \wedge \ell(X)$$

and, moreover,

$$(26) \quad \begin{aligned} L_{\nabla_X^E} D(Y) - L_{\nabla_Y^E} D(X) &= D(\llbracket X, Y \rrbracket), \\ L_{\nabla_X^E} \ell(Y) + i_{\rho(Y)} D(X) &= \ell(\llbracket X, Y \rrbracket), \\ i_{\rho(X)} \ell(Y) + i_{\rho(Y)} \ell(X) &= 0, \end{aligned}$$

for all $X, Y \in \Gamma(A)$. There is a difference in signs between the above definition and the original one in [Crainic et al. 2015]. The original definition is recovered by replacing $D \rightarrow -D$. We chose the sign convention which makes formulas simpler in the present graded context.

Spencer operators are the infinitesimal counterparts of multiplicative vector valued forms on Lie groupoids. When the vector bundle is a trivial line bundle, they reduce to the IM forms of Bursztyn and Cabrera [2012] (see also [Bursztyn et al. 2009] for the 2-form case). Hence the result of this section is the expected generalization of the following (well) known facts:

- Jacobi manifolds can be understood either as infinitesimal counterparts of contact Lie groupoids [Crainic and Salazar 2015] or as degree-one contact $\mathbb{N}Q$ -manifolds [Mehta 2013; Grabowski 2013].
- Poisson manifolds can be understood either as infinitesimal counterparts of symplectic Lie groupoids [Weinstein 1987] or as degree-one symplectic $\mathbb{N}Q$ -manifolds [Roytenberg 2002].

Theorem 36. *There is a one-to-one correspondence between*

- *degree-one \mathbb{N} -manifolds equipped with a degree-zero $\mathbb{N}Q$ -vector bundle \mathcal{E} and a degree-one compatible \mathcal{E} -valued differential k -form, and*

- *Lie algebroids equipped with a representation (E, ∇^E) and an E -valued k -Spencer operator.*

Proof. Let \mathcal{M} be a degree-one \mathbb{N} -manifold, and let $(\mathcal{E}, \mathbb{Q})$ be a degree-zero $\mathbb{N}Q$ -vector bundle over it. In particular $\mathcal{E} = \mathcal{M} \times_M E$ for a nongraded vector bundle $E \rightarrow M$. Let $(T^*M, \rho, \llbracket -, - \rrbracket)$ and (E, ∇^E) be the Lie algebroid and the Lie algebroid representation corresponding to \mathbb{Q} . Finally, let ω be a degree-one E -valued k -form on \mathcal{M} . Then, ω is compatible with \mathbb{Q} if and only if (16), (17), and (18) are satisfied, for all $X, Y \in \mathfrak{X}_{-1}(\mathcal{M}) \simeq \Gamma(A)$. Denote by (D, ℓ) the Spencer data corresponding to ω . Then

$$\begin{aligned} A(X, Y) &= D(\llbracket X, Y \rrbracket) - L_{\nabla_X^E} D(Y) + L_{\nabla_Y^E} D(X), \\ B(X, Y) &= \ell(\llbracket X, Y \rrbracket) + i_{\rho(X)} D(Y) + L_{\nabla_Y^E} \ell(X), \\ C(X, Y) &= i_{\rho(X)} \ell(Y) + i_{\rho(Y)} \ell(X). \end{aligned}$$

Concluding, ω is compatible with \mathbb{Q} if and only if (D, ℓ) is an E -valued k -Spencer operator on the Lie algebroid A . □

5.2. Degree one multisymplectic $\mathbb{N}Q$ -manifolds. We conclude this section specializing to degree-one multisymplectic $\mathbb{N}Q$ -manifolds. Let k be a positive integer. Recall that a k -plectic manifold (see, for instance, [Rogers 2012], see also [Cantrijn et al. 1999] for more details on multisymplectic geometry) is a manifold N equipped with a k -plectic structure, i.e., a closed $(k + 1)$ -form ω which is nondegenerate in the sense that the vector bundle morphism $TN \rightarrow \wedge^k T^*N$, $X \mapsto i_X \omega$ is an embedding. As expected, degree-one multisymplectic $\mathbb{N}Q$ -manifolds are equivalent to Lie algebroids equipped with an *IM multisymplectic structure*, also called a *higher Poisson structure* in [Bursztyn et al. 2015]. The latter are infinitesimal counterparts of multisymplectic groupoids. Specifically, an *IM k -plectic structure* on a Lie algebroid $(A, \llbracket -, - \rrbracket, \rho)$ (see [Bursztyn et al. 2015]) is a $C^\infty(M)$ -linear map $\ell : A \rightarrow \Omega^k(M)$ such that

$$(27) \quad i_{\rho(X)} \ell(Y) + i_{\rho(Y)} \ell(X) = 0,$$

$$(28) \quad L_{\rho(X)} \ell(Y) - i_{\rho(Y)} d\ell(X) = \ell(\llbracket X, Y \rrbracket),$$

for all $X, Y \in \Gamma(A)$, and, moreover,

$$\ker \ell := \{a \in A : \ell(a) = 0\} = 0, \quad (\text{im } \ell)^\circ := \{\zeta \in TM : i_\zeta \circ \ell = 0\} = 0.$$

Definition 37. A degree- n *k -plectic \mathbb{N} -manifold* is a degree- n \mathbb{N} -manifold \mathcal{M} equipped with a degree- n *k -plectic structure*, i.e., a closed $(k + 1)$ -form which is *nondegenerate* in the sense that the degree- n vector bundle morphism $T\mathcal{M} \rightarrow S^k T^*[-1]\mathcal{M}$, $X \mapsto i_X \omega$ is an embedding. A *k -plectic $\mathbb{N}Q$ -manifold* of degree- n is an $\mathbb{N}Q$ -manifold equipped with a compatible k -plectic structure.

Example 38. Let M be an ordinary (nongraded) manifold. The degree- n \mathbb{N} -manifold $\mathcal{M} = (\wedge^k T^*)[n]M$ comes equipped with the obvious tautological, degree- n k -form ϑ . Consider the degree- n $(k + 1)$ -form $\omega = d\vartheta$. It is a degree- n k -plectic structure. Negatively graded vector fields on \mathcal{M} identify with k -forms on M and it is easy to see, along similar lines as in Example 14, that the Spencer data (D, ℓ) of ω identify with $(-)^n$ times the exterior differential $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ and the identity $\text{id} : \Omega^k(M) \rightarrow \Omega^k(M)$ respectively.

Theorem 39. Degree one k -plectic $\mathbb{N}Q$ -manifolds are in one-to-one correspondence with Lie algebroids equipped with an IM k -plectic structure.

Proof. Let \mathcal{M} be a degree-one \mathbb{N} -manifold, ω a degree-one $(k + 1)$ -form on it and let (D, ℓ) be the corresponding Spencer data. In particular, $\mathcal{M} = A[1]$ for some vector bundle $A \rightarrow M$. Moreover, ω is closed if and only if $i_X d\omega = 0$ for all negatively graded vector fields X on \mathcal{M} . Indeed, from $i_X d\omega = 0$ it also follows that $L_X d\omega = 0$. In other words, $d\omega = 0$ if and only if the diagram

$$\begin{array}{ccc}
 \Gamma(A) & \xrightarrow{D} & \Omega^{k+1}(M) \\
 \ell \downarrow & \nearrow -d & \\
 \Omega^k(M) & &
 \end{array}$$

commutes. Conversely, a $C^\infty(M)$ -linear map $\ell : \Gamma(A) \rightarrow \Omega^k(M)$ uniquely determines a closed degree-one $(k + 1)$ -form on \mathcal{M} whose Spencer data are $(-d \circ \ell, \ell)$. Concluding, degree-one \mathbb{N} -manifolds equipped with a closed $(k + 1)$ -form are equivalent to vector bundles $A \rightarrow M$ equipped with a linear map $\ell : \Gamma(A) \rightarrow \Omega^k(M)$.

Now, let Q be a homological vector field on \mathcal{M} and let $(A, \rho, \llbracket -, - \rrbracket)$ be the corresponding Lie algebroid. The $(k + 1)$ -form ω is compatible with Q if and only if $(-d \circ \ell, \ell)$ is a $(k + 1)$ -Spencer operator, i.e., ℓ fulfills (27) and (28) (Equation (26) then follows from $D = -d \circ \ell$).

Finally, we need to characterize nondegeneracy of the closed form ω in terms of ℓ . Recall that M can be understood as a submanifold in \mathcal{M} via the “zero section” of $\mathcal{M} \rightarrow M$, and the vector bundle morphism $\Gamma : T\mathcal{M} \rightarrow S^k T^*[-1]\mathcal{M}$, $X \mapsto i_X \omega$, restricts to a vector bundle morphism $\Gamma|_M : T\mathcal{M}|_M \rightarrow S^k T^*[-1]\mathcal{M}|_M$. Now, there are canonical identifications $T\mathcal{M}|_M = TM \oplus A[1]$, and

$$S^k T^*[-1]\mathcal{M}|_M = \bigoplus_{i+j=k} \wedge^i T^*M \otimes S^j A^*[-1].$$

It follows from $|\omega| = 1$ that $\Gamma|_M$ does actually take values in $\wedge^{k-1} T^*M \otimes A^*[-1] \oplus \wedge^k T^*M$. More precisely, it identifies with the pair of vector bundle morphisms

$$A[1] \rightarrow \wedge^k T^*M, \quad X \mapsto \ell(X).$$

and

$$TM \rightarrow \wedge^{k-1} T^*M \otimes A^*[-1], \quad Z \mapsto i_Z \circ \ell.$$

Consequently, $\ker \ell$ and $(\operatorname{im} \ell)^\circ$ are trivial if and only if $\Gamma|_M$ is an embedding. It remains to show that ω is nondegenerate provided $\Gamma|_M$ is an embedding. This is easily seen, for instance, in local coordinates: let x^i be coordinates in M and z^a be (degree-one) fiber coordinates in $A[1] \rightarrow M$. Locally,

$$\omega = \omega_{a|i_1 \dots i_k} dz^a dx^{i_1} \dots dx^{i_k} + \omega'_{a|i_1 \dots i_{k+1}} z^a dx^{i_1} \dots dx^{i_{k+1}}.$$

In the basis $\{ \partial/\partial z^a \mid \partial/\partial x^i \}$ of $\mathfrak{X}(\mathcal{M})$ and $\{ dx^{i_1} \dots dx^{i_k} \mid dz^a dx^{i_1} \dots dx^{i_{k-1}} \mid \dots \}$ of $\Omega^k(\mathcal{M})$, the vector bundle morphism Γ is represented by the matrix

$$\left(\begin{array}{c|cc} \omega_{a|i_1 \dots i_k} & 0 & \dots \\ \hline * & k\omega_{a|wei_1 \dots i_k} & \dots \end{array} \right)$$

and $\Gamma|_M$ is represented by the same matrix with the lower-left block set to zero. This concludes the proof. □

Appendix A: Locally conformal symplectic manifolds revisited

We refer to [Vaisman 1985] for details about standard locally conformal symplectic (lcs) structures. Here, we present a slightly more intrinsic approach to them (A. M. Vinogradov, personal communication, 2014; see also [Vitagliano 2015a, Section 3]). Let M be a smooth manifold.

Definition 40. An abstract lcs structure on M is a triple (L, ∇, ω) , where $L \rightarrow M$ is a line bundle, ∇ is a flat connection in L , and ω is a nondegenerate L -valued 2-form on M such that $d_\nabla \omega = 0$, where $d_\nabla : \Omega(M, L) \rightarrow \Omega(M, L)$ is the de Rham differential of ∇ . A manifold equipped with an abstract lcs structure is an abstract lcs manifold.

Example 41. Let (L, ∇, ω) be an abstract lcs structure on M . If $L = M \times \mathbb{R}$ is the trivial line bundle, then ∇ is the same as a closed 1-form on M , specifically, the connection 1-form $\phi := -d_\nabla 1 \in \Omega^1(M)$. Moreover, ω is a standard (nondegenerate) 2-form on M and it is easy to see that (ϕ, ω) is a standard lcs structure, i.e., $d\omega = \phi \wedge \omega$. In particular, if $\phi = 0$, then ω is a symplectic structure.

The word “abstract” in Definition 40 refers to the fact that ω takes values in an “abstract” line-bundle L , as opposed to the concrete, trivial line bundle $M \times \mathbb{R}$. Similarly, one can define “abstract” locally conformal Poisson manifolds (see

below) and, more generally, “*abstract*” *Jacobi manifolds*. An *abstract Jacobi structure* (called a Jacobi bundle in [Marle 1991]) on a manifold M is a line bundle L equipped with a Lie bracket $\{-, -\}$ on $\Gamma(L)$ which is a first order differential operator in each entry (see, e.g., [Crainic and Salazar 2015] for details). Abstract Jacobi manifolds were introduced by Kirillov [1976] under the name *local Lie algebras with one dimensional fibers*. An abstract lcs structure (L, ∇, ω) on M determines an abstract Jacobi structure $(L, \{-, -\})$ as follows. First of all, by nondegeneracy, ω establishes an isomorphism $TM \rightarrow T^*M \otimes L$, $X \mapsto i_X \omega$. Denote by $\sharp : T^*M \otimes L \rightarrow TM$ the inverse isomorphism and, for $\lambda \in \Gamma(L)$, put $X_\lambda := \sharp(d_\nabla \lambda) \in \mathfrak{X}(M)$. Finally, put

$$\{\lambda, \mu\} := \omega(X_\lambda, X_\mu) = \nabla_{X_\lambda} \mu,$$

$\lambda, \mu \in \Gamma(L)$. Clearly, $\{-, -\}$ is a first order differential operator in each entry. Moreover, the Jacobi identity is equivalent to $d_\nabla \omega = 0$. Thus, $(L, \{-, -\})$ is an abstract Jacobi structure on M . Notice that there exists a unique linear morphism $P : \wedge^2(T^*M \otimes L) \rightarrow L$ such that

$$\{\lambda, \mu\} = P(d_\nabla \lambda, d_\nabla \mu), \quad \text{for all } \lambda, \mu \in \Gamma(L).$$

Example 42. Let $L = M \times \mathbb{R}$ so that (L, ∇, ω) is the same as a standard lcs structure (ϕ, ω) . Then, for $f, g \in C^\infty(M) = \Gamma(L)$, X_f is implicitly defined by

$$i_{X_f} \omega = df - f\phi,$$

and

$$\{f, g\} := \omega(X_f, X_g) = X_f(g) - g\phi(X_f).$$

In particular, if $\phi = 0$, then P is the Poisson bivector determined by the symplectic structure ω .

More generally, Let M be a smooth manifold, (L, ∇) a line bundle over M equipped with a flat connection, and let $P : \wedge^2(T^*M \otimes L) \rightarrow L$ be a linear morphism. One can then define a bracket $\{-, -\}_P$ in $\Gamma(L)$ by putting

$$\{\lambda, \mu\}_P = P(d_\nabla \lambda, d_\nabla \mu),$$

$\lambda, \mu \in \Gamma(L)$.

Definition 43. An *abstract locally conformal Poisson structure* on M is a triple (L, ∇, P) , where $L \rightarrow M$ is a line bundle, ∇ is a flat connection in L , and P is a linear morphism $P : \wedge^2(T^*M \otimes L) \rightarrow L$ such that $\{-, -\}_P$ is a Lie bracket. A manifold equipped with an abstract locally conformal Poisson structure is an *abstract locally conformal Poisson manifold*.

Thus, abstract lcs manifolds are abstract locally conformal Poisson manifolds (much as standard symplectic manifolds are standard Poisson manifolds), but the latter are more general.

Example 44. Let (L, ∇, P) be an abstract locally conformal Poisson structure on M . If $L = M \times \mathbb{R}$ is the trivial line bundle, and $\phi := -d_\nabla 1 \in \Omega^1(M)$ is the connection 1-form, then P is a standard bivector on M and a lengthy but straightforward computation shows that (ϕ, P) is a *locally conformal Poisson structure* in the sense of [Vaisman 2007], i.e., $[P, P]_{\text{ns}} = i_\phi P \wedge P$ (where $[-, -]_{\text{ns}}$ is the Nijenhuis–Schouten bracket of multivectors). In particular, if $\phi = 0$, then P is a Poisson structure.

Finally, notice also that abstract locally conformal Poisson manifolds are abstract Jacobi manifolds (of a special kind).

Appendix B: Lie algebroids and their representations

Recall that a *Lie algebroid* over a manifold M is a vector bundle $A \rightarrow M$ equipped with 1) a $C^\infty(M)$ -linear map $\rho : \Gamma(A) \rightarrow \mathfrak{X}(M)$ called the *anchor*, and 2) a Lie bracket $\llbracket -, - \rrbracket$ on $\Gamma(A)$ such that

$$\llbracket X, fY \rrbracket = \rho(X)(f)Y + f\llbracket X, Y \rrbracket, \quad X, Y \in \Gamma(A), \quad f \in C^\infty(M).$$

Example 45. The tangent bundle TM is a Lie algebroid with Lie bracket given by the commutator of vector fields and anchor given by the identity.

Let $A \rightarrow M$ be a Lie algebroid. A *representation* of A is a vector bundle $E \rightarrow M$ equipped with a *flat A-connection* ∇^E , i.e., a $C^\infty(M)$ -linear map

$$\nabla^E : \Gamma(A) \rightarrow \Gamma(DE), \quad X \mapsto \nabla_X^E,$$

such that the symbol of the derivation ∇_X^E is $\rho(X)$, and $[\nabla_X^E, \nabla_Y^E] = \nabla_{\llbracket X, Y \rrbracket}^E$, for all $X, Y \in \Gamma(A)$. Let (E, ∇^E) be a representation of A . The graded vector space $\Gamma(\wedge^* A^* \otimes E)$ of alternating, $C^\infty(M)$ -multilinear, $\Gamma(E)$ -valued forms on $\Gamma(A)$ is naturally equipped with a homological operator d_E given by the following *Chevalley–Eilenberg formula*:

$$(d_E \varphi)(X_1, \dots, X_{k+1}) \\ := \sum_i (-)^{i+1} \nabla_{X_i}^E (\varphi(\dots, \widehat{X}_i, \dots)) + \sum_{i < j} (-)^{i+j} \varphi(\llbracket X_i, X_j \rrbracket, \dots, \widehat{X}_i, \dots, \widehat{X}_j, \dots),$$

where $\varphi \in \Gamma(\wedge^k A^* \otimes E)$ is an alternating form with k -entries, $X_1, \dots, X_{k+1} \in \Gamma(A)$, and a hat ($\widehat{}$) denotes omission.

Example 46. Let ∇ be a standard flat connection in a vector bundle E . Then (E, ∇) is a representation of the Lie algebroid TM and the de Rham operator d_∇ of ∇ is its associated homological operator.

Example 47. Let $A \rightarrow M$ be a Lie algebroid. Clearly $(M \times \mathbb{R}, \rho)$ is a canonical representation of A . In particular, $\Gamma(\wedge^* A)$ is equipped with a homological operator (in fact a derivation) which we denote by d_A .

Example 48. Let $(L, \{-, -\})$ be an abstract Jacobi structure on a manifold M . There is a unique Lie algebroid $(J^1 L, \rho, \llbracket -, - \rrbracket)$ such that $\llbracket j^1 \lambda, j^1 \mu \rrbracket = j^1 \{\lambda, \mu\}$, and $\rho(j^1 \lambda)$ is the symbol of the first order differential operator (in fact a derivation) $\{\lambda, -\}$, where $\lambda, \mu \in \Gamma(L)$. Moreover, there is a unique representation (L, ∇^L) of $J^1 L$ such that $\nabla_{j^1 \lambda}^L \mu = \{\lambda, \mu\}$. In particular,

$$(29) \quad \llbracket j^1 \lambda, j^1 \mu \rrbracket = j^1(\nabla_{j^1 \lambda}^L \mu).$$

Conversely, let $(J^1 L, \rho, \llbracket -, - \rrbracket)$ be a Lie algebroid equipped with a representation (L, ∇^L) such that (29) holds. For $\lambda, \mu \in \Gamma(L)$ put $\{\lambda, \mu\} := \nabla_{j^1 \lambda}^L \mu$. Then $(L, \{-, -\})$ is an abstract Jacobi structure on M . This shows that abstract Jacobi structures $(L, \{-, -\})$ are equivalent to Lie algebroids $(J^1 L, \rho, \llbracket -, - \rrbracket)$ equipped with a representation (L, ∇^L) such that (29) holds.

Example 49. Let $\{-, -\}$ be a Poisson structure on a manifold M . There is a unique Lie algebroid $(T^*M, \rho, \llbracket -, - \rrbracket)$ such that $\llbracket df, dg \rrbracket = d\{f, g\}$, and $\rho(df)$ is the Hamiltonian vector field of f , where $f, g \in C^\infty(M)$. In particular,

$$(30) \quad \llbracket df, dg \rrbracket = d(\rho(df)(g)) \quad \text{and} \quad \rho(df)(g) + \rho(dg)(f) = 0.$$

Conversely, let $(T^*M, \rho, \llbracket -, - \rrbracket)$ be a Lie algebroid such that (30) holds. For $f, g \in C^\infty(M)$ put $\{f, g\} := \rho(df)(g)$. Then $\{-, -\}$ is a Poisson structure on M . This shows that Poisson structures are equivalent to Lie algebroids $(T^*M, \rho, \llbracket -, - \rrbracket)$ such that (30) holds.

Example 50. Let (L, ∇, ω) be an abstract locally conformal Poisson structure on a manifold M (see the previous appendix). There is a unique Lie algebroid $(T^*M \otimes L, \rho, \llbracket -, - \rrbracket)$ such that $\llbracket d_\nabla \lambda, d_\nabla \mu \rrbracket = d_\nabla \{\lambda, \mu\}$, and $\rho(d_\nabla \lambda)$ is the symbol of the first order differential operator $\{\lambda, -\}$, where $\lambda, \mu \in \Gamma(L)$. Moreover, there is a unique representation (L, ∇^L) of $T^*M \otimes L$ such that $\nabla_{d_\nabla \lambda}^L \mu = \{\lambda, \mu\}$. In particular,

$$(31) \quad \llbracket d_\nabla \lambda, d_\nabla \mu \rrbracket = d_\nabla(\nabla_{d_\nabla \lambda}^L \mu) \quad \text{and} \quad \nabla_{d_\nabla \lambda}^L \mu + \nabla_{d_\nabla \mu}^L \lambda = 0.$$

Conversely, let $(T^*M \otimes L, \rho, \llbracket -, - \rrbracket)$ be a Lie algebroid equipped with a representation (L, ∇^L) such that (31) holds. For $\lambda, \mu \in \Gamma(L)$ put $\{\lambda, \mu\} := \nabla_{d_\nabla \lambda}^L \mu$. Then $(L, \nabla, \{-, -\})$ is a locally conformal Poisson structure on M . Thus locally

conformal Poisson structures are equivalent to Lie algebroids $(T^*M \otimes L, \rho, \llbracket -, - \rrbracket)$ such that (31) holds.

Acknowledgement. I thank the anonymous referee for carefully reading the first manuscript and for her/his suggestions to improve the readability of the paper.

References

- [Bonavolontà and Poncin 2013] G. Bonavolontà and N. Poncin, “On the category of Lie n -algebroids”, *J. Geom. Phys.* **73** (2013), 70–90. [MR](#) [Zbl](#)
- [Bruce 2011] A. J. Bruce, “From L_∞ -algebroids to higher Schouten/Poisson structures”, *Rep. Math. Phys.* **67**:2 (2011), 157–177. [MR](#) [Zbl](#)
- [Bursztyn 2013] H. Bursztyn, “A brief introduction to Dirac manifolds”, pp. 4–38 in *Geometric and topological methods for quantum field theory*, edited by A. Cardona et al., Cambridge Univ. Press, 2013. [MR](#) [Zbl](#) [arXiv](#)
- [Bursztyn and Cabrera 2012] H. Bursztyn and A. Cabrera, “Multiplicative forms at the infinitesimal level”, *Math. Ann.* **353**:3 (2012), 663–705. [MR](#) [Zbl](#)
- [Bursztyn et al. 2009] H. Bursztyn, A. Cabrera, and C. Ortiz, “Linear and multiplicative 2-forms”, *Lett. Math. Phys.* **90**:1-3 (2009), 59–83. [MR](#) [Zbl](#)
- [Bursztyn et al. 2015] H. Bursztyn, A. Cabrera, and D. Iglesias, “Multisymplectic geometry and Lie groupoids”, pp. 57–73 in *Geometry, mechanics, and dynamics*, edited by D. E. Chang et al., Fields Inst. Commun. **73**, Springer, New York, 2015. [MR](#) [Zbl](#) [arXiv](#)
- [Cantrijn et al. 1999] F. Cantrijn, A. Ibort, and M. de León, “On the geometry of multisymplectic manifolds”, *J. Austral. Math. Soc. Ser. A* **66**:3 (1999), 303–330. [MR](#) [Zbl](#)
- [Cattaneo and Schätz 2011] A. S. Cattaneo and F. Schätz, “Introduction to supergeometry”, *Rev. Math. Phys.* **23**:6 (2011), 669–690. [MR](#) [Zbl](#)
- [Courant 1990] T. J. Courant, “Dirac manifolds”, *Trans. Amer. Math. Soc.* **319**:2 (1990), 631–661. [MR](#) [Zbl](#)
- [Crainic and Salazar 2015] M. Crainic and M. A. Salazar, “Jacobi structures and Spencer operators”, *J. Math. Pures Appl.* (9) **103**:2 (2015), 504–521. [MR](#) [Zbl](#)
- [Crainic et al. 2015] M. Crainic, M. A. Salazar, and I. Struchiner, “Multiplicative forms and Spencer operators”, *Math. Z.* **279**:3-4 (2015), 939–979. [MR](#) [Zbl](#)
- [Getzler 2010] E. Getzler, “Higher derived brackets”, preprint, 2010. [arXiv](#)
- [Grabowski 2013] J. Grabowski, “Graded contact manifolds and contact Courant algebroids”, *J. Geom. Phys.* **68** (2013), 27–58. [MR](#) [Zbl](#)
- [Jotz and Ortiz 2011] M. Jotz and C. Ortiz, “Foliated groupoids and their infinitesimal data”, preprint, 2011. [arXiv](#)
- [Kirillov 1976] A. A. Kirillov, “Local Lie algebras”, *Uspehi Mat. Nauk* **31**:4(190) (1976), 57–76. In Russian; translated in *Russian Math. Surveys* **31**:4 (1976), 55–75. [MR](#)
- [Kjeseth 2001a] L. Kjeseth, “A homotopy Lie–Rinehart resolution and classical BRST cohomology”, *Homology Homotopy Appl.* **3**:1 (2001), 165–192. [MR](#) [Zbl](#)
- [Kjeseth 2001b] L. Kjeseth, “Homotopy Rinehart cohomology of homotopy Lie–Rinehart pairs”, *Homology Homotopy Appl.* **3**:1 (2001), 139–163. [MR](#) [Zbl](#)
- [Marle 1991] C.-M. Marle, “On Jacobi manifolds and Jacobi bundles”, pp. 227–246 in *Symplectic geometry, groupoids, and integrable systems* (Berkeley, 1989), edited by P. Dazord and A. Weinstein, Math. Sci. Res. Inst. Publ. **20**, Springer, New York, 1991. [MR](#) [Zbl](#)

- [Mehta 2006] R. A. Mehta, *Supergroupoids, double structures, and equivariant cohomology*, Ph.D. thesis, University of California, Berkeley, 2006, <http://search.proquest.com/docview/305363751>. [MR](#)
- [Mehta 2013] R. A. Mehta, “Differential graded contact geometry and Jacobi structures”, *Lett. Math. Phys.* **103**:7 (2013), 729–741. [MR](#) [Zbl](#)
- [Rogers 2012] C. L. Rogers, “ L_∞ -algebras from multisymplectic geometry”, *Lett. Math. Phys.* **100**:1 (2012), 29–50. [MR](#) [Zbl](#)
- [Roytenberg 2002] D. Roytenberg, “On the structure of graded symplectic supermanifolds and Courant algebroids”, pp. 169–185 in *Quantization, Poisson brackets and beyond* (Manchester, 2001), edited by T. Voronov, Contemp. Math. **315**, Amer. Math. Soc., Providence, RI, 2002. [MR](#) [Zbl](#)
- [Salazar 2013] M. A. Salazar, *Pfaffian groupoids*, Ph.D. thesis, Universiteit Utrecht, 2013, <http://dspace.library.uu.nl/bitstream/handle/1874/275521/salazar-pinzon.pdf>.
- [Sati et al. 2012] H. Sati, U. Schreiber, and J. Stasheff, “Twisted differential string and fivebrane structures”, *Comm. Math. Phys.* **315**:1 (2012), 169–213. [MR](#) [Zbl](#)
- [Sheng and Zhu 2011] Y. Sheng and C. Zhu, “Higher Extensions of Lie Algebroids and Application to Courant Algebroids”, preprint, 2011. [arXiv](#)
- [Vaisman 1985] I. Vaisman, “Locally conformal symplectic manifolds”, *Internat. J. Math. Math. Sci.* **8**:3 (1985), 521–536. [MR](#) [Zbl](#)
- [Vaisman 2007] I. Vaisman, “Dirac structures and generalized complex structures on $TM \times \mathbb{R}^h$ ”, *Adv. Geom.* **7**:3 (2007), 453–474. [MR](#) [Zbl](#)
- [Vitagliano 2014] L. Vitagliano, “On the strong homotopy Lie–Rinehart algebra of a foliation”, *Commun. Contemp. Math.* **16**:6 (2014), 1450007, 49. [MR](#) [Zbl](#)
- [Vitagliano 2015a] L. Vitagliano, “Dirac–Jacobi Bundles”, preprint, 2015. To appear in *J. Sympl. Geom.* [arXiv](#)
- [Vitagliano 2015b] L. Vitagliano, “Representations of homotopy Lie–Rinehart algebras”, *Math. Proc. Cambridge Philos. Soc.* **158**:1 (2015), 155–191. [MR](#)
- [Voronov 2010] T. T. Voronov, “ Q -manifolds and higher analogs of Lie algebroids”, pp. 191–202 in *XXIX Workshop on Geometric Methods in Physics*, edited by P. Kielanowski et al., AIP Conf. Proc. **1307**, Amer. Inst. Phys., Melville, NY, 2010. [MR](#) [Zbl](#) [arXiv](#)
- [Weinstein 1987] A. Weinstein, “Symplectic groupoids and Poisson manifolds”, *Bull. Amer. Math. Soc. (N.S.)* **16**:1 (1987), 101–104. [MR](#) [Zbl](#)
- [Zambon and Zhu 2012] M. Zambon and C. Zhu, “Distributions and quotients on degree 1 NQ-manifolds and Lie algebroids”, *J. Geom. Mech.* **4**:4 (2012), 469–485. [MR](#) [Zbl](#) [arXiv](#)

Received April 14, 2015. Revised January 14, 2016.

LUCA VITAGLIANO
 DEPARTMENT OF MATHEMATICS
 UNIVERSITÀ DEGLI STUDI DI SALERNO
 & ISTITUTO NAZIONALE DI FISICA NUCLEARE
 GC SALERNO
 VIA GIOVANNI PAOLO II N° 123
 I-84084 FISCIANO (SA)
 ITALY
lvitagliano@unisa.it

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.


The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic.

Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 283 No. 2 August 2016

The fundamental theorem of tropical differential algebraic geometry	257
FUENSANTA AROCA, CRISTHIAN GARAY and ZEINAB TOGHANI	
A simple solution to the word problem for virtual braid groups	271
PAOLO BELLINGERI, BRUNO A. CISNEROS DE LA CRUZ and LUIS PARIS	
Completely contractive projections on operator algebras	289
DAVID P. BLECHER and MATTHEW NEAL	
Invariants of some compactified Picard modular surfaces and applications	325
AMIR DŽAMBIĆ	
Radial limits of bounded nonparametric prescribed mean curvature surfaces	341
MOZHGAN (NORA) ENTEKHABI and KIRK E. LANCASTER	
A remark on the Noetherian property of power series rings	353
BYUNG GYUN KANG and PHAN THANH TOAN	
Curves with prescribed intersection with boundary divisors in moduli spaces of curves	365
XIAO-LEI LIU	
Virtual rational Betti numbers of nilpotent-by-abelian groups	381
BEHROOZ MIRZAI and FATEMEH Y. MOKARI	
A planar Sobolev extension theorem for piecewise linear homeomorphisms	405
EMANUELA RADICI	
A combinatorial approach to Voiculescu's bi-free partial transforms	419
PAUL SKOUFRANIS	
Vector bundle valued differential forms on $\mathbb{N}\mathbb{Q}$ -manifolds	449
LUCA VITAGLIANO	
Discriminants and the monoid of quadratic rings	483
JOHN VOIGHT	