

ORIGINAL ARTICLE

Vertical segregation and phylogenetic characterization of ammonia-oxidizing Archaea in a deep oligotrophic lake

Jean-Christophe Auguet, Xavier Triadó-Margarit, Natalya Nomokonova, Lluís Camarero and Emilio O Casamayor

Limnological Observatory of the Pyrenees (LOOP)-Biogeodynamics & Biodiversity Interactions Group, CEAB-CSIC, Centre d'Estudis Avançats de Blanes – CSIC, Girona, Spain

Freshwater habitats have been identified as one of the largest reservoirs of archaeal genetic diversity, with specific lineages of ammonia-oxidizing archaea (AOA) populations different from soils and seas. The ecology and biology of lacustrine AOA is, however, poorly known. In the present study, vertical changes in archaeal abundance by CARD-FISH, quantitative PCR (qPCR) analyses and identity by clone libraries were correlated with environmental parameters in the deep glacial high-altitude Lake Redon. The lake is located in the central Spanish Pyrenees where atmospheric depositions are the main source of reactive nitrogen. Strong correlations were found between abundance of thaumarchaeotal 16S rRNA gene, archaeal *amoA* gene and nitrite concentrations, indicating an ammonium oxidation potential by these microorganisms. The bacterial *amoA* gene was not detected. Three depths with potential ammonia-oxidation activity were unveiled along the vertical gradient, (i) on the top of the lake in winter–spring (that is, the 0 °C slush layers above the ice-covered sheet), (ii) at the thermocline and (iii) the bottom waters in summer–autumn. Overall, up to 90% of the 16S rRNA gene sequences matched Thaumarchaeota, mostly from both the Marine Group (MG) 1.1a (*Nitrosoarchaeum*-like) and the sister clade SAGMGC – 1 (*Nitrosotalea*-like). Clone-libraries analysis showed the two clades changed their relative abundances with water depth being higher in surface and lower in depth for SAGMGC – 1 than for MG 1.1a, reflecting a vertical phylogenetic segregation. Overall, the relative abundance and recurrent appearance of SAGMGC – 1 suggests a significant environmental role of this clade in alpine lakes. These results expand the set of ecological and thermal conditions where Thaumarchaeota are distributed, unveiling vertical positioning in the water column as a key factor to understand the ecology of different thaumarchaeotal clades in lacustrine environments.

The ISME Journal (2012) 6, 1786–1797; doi:10.1038/ismej.2012.33; published online 12 April 2012

Subject Category: microbial ecology and functional diversity of natural habitats

Keywords: Archaea; AOA; lakes; alpine

Introduction

Freshwater habitats have recently been proposed as one of the largest reservoirs of archaeal genetic diversity (Llirós *et al.*, 2008; Auguet *et al.*, 2010). Some of the representative freshwater lineages identified may represent potential key archaeal groups in terms of distribution and ecological processes (Auguet *et al.*, 2010; Barberán *et al.*, 2011). Unfortunately, at present, only a few examples provide enough information to gain clues on the ecological significance of these

uncultured microorganisms, such as the ammonia-oxidizing populations within the thaumarchaeotal Marine Group (MG) 1.1a. The genomes of widespread and abundant representatives of the domain Archaea encode for the ammonia mono-oxygenase enzyme (Venter *et al.*, 2004; Treusch *et al.*, 2005). The isolated Thaumarchaeota *Nitrosopumilus maritimus* (Konneke *et al.*, 2005) have shown chemolithoautotrophic growth fueled with ammonia as energy source, initiating further research on the role of archaea in the global nitrogen cycle.

In aquatic ecosystems, the phylogenies of both *amoA* and 16S rRNA genes indicate that most ammonia-oxidizing archaea (AOA) belong to a few specific lineages, mainly the MG 1.1a and the deeply branching thaumarchaeotal clade pSL12 (Mincer *et al.*, 2007). These lineages are distantly related to those associated with soil or sediment

Correspondence: EO Casamayor, Limnological Observatory of the Pyrenees (LOOP)-Biogeodynamics & Biodiversity Interactions Group, CEAB-CSIC, Centre d'Estudis Avançats de Blanes – CSIC, C/ d'accès a la Cala St. Francesc, 14, Blanes, Girona 17300, Spain. E-mail: casamayor@ceab.csic.es

Received 22 December 2011; revised 6 March 2012; accepted 6 March 2012; published online 12 April 2012

(Auguet *et al.*, 2011). Despite the fact that only a handful of freshwater environments have been investigated so far, AOA have been consistently detected in rivers (Galand *et al.*, 2006; Herfort *et al.*, 2009; Merbt *et al.*, 2011), the Great African Lake Kivu (Llirós *et al.*, 2010), groundwater (Reed *et al.*, 2009), drinking waters (van der Wielen *et al.*, 2009), freshwater sediments (Herrmann *et al.*, 2008) and alpine lakes (Auguet *et al.*, 2011). These studies have significantly expanded the archaeal *amoA* phylogenetic tree showing lacustrine *amoA* gene sequences grouped in coherent phylogenetic clusters, and suggesting that freshwater habitats contain archaeal nitrifiers different from soil and marine habitats.

Spatio-temporal changes in the distribution of planktonic freshwater AOA are, however, poorly documented (Pouliot *et al.*, 2009; Llirós *et al.*, 2010; Auguet *et al.*, 2011) in contrast to what is known in the marine realm. The characteristic AOA abundance peak observed in the oxygen-minimum zone and dimly lit oceanic waters are often associated to maxima in nitrite concentrations, suggesting active ammonia oxidation in these areas. Significant correlations found between archaeal *amoA* transcripts abundance and nitrite maxima in the oxygen-minimum zone of the Black Sea (Lam *et al.*, 2007), the Peruvian upwelling (Lam *et al.*, 2009), and upwelled waters north of the equator (Church *et al.*, 2010) have been reported. Consistent differences have been observed in the sea for vertical AOA abundance, identity and activity (Francis *et al.*, 2005; Hallam *et al.*, 2006; Mincer *et al.*, 2007; Beman *et al.*, 2008; Agogué *et al.*, 2008; Church *et al.*, 2010). In an interannual study of surface waters in the Mediterranean sea, the abundance of AOA showed a consistent seasonal pattern, and *amoA* gene relative abundance was positively correlated to nitrate concentrations (Galand *et al.*, 2010). From the few studies carried out in planktonic freshwater, both strong seasonal dynamics in AOA observed in surface waters of shallow alpine lakes, which were well explained by temporal changes in ammonium and nitrite concentrations (Auguet *et al.*, 2011), and the maximal abundance of AOA in the oxycline/nitracline of chemically stratified lakes (Pouliot *et al.*, 2009; Llirós *et al.*, 2010), suggest a close link between freshwater Thaumarchaeota and the nitrogen cycling. In a recent investigation (Auguet *et al.*, 2011), we studied the temporal changes in the archaeal assemblages inhabiting the surface waters of three connected shallow and well-mixed Pyrenean lakes (Redó AigüesTortes (AT), Llong and Llebreta in the AigüesTortes valley) with a very low water-residence time (see Table 1 in Auguet and Casamayor, 2008). In the present study, we followed the vertical changes of AOA along the water column in the deep glacial stratified alpine Lake Redon (three orders of magnitude higher water-residence time, Auguet and Casamayor, 2008) during one year unveiling potential archaeal nitrification hotspots

Table 1 Diversity indicators of archaeal assemblages in Lake Redon based on OTUs defined by the 16S rRNA (at 3% cutoff) and *amoA* (5% cutoff) genes.

Clone libraries	Clone numbers	OTU	Coverage(%)	Chao1	PD
<i>16S rRNA</i>					
Slush-April	24	4	87	7	0.78 ± 0.05
20m-April	25	8	86	10	0.92 ± 0.08
60m-April	26	13	73	18	1.03 ± 0.11
SML-Sep	29	8	89	9	0.48 ± 0.02
20m-Sep	29	13	69	31	1.05 ± 0.10
60m-Sep	29	16	59	49	1.46 ± 0.15
<i>amoA</i>					
Slush-April	24	6	91	6	0.86 ± 0.09
20m-April	18	5	89	6	0.79 ± 0.05
60m-April	30	10	83	13	0.64 ± 0.06
SML-Sep	5	4	40	5	0.59 ± 0.10
20m-Sep	27	6	92	6	0.35 ± 0.08
60m-Sep	27	5	96	5	0.39 ± 0.11

Abbreviations: OTUs, operational taxonomic units; PD, phylogenetic diversity; rRNA, ribosomal RNA; SML, surface microlayer.

along the water column and vertical phylotype segregation.

Materials and methods

Study site, sampling and general analyses

Lake Redon is a deep-glacial cirque lake located in the Conanglés valley, central Spanish Pyrenees (42°38'34"N, 0°46'13"E, altitude 2240 m, maximum depth 73 m, surface 0.24 km²) extensively studied in the last 30 years by limnologists (see Catalan *et al.*, 2006 for more details) and recently by bacterial ecologists (Hervàs and Casamayor 2009; Llorens-Marès *et al.*, 2012). This lake is completely different from Lake Redó AT located in the Aigües Tortes National Park, a shallow (12 m depth) well-mixed lake with a very low water-residence time (see more details in Auguet and Casamayor, 2008; Auguet *et al.*, 2011). Local precipitation is the sole source of water for Lake Redon, as there are no permanent rivers in the basin, and the lake has a small catchment area of c. 1.5 km². The lake is usually ice-covered for about 6 months of the year. The thickness of the ice cover reaches several meters during its maximum in the month of April with dynamic changes in composition of the ice bacterial assemblages (Llorens-Marès *et al.*, 2012). Lake Redon has a dimictic regime, with mixing periods in spring and in autumn. The lake is oligotrophic owing to the sparse vegetation and poorly developed soils in the small catchment area, and atmospheric deposition is the main source of reactive nitrogen (Catalan *et al.*, 1994).

Samples for this study were collected monthly between May 2007 and April 2008 from six selected

depths (air–water surface microlayer (SML), 2, 10, 20, 35 and 60 m) at the deepest point of the lake using a 2-l Niskin bottle. These depths covered different limnological characteristics of the lake. The lake was frozen from November to April and inaccessible during January and February (data for these two months are missing in the graphs). The first 400 μm of the SML were collected with a nylon screen sampler (Auguet and Casamayor, 2008) during the ice-free period. In March and April 2008, we sampled the slush (that is, mixture layer of snow and lake water formed within the ice cover) as previously reported (Llorens-Marès *et al.*, 2012) and samples were processed in the laboratory within 1–2 h.

Samples for dissolved organic carbon (DOC) were filtered at the sampling site through a pre-combusted Whatman GF/F filter (Whatman Ltd, Kent, UK) and stored in the dark at 4 °C until analyzed (between 3 and 5 h after sampling) by high temperature combustion/direct injection into a Shimadzu TOC 5000 IR gas analyzer (Shimadzu Co., Kyoto, Japan). Sample stability and blank tests were run to check that the storage did not affect DOC measurements. Ammonium was colorimetrically determined according to Solórzano (1969). Nitrate was measured by capillary electrophoresis (model Quanta-4000, Waters Co., Milford, MA, USA) (Doble and Haddad, 1999). Nitrite was determined spectrophotometrically at 560 nm after the reaction in H_3PO_4 acid medium with a solution of sulfanilamide and *N*-(1-naphthyl) ethylene diamine dihydrochloride. Chlorophyll *a* was spectrophotometrically determined using water samples (0.5 ml) collected on Whatmann GF/F glass fiber filters and extracted with acetone (90%, v/v). Prokaryotic cells were collected in triplicate on 0.2- μm pore-size polycarbonate filters and processed for DAPI and CARD-FISH counts with probes EUB338 for bacterial and ARCH915 for archaeal cells as previously reported (Auguet and Casamayor, 2008).

Genetic analyses

Water samples were pre-filtered *in situ* through a 40 μm pore-size net, to exclude large zooplankton and algae, and 300–500 ml were subsequently filtered on 0.2 μm polycarbonate membranes (47 mm diameter, Nuclepore, Whatman Ltd). The filter was stored at –20 °C until further processing. Filters were incubated with lysozyme, proteinase K, and sodium dodecyl sulfate in lysis buffer (40 mM EDTA, 50 mM Tris pH 8.3, 0.75 M sucrose), and DNA was phenol extracted as previously reported (Demergasso *et al.*, 2008).

We initially used denaturing gradient gel fingerprinting (DGGE) on the whole samples set for a rapid screening on the vertical shifts in archaeal 16S rRNA gene diversity (see Auguet *et al.*, 2011 for specific PCR and DGGE conditions). Next, according to the shifts observed in the fingerprints, we selected six

samples to describe the phylogenetic composition and vertical changes by PCR cloning of both the 16S rRNA (general archaeal primer set 21f-958r, 937 bp fragment length) and the archaeal *amoA* (primer set Arch-*amoA*F – Arch-*amoA*R, 635 bp fragment length) genes following previously reported conditions (Auguet *et al.*, 2011 and see also the original references for the primers therein). PCR products were purified with the QIA quick PCR Purification kit (Qiagen, Hilden, Germany) and cloned with the TOPO TA cloning kit (Invitrogen, Life Technologies Co., Carlsbad, CA, USA) following the manufacturer's instructions. The presence of inserts was checked by ampicillin resistance and blue/white selection on LB plates supplemented with ampicillin 100 $\mu\text{g ml}^{-1}$ and X-gal 40 $\mu\text{g ml}^{-1}$ as previously reported (Ferrera *et al.*, 2004). Single-strand sequencing reactions were carried out using external facilities (<http://www.macrogen.com>).

Phylogenetic analysis

The 16S rRNA gene sequences were automatically aligned with the NAST aligner (DeSantis *et al.*, 2006), clustered at 97% identity threshold, and imported into the Greengenes database (<http://greengenes.lbl.gov/>) based on the ARB package (Ludwig *et al.*, 2004; <http://www.arb-home.de>). The Archaea base frequency ARB filter was applied to exclude highly variable positions before sequences were added using the ARB parsimony insertion tool to the original Greengenes tree provided by default.

The *amoA* gene sequences were manually checked with BioEdit (Hall, 1999) and submitted for matching in the protein database using translated nucleotide sequences (BLASTX, www.ncbi.nlm.nih.gov; Altschul *et al.*, 1990) to check for protein identity. Next, sequences were clustered at 99% identity in nucleotides with CD-hit (Li and Godzik, 2006). Multiple sequence alignment, phylogenetic inference by maximum likelihood and calculation of *amoA* gene identity matrices was carried out as recently reported (Auguet *et al.*, 2011).

Quantitative PCR (qPCR)

Abundances of thaumarchaeotal (both MG 1.1a and SAGMGC – 1) 16S rRNA and archaeal *amoA* genes were determined by quantitative real-time PCR amplification. The qPCR assays were run in triplicate on 96 well white qPCR plates with adhesive seals (Bio-Rad, Hercules, CA, USA) in a DNA Engine thermal cycler (Bio-Rad) equipped with a Chromo 4 Real-Time Detector (Bio-Rad) using primers sets (MCGI-391f/MCGI-554r and AOA-*amoA*-f/AOA-*amoA*-r, respectively) and conditions previously reported (Auguet *et al.*, 2011). Primers MCGI-391f/MCGI-554r targeted both MG 1.1a and SAGMGC-1. Additionally, the bacterial *amoA* gene was targeted with primers *amoA*-1F (5'-GGGTTTCT

ACTGGTGGT-3') and *amoA*-2R (5'-CCCCTCKGSAA AGCCTTCTTC-3') (Rotthauwe *et al.*, 1997). However, no amplification for the *amoA* gene of bacteria was obtained with the set of samples tested. Overall, average efficiencies for all quantification reactions ranged from 86% for the archaeal *amoA* to 91% for the 16S rRNA gene with *R* values >0.99. Specificity of reactions was confirmed by melting curve analyses and after looking for unspecific PCR products such as primer dimers or gene fragments of unexpected length in agarose gel electrophoresis.

Diversity indices and statistical analyses

Phylogenetic diversity (PD) was calculated as the sum of the branch lengths associated with the 16S rRNA gene sequences for each sample (Faith, 1992). To correct the unequal number of sequences, we calculated the mean phylogenetic diversity of 1000 randomized subsamples (Barberán and Casamayor, 2010). Chao diversity indices were calculated using the FastGroupII server (http://fastgroup.sdsu.edu/fg_tools.htm). Coverage of libraries was calculated using the diversity tool (<http://www.aslo.org/lomethods/free/2004/0114a.html>) developed by Kemp and Aller (2004). Distance matrices for 16S rRNA and *amoA* genes were constructed with UniFrac (<http://bmf.colorado.edu/unifrac/>) (Lozupone *et al.*, 2006). To assess the sources of variation in the 16S rRNA and *amoA* UniFrac matrices, we carried out permutational multivariate analysis of variance based on 1000 permutations (McArdle, 2001) using the function *adonis* in vegan package (Oksanen *et al.*, 2008). Spearman's rank (r_s) correlations were run to investigate the relationship between environmental parameters and gene abundance. All statistical analyses were run in the R environment (<http://www.r-project.org/>) with the *pvclust* package for cluster analysis.

Nucleotide sequence accession numbers

Sequences were deposited in GenBank under the accession numbers HE589603 to HE589764 for the 16S rRNA gene and HE598566 to HE598696 for the *amoA* gene (see also Supplementary Table S2 for detailed assignment). Gene sequences were deposited following MIENS (Minimal Information about an Environmental Study Standard) and meta-data have the accession numbers ERS054479–ERS054484.

Results

The complete field data set of the vertical changes in temperature, oxygen, DOC, nitrogen compounds and prokaryotes abundance during the annual study can be found in Supplementary Material (Table S1). Water temperatures ranged from 0 (in the slush in winter-spring) to 13 °C (in the summer surface layers). In July–September, the water column was

thermally stratified with the thermocline located between 10 and 35 m. During the ice-cover period, the lake was inversely stratified (warmer water downward). Below 35 m the water column was thermally stable at 4 °C all over the year. Oxygen profiles ranged from 170 to 400 µM throughout the year. Oxygen saturation was found in most of the samples and the lowest concentration was detected close to the bottom in October and April. Chlorophyll *a* (Chl *a*) concentrations ranged from 0.07 to 4.44 µg l⁻¹ and were consistently higher in summer with a deep chlorophyll maximum at 35 m, and a secondary-abundance peak at 10 m. DOC ranged from 12 to 100 µM with higher concentrations in surface waters in summer. Concentrations of ammonium, nitrate and nitrite ranged from 0.08 to 4.55, 5.8 to 11.2 and 0.02 to 0.10 µM, respectively, nitrate being the predominant form of inorganic nitrogen with maximal concentrations in summer around the thermocline. Ammonium concentrations showed higher concentration in bottom waters than in surface with the highest concentration found at the transition from summer to autumn. We also observed ammonium accumulation in the slush layers. Nitrite concentrations showed two maxima at 20 and 60 m during the ice-free period, and an additional maximum in the slush layers during the ice-covered period. Cell abundance ranged from 3 × 10⁵ to 1 × 10⁶ bacterial cells per ml, and from 2 × 10³ to 1 × 10⁵ archaeal cells per ml. These cells abundances were within the same range as those previously reported for Pyrenean lakes (Auguet and Casamayor, 2008).

The DGGE analysis of the archaeal 16S rRNA gene produced a reproducible genetic fingerprint for each sample in different runs. The number of DGGE bands for the whole set of samples ranged between 5 and 12 per sample. Similarity and permutational multivariate analysis of variance analyses made on the manually assigned fingerprint showed 'depth' as a significant source of variation ($R^2 = 0.31$, $P < 0.01$), particularly during the summer period (Figure 1a). Here, we identified two different clusters of samples according to the archaeal community composition, (i) surface samples up to 20 m, and (ii) samples from 20 m to the bottom of the lake. Samples also clustered according to the temporal variations ($R^2 = 0.57$, $P < 0.001$) into two main data sets, one for the freezing–melting period, and another for summer as previously reported in other Pyrenean lakes (Auguet *et al.*, 2011). In fact, the spatio-temporal variations in archaeal diversity mirrored the most significant sources of variation in environmental parameters as determined by a hierarchical clustering and permutational multivariate analysis of variance analysis of the whole environmental data set (Figure 1b) ('depth' $R^2 = 0.22$, $P < 0.01$; 'month' $R^2 = 0.65$, $P < 0.001$). On the basis of these results, we selected samples representative of each period (that is, April and September) and each depth for more detailed genetic analyses




Figure 1 (a) Spatio-temporal clustering according to the similarity matrix generated from archaeal 16S rRNA gene DGGE fingerprints and UPGMA clustering method. Code indicates month.depth. Six selected samples (bold labeling) were further studied by cloning and qPCR analysis. (b) Samples clustering according to changes in environmental parameters (chlorophyll *a*, DOC, nutrients, oxygen concentrations, cells abundance and temperature) based on a hierarchical ascendant classification analysis.

by qPCR and cloning of both 16S rRNA and *amoA* genes.

In April, the lake was thermally mixed with a constant water column temperature of 4 °C below the ice cover. DOC, nitrite, ammonium, Chl *a* and oxygen concentrations gradually decreased with depth although ammonium concentration remarkably increased from 35 to 60 m (Figure 2). The cell abundances (CARD-FISH counts) and quantitative distribution of specific archaeal genes showed little variations with depth, ranging from 3.3×10^2 to 6.7×10^2 16S rRNA gene copies ml⁻¹ and from 1.8×10^3

to 5.6×10^3 *amoA* gene copies ml⁻¹ (Figure 3, black labels for April). Interestingly, the slush layer showed higher nutrients concentrations (particularly enriched in nitrite) and cell abundances than the water column below (Figure 3). This was particularly true for archaea. In the slush, we observed the highest proportion of positive CARD-FISH hybridized archaeal cells (14.5% of total DAPI counts), and the highest concentration of MG 1.1a + SAGMGC – 1 16S rRNA (4.3 times the average value in the water column) and archaeal *amoA* (times 3.5) genes (Figure 3, right panels). In September the lake




Figure 2 Vertical profiles of temperature, DOC, Nitrogen species, Chl *a* and oxygen, along the water column of Lake Redon in April (black symbols) and September (white symbols). The slush sample was sampled in April and the SML (0 m) in September.




Figure 3 Vertical profiles of (i) the percentage of archaea (% ARCH915) versus total cell counts, (ii) the abundance of bacterial and archaeal cells as determined by CARD-FISH and (iii) the concentrations of MG 1.1a + SAGMGC-1 16S rRNA and archaeal *amoA* gene copies as determined by qPCR, along the water column of Lake Redon in April (black symbols) and September (white symbols). The slush sample was sampled in April and the SML (0 m) in September. Standard deviations < 10% applies for all the data.

was thermally stratified and three differentiated water masses were observed along the vertical axis, that is, epilimnion (0–10 m), metalimnion (thermocline, 10–40 m) and hypolimnion (40 m bottom) (Figure 2). The epilimnion showed the lowest proportion of (i) positively hybridized archaeal cells, (ii) cell abundances and (iii) specific archaeal genes (Figure 3, white labels for September). The microbial parameters markedly increased in the metalimnion with the archaeal abundance peak found at 20 m matching the peak of nitrite concentration (Figure 2, Supplementary Table S1). The hypolimnion was enriched in different nitrogen compounds, and the abundances for MG 1.1a + SAGMGC-1 16S rRNA and archaeal *amoA* genes showed a second abundance peak at the bottom of the lake (Figure 3). Overall, only nitrite significantly explained the vertical variability of specific archaeal

gene abundances ($P < 0.05$) after multiple linear regression analysis. In addition, the abundances of the 16S rRNA and *amoA* genes were significantly correlated ($r_s = 0.88$, $P < 0.01$, $n = 12$, slope = 4.8).

As a whole, the qPCR results indicated three potential ammonia oxidation hotspots in the lake, (i) on the top (slush) in winter–spring, (ii) at the thermocline and (iii) at the bottom in summer–autumn. A similar vertical distribution between MG 1.1a + SAGMGC-1 16S rRNA and archaeal *amoA* genes was also observed. In order to unveil the phylogenetic composition for archaeal 16S rRNA and *amoA* genes, and the vertical changes in identity, six samples from three selected depths (top, 20 m, and 60 m) were cloned for April and September. Most of the 16S rRNA gene sequences matched Thaumarchaeota (close to 90%) and only a few Euryarchaeota were detected (Figure 4).




Figure 4 Different analyses carried out with sequence data from the six 16S rRNA clone libraries of Lake Redon in April and September. (a) Hierarchical clustering analysis (UPGMA algorithm with Jackknife supporting values, 100 replicates). Distances between clusters expressed in UniFrac units (distance 0 means identical environments, distance 1 means two environments containing mutually exclusive lineages). (b) Archaeal community structure (relative abundances). Thaumarchaeota Group 1.1a (white label) and SAGMAGC-1 (gray label) are the dominant groups. Other minor thaumarchaeotal/crenarchaeotal groups detected are shown together (black label), as well as the euryarchaeotal groups (shaded label). (c) Collapsed maximum parsimony tree showing the phylogenetic position of the archaeal groups detected. See more details in Figure 5. Scale bar, 10% estimated divergence.

Thaumarchaeota were spread in the clusters MG 1.1a, SAGMAGC-1, pSL12 (mostly found at the bottom waters during the ice-cover period), the soil group 1.1b (found at the SML), and a few sequences from the crenarchaeotal group C2. Overall, the members of SAGMAGC-1 and MG 1.1a accounted for most archaeal 16S rRNA gene sequences (>80%). The euryarchaeotal sequences detected, essentially belonged to the widespread clades HV-Fresh (also named LDS, for Lake Dagow sediment) and PISA1 (also named RC V, for Rice Cluster V) (Auguet *et al.*, 2010; Barberán *et al.*, 2011).

Interestingly, the archaeal assemblages from the two most peculiar samples of this study (that is, winter slush and summer SML) were largely dominated (c. 85–90% of the sequences) by SAGMAGC-1 thaumarchaeotal sequences rooted with the MG 1.1a (89% identity between both the clades) (Figure 4). SAGMAGC-1 was also detected as the predominant clade along the whole water column for most of the samples. Remarkably, sequences from the MG 1.1a were not detected in the slush and SML layers, and its contribution was larger in the summer-stratified hypolimnion. Most of the MG 1.1a sequences were closely related (>97% identity in the 16S rRNA gene sequence) to *Nitrosoarchaeum limnia* (Figure 5, upper clade), a novel AOA recently enriched from low-salinity sediments (Blainey *et al.*, 2011). SAGMAGC-1 sequences, in turn, were closely related (between 99 and 94% identity) to the AOA *Nitrosotalea devanattera* (Lehtovirta-Morley *et al.*, 2011) and to environmental thaumarchaeotal sequences previously found in lakes, aquifers, rhizosphere and rice-paddy soil (Figure 5, lower clade).

For the archaeal *amoA* gene, 130 clones were sequenced (Table 1), which ended in 17 unique operational taxonomic units (OTUs, 5% cutoff ranging from 50 to 94% nucleotide identity).

The recovered *amoA* genes were shared between 60 and 88% nucleotide identity with *Nitrosopumilus maritimus*. Interestingly, we noticed that, within a single lake sample, we could find OTUs diverging up to 38% in nucleotide sequence. Five OTUs represented $80 \pm 20\%$ of the sequences, that is, OTUs numbers 2, 4, 8, 9 and 12 (Figure 6, panels b and c), and up to 98% of the *amoA* gene sequences fell into two previously reported freshwater clusters obtained from Lake Llebrete, a shallow alpine lake belonging to the Limnological Observatory of the Pyrenees (Supplementary Figure. S1). Most of the sequences matched the soil/fresh-1 group (cluster A, in Auguet *et al.*, 2011). In turn, OTU-12 (very abundant at the bottom of the lake in winter-spring, and responsible for splitting this sample from the remaining April samples, Figure 6a) fell into the fresh/low-salinity group (cluster C), which has 95% identity with the *amoA* sequence of *N. limnia*. Probably, OTU-12 belongs to those Archaea from the 1.1a group which were closer to *N. limnia*. Overall, community similarity analysis of the 16S rRNA and *amoA* gene libraries showed consistent vertical differences in the composition of the archaeal sequences ($P < 0.05$, permutational multivariate analysis of variance test). Despite the fact that relative abundances in clone libraries cannot be translated to real abundances, and truly quantitative methods are needed to substantiate and further explore these findings, the set of samples were treated in exactly the same way and, therefore, any bias should apply the same and the comparison would still be valid. In addition, the quite limited number of clones analyzed per sample (Table 1) is in agreement with the high coverage values found (>77% coverage on average for the 16S rRNA gene, and 82% coverage on average for the *amoA* gene, Table 1), and the low number of DGGE bands found for each sample (5–12), indicating a low archaeal richness.

Discussion

Recently described phylogenetic lineages within the domain Archaea are significant components in freshwater plankton (Auguet and Casamayor, 2008; Barberán *et al.*, 2011). Archaeal 16S rRNA and *amoA* genes had been previously reported to be abundant and genetically diverse in surface waters of alpine lakes (Auguet and Casamayor, 2008; Auguet *et al.*, 2011). To understand better, the ecology of these microorganisms, we investigated the relative abundance, distribution and phylogenetic composition along the vertical gradient in a deep oligotrophic alpine lake. The recurrent dominance of AOA lineages and the strong correlations detected between abundances of MG 1.1a + SAGMGC-1 16S rRNA and archaeal *amoA* genes, and nitrite concentrations both in surface waters (Auguet *et al.*, 2011), and along the vertical gradient (present study) suggest a pivotal role of archaea in the N biogeochemistry of oligotrophic high-altitude lakes. In agreement, we did not find either ammonia-oxidizing bacteria (AOB) after specifically looking for the bacterial *amoA* gene, or β - and γ -proteobacterial nitrifiers when looking at the bacterial 16S rRNA gene clone libraries (Hervàs and Casamayor, 2009). This result was expected as ammonium concentrations in the water are mostly below the AOB affinity threshold ($>1 \mu\text{M NH}_4^+$, Bollmann *et al.*, 2002). Therefore, ultra-oligotrophic waters from mountain areas may constitute an ideal habitat for AOA as the low ammonium concentrations *in situ* would select for AOA because of their higher affinity to ammonium ($K_m 130 \text{ nM NH}_4^+$, Martens-Habbena *et al.*, 2009) as compared with AOB.

Obviously, caution is needed before a direct link between *amoA* gene abundance and nitrification activity can be established in absence of bulk nitrification measurements. In some cases it has been reported that the archaeal *amoA* gene could be an evolutionary relic that is not functional anymore (Mussmann *et al.*, 2011). Nonetheless, in Lake Redon the match found between the vertical abundance of nitrite and peaks in the abundance of both *amoA* and 16S rRNA genes strongly suggests potential active ammonium oxidation zones related to archaea. Intriguingly, abundances of the 16S rRNA and the *amoA* genes differ markedly. Studies in marine environments reported an archaeal *amoA* to 16S ratio of 0.5:1 and lower (De Corte *et al.*, 2009). In Lake Redon ratios of 5:1–8:1 were observed, suggesting that either not all 16S genes of *amoA* gene containing organisms were detected with the qPCR assay or that the archaea present in the lake have a higher copy number of *amoA* genes per genome.

Interestingly, we unveiled ammonium oxidation zones changing vertically with seasons. In summer–autumn, nitrification at the bottom of the lake could be related to the ammonium produced in the




Figure 5 Maximum-parsimony tree of the 16S rRNA gene sequences from the Marine 1.1a and SAGMGC-1 groups obtained along the water column of Lake Redon (in bold label) in April and September. GenBank accession numbers are shown for each sequence. Scale bar, 10% estimated divergence.




Figure 6 Different analyses carried out with sequence data from the six *amoA* clone libraries of Lake Redon in April and September as in Figure 4. (a) Hierarchical clustering analysis. (b) AOA community structure presented by relative abundances of the five most represented OTUs (i.e., OTUs 2, 4, 8, 9 and 12, see labels and number of sequences within each group in panel c) whereas the remaining OTUs are condensed (black bars). (c) Collapsed maximum likelihood tree inferred by RaxML and based on 579 nucleotides positions. See more details in Supplementary Figure. S1. Scale bar, 10% estimated divergence.

sediment by microbial decomposition of organic material from early summer phytoplankton blooms (Catalan *et al.*, 2006), and the oxygen-minimum zone observed. In winter–spring, microbial activities and biomasses are by far larger in the slush layers generated by episodic events of snow deposition, flooding by lake water, melting, freezing and atmospheric depositions (that is rain or snow) (Felip *et al.*, 1995) than in the water column. Atmospheric depositions are the main source of reactive nitrogen in remote mountain catchments (Catalan *et al.*, 1994), particularly ammonium with concentrations in the rain up to 30fold higher than mean values measured in the lake water (Auguet *et al.*, 2011). High influence of airborne dust nutrients has also been reported (Hervàs *et al.*, 2009; Reche *et al.*, 2009) that may fuel AOA nitrification activity. These results reflect the wide set of ecological and thermal conditions where AOA are distributed, ranging from hot springs (Zhang *et al.*, 2008) to the ice layers reported here. Intriguingly, the presence of a potential nitrification zone at 20 m is more difficult to understand because both photoinhibition (Merbt *et al.*, 2012) and preference for a dark environment (Church *et al.*, 2010; Merbt *et al.*, 2011) has been recently reported for some types of AOA.

Another interesting finding of this work is the high relative abundance and high frequency of appearance of the SAGMGC – 1 Thaumarchaeota, a sister clade to the marine 1.1 a clade, originally discovered in South Africa gold mines (Takai *et al.*, 2001). This result suggests a significant environmental role for the SAGMGC – 1 clade in alpine lakes. Previous reports showed that the members of the SAGMGC – 1 clade to be mainly found in freshwater systems, for example, alpine lakes (Auguet and Casamayor, 2008; Auguet *et al.*, 2011) groundwater (Shimizu *et al.*, 2007), aquifers (López-Archilla *et al.*, 2007) and terrestrial habitats such as rhizospheres (Herrmann *et al.*, 2008) and rice-paddy soils (Sakai *et al.*, 2007). Whereas members of the MG 1.1a have been largely implicated in

nitrification (Konneke *et al.*, 2005; Hallam *et al.*, 2006; Blainey *et al.*, 2011), for the SAGMGC – 1 clade only a very recent study indicates that a cultured strain from an acid soil, *Nitrosotalea devanattera*, codes for the *amoA* gene (Lehtovirta-Morley *et al.*, 2011). Thus, several lines of evidence may support ammonium oxidation potential in freshwater SAGMGC – 1. First, SAGMGC-1 belongs to the Thaumarchaeota phylum in which all the cultivated or enriched members contain the *amoA* gene, and more precisely are closely related to the AOA *N. devanattera*. Second, we noticed substantial abundance of *amoA* gene in a sample (that is, the slush), where the only thaumarchaeal group observed by 16S rRNA gene sequencing was SAGMGC – 1. And third, the identity of the *amoA* gene found in the slush sample was associated with the previously reported soil/fresh 1 – *amoA* cluster A (Auguet *et al.*, 2011). The *amoA* cluster A is mainly formed by sequences from freshwater origin not yet affiliated to any 16S rRNA-defined population and interestingly fits with the *Nitrosotalea* subcluster 1 (Pester *et al.*, 2012). Unfortunately, these are indirect findings yet and further work and interdisciplinary approaches are needed to probe that the freshwater SAGMGC – 1 members are AOA.

Field studies may provide still helpful insights into the ecology of uncultured SAGMGC – 1. In a previous work carried out in 10 alpine lakes, we unveiled the neuston as a selective environment where members of SAGMGC – 1 are naturally enriched (Auguet and Casamayor, 2008), in agreement with the result found here. Remarkably, in an annual survey in three connected shallow alpine lakes made in parallel to the present work, we found surface archaeal assemblages dominated by the MG1.1a all around the year (Auguet *et al.*, 2011), although SAGMGC – 1 was also detected. In the deep Lake Redon, the two clades segregated inversely along the water column with the relative abundance of SAGMGC – 1 vertically decreasing whereas that of MG 1.1a increased. Although caution is needed to translate relative abundances

of clones in the clone library to real cell abundance distributions, data suggest that the vertical gradient is as a key factor to understand the ecology of both the groups. In fact, the community similarity analysis indicated archaeal assemblage composition to be significantly related to depth. There are, however, many physical, chemical and biological variables that change with depth, and high-altitude oligotrophic lakes are especially affected by high irradiances and UV penetration, with varying effects associated with altitude (Catalan *et al.*, 2006). It has been reported that parameters such as DOC and oxygen concentrations are potential environmental factor shaping the ecological niches of AOA (Erguder *et al.*, 2009). The recently reported photo-inhibition in AOA (Merbt *et al.*, 2012) may provide additional hypothesis to explore differential tolerances to irradiance among clades, as diverse marine planktonic AOA ecotypes (that is, shallow and deep) (Francis *et al.*, 2005; Hallam *et al.*, 2006; Mincer *et al.*, 2007; Beman *et al.*, 2008) may be related to the differential resistance and adaptations to light. The vertical distribution of the MG 1.1a observed in Lake Redon was most closely related to that found for this clade in oceans (that is, below the euphotic zone) and may be associated to a lower tolerance to UV radiations and high light intensities. Conversely, the presence of SAGMGC – 1 in surface waters and particularly at the air–water interface suggests a better adaptation to solar irradiance. Nevertheless, different metabolic adaptations and environmental distribution may be contained in the different clusters observed within each clade that would require finer phylogenetic resolution. Altogether, both altitudinal and vertical gradients in high mountain lake districts provide an excellent framework to further explore the ecology and biology of the idiosyncratic AOA populations.

Acknowledgements

We are thankful to the Centre de Recerca d'Alta Muntanya (CRAM), Universitat de Barcelona, Vielha for laboratory facilities. C Gutiérrez-Provecho, A Barberán, M Bacardit and L Alonso-Sáez are acknowledged for field and laboratory assistance and J Catalan for scientific feedback and support. We are also thankful for the constructive comments provided by the reviewers and the editor. This research was supported by grants CRENYC CGL2006-12058 and PIRENA CGL2009-13318 to EOC, and CONSOLIDER grant GRACCIE CSD2007-00067 from the Spanish Office of Science and Innovation (MICINN). JCA benefits from a Juan de la Cierva postdoctoral fellow (MICINN).

References

Agogué H, Brink M, Dinasquet J, Herndl GJ. (2008). Major gradients in putatively nitrifying and non-nitrifying Archaea in the deep North Atlantic. *Nature* **456**: 788–792.

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic Local Alignment Search Tool. *J Mol Biol* **215**: 403–410.
- Auguet JC, Barberán A, Casamayor EO. (2010). Global ecological patterns in uncultured Archaea. *ISME J* **4**: 182–190.
- Auguet JC, Casamayor EO. (2008). A hotspot for cold crenarchaeota in the neuston of high mountain lakes. *Environ Microbiol* **10**: 1080–1086.
- Auguet JC, Nomokonova N, Camarero L, Casamayor EO. (2011). Seasonal changes of freshwater ammonia-oxidizing archaeal assemblages and nitrogen species in oligotrophic alpine lakes. *Appl Environ Microbiol* **77**: 1937–1945.
- Barberán A, Casamayor EO. (2010). Global phylogenetic community structure and beta-diversity patterns of surface bacterioplankton metacommunities. *Aquat Microb Ecol* **59**: 1–10.
- Barberán A, Fernandez-Guerra A, Auguet J-C, Galand PE, Casamayor EO. (2011). Phylogenetic ecology of widespread uncultured clades of the Kingdom Euryarchaeota. *Mol Ecol* **20**: 1988–1996.
- Beman JM, Popp BN, Francis CA. (2008). Molecular and biogeochemical evidence for ammonia oxidation by marine Crenarchaeota in the Gulf of California. *ISME J* **2**: 429–441.
- Blainey PC, Mosier AC, Potanina A, Francis CA, Quake SR. (2011). Genome of a low-salinity ammonia-oxidizing archaeon determined by single-cell and metagenomic analysis. *PLoS One* **6**: e16626.
- Bollmann A, Bar-Gilissen MJ, Laanbroek HJ. (2002). Growth at low ammonium concentrations and starvation response as potential factors involved in niche differentiation among ammonia-oxidizing bacteria. *Appl Environ Microbiol* **68**: 4751–4757.
- Catalan J, Camarero L, Gacia E, Ballesteros E, Felip M. (1994). Nitrogen in the Pyrenean lakes (Spain). *Hydrobiologia* **274**: 17–27.
- Catalan J, Camarero L, Felip M, Pla S, Ventura M, Buchaca T *et al.* (2006). High mountain lakes: extreme habitats and witnesses of environmental changes. *Limnetica* **25**: 551–584.
- Church MJ, Wai B, Karl DM, DeLong EF. (2010). Abundances of crenarchaeal *amoA* genes and transcripts in the Pacific Ocean. *Environ Microbiol* **12**: 679–688.
- De Corte D, Yokokawa T, Varela MM, Agogue H, Herndl GJ. (2009). Spatial distribution of bacteria and Archaea and *amoA* gene copy numbers throughout the water column of the eastern Mediterranean Sea. *ISME J* **3**: 147–158.
- Demergasso C, Escudero L, Casamayor EO, Chong G, Balagué V, Pedrós-Alió C. (2008). Novelty and spatio-temporal heterogeneity in the bacterial diversity of hypersaline Lake Tebenquiche (Salar de Atacama). *Extremophiles* **12**: 491–504.
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K *et al.* (2006). Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* **72**: 5069–5072.
- Doble P, Haddad PR. (1999). Indirect photometric detection of anions in capillary electrophoresis. *J Chromatogr* **A834**: 189–212.
- Erguder T.H, Boon N, Wittebolle L, Marzorati M, Verstraete W. (2009). Environmental factors shaping the ecological niches of ammonia-oxidizing Archaea. *FEMS Microbiol Rev* **33**: 855–869.

- Faith DP. (1992). Conservation evaluation and phylogenetic diversity. *Biol Conserv* **61**: 1–10.
- Felip M, Sattler B, Psenner R, Catalan J. (1995). Highly active microbial communities in the ice and snow cover of high mountain lakes. *Appl Environ Microbiol* **61**: 2394–2401.
- Ferrera I, Massana R, Casamayor EO, Balague V, Sanchez O, Pedros-Alio C *et al.* (2004). High-diversity biofilm for the oxidation of sulfide-containing effluents. *Appl Microbiol Biotechnol* **64**: 726–734.
- Francis CA, Roberts KJ, Beman JM, Santoro AE, Oakley BB. (2005). Ubiquity and diversity of ammonia-oxidizing Archaea in water columns and sediments of the ocean. *Proc Natl Acad Sci USA* **102**: 14683–14688.
- Galand PE, Lovejoy C, Vincent WF. (2006). Remarkably diverse and contrasting archaeal communities in a large arctic river and the coastal Arctic Ocean. *Aquat Microb Ecol* **44**: 115–126.
- Galand PE, Gutierrez-Provecho C, Massana R, Gasol JM, Casamayor EO. (2010). Inter-annual recurrence of archaeal assemblages in the coastal NW Mediterranean Sea (Blanes Bay Microbial Observatory). *Limnol Oceanogr* **55**: 2117–2125.
- Hall TA. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* **41**: 95–98.
- Hallam SJ, Mincer TJ, Schleper C, Preston CM, Roberts K, Richardson PM *et al.* (2006). Pathways of carbon assimilation and ammonia oxidation suggested by environmental genomic analyses of marine Crenarchaeota. *PLoS Biol* **4**: 520–536.
- Herfort L, Kim JH, Coolen MJL, Abbas B, Schouten S, Herndl GJ *et al.* (2009). Diversity of Archaea and detection of crenarchaeotal *amoA* genes in the rivers Rhine and Tet. *Aquat Microb Ecol* **55**: 189–201.
- Herrmann M, Saunders AM, Schramm A. (2008). Archaea dominate the ammonia-oxidizing community in the rhizosphere of the freshwater macrophyte *Littorellau-niflora*. *Appl Environ Microbiol* **74**: 3279–3283.
- Hervàs A, Casamayor EO. (2009). High similarity between bacterioneuston and airborne bacterial community compositions in a high mountain lake area. *FEMS Microbiol Ecol* **67**: 219–228.
- Hervàs A, Camarero L, Reche I, Casamayor EO. (2009). Viability and potential for immigration of airborne bacteria from Africa that reach high mountain lakes in Europe. *Environ. Microbiol* **11**: 1612–1623.
- Kemp PF, Aller JY. (2004). Estimating prokaryotic diversity: when are 16S rDNA libraries large enough? *Limnol Oceanogr-Meth* **2**: 114–125.
- Konneke M, Bernhard AE, de la Torre JR, Walker CB, Waterbury JB, Stahl DA. (2005). Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* **437**: 543–546.
- Lam P, Jensen MM, Lavik G, McGinnis DF, Muller B, Schubert CJ *et al.* (2007). Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea. *Proc Natl Acad Sci USA* **104**: 7104–7109.
- Lam P, Lavik G, Jensen MM, van de Vossenberg J, Schmid M, Woebken D *et al.* (2009). Revising the nitrogen cycle in the Peruvian oxygen minimum zone. *Proc Natl Acad Sci USA* **106**: 4752–4757.
- Lehtovirta-Morley LE, Stoecker K, Vilcinskis A, Prosser JI, Nicol GW. (2011). Cultivation of an obligate acidophilic ammonia oxidizer from a nitrifiant acid soil. *Proc Natl Acad Sci USA* **108**: 15892–15897.
- Li W, Godzik A. (2006). Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.
- Llirós M, Casamayor EO, Borrego CM. (2008). High archaeal richness in the water column of a freshwater sulfurous karstic lake along an interannual study. *FEMS Microbiol. Ecol* **66**: 331–342.
- Llirós M, Gich F, Plasencia A, Auguet JC, Darchambeau F, Casamayor EO *et al.* (2010). Vertical distribution of ammonia oxidizing crenarchaeota and methanogens in the epipelagic waters of Lake Kivu. *Appl Environ Microbiol* **76**: 6853–6863.
- Llorens-Marès T, Auguet JC, Casamayor EO. (2012). Winter to Spring changes in the slush bacterial community composition of a high mountain lake (Lake Redon, Pyrenees). *Environ Microbiol Rep* **4**: 50–56.
- López-Archilla AI, Moreira D, Velasco S, López-García P. (2007). Archaeal and bacterial community composition of a pristine coastal aquifer in Doñana National Park, Spain. *Aquat Microb Ecol* **47**: 123–139.
- Lozupone C, Hamady M, Knight R. (2006). UniFrac—An online tool for comparing microbial community diversity in a phylogenetic context. *BMC Bioinformatics* **7**: 371.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar *et al.* (2004). ARB: a software environment for sequence data. *Nucl Acids Res* **32**: 1363–1371.
- Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR, Stahl DA. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* **461**: 976–979.
- McArdle B. (2001). Fitting multivariate models to community data: a comment on distance-based redundancy analysis. *Ecology* **82**: 290–297.
- Merbt S, Auguet JC, Casamayor EO, Martí E. (2011). Biofilm recovery in a wastewater treatment plant-influenced stream and spatial segregation of ammonia-oxidizing microbial populations. *Limnol Oceanogr* **56**: 1054–1064.
- Merbt S, Stahl DA, Casamayor EO, Martí E, Nicol GW, Prosser JI. (2012). Differential photoinhibition of bacterial and archaeal ammonia oxidation. *FEMS Microbiol Lett* **327**: 41–46.
- Mincer TJ, Church MJ, Taylor LT, Preston C, Kar DM, DeLong EF. (2007). Quantitative distribution of presumptive archaeal and bacterial nitrifiers in Monterey Bay and the North Pacific Subtropical Gyre. *Environ Microbiol* **9**: 1162–1175.
- Mussmann M, Brito I, Pitcher A, Sinninghe Damsté JS, Hatzenpichler R, Richter A *et al.* (2011). Thaumarchaeotes abundant in refinery nitrifying sludges express *amoA* but are not obligate autotrophic ammonia oxidizers. *Proc Natl Acad Sci USA* **108**: 16771–16776.
- Oksanen J, Kindt R, Legendre P, O'hara B, Simpson GL, Stevens MH. (2008). *Vegan: community Ecology Package, version 1*: 11–14; <http://vegan.r-forge.r-project.org>.
- Pester M, Rattei T, Flechl S, Gröngröft A, Richter A, Overmann J *et al.* (2012). AmoA-based consensus phylogeny of ammonia-oxidizing archaea and deep sequencing of *amoA* genes from soils of four different geographic regions. *Environ Microbiol* **14**: 525–539.

- Pouliot J, Galand PE, Lovejoy C, Vincent WF. (2009). Vertical structure of archaeal communities and the distribution of ammonia monooxygenase A gene variants in two meromictic High Arctic lakes. *Environ Microbiol* **11**: 687–699.
- Reche I, Ortega-Retuerta E, Romera O, Pulido-Villena E, Morales-Baquero R, Casamayor EO. (2009). Effect of Saharan dust inputs on bacterial activity and community composition in Mediterranean lakes and reservoirs. *Limnol Oceanogr* **54**: 869–879.
- Reed D.W, Smith JM, Francis CA, Fujita Y. (2009). Responses of ammonia-oxidizing bacterial and archaeal populations to organic nitrogen amendments in low-nutrient groundwater. *Appl Environ Microbiol* **76**: 2517–2523.
- Rotthauwe JH, Witzel KP, Liesack W. (1997). The ammonia monooxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Appl Environ Microbiol* **63**: 4704–4712.
- Sakai S, Imachi H, Sekiguchi Y, Ohashi A, Harada H, Kamagata Y. (2007). Isolation of key methanogens for global methane emission from rice paddy fields: a novel isolate affiliated with the clone cluster rice cluster I. *Appl Environ Microbiol* **3**: 4326–4331.
- Shimizu S, Akiyama M, Naganuma T, Fujioka M, Nako M, Ishijima Y. (2007). Molecular characterization of microbial communities in deep coal seam groundwater of northern Japan. *Geobiology* **5**: 423–433.
- Solórzano L. (1969). Determination of ammonia in natural waters by the phenylhypochlorite method. *Limnol Oceanogr* **14**: 799–801.
- Takai K, Moser DP, DeFlaun M, Onstott TC, Fredrickson JK. (2001). Archaeal diversity in waters from deep South African gold mines. *Appl Environ Microbiol* **67**: 5750–5760.
- Treusch AH, Leininger S, Kletzin A, Schuster SC, Klenk HP, Schleper C. (2005). Novel genes for nitrite reductase and Amo-related proteins indicate a role of uncultivated mesophilic crenarchaeota in nitrogen cycling. *Environ Microbiol* **7**: 1985–1995.
- van der Wielen P, Voost S, van der Kooij D. (2009). Ammonia-oxidizing bacteria and archaea in groundwater treatment and drinking water distribution systems. *Appl Environ Microbiol* **75**: 4687–4695.
- Venter JC, Remington K, Heidelberg JF, Halpern AL, Rusch D, Eisen JA *et al*. (2004). Environmental genome shotgun sequencing of the Sargasso Sea. *Science* **304**: 66–74.
- Zhang CL, Ye Q, Huang Z, Li W, Chen J, Song Z *et al*. (2008). Global occurrence of archaeal *amoA* genes in terrestrial hot springs. *Appl Environ Microbiol* **74**: 6417–6426.

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)