

Vibrational Spectra of Tetrafluoroethylene and Tetrachloroethylene^{1, 2}

D. E. Mann, Nicolo Acquista, and Earle K. Plyler

The infrared spectra of gaseous tetrafluoroethylene from 22 to 52 μ and liquid tetrachloroethylene from 3 to 52 μ have been determined. Several hitherto unobserved bands were found. New and satisfactory assignments, for which a detailed discussion is given, have been achieved. The out-of-plane force constants are discussed and used to predict the wagging and torsion frequencies of tetrabromoethylene. Tables of the thermodynamic functions for tetrafluoro- and tetrachloroethylene are presented.

1. Introduction

As part of a more general investigation of the vibrational spectra and force constants of halogenated ethylenes, it was found desirable to reexamine the assignments for tetrafluoroethylene and tetrachloroethylene.³ Despite a considerable body of earlier work, several fundamentals in each molecule have remained in doubt or unobserved. The availability of a cesium iodide prism [1]⁴ has made convenient the extension of the observable spectral range to about 52 μ , with the result for TFE and TCE that several hitherto unsuspected and otherwise unattainable bands have been revealed. These new data, in conjunction with the recently determined value of the entropy of TFE [2] and some preliminary force-constant calculations, indicate the need, and provide the basis, for achieving more satisfactory assignments.

Although TCE has been subjected to much more spectroscopic investigation, analysis, and discussion than TFE, its assignment has remained the more questionable. This seems to have been due, at least in part, to the greater inaccessibility of the low-frequency infrared-active fundamentals of TCE. Its successful analysis may also have been impeded by the lack of adequate data for other molecules closely enough related to TCE to render some correlations useful. Tetrafluoroethylene bears a sufficiently close resemblance, both spectrally and structurally, to its chlorine analogue to make it profitable to consider them together. Moreover, the entropy data lend additional credibility to the present TFE assignment, so that its use as a guide may be at least partially justified.

2. Experimental Methods

The general method of measurement has been described previously [3, 4]. To ensure adequate resolution over the range 3 to about 52 μ , prisms of NaCl, KBr, and CsI were employed in appropriate regions. The first two were used in Perkin-Elmer model 21 and Baird Associates instruments, respectively, and the

cesium iodide prism was used with a Perkin-Elmer model 12C spectrometer.

A pure sample of gaseous TFE was examined only in the region 22 to 52 μ because the earlier investigation by Nielsen, Claassen, and Smith [5] of its infrared spectrum between 2 and 22 μ is satisfactory. Recent studies of the infrared spectrum of TCE have been made by Plyler [6], who reported four bands in the region 23 to about 40 μ , and by Bernstein [7], who investigated the region between 2.5 and 25 μ . It was thought desirable to reexamine the previously studied range, using a highly purified sample of TCE, as well as to extend it to 52 μ . There are some discrepancies between the spectrum presented here and that displayed in [7]. These may be largely due to differences in purity and resolution. The sample of TCE used in this work had been specially purified by distillation through a 40-plate Pirox-Glover spinning band still, and by passage through a 10-in. column of silica gel.

The spectral curves for TFE and TCE are displayed in figures 1 and 2, respectively. A complete tabulation of the observed data for the chloro compound, together with its assignment, is given in table 1.

3. Discussion

3.1. Spectra of Tetrafluoroethylene

The infrared spectrum of gaseous TFE has been examined by Torkington and Thompson [8], and more thoroughly by Nielsen, Claassen, and Smith [5] from 2 to 22 μ . The region between 22 and 37 μ was covered by Plyler with the aid of a KRS-5 prism [5]. Raman spectra of gaseous TFE, including depolarization factors, were also obtained by Nielsen, Claassen, and Smith. Monfils and Duchesne [9] have reported Raman spectra for the liquid but did not give polarization data.

The infrared band at 406 cm^{-1} observed by Plyler, was reported in [5] to have a type C contour. From figure 1 it is evident that this observation is confirmed despite the appearance of an overlapped band in the *R* branch. Apart from the very weak peak adjacent to the *P* branch of the 406 cm^{-1} band, the intermediate region is quite barren until the strong absorption around 218 cm^{-1} is reached. The resolu-

¹ Presented before the American Physical Society at its meeting in Washington, D. C., on May 1, 1953.

² This work has been supported in part by the ONR under contract NAomr 112-51.

³ Hereafter in this paper tetrafluoroethylene, $\text{F}_2\text{C}=\text{CF}_2$, will be abbreviated to TFE, and tetrachloroethylene, $\text{Cl}_2\text{C}=\text{CCl}_2$, will be contracted to TCE.

⁴ Figures in brackets indicate the literature references at the end of this paper.

TABLE 1. Infrared and Raman spectra of Cl₂C:CCl₂ (liquid)

Raman		Infrared ^a		Interpretation
ν	(I, ρ)	ν	I	
cm^{-1}		cm^{-1}		
218	(00,-)	224	w	$2 \times 110 = 220(A_g)$ $\nu_7(b_{1u}; \beta\text{-CCl}_2)$
235	(10,0.62)	288	w	$\nu_3(a_g; \delta\text{-CCl}_2)$
		310	vw	$\nu_{12}(b_{3u}; \delta\text{-CCl}_2)$
		328	vw	$288 + \nu_i - \nu_i$
		345	vw	$777 - 447 = 330(B_{3u})$ $(\nu_6)^b$
^b 347	(4,0.83)	405	vw	$\nu_8(b_{1g}; \rho\text{-CCl}_2)$ $176 + 235 = 411(B_{2u})$
		433	vw	$777 - 347 = 430(B_{2u})$
447	(15,0.14)	465	vw	$\nu_2(a_g; \nu\text{-CCl})$ $2 \times 235 = 470(A_g)$
464	(1/2,-)			$224 + 235 = 459(B_{1u}); 908 - 447 = 461(B_{2u})$
512	(4,0.77)	513	vw	$\nu_8(b_{2g}; \beta\text{-CCl}_2)$ $176 + 347 = 523(B_{3u})$
		526	vw	$235 + 288 = 523(B_{3u})$
		540	vw	$777 - 235 = 542(B_{3u})$
		559	vw	$908 - 347 = 561(B_{3u})$
574	(0,-)	620	vw	$2 \times 288 = 576(A_g)$ $176 + 447 = 623(B_{2u})$
631	(00,-)	641	vw	$908 - 288 = 620(B_{1g})$ $288 + 347 = 635(B_{2u})$
		679	w	$224 + 447 = 671(B_{1u})$
726	(00,-)	755	s	$2 \times 347 = 694(A_g); 235 + 512 = 747(B_{2g})$
		777	vs	$224 + 512 = 736(B_{3u}); 288 + 447 = 735(B_{3u})$ $\nu_{11}(b_{3u}; \nu\text{-CCl})$
784	(00,-)	800	s	$347 + 447 = 794(B_{1g})$ $288 + 512 = 800(B_{1u})$
		908	vvs	$\nu_9(b_{2u}; \nu\text{-CCl})$
		979	m	$224 + 235 + 512 = 971(B_{3u})$
		994	m	$235 + 777 = 1012(B_{3u})$
1000	(00,-)	1102	m	$\nu_5(b_{1g}; \nu\text{-CCl})$ $2 \times 512 = 1024(A_g)$
1025	(1,p)	1121	m	$224 + 347 + 512 = 1083(B_{2u})$
		1142	m	$347 + 777 = 1124(B_{2u})$
		1171	w	$235 + 908 = 1143(B_{2u})$ $176 + 1000 = 1176(B_{3u})$
		1200	w	$224 + 447 + 512 = 1183(B_{3u})$
		1222	w	$447 + 777 = 1224(B_{3u})$
		1253	w	$347 + 908 = 1255(B_{3u})$
		1355	m	$447 + 908 = 1355(B_{2u})$
		1392	vw	$176 + 235 + 1000 = 1411(B_{3u})$
1441	(0?, -)	1481	vw	$447 + 1000 = 1447(B_{1g})$ $2 \times 288 + 908 = 1484(B_{2u})$
1571	(10,0.36)	1577	w	$\nu_1(a_g; \nu\text{-CC})$ $347 + 447 + 777 = 1571(B_{2u})$
		1733	vw	$288 + 447 + 1000 = 1735(B_{2u})$
		1751	w	$176 + 1571 = 1747(B_{2u})$
		1773	w	$777 + 1000 = 1777(B_{2u})$
		1795	vw	$224 + 1571 = 1795(B_{1u})$
1819	(0,-)	1862	w	$2 \times 908 = 1816(A_g)$ $288 + 1571 = 1859(B_{3u})$
		1887	w	$908 + 1000 = 1908(B_{3u})$
1998	(1/2,-)	2475	w	$2 \times 1000 = 2000(A_g)$ $908 + 1571 = 2479(B_{2u})$
		2717	vw	$235 + 908 + 1571 = 2714(B_{2u})$
		2755	vw	$777 + 2 \times 1000 = 2777(B_{3u})$
		2890	vw	$908 + 2 \times 1000 = 2908(B_{2u})$
		3448	vw	$908 + 1000 + 1571 = 3479(B_{3u})$




FIGURE 1. Infrared spectrum of gaseous $F_2C:CF_2$ from 22 to 52 μ .




FIGURE 2. Infrared spectrum of liquid $Cl_2C:CCl_2$ from 2 to 52 μ .

^a The infrared bands at 176 and 152 cm^{-1} observed by Pitzer (see footnote 6) may be interpreted as $\nu_{10}(b_{2g}; \rho-CCl_2)$, and $\nu_{10} + \nu_2 - \nu_4$, respectively. Before Professor Pitzer's results were made known to present authors, the ν_{10} fundamental used in this table had been assigned the value ~ 170 cm^{-1} . This has now been corrected to 176 cm^{-1} to accord with the new data.

^b For lack of a better explanation, the 347 cm^{-1} infrared band is interpreted as ν_8 appearing in virtue of the partial breakdown of selection rules in the liquid. It should be noticed that there are a number of other coincidences between infrared and Raman frequencies: 464 and 465, 512 and 513, 1000 and 994, 1571 and 1577. Fortunately, these can be explained otherwise.

tion of the cesium iodide prism is not sufficient to reveal much structure in this region, and the bumps on the low-frequency shoulder should be regarded with some scepticism. Though they appear to be real, they may derive at least in part from some uncertainty in the background-absorption correction.

The band at 218 cm^{-1} undoubtedly represents the hitherto unobserved b_{2u} rocking fundamental. It had been placed at 245 cm^{-1} by Nielsen, Claassen, and Smith on the basis of their interpretation of several combination bands. A value of 250 cm^{-1} was deduced by Monfils and Duchesne [9] from a force-constant analysis. Inasmuch as 406 and 558 cm^{-1} have been unambiguously established as representing the b_{1u} wagging and b_{3u} deformation fundamentals, respectively, only the inactive torsion and the Raman-active vibrations need be considered in order to complete the assignment.

The polarization measurements of Nielsen, Claassen, and Smith leave no doubt as to the correctness of their assignment of 394, 778, and 1872 cm^{-1} to the totally symmetric (a_g) vibrations. Moreover, their interpretation of the $503\text{--}517\text{ cm}^{-1}$ doublet as the b_{2g} fundamental, and their deduction of a frequency around 550 cm^{-1} for the lower of the b_{1g} vibrations is supported by the work of Monfils and Duchesne [9] on the Raman spectrum of liquid TFE. Although the latter workers originally questioned [9] the assignment by Nielsen, Claassen, and Smith of a very feeble line at about 1340 cm^{-1} to b_{1g} , they appear now [10] to have accepted this interpretation. The present analysis also favors 1340 cm^{-1} for the higher b_{1g} vibration. Only the inactive torsion frequency remains unassigned.

3.2. Torsion Frequency of Tetrafluoroethylene

In order to obtain a rough estimate of the torsion frequency, the assumption was made that the force constant for the twisting motion in TFE is the same as that obtained by Arnett and Crawford [11] for ethylene, $\text{H}_2\text{C}=\text{CH}_2$.⁵ In units of $10^{-11}\text{ erg/radian}^2$ the constant obtained by Arnett and Crawford is $F_\tau=0.269$. With the distances and angles appropriate to TFE [13], this leads to a value of 199 cm^{-1} for the torsion frequency. Hence, the interesting possibility exists that one of the peaks around 205 cm^{-1} in the infrared spectrum of TFE may arise from the torsion vibration made active by Coriolis interaction with the neighboring b_{2u} fundamental at 218 cm^{-1} . Attempts to determine the torsion frequency directly from the observed combination bands tend to support a value in the range 190 to 200 cm^{-1} . Fortunately, the entropy of TFE has been determined very recently [2] and provides valuable evidence in favor of the complete assignment, while placing the torsion frequency at 190 cm^{-1} . Small inaccuracies in the fundamental frequencies, together with the experimental uncertainties in the entropy value, may account for the slight discrepancy. Because of the lack of stronger evidence favoring the higher value, 190 cm^{-1} was

adopted for the torsion frequency. The corresponding force constant then becomes $F_\tau=0.245$. The final and complete assignment is given in table 3.

3.3. Spectra of Tetrachloroethylene

The infrared spectrum of liquid TCE was first studied by Coblenz [14]. Bonino examined a small portion of its spectrum in 1925 [15], and Spence and Easley observed the region between 0.8 and 3μ but found no bands [16]. Another early study of its near infrared spectrum was made by Freymann [17]. In 1934 Wu reported four bands, $755(\text{m})$, $782(\text{s})$, $802(\text{s})$, $913(\text{s})$, for gaseous TCE [18]. A few years later Duchesne and Parodi reported infrared bands at 332 and 387 cm^{-1} [19]. Bernstein has observed the spectrum of TCE from 2.5 to 25μ [7], and Plyler has extended the range to about 40μ [6]. In the present work the entire region 3 to 52μ has been investigated.⁶

The Raman spectrum of liquid TCE has been extensively investigated [20 to 31]. With the exception of the recent work of Sanyal [31], the available data have been considerably augmented and carefully reviewed by Wittek [29]. Summaries have also been given by Kohlrusch [32], Wu [33], and Herzberg [34]. The frequencies, relative intensities, and depolarization factors⁷ given by Wittek are: $218(00,-)$, $235(10,0.62)$, $347(4,0.83)$, $447(15,0.14)$, $464(\frac{1}{2},-)$, $512(4,0.77)$, $574(0,-)$, $631(00,-)$, $726(00,-)$, $784(00,-)$, $1000(00,-)$, $1025(1,\text{p})$, $1441(0?,-)$, $1571(10,0.36)$, $1819(0,-)$, $1998(\frac{1}{2},-)$. Sanyal [31] reported for liquid TCE $238(7,\text{dp})$, $346(4,\text{dp})$, $382(\text{Ob},-)$, $450(9,\text{p})$, $516(3,\text{dp})$, $1572(10,\text{dp})$; and for the solid at -150° C , he found $242(2,-)$, $346(1,-)$, $380(0,-)$, $454(1,-)$, $1576(58,-)$. The line at 382 cm^{-1} found by Sanyal was also reported by Wu [27] but was shown by Wittek to be spurious. Moreover, the depolarized character ascribed by Sanyal to the 1572 cm^{-1} line is in disagreement with all prior polarization measurements [25 to 27, 29]. Wittek's results are used in this paper.

The infrared spectrum of liquid TCE is shown in figure 2. The wave numbers and relative intensities of the observed bands, together with Wittek's Raman data and the present interpretation, are given in table 1.

3.4. Interpretation of the Vibrational Spectra Tetrachloroethylene

The Raman data for TCE clearly prescribe the three a_{1g} fundamentals: 235 , 447 , and 1571 cm^{-1} . These correspond approximately to the totally symmetric CCl_2 deformation ($\delta\text{-CCl}_2$), CCl stretching ($\nu\text{-CCl}$), and CC stretching ($\nu\text{-CC}$), respectively.⁸

⁶ After the work described in this paper was completed a private communication was received from K. S. Pitzer stating that the infrared spectrum of TCE had been observed in his laboratory down to about 135 cm^{-1} . Outside of the range used in the present investigation, two bands, a strong one at 176 cm^{-1} and a somewhat weaker one at 152 cm^{-1} , were observed.

⁷ The depolarization factors are averages of the results given by Wittek for four separate determinations. For the 1025 cm^{-1} line, the data are too few to justify averaging but suffice to indicate its character.

⁸ See [4] for a description of the notation used here to represent the various modes.

⁵ For brief mention of this point, see the paper by Torkington on TCE [12].

There remain three Raman-active fundamentals: a CCl stretching and CCl₂ rocking (ρ -CCl₂) in b_{1g} , and the out-of-plane CCl₂ wagging (β -CCl₂) in b_{2g} . Apart from the lines already assigned to a_{1g} vibrations, only 347 and 512 have appreciable intensity. As they are depolarized they cannot be overtones, nor, indeed, is it reasonable to suppose they arise from sum bands. In fact, the weak polarized line at 1025 cm⁻¹ is most readily interpreted as 2×512. As fundamentals, they must be assigned in some manner to the b_{1g} and b_{2g} species. Analogy with TFE suggests that the b_{1g} stretching frequency in TCE is high, on a par with the infrared-active stretchings at 777 and 908 cm⁻¹, and that it would appear only feebly in the Raman spectrum. Hence, 347 and 512 may be assigned to the b_{1g} rocking and b_{2g} wagging modes, though it is not yet clear in what order. For the b_{1g} stretching one of the very weak lines at 726, 784, or 1000 cm⁻¹ may be chosen. The repeated occurrence of the latter in combinations and its usefulness in explaining the line at 1998 as 2×1000 lends considerable support to its interpretation as a fundamental.

It is now convenient to consider the infrared-active vibrations. The very strong bands at 777 and 908 cm⁻¹ may at once be assigned to the CCl stretchings in b_{2u} and b_{3u} , but in the absence of characteristic envelopes, it is not possible to determine uniquely their correspondence with the species. The infrared spectrum of TFE suggests that the b_{1u} wagging and the remaining b_{2u} and b_{3u} modes of TCE give rise to moderately strong bands. Figure 2 shows that between the intense cluster around 777 cm⁻¹ and the lower limit of the present range of observation, 190 cm⁻¹, there appear only two relatively strong bands, viz., those at 288 and 224 cm⁻¹. It will be recalled that the b_{2u} , b_{1u} , and b_{3u} angular modes in TFE occur at 558, 406, and 218 cm⁻¹. This suggests that the TCE rocking frequency occurs well below the 224 cm⁻¹ band. At the same time, however, this analogy with TFE requires the assignment of 288 cm⁻¹ to the b_{3u} deformation and 224 cm⁻¹ to the b_{1u} wagging motion. The attempt to infer the frequency of the b_{1u} rocking from the observed combination bands leads to a value around 170 cm⁻¹.⁹

Although it seems clear that aside from the torsion vibration all the fundamental frequencies are now known, it is still necessary to determine unequivocally the species to which the pairs 347, 512 and 777, 908 belong. An analysis of the force constants for the b_{1g} block leads to the conclusion that the frequency 512 cm⁻¹ is far too high to be ascribed to the rocking mode. It must, therefore, correspond to the b_{2g} wagging, whereas 347 may be satisfactorily interpreted as the b_{1g} rocking frequency. Contrary to the objections raised by Torkington [12], this assignment leads to reasonable force constants for the out-of-plane bending modes. In table 2 are

given the principal and interaction constants for the wagging motions in ethylene, TFE, and TCE. As was to be expected both H and h are much larger for TFE than for ethylene. For TCE, on the other hand, the principal constant H shows a marked decrease from its value for TFE, approaching, in fact, that for ethylene. Most interesting, however, is the zero interaction constant for TCE. Indeed, it may be anticipated that h also vanishes for the next higher member of the series, viz., tetrabromoethylene (TBE). Moreover, it is reasonable to expect that for TBE the constant H will be close to its value for TCE, being perhaps slightly smaller. If, then, the TCE constants are transferred to TBE, the b_{1u} and b_{2g} frequencies for the latter are found to be 200 and 489 cm⁻¹, respectively. The Raman data given for TBE by Köhler [32] include a weak line at 463 cm⁻¹, which he has assigned to the b_{2g} wagging mode. If this interpretation of the 463 cm⁻¹ is correct and h is taken to be zero, then H becomes 0.238, which is to be compared with 0.266 for TCE. It is, moreover, now possible to predict the value of the b_{1u} wagging frequency in TBE. With $H=0.238$, $h=0$, this vibration is calculated to occur at 189 cm⁻¹.

TABLE 2. Frequencies and force constants for wagging modes of H₂C:CH₂, F₂C:CF₂, and Cl₂C:CCl₂

Constants	H ₂ C:CH ₂ ^a	F ₂ C:CF ₂	Cl ₂ C:CCl ₂
$\nu_7(b_{1u})$ -----	949.2	406	224
$\nu_8(b_{2g})$ -----	943	508	512
$^b H$ -----	0.230	0.326	0.266
$^b h$ -----	.033	.127	.000

^a The data for ethylene are taken from the paper by Arnett and Crawford [11].
^b The principal constant H and the interaction constant h are given here in units of 10⁻¹¹ erg/radian². The actual symmetrized constant for the b_{1u} motion is $(H+h)$, and that for b_{2g} is $(H-h)$.

The correlation of the 777 and 908 cm⁻¹ bands with the species b_{2u} and b_{3u} is hindered by the absence of the usual aids to assignment. Analogy with TFE suggests that 908 cm⁻¹ refers to the b_{2u} stretching mode and 777 cm⁻¹ to that in b_{3u} . Because the remainder of the assignment is fairly well established, a force-constant calculation is justified. As yet unpublished results obtained by D. E. Mann for TCE, with the aid of a Urey-Bradley type of potential function [35], indicate that the suggested analogy with the TFE assignment is probably correct.¹⁰

All that is needed now to complete the assignment is a value for the torsion frequency. If we use for TCE the torsion constant previously determined for TFE, the frequency is calculated to be 106 cm⁻¹. This provides a plausible explanation for the very weak Raman line at 218 cm⁻¹ as the overtone of the torsion fundamental. To avoid imputing too much significance to this calculation, the value of the frequency has been rounded to 110 cm⁻¹. The final assignment is given in table 3.

¹⁰ It is worth pointing out, however, that some modification of the simple Urey-Bradley field [35] is needed. In particular, the addition of a constant that describes the effect of the interaction of the CCl₂ groups in the rocking modes seems desirable.

⁹ The results quoted in footnote 6 provide encouraging support for this prediction.

TABLE 3. Fundamental vibration frequencies for F₂C:CF₂ and Cl₂C:CCl₂

Species (V _i)	Number	Designation	F ₂ C:CF ₂	Cl ₂ C:CCl ₂
a _g -----	1, 2, 3	ν-CC	cm ⁻¹ 1872	cm ⁻¹ 1571
		ν-CX	778	447
		δ-CX ₂	394	235
a _u -----	4, 5	τ	190	100
		ν-CX	1340	1000
b _{1g} -----	6, 7	ν-CX ₂	551	347
		β-CX ₂	406	224
b _{2g} -----	8, 9	β-CX ₂	508	512
		ν-CX	1337	908
b _{2u} -----	10, 11	ν-CX ₂	218	176
		ν-CX	1186	777
b _{3u} -----	12	δ-CX ₂	558	288

^a See footnote 6.

4. Thermodynamic Functions

The thermodynamic functions to the rigid-rotator, harmonic-oscillator approximation were computed for TFE and TCE in their ideal gaseous state. The molecular constants used were taken from [13] and [36], but the assignments were those given in table 3. The results given in tables 4 and 5 are taken from the more complete tables calculated with the aid of SEAC.

TABLE 4. Calculated thermodynamic functions in dimensionless units for F₂C:CF₂ as an ideal gas

T	C _p ^o /R	(H ^o -E ₀ ^o)/RT	-(F ^o -E ₀ ^o)/RT	S ^o /R
°K				
197.53	7.787	5.467	27.013	32.480
300	9.707	6.607	29.526	36.133
400	11.054	7.559	31.361	39.120
500	12.073	8.364	33.336	41.701
600	12.849	9.050	34.924	43.974
700	13.441	9.637	36.364	46.001
800	13.895	10.142	37.685	47.827
900	14.247	10.580	38.905	49.485
1,000	14.523	10.961	40.040	51.001
1,100	14.741	11.295	41.101	52.395
1,200	14.917	11.590	42.096	53.686
1,300	15.059	11.851	43.034	54.886
1,400	15.176	12.085	43.921	56.006
1,500	15.273	12.294	44.762	57.056

TABLE 5. Calculated thermodynamic functions in dimensionless units for Cl₂C:CCl₂ as an ideal gas ^a

T	C _p ^o /R	(H ^o -E ₀ ^o)/RT	-(F ^o -E ₀ ^o)/RT	S ^o /R
°K				
300	11.519	8.054	33.292	41.346
^b 394.18	12.623	9.021	35.622	44.643
400	12.679	9.074	35.755	44.829
500	13.479	9.880	37.869	47.750
600	14.043	10.530	39.730	50.260
700	14.449	11.062	41.395	52.457
800	14.747	11.505	42.902	54.407
900	14.971	11.878	44.279	56.157
1,000	15.142	12.196	45.547	57.744
1,100	15.275	12.471	46.723	59.193
1,200	15.380	12.709	47.818	60.527
1,300	15.464	12.918	48.844	61.762
1,400	15.533	13.102	49.808	62.910
1,500	15.590	13.266	50.718	63.984

^a The entries in this table have been corrected for the change of μ₁₀ from 170 to 176 cm⁻¹.

^b The boiling point of TCE at 1 atmosphere is given by Dreisbach and Shrader as 121.02° C [37]. The ice point, as well as the other basic constants needed for the computation of these tables, are taken from [38].

The authors are indebted to Prof. K. S. Pitzer for communicating to them the infrared spectrum of TCE below 190 cm⁻¹, and also acknowledge several helpful discussions with Dr. Takehiko Shimanouchi.

5. References

- [1] Earle K. Plyler and Nicolo Acquista, *J. Opt. Soc. Am.* **43**, 212 (1953).
- [2] G. T. Furukawa, R. E. McCoskey, and M. L. Reilly, *J. Research NBS* **51**, 69 (1953) RP2423.
- [3] E. K. Plyler, R. Stair, and C. J. Humphreys, *J. Research NBS* **38**, 211 (1947) RP1769.
- [4] D. E. Mann, N. Acquista, and Earle K. Plyler, *J. Chem. Phys.* **21**, 1949 (1953).
- [5] J. R. Nielsen, H. H. Claassen, and D. C. Smith, *J. Chem. Phys.* **18**, 812 (1950).
- [6] E. K. Plyler, *J. Chem. Phys.* **16**, 1008 (1948).
- [7] H. J. Bernstein, *J. Chem. Phys.* **18**, 478 (1950).
- [8] P. Torkington and H. W. Thompson, *Trans. Faraday Soc.* **41**, 236 (1945).
- [9] A. Monfils and J. Duchesne, *J. Chem. Phys.* **18**, 1415 (1950).
- [10] A. Monfils and J. Duchesne, *Compt. rend.* **236**, 685 (1953).
- [11] R. L. Arnett and B. L. Crawford, Jr., *J. Chem. Phys.* **18**, 118 (1950).
- [12] P. Torkington, *Trans. Faraday Soc.* **45**, 445 (1949).
- [13] I. L. Karle and J. Karle, *J. Chem. Phys.* **18**, 963 (1950).
- [14] W. W. Coblentz, *Investigations of infrared spectra* (Publ. Carnegie Inst. of Wash., D. C. No. 35, Part I (1905)).
- [15] G. B. Bonino, *Gazz. chim. ital.* **55**, 335 (1925).
- [16] B. J. Spence and M. A. Easley, *Phys. Rev.* **34**, 730 (1929).
- [17] R. Freymann, *Compt. rend.* **194**, 1471 (1932).
- [18] T-Y. Wu, *Phys. Rev.* **46**, 465 (1934).
- [19] J. Duchesne and M. Parodi, *Nature* **144**, 382 (1939).
- [20] P. Pringsheim and B. Rosen, *Z. Physik*, **50**, 741 (1928).
- [21] A. Dadieu and K. W. F. Kohlrausch, *Sitz. ber. Akad. Wiss. Wien, Math.-naturw. Kl. [I]* **138**, 635 (1929).
- [22] A. Dadieu and K. W. F. Kohlrausch, *Monatsh. Chem.* **55**, 58 (1930).
- [23] V. N. Thatte and S. M. Shahane, *Indian J. Phys.* **6**, 155 (1931).
- [24] C. S. Morris, *Phys. Rev.* **38**, 141 (1931).
- [25] L. Simons, *Soc. Sci. Fennica, Commentationes Phys.-Math.* **6**, No. 13 (1932).
- [26] F. Heidenreich, *Z. Physik* **97**, 277 (1935).
- [27] T-Y. Wu, *J. Chinese Chem. Soc.* **4**, 402 (1936).
- [28] M. Kowalewska, *Acta Phys. Polon.* **7**, 279 (1938).
- [29] H. Wittek, *Z. physik. Chem. B* **48**, 1 (1940).
- [30] D. H. Rank, N. Sheppard, and G. J. Szasz, *J. Chem. Phys.* **16**, 698 (1948).
- [31] S. B. Sanyal, *Indian J. Phys.* **24**, 151 (1950).
- [32] K. W. F. Kohlrausch, *Ramanspektren* (Becker and Esler, Leipzig, 1943).
- [33] T-Y. Wu, *Vibrational spectra and structure of polyatomic molecules* (J. W. Edwards, Ann Arbor, Mich., 1946).
- [34] G. Herzberg, *Infrared and Raman spectra of polyatomic molecules* (D. van Nostrand Co., Inc., New York, N. Y., 1945).
- [35] Y. Morino, K. Kuchitsu, and T. Shimanouchi, *J. Chem. Phys.* **20**, 726 (1952).
- [36] I. L. Karle and J. Karle, *J. Chem. Phys.* **20**, 63 (1952).
- [37] R. R. Dreisbach and S. A. Shrader, *Ind. Eng. Chem.* **41**, 2879 (1949).
- [38] F. D. Rossini, F. T. Gucker, Jr., H. L. Johnston, Linus Pauling, and G. W. Vinal, *J. Am. Chem. Soc.* **74**, 2699 (1952).

WASHINGTON, October 2, 1953.