



Virucidal efficacy of chlorhexidine: a systematic review

Matheus dos Santos Fernandez¹ · Maria Izabel Florindo Guedes² · Gerson Pedro José Langa³ · Cassiano Kuchenbecker Rösing³ · Juliano Cavagni³ · Francisco Wilker Mustafa Gomes Muniz⁴ 

Received: 10 May 2021 / Accepted: 26 September 2021 / Published online: 12 October 2021
© The Society of The Nippon Dental University 2021

Abstract

This study aimed to systematically review the literature about the virucidal efficacy of CHX in comparison to other substances used in the oral cavity. Electronic searches were performed in four databases (PubMed, Scopus, Embase, and Web of Science). Only studies that presented the following characteristics were included: (1) verified virucidal efficacy of CHX against Herpes Simplex Type-1 (HSV-1), any Influenza, or any human coronavirus (HCoV); and (2) compared the virucidal efficacy of CHX with essential oils (Listerine[®]), quaternary ammonium compounds, povidone-iodine, hydrogen peroxide, negative control substance, and absence of therapy. Two researchers independently selected the studies, extracted data and evaluated the risk of bias. A narrative data synthesis was used. Twenty-five studies were included, of which 21 were in vitro and four were randomized clinical trials (RCT). Studies assessed the virucidal efficacy of CHX against Herpes Simplex Type-1 (HSV-1) (10 studies), Influenza A (InflA) (4 studies), human coronavirus (HCoV) (4 studies) and Severe Acute Respiratory Syndrome-Related Coronavirus (SARS-CoV-2) (11 studies). Most studies demonstrated that CHX has a positive virucidal efficacy against HSV-1 and InflA strains. However, lower efficacy was shown to InflA strain in comparison to povidone-iodine. Lower to none virucidal efficacy of CHX is expected for HCoV and SARS-CoV-2 strains for in vitro studies. Three RCT demonstrated that CHX was able to significantly reduce the viral load of SARS-CoV-2 for a short period. CHX may present an interesting virucidal efficacy against HSV-1 and InflA viruses. CHX also presents transient efficacy against SARS-CoV-2 when used as a mouthwash.

Keywords Antiseptics · Chlorhexidine · Mouthrinses · Mouthwashes · Virus

Introduction

Chlorhexidine (CHX) is widely used as an antiseptic formulation in dental practice [1]. CHX is a dicationic biguanide, and its use in children and adults has provided fast-acting and excellent safety [2, 3]. This product is commercially

available in various concentrations and presentations. In contrast to other antiseptic agents, the residual antimicrobial activity of CHX is unaffected by the presence of body fluids and blood [4, 5].

Many studies have established a clinically relevant antiplaque and antigingivitis efficacy of CHX in a dental clinical setting [6, 7]. For these reasons, CHX is recommended as the gold standard mouthrinse for chemical control of supragingival biofilm. Other studies have also reported positive results of CHX in reducing the proliferation of bacterial species associated with periodontal disease [6, 8]. This product also reduces the levels of halitosis-related bacteria colonizing the tongue's dorsal surface [9, 10]. However, the adverse events of CHX limit the long-term use of this mouthwash, which includes taste alteration, formation of supragingival calculus, soft tissue lesions in young patients, allergic responses, and staining of teeth and soft tissues [11, 12].

CHX is also suggested as a preprocedural mouthrinse to control the infection and reduce the number of bacteria in

✉ Francisco Wilker Mustafa Gomes Muniz
wilkermustafa@gmail.com

¹ School of Dentistry, Federal University Pelotas (UFPel), Rua Gonçalves Chaves, 457, Pelotas, RS 96015-560, Brazil

² Biotechnology and Molecular Biology Laboratory, State University of Ceará (UECE), Av. Dr. Silas Munguba, 1700, Fortaleza, CE 60714-903, Brazil

³ Department of Periodontology, Federal University Rio Grande do Sul (UFRGS), Rua Ramiro Barcelos, 2492, Porto Alegre, RS 90035-003, Brazil

⁴ Department of Periodontology, Federal University Pelotas (UFPel), Rua Gonçalves Chaves, 457, Pelotas, RS 96015-560, Brazil

aerosol [13, 14] since important effects in a broad spectrum of Gram-positive and Gram-negative bacteria were demonstrated [2, 3, 15]. Therefore, considering that CHX may also be effective against lipid-enveloped microorganisms, different authors have also investigated the effectiveness of CHX against viral strains. Therefore, CHX may be a viable candidate to prevent or treat virus infection in the oral cavity, but no synthesized information is available about the virucidal efficacy of CHX. Therefore, this study aimed to systematically review the literature about the virucidal efficacy of CHX in viruses that affects the oral cavity.

Materials and methods

The report of this systematic review is based on the Preferred Reporting Items for Systematic Reviews and Metaanalyses (PRISMA) 2020 statement [16].

Focused question

The present study is based on the following focused question: “Does chlorhexidine have significant virucidal effect, as compared to other solutions or a placebo solution, in viruses affecting the oral cavity?”. Therefore, the PICO question for this systematic review was defined as follows:

P: Any population and in vitro assays involving viruses that affect the oral cavity.

I: Use of chlorhexidine in any concentration and formulation.

C: Absence of therapy, placebo solution, hydrogen peroxide, povidone-iodine, any quaternary ammonium compounds or essential oils (Listerine®).

O: Any assessment of virucidal efficacy in the following virus: Herpes Simplex Type-1 (HSV-1), Influenza A, Human coronavirus (HCoV) and Severe Acute Respiratory Syndrome-Related Coronavirus (SARS-CoV-2) strains.

Search strategy and selection of studies

All articles were selected from the following electronic databases: PubMed, Web of Science, Embase and Scopus. Sensitivity analysis strategies were developed for each database using a combination of free terms and, whenever applicable, indexed specifically to each database. The search strategy was performed on July 9th, 2021. Hand searching of the reference lists of all studies included in the search strategy and in the reference list of previously published reviews was performed [14, 17–19]. The full search strategy for all databases can be found in Appendix 1. A search for grey literature was also performed on Google Scholar database

using an adapted search strategy. Only the first 100 studies were screened for eligibility on this database.

Studies that performed any in vitro microbiological analysis that used cells or surfaces contaminated by these viral agents were also screened for eligibility, as well as clinical trials that assessed the virucidal efficacy of CHX. The exclusion criteria were review articles, case reports, letters to the editor and observational studies. Studies that associated CHX with another therapy in the same group, those who evaluated other viruses or those studies without the above-mentioned control groups were also excluded. No restriction of language or date of publication were applied.

Results of literature searches were uploaded in Endnote® X9 software (Thomson Reuters, New York, NY, USA), and duplicate records were removed. Two researchers (FWMGM and GPJL) independently screened all titles and abstracts considering the abovementioned eligibility criteria. The same two researchers independently assessed the full text of the included studies, and discrepancies were solved by a third researcher (CKR). Regarding study selection, kappa coefficients between the two researchers were 0.976 and 0.957 for title/abstract and full-text selection, respectively.

Data extraction

Data extraction was conducted using a standardized Excel spreadsheet (Microsoft®) specifically developed for this study. The following data were extracted: author, year, country, study design, CHX and control solution concentrations, the form of administration, contact time of all tested solutions, virus assessed, and origin of the virus. In addition, the test used to determine the virucidal efficacy, results of virus inactivation in the CHX group, results of virus inactivation in the control groups, main results, and other important observations were also evaluated. Two researchers extracted all data independently (FWMGM and MSF), and a third reviewer was involved only in case of discrepancy (MIFG).

Risk of bias and synthesis of evidence

The risk of bias of all included studies was assessed independently by two researchers (FWMGM and MSF), following the criteria proposed by the *Joanne Briggs Institute* [20]. When a consensus was not possible, a third researcher was involved in this process (GPJL). The checklist was adapted according to the statements proposed by CRIS Guidelines (Checklist for Reporting *In-Vitro* Studies) [21], which suggests evaluating factors such as the randomization process, blinding and statistical analysis. The tool has ten questions, two of which were not considered for the present study because they do not agree with the risk of bias analysis for

in vitro studies. Therefore, the following questions were evaluated:

- (1) Was the assignment to treatment groups truly random?
- (2) Was allocation to treatment groups concealed from the allocator?
- (3) Were those assessing the outcomes blind to the treatment allocation?
- (4) Were control and treatment groups comparable at entry?
- (5) Were groups treated identically other than for the named interventions?
- (6) Were outcomes measured in the same way for all groups?
- (7) Were outcomes measured in a reliable way?
- (8) Was appropriate statistical analysis used?

Each question was answered and when enough information was available, a “Yes” answer was given, which is equivalent to a low risk of bias. In case of non-existent information, the “No” answer was given for this criterion, the equivalent of a high risk of bias. The “not clear” response was attributed when a high or low risk of bias could not be classified. For each “Yes” attributed to each criterion, the study received one point.

For randomized clinical trials (RCT), the Cochrane risk of bias tool for randomized trials (RoB 2) was used [22]. The tool evaluates the risk of bias considering six different domains: random sequence generation, deviations from the intended interventions, missing outcomes, measurement of the outcome and selective outcome reporting.

Due to the high heterogeneity among the included studies, no meta-analysis could be performed. Therefore, data synthesis is presented in a qualitative way.

Results

The searches performed in the electronic databases presented 2,662 potentially relevant studies [PubMed ($n = 1,225$); Scopus ($n = 1741$); EMBASE ($n = 1562$); Web of Science ($n = 351$)]. From these, 196 full-texts were read, of which 170 were excluded. Therefore, 26 studies were included in this systematic review. Seven evaluated the virucidal effect of CHX only against HSV-1 [23–29] and three studies verified the effect on HSV-1 and Influenza A [30–32]. In addition, another four studies investigated the virucidal effect from CHX in HCoV strains [33–36]. Ten studies assessed the virucidal efficacy of CHX against SARS-CoV-2 [37–46]. One study

assessed the virucidal efficacy of CHX against SARS-CoV-2 and Influenza A [47]. Figure 1 shows the flow-chart of study inclusion. In general, the included studies were published between 1972 and 2021.

Risk of bias

All in vitro studies presented an unclear risk of bias for randomization, allocation concealment, and blinding of the evaluators (Fig. 2) [23–38, 40, 44–47]. Based on RoB 2 assessment, overall low risk of bias was detected in two studies [41, 42]. The other RCTs presented some concerns [43] and high [39] risk of bias. Figure 3 shows the risk of bias assessment for these studies.


Efficacy of CHX against HSV-1

Table 1 summarizes the main findings of all studies that evaluated the virucidal effect of CHX against HSV-1. In these assays, the concentration of CHX varied from 0.001% [26, 27] to 2.5% [25]. The main method of administration of CHX was direct contact with cells contaminated by HSV-1, and the time of exposure to the virus varied significantly among the included studies.

All studies presented a control group that could be composed of contaminated cells that were exposed to antiseptic solutions [26–28, 30], samples exposed to a placebo solution, without an active agent [31], or to sterile phosphate-buffered saline [23]. Other substances were also used, such as essential oils [24], povidone-iodine [25, 29, 32], hydrogen peroxide [25], and quaternary ammonium compounds, Ethylenediaminetetraacetic acid (EDTA) [29]. Most of the viral samples used in the studies were obtained from culture collection organizations, cultivation from other laboratories and distributed standard reference microorganisms [23–28, 31, 32]. Two studies did not report the origin of the viral samples used in their experiments [29, 30].

All studies were in vitro, and the most used test to verify the virucidal efficacy of CHX against viral agents was minimum inhibitory concentration by viral titration [25, 26, 28–32] and colony-forming units [24, 27], followed by the sensitivity of tissue culture cells [23]. Regardless of the CHX’s concentration, significant virucidal efficacy was demonstrated in studies that used viral titration reduction tests when compared to a control group of contaminated cells that did not receive any antiseptic solution [28, 30] and to those exposed to a placebo solution [31]. Other studies that used the same methodology demonstrated that CHX showed low [26] or uncertain [25, 29, 32] efficacy to inhibit HSV-1 viral replication when compared to cells that had no contact with disinfectants [26] or exposed to povidone-iodine [25,

Fig. 1 Flow chart of studies selection according to PRISMA statement



29, 32], respectively. The study that assessed the sensitivity of tissue culture cells demonstrated that CHX has superior virucidal efficacy compared to a sterile phosphate-buffered saline solution [23].

Moreover, other studies that evaluated the effect of CHX compared to uncontaminated cells and exposed to sterile phosphate-buffered saline by means of colony-forming units and sensitivity of tissue culture cells found that CHX is a viable alternative for the reduction of HSV-1 viral load [24, 27]. Regarding the virucidal effect of CHX compared to essential oils, both solutions showed similar and effective anti-HSV-1 properties [24].

Efficacy of CHX against Influenza A and coronavirus (HCoV and SARS-CoV-2)

Five studies evaluated the virucidal effect of CHX against Influenza A strains [30–33, 47] (Table 2). Of these, three studies also verified the virucidal efficacy of the antiseptic against HSV-1. These studies were previously characterized

[30–32]. One study [33] verified the viral sensitivity by analyzing the detection inoculation of samples in cell culture and nested multiplex RT-PCR. For this, cells infected by the viruses were exposed to CHX, while another group of infected cells did not receive solutions with disinfectant agents [33]. Disinfectant effectiveness of CHX was assessed by titrating Influenza A on a model using skin tissues [47]. In vitro inactivation of Influenza A was also tested in this study [47].

In general, studies demonstrated the effectiveness of CHX in reducing the viral load of Influenza A after 30 s [31, 33] and after 10 min of contact [30], when compared to groups of contaminated cells that had no contact with disinfectants [30, 33] or exposed to CHX without the active agent [31]. Notwithstanding, one study [32] demonstrated that, compared to povidone-iodine solution and gargle, CHX might also be a viable alternative against the Influenza A virus [32]. Another study demonstrated a low virucidal efficacy of CHX against Influenza A viruses on both skin and in vitro models [47].

Author (Year)	Joanne Briggs Institute (JBI)							
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Ansaldi et al., 2004	?	?	?	+	+	+	+	-
Bailey & Longson, 1972	?	?	?	+	+	+	+	-
Baqui et al., 2001	?	?	?	+	+	+	+	-
Bernstein et al., 1990	?	?	?	+	+	+	+	-
Davies et al., 2021	?	?	?	+	+	+	+	-
Damery & Cremieux, 1989	?	?	?	+	+	+	+	-
Geller et al., 2009	?	?	?	+	+	+	+	+
Hirose et al., 2021	?	?	?	+	+	+	+	-
Imai et al., 2021	?	?	?	+	+	+	+	-
Jain et al., 2021	?	?	?	+	+	+	+	-
Kawana et al., 1997	?	?	?	+	+	-	?	+
Komine et al., 2021	?	?	?	+	-	+	+	-
Meister et al., 2020	?	?	?	+	+	+	+	-
Okunishi et al., 2009	?	?	?	+	+	+	?	-
Park & Park, 1989	?	?	?	+	+	+	+	-
Park et al., 1991	?	?	?	+	+	+	+	-
Sattar et al., 1989	?	?	?	+	+	+	+	+
Steinhauer et al., 2021	?	?	?	+	-	+	+	-
Shinkai & Yoshino, 1974	?	?	?	+	+	+	+	-
Tyler & Ayliffe, 1987	?	?	?	+	+	+	+	-
Xu et al., 2020	?	?	?	+	+	+	+	+

JBI - Critical appraisal checklist for <i>In-Vitro</i> Studies	
(1) Was the assignment to treatment groups truly random?	
(2) Was allocation to treatment groups concealed from the allocator?	
(3) Were those assessing the outcomes blind to the treatment allocation?	
(4) Were control and treatment groups comparable at entry?	
(5) Were groups treated identically other than for the named interventions?	
(6) Were outcomes measured in the same way for all groups?	
(7) Were outcomes measured in a reliable way?	
(8) Was appropriate statistical analysis used?	

Legend	
+	Yes
-	No
?	Unclear

Fig. 2 Risk of bias of the in vitro included studies

The resistance of HCoV against the virucidal action of CHX was evaluated in four studies [33–36]. Minimum inhibitory concentration by viral titration was the main test to verify the virucidal effect of the antiseptic on cells [34] and stainless steel disc surfaces [35] contaminated with the virus, except for one study [33]. One study assessed the viral titers reduction after exposure to CHX [36].

Detection analyses by inoculation of samples in cell culture demonstrated no virucidal efficacy after 30 s of contact with CHX when a culture of HCoV was used. Similar results were detected, in another study, after 15 s, 30 s, and 60 s [36]. However, a different result was obtained when RT-PCR was applied, as CHX inactivated the viruses after 30 s [33]. Nevertheless, the results of viral titration tests in plaque-forming units reflect that CHX has moderate anti-HCoV activity [34] or is ineffective [35] when compared to the results of contaminated samples not exposed to disinfectants (Table 2).

Eleven included studies verified the antiviral efficacy of CHX on SARS-CoV-2 strains, of which seven were in vitro studies [37, 38, 40, 44–47], and four were RCT [39–42]. One study [37] verified the viral infectivity of SARS-CoV-2 after contact with two different products based on 0.02% CHX (Chlorhexamed Forte® and Dynexidine Forte®). Through viral titers determined upon titration on Vero cells, the study demonstrated that CHX (regardless of the tested product) has not been able to significantly reduce the viral infectivity of the three strains of SARS-CoV-2 in comparison to other evaluated solutions (hydrogen peroxide, povidone-iodine and essential oils) [37]. Moreover, CHX was not able to significantly reduce the viral titers in two other studies, using 0.12% [45] or 0.2% diluted or not in alcohol [40]. Conversely, these studies showed that essential oils (Listerine)

Fig. 3 Risk of bias assessment of included the randomized controlled trials

	Randomization process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported result	Overall bias	
Eduardo et al., 2021	!	●	+	●	+	●	LEGEND + Low risk ! Some concerns ● High risk
Elzien et al., 2021	+	+	+	+	+	+	
Huang et al., 2016	+	+	+	+	+	+	
Seneviratne et al., 2020	!	+	+	+	+	!	

Table 1 General characteristics of included studies that verified the virucidal effect of chlorhexidine (CHX) against Herpes Simplex Type-1 (HSV-1) in this review (*n* = 10)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was administered? Time of contact	Virus assessed (origin)	Main results
Bailey 1972 [23] United Kingdom	0.02% CHX in direct contact with contaminated cells 90'	Negative control: sterile phosphate-buffered saline solution. The substance was in direct contact with contaminated cells 90'	HSV-1 Oral strain SUE	Surviving virus is presented, respectively, for CHX and control groups TCID ₅₀ /ml ² : < 1 and 1 × 10 ⁶ PFU/ml ³ : < 1 and 1.2 × 10 ⁵
Baqui 2001 [24] United States of America	0.12% and 0.2% CHX in direct contact with contaminated cells 30"	Essential oil-containing mouthrinses (LA and TLA) Direct contact with contaminated cells 30"	HSV-1 McIntyre strain	The antiviral effects of LA and TLA were found to be similar and also the antiviral effect of 0.12% and 0.2% CHX were also found to be comparable Inhibition of the syncytia formation or the cytopathic effect and inhibition of the plaque formation for HSV-1 on Vero cell monolayers
Bernstein 1990 [31] United States of America	0.12% CHX in direct contact with contaminated cells 30", 5', and 15'	Negative control substance: similar to CHX, but without the active ingredient. The substance was in direct contact with contaminated cells 30", 5', and 15'	HSV-1 Strain/ Bethesda 1/85	Infectivity of HSV-1 (0.12% CHX and 0.2% CHX): undiluted: no growth; dilution 1:2—1 to 10 plaques/well; dilution 1:4, 1:8, 1:16, 1:32, 1:64. All presented +30 plaques/well Infectivity of HSV-1 (LA/TLA): undiluted: no growth; dilution 1:2: no growth; dilution 1:4: 1—10 plaque/well; dilution 1:8, 1:16, 1:32, 1:64 All presented +30 plaques/well Virus titers in plaque-forming units, PFU/ml. Values are presented, respectively, for CHX and control groups: CHX demonstrated an effective virucidal effect against HSV-1 after 30' of contact 30': 97% and 0% 5': 99.9% and 1% 15': > 99.9% and 1%
Damery 1989 [25] France	0.5% CHX in direct contact with contaminated cells 15', 30' and 60'	PVP-I (20%) HP (10%) Direct contact with virus-contaminated cells 15', 30' and 60'	HSV-1 C.H.U. Toulouse- Rangueil	After 60 min of contact, all solutions were effective in reducing viral load Reduction in virus titer in log IU/ml: CHX: 15': 4.73; 30': 4.26; 60': 3.36 Povidone-iodine: 15': 4.95; 30': 4.16; 60': < 3 Hydrogen peroxide: 15': < 4; 30': < 4; 60': < 4

Table 1 (continued)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was administered? Time of contact	Virus assessed (origin)	Main results
Kawana 1997 [32] Japan	0.05%, 0.1%, 0.5% CHX in direct contact with contaminated cells	PVP-I (solution): 0.05%, 0.1%, 0.5%, 1% and 5%; gargle: 0.07%, 0.1%, 0.5%, 1%; cream 0.05%	HSV-1 A/Kitakyushu/159/93	Reduction in virus titer after treatment: CHX: uncertain virucidal effect PVP-I solution and PVP-I gargle: virucidal effect clearly effective PVP-I cream: not performed
Okunishi 2009 [30] Japan	NR 0.5% and 0.1% CHX in direct contact with contaminated cells	NR Negative control: Contaminated cells did not receive disinfectant solutions Not applied	HSV-1 NR	CHX showed an efficient virucidal effect Log ₁₀ reduction of virus titer calculated as log ₁₀ TCID ₅₀ of control—Log ₁₀ TCID ₅₀ of CHX: 0.5% CHX reduced the viability titer to below the detection limit after just 1 min
Park 1989 [27] United States of America	0.2%, 0.1%, 0.05%, 0.001%, 0.0008% and 0.0005% CHX in direct contact with contaminated cells	Negative control: contaminated cells did not receive disinfectant solutions Not applied	HSV-1 F-strain—American Type Culture	Various concentrations of CHX, even if minimal, are effective in decreasing the viral load of HSV-1 The viral titers were then determined from the supernatant by use of a plaque assay technique At any concentration used, the reduction of HSV-1 by CHX was significantly greater than the control group. CHX > 0.001% showed a cytotoxicity index
	Baseline, 10", 20" and 60"			The cytolytic activity of the virus was evident in the presence of 0.5% ($p < 0.05$), 0.1% ($p < 0.02$) or 0.2% CHX ($p < 0.02$) in the culture medium

Table 1 (continued)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was (were) administered? Time of contact	Virus assessed (origin)	Main results
Park 1991 [26] United States of America	1, 5, 8, 10 and 20 µg/ml CHX in direct contact with contaminated cells	Negative control: Contaminated cells did not receive disinfectant solutions Not applied	HSV-1 F-strain—American Type Culture Collection	CHX showed low efficacy in inhibiting viral replication Reduction in virus titer after treatment: CHX has the ability to slightly reduce viral titration but does not significantly decrease from 10 ² to less than 10 ¹
Shinkai 1974 [28] Japan	NR 50, 100 and 200 µg/ml CHX in direct contact with contaminated cells	Negative control: Contaminated cells did not receive disinfectant solutions	HSV-1 F-strain	At 0 °C, the virus was not activated by CHX. At 37 °C, CHX was able to inactivate the viruses The sensitivity was expressed by titer reduction in the log of test mixture minus that of control mixture as estimated after 60' incubation: 37 °C and 200 µg/ml: 1.1; 37 °C and 100 µg/ml: 0.3; 37 °C and 50 µg/ml: 0.1; 0 °C and 200 µg/ml: 0
Tyler 1987 [29] United Kingdom	60' 0.50% CHX in direct contact with contaminated cells	Not applied PVP-I (10%; 1%) Quaternary ammonium—EDTA solution (8%)	HSV-1 NR	Counts were expressed as the number of PFU/ml of the original inoculum: Reductions were similar or lower than the controls were obtained with CHX CHX: 1': 0.4±0.2; 5': 1.4±0.3; 10': 1.1 Povidone-iodine: 1': 0.3±0.1; 5': 1.9±0.4; 10': 2.5±0.6 EDTA solution: 1': 0.6±0.5; 5': 0.1; 10': 0.3
	1', 5' and 10'	1', 5' and 10'		

NA not applicable, NR not reported, PVP-I povidone-iodine, HP hydrogen peroxide

Table 2 General characteristics of included studies that verified the virucidal effect of CHX against Influenza A, Human coronavirus (HCoV) and Severe Acute Respiratory Syndrome-Related Coronavirus (SARS-CoV-2) strains in this review ($n = 18$)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was (were) administered? Time of contact	Virus assessed (origin)	Main results
Ansaldo 2004 [33] Italy	1% CHX in direct contact with contaminated cells 30', 1', 2', 5', 15', and 30'	Negative control: contaminated cells did not receive disinfectant solutions NR	Influenza A and HCoV NR	Infectivity, detected by inoculation of samples in suitable cell culture; genome integrity, detected by nested RRT-PCR for SARS-CoV and multiplex nested RRT-PCR Influenza virus and SARS-CoV RNA are still detectable after 30' of contact time with 1% CHX Last contact time resulting positive by cell culture and PCR for CHX: Influenza (culture): 30' Influenza (PCR): 30' SARS-CoV (culture): negative after 30' SARS-CoV (PCR): 30' 1% CHX in 2 ml cell culture medium did not significantly damage the cell monolayer
Bernstein 1990 [31] United States of America	0.12% CHX in direct contact with contaminated cells 30', 5', and 15'	Negative control substance: similar to CHX, but without the active ingredient. The substance was in direct contact with contaminated cells 30', 5', and 15'	Influenza A Strain/Bethesda 1/85	Virus titers in plaque-forming units, PFU/ml. Values are presented, respectively, for CHX and control groups: CHX demonstrated an effective virucidal effect against Influenza A after 30' of contact 30': 93% and 0% 5': >98% and 1% 15': >98% and 1%
Davies et al. 2021 [40] United Kingdom	Product 1—CHX 0.2%—formulation contains ethanol; Product 1—CHX 0.2%—alcohol-free formulation CHX in direct contact with contaminated cells 1'	HP (1.5%) PVP-I (0.58%) Essential oils mouthrinse (Listerine total care)	SARS-CoV-2 England 2 strain	Log ₁₀ reduction of virus titer calculated as log ₁₀ TCID ₅₀ of control—log ₁₀ TCID ₅₀ of CHX TCF unconcentrated—mean (95% CI)
Eduardo et al. 2021 [41] Brazil	0.12% CHX in direct contact with contaminated saliva (mouthwash) 30'	Placebo—distilled water; CPC (0.075%) + zinc (Zn) lactate (0.28%) HP (1.5%) Antiseptic agents in direct contact with contaminated saliva (mouthwash)	SARS-CoV-2 Saliva from contaminated patients	Product 1—CHX (0.2%) formulation ethanol: 0.5 (0.1–0.9) Product 2—CHX (0.2%) alcohol-free formulation: 0.2 (-0.2–0.7) HP (1.5%): 0.2 (-0.1–0.5) PVP-I (0.58%): ≥ 4.1 (3.8–4.4) Essential oils mouthrinse (Listerine total care): ≥ 4.1 (3.8–4.4) RRT-PCR cycle threshold (Ct). RRT-PCR assessed before rising, immediately after, 30' and 60' after. Mean Ct value for each experimental group was compared to the baseline value. Comparisons among groups were also performed CHX: significantly lower viral load when compared to baseline after 30' and 60' CPC + Zn: significantly lower viral load when compared to baseline and immediately after rising HP: significantly lower viral load when compared to baseline and immediately after, 30', and 60' CPC + Zn and HP promoted higher reductions of viral load when compared to CHX
		Placebo: 1'' CPC + Zn: 30' HP: 1''		

Table 2 (continued)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was administered? Time of contact	Virus assessed (origin)	Main results
Elzein et al. 2021 [42] Lebanon	0.2% CHX in direct contact with contaminated saliva (mouthwash) 30' 0.05% CHX in direct contact with contaminated cells 5', 15', 30' and 60'	Placebo—distilled water PVP-I (1%) Antiseptic agents in direct contact with contaminated saliva (mouthwash) 30' Negative control: contaminated cells did not receive disinfectant solutions 5', 15', 30' and 60'	SARS-CoV-2 Saliva from contaminated patients HCoV VR 70—American Type Culture Collection	Saliva samples were collected before and 5 min after rinsing. Subsequently, SARS-CoV-2 RRT-PCR was performed. The change in cycle threshold (delta Ct) values of salivary SARS-CoV-2 were calculated CHX: significantly lower viral load was detected in comparison to baseline. No significant difference was found between the delta Ct of patients using CHX and PVP-I solutions PVP-I (1%): significantly lower viral load was detected in comparison to baseline Infected wells were counted and viral titers or 50% cell culture infective dose (TCID ₅₀) were estimated: CHX showed a moderate anti-HCoV 229E activity but insufficient to be antiseptic Log ₁₀ reduction of viral titers: 5': 0.8 ± 0.7 (10 ⁻⁴ mol/L) and 1.4 ± 0.8 (10 ⁻³ mol/L); 15': 0.5 ± 0.4 (10 ⁻⁴ mol/L) and 2.1 ± 0.4 (10 ⁻³ mol/L); 30': 1.4 ± 1.5 (10 ⁻⁴ mol/L) and 2.4 ± 0.6 (10 ⁻³ mol/L); 60': 2.1 ± 1.2 (10 ⁻⁴ mol/L) and 3.0 ± 0.2 (10 ⁻³ mol/L) The measurement limits of the titers of Influenza A and SARS-CoV-2 were 10 ¹ focus-forming units/ml and 10 ^{0.5} 50% tissue culture infectious dose (TCID ₅₀ /ml), respectively
Geller 2009 [34] France	CHX in direct contact with contaminated cells 5', 15', 30' and 60'	Negative control: contaminated cells did not receive disinfectant solutions NA	Influenza A and SARS-CoV-2 NR	Influenza A: CHX presented low virucidal efficacy in both in vitro and on skin models SARS-CoV-2: CHX presented low virucidal efficacy in both in vitro and on skin models The disinfection efficacy of CHX in SARS-CoV-2 was slightly greater than that in Influenza A
Hirose et al. 2021 [47] Japan	0.2% and 1.0% CHX in direct contact with contaminated cells 5', 15' and 60'	Negative control: contaminated patients did not receive disinfectant solutions NA	SARS-CoV-2 Saliva from contaminated patients	After 4 days of CHX administration, the oropharynx was swabbed and tested for the presence of SARS-CoV-2 by rRT-PCR (1) Most individuals in the group that received CHX intervention protocol were considered negative for SARS-CoV-2 in the oropharynx (62.1%; n = 41). In the control group, only 3 (5.5%) were negative for SARS-CoV-2 in the oropharynx (p < 0.01) (2) Among individuals who used CHX rinse, 80 (86.0%) of patients were negative for the presence of SARS-CoV-2, while 5 (6.2%) of the control patients were negative for the virus (p < 0.01)
Huang et al. 2021 [43] United States of America	0.12% CHX in direct contact with contaminated saliva (1) Only mouthwash (n = 66); (2) Mouthwash and 1.5 mL of CHX spray (n = 93) 30' twice a day for 4 days	Negative control: contaminated patients did not receive disinfectant solutions. Two negative control groups were involved: (1 - n = 55) and (2 - n = 80) NA	SARS-CoV-2 HCoV VR-1558—American Type Culture Collection	Viral titers (log ₁₀ TCID ₅₀ /ml) were measured by quantal tests of six wells per dilution CHX (0.5%) was ineffective against two coronavirus strains The virucidal efficacy of CHX against HCoV through the suspension test [mean log ₁₀ reduction (95% CI)] after 15', 30' and 60 were, respectively, 0.60 (0.34), 0.75 (0.35), and 0.75 (0.38)
Imai et al. 2021 [36] Japan	0.1% and 0.5% CHX in direct contact with contaminated cells 15', 30', and 60'	Negative control: contaminated cells did not receive disinfectant solutions NA	HCoV VR-1558—American Type Culture Collection	

Table 2 (continued)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was administered? Time of contact	Virus assessed (origin)	Main results
Jain et al. 2021 [44] India	0.2% and 0.12% CHX in direct contact with contaminated cells 30' and 60'	PVP-I (1%) Antiseptic agents in direct contact with contaminated cells 30' and 60'	SARS-CoV-2 NR	Analysis of the virus inactivation was based on the quantification of viral RNA (Cycle threshold [Ct]) present in the culture supernatant using rRT-PCR CHX and PVP-I were able to inactivate SARS-CoV-2. 0.2% CHX performed better than the other substances 30'; 60' = relative Ct change log ₁₀ reduction (standard deviation) CHX (0.2%): 12.5 (0.5); 13 (0) CHX (0.12%): 10.5 (0.5); 11 (1.0) PVP-I: 9.5 (0.5); 11 (2)
Kawana [32] 1997 Japan	0.05%, 0.1%, 0.5% CHX in direct contact with contaminated cells 30' and 60'	Povidone-iodine (solution: 0.05%, 0.1%, 0.5%, 1% and 5%; gargle: 0.07%, 0.1%, 0.5%, 1%; cream 0.05%) NR	Influenza A A/Kitakyushu/159/93	Reduction in virus titer after treatment: CHX: effective virucidal effect for Influenza A PVP-I solution and PVP-I gargle: virucidal effect clearly effective for Influenza A PVP-I cream: not performed
Komine et al. 2021 [45] Japan	0.12% CHX in direct contact with contaminated cells 30'	Negative control: contaminated cells did not receive disinfectant solutions CPC toothpaste (0.05%) CPC mouthwash (0.05%) CPC spray (0.3%) CPC mouthwash (0.075%) CPC mouthwash (0.04%) NR	SARS-CoV-2	Log ₁₀ reduction of virus titer (PFU/ml) CHX (0.12%) did not show a sufficient inactivation effect against SARS-CoV-2; inactivation effectiveness = 42.5%
Meister [37] 2020 Germany	(1) Product B—0.2% (2) Product D—0.2% CHX in direct contact with contaminated cells 30"	Hydrogen peroxide (1.5%) Polyvidone-iodine (1%) Essential oils Antiseptic agents in direct contact with contaminated cells 30"	JPN/TY/ WK-521 strain— National Institute of Infectious Disease SARS-CoV-2 UKEssen strain	CPC toothpaste [0.05% (1/4 slurry): 0.0125%]: 99.94% CPC mouthwash (0.05%): 99.994% CPC spray (0.3%): > 99.96%; CPC mouthwash (0.075%): > 99.995% CPC mouthwash (0.04%): > 99.996% Viral titers were determined upon titration on Vero cells: virucidal activities could be observed with log reduction factors ranging between 0.3 and 1.78 Although CHX demonstrated mild virucidal activities, it did not significantly reduce the viral infectivity of the three strains of SARS-CoV-2
				Active compound: strain 1/strain 2/strain 3 (1) CHX 0.2%: 1.00/0.78/1.17 (2) CHX 0.2%: 0.50/0.56/0.50 Hydrogen peroxide: 0.78/0.68/0.73 Polyvidone-iodine: ≥ 3.11/≥ 2.78/≥ 2.61 Essential oils: ≥ 3.11/≥ 2.78/≥ 2.61

Table 2 (continued)

Author (year) Country	CHX concentration; how CHX was administered? Time of contact	Control group (concentration); how product(s) was (were) administered? Time of contact	Virus assessed (origin)	Main results
Okunishi 2009 [30] Japan	0.5% and 0.1% CHX in direct contact with contaminated cells 15', 30', and 60'	Negative control: contaminated cells did not receive disinfectant solutions	Influenza A NR	Log ₁₀ reduction of virus titer calculated as log ₁₀ TCID ₅₀ of control—log ₁₀ TCID ₅₀ of CHX CHX showed virucidal efficacy against Influenza A 0.5% and 0.1% CHX failed to show virucidal efficacy against Influenza A at any exposure time lower than 5 min CHX decreased only 45.1% of Influenza A titers after 10 min of exposure
Sattar 1989 [35] Canada	0.008% and 0.08% CHX applied in the surface test (stainless steel disks) 1'	Povidone-iodine (10—1% iodine) Quaternary ammonium (Dimethyl benzyl ammonium chloride)—50% C14, 40% C12, 10% C16—0.04% applied in the surface test (stainless steel disks)	HCoV NR	The criterion of efficacy (PFU/disk) for the disinfectants was ≥ 3 log ₁₀ reduction in the number of infectious virus units (reduction of virus titer by > 99.9%); CHX was ineffective in reducing viral replication CHX: no Povidone-iodine: yes Quaternary ammonium: no
Seneviratne 2020 [39] China	0.02% CHX in direct contact with contaminated saliva 5', 120', and 240'	Povidone-iodine (0.5%) Cetylpyridinium chloride (0.075%) Antiseptic agents in direct contact with contaminated saliva	SARS-CoV-2 UKEssen strain	RRT-PCR cycle threshold (Ct) The relative change in the doubling of the cycle limit values in patients in the mouthwash and water group demonstrated no significant difference in viral reduction in the CHX group than the others (<i>p</i> > 0.05)
Steinhauer et al. 2021 [46] Germany	0.1% and 0.2% CHX in direct contact with contaminated cells CHX 0.1%: 5', 10', and 5'' CHX 0.2%: 1' and 5''	Negative control: contaminated cells did not receive disinfectant solutions NA	SARS-CoV-2 NR	Viral titers were determined upon limited end-point titration on Vero E6 cells. Tissue culture infectious dose 50% (TCID ₅₀ /ml) was calculated—Log ₁₀ reduction of virus titer Both formulations based on CHX were found to have limited efficacy against SARS-CoV-2 CHX 0.1%: reduced the virus titer even at a prolonged contact time of 10 min by < 1 log ₁₀ CHX 0.2%: reduced SARS-CoV-2 within a contact time of 1 min as well as at a prolonged contact time of 5 min when tested by < 1 log ₁₀
Xu 2020 [38] United States of America	0.12% CHX in direct contact with contaminated saliva 20'	HP (1.5%) PVP-I (1%) Essential oils mouthrinse	SARS-CoV-2	Plaque assays determined virus titers Viruses that came into contact with CHX (50%) completely lost their infectivity. Treatment with 5% essential oils or CHX had a moderate antiviral effect; Hydrogen peroxide and povidone-iodine had greater inhibitory effects on viruses than CHX or essential oils. In general, the effect of CHX on the viruses after removal of the mouth rinse during the infection, 5% (v/v) CHX had only a moderate effect, reducing infection by 35–55%

NA not applicable, NR not reported, PVP-I povidone-iodine, HP hydrogen peroxide, CPC cetylpyridinium chloride

[40], povidone-iodine [40], and cetylpyridinium chloride were able to significantly SARS-CoV-2 strains [45].

Another study [38] investigated the infectivity of viral strains present in saliva after exposure to CHX solution. It was demonstrated that those viruses that came into contact with CHX (1.5%) completely lost their infectivity. In addition, treatment with 5% CHX had a moderate antiviral effect. Hydrogen peroxide and povidone-iodine had greater inhibitory effects on viruses than CHX. In general, CHX significantly blocked viral infectivity.

The efficacy of CHX against SARS-CoV-2 was also demonstrated in lower concentrations, such as 0.2%, inactivating > 99.9% of the viruses after 30 s and 60 s [44]. However, different results were also detected in the literature, as low [46, 47] virucidal efficacy was also seen in three other studies, using 0.1% to 1.0% CHX.

Still, in this context, three recent RCTs evaluated the efficacy of CHX in reducing the salivary SARS-CoV-2 viral load in patients diagnosed with coronavirus disease 2019 (COVID-19) [39, 41, 42]. One study assessed the reduction of the viral load of SARS-CoV-2 in the oropharynx [43]. It is important to highlight that these studies included patients with no need for nasogastric or endotracheal intubation. The time-lapse between COVID-19 diagnosis and inclusion in the trial was short. The reduction in viral load was measured using the rRT-PCR Cycle threshold (Ct) technique. Ct values are proportional inversely to viral load in this technique and can provide an indirect method of quantifying the copy number of viral RNA in the sample. It was observed that CHX demonstrated heterogeneous virucidal efficacy against SARS-CoV-2 present in saliva in one study [39]. After a single-use, the antiseptic failed to significantly reduce viral load when compared to the viral reduction observed in groups exposed to povidone-iodine, cetylpyridinium chloride and water [39].

Conversely, another study demonstrated a significantly lower viral load in the CHX group when compared to a control group (distilled water) [42]. This study also showed no significant difference when CHX and povidone-iodine were compared [42]. Another study also showed that a single rinse with CHX significantly reduced the viral load of SARS-CoV-2 after 30 min and 60 min [41]. However, this study showed that higher reductions in viral load were detected in individuals that rinsed with hydrogen peroxide or with cetylpyridinium chloride + zinc mouthwashes [41].

The study that assessed the reduction in viral load of SARS-CoV-2 in the oropharynx compared rinsing with CHX to standard care only [43]. A combination of spray and mouthwash with CHX was also compared to a standard care. In both test groups, rinse with CHX was performed for 4 days, and, in comparison to the control group, a number significantly higher of patients with negative detection of SARS-CoV-2 in the oropharynx was detected [43]. It

is important to highlight that all RCT reported no adverse events after a single use or after 4 days use of CHX.

Discussion

The present study aimed to systematically review the literature about the virucidal efficacy of CHX for some strains of viruses. Based on *in vitro* studies, it was demonstrated that CHX might reduce the levels of HSV-1 and Influenza A viruses, but lower to none *in vitro* effect may be expected to HCoV and SARS-CoV-2. In addition, a transient efficacy of CHX may be expected in patients diagnosed with COVID-19. It must be acknowledged that, among the included studies, the standards of testing such products were considerably heterogeneous, and the combination of data from different studies with the aforementioned draws peril for veritable comparisons. It may be hypothesized that the biophysiological mechanisms for the virucidal effect of CHX are the lysis of viral envelope, deterioration of nucleotide carbon chains, and impact on the inactivation and/or blocking of viral proteins [17, 24, 38], which may explain the efficacy against all the viruses tested in the present study.

The antiplaque and antigingivitis efficacy of CHX has been widely proven in the literature [6, 7]. In addition to these important effects, CHX can also reduce bacteria levels in aerosol [14]. The use of CHX is not restricted to Dentistry, as the literature demonstrates a reduction in the incidence of ventilator-associated pneumonia in adults [48] and prevention of infection in clean surgeries in adults [49]. In addition, vaginal cleansing with CHX demonstrated lower morbidity rates in post-cesarean [50]. Despite this knowledge, the present study is the first systematic review to assess the virucidal efficacy of CHX on human viruses affecting the oral cavity.

Greater efficacy of CHX against HSV-1 viruses was demonstrated in the present study. The clinical feature of HSV-1 infection is characterized by the appearance of vesicles on the skin or mucous membranes of the mouth and lips. This virus is mainly transmitted by oral-to-oral contact, but the occurrence of these lesions in the genital area may also be detected. However, this is highly attributed to the infection of HSV-2 [51]. HSV-1 structure is relatively large, presenting a linear DNA genome wrapped in a lipid bilayer, the envelope. Therefore, it may be speculated that the antiviral efficacy of CHX on enveloped viruses may be similar to the one observed in the cytoplasmic membrane of bacteria [52].

It is estimated that approximately 66.6% in those aged 0–49 years are living with HSV-1 in the world [53]. The literature reports that topical antiviral agents showed no efficacy in preventing herpes simplex labialis in individuals of all ages [54]. This is of utmost importance as HSV-1 infections may be self-limited. However, the HSV-1 vesicles

are very painful and may delay dental treatment when clinically detected. As an interesting effect of CHX was detected against this virus, further clinical trials are needed to establish these findings.

Another virus whose effect of CHX has been studied is Influenza A that is responsible for a disease transmitted through air, which causes fever, fatigue and sore throat. This virus is from the Orthomyxoviridae family and it has a negative sense RNA. Although infections with this virus are most common among birds [55], it also affects humans, which originated the 2009 pandemic of H1N1 [56]. A high hospitalization rate and hospitalization fatality risk are associated with these viruses, especially in developing countries [56].

The literature has demonstrated that saliva samples may be an excellent predictive tool for H1N1 [57], making mouthrinses very important to decrease contaminated aerosol with this virus. Only three included studies assessed the virucidal efficacy of CHX against Influenza A, and all of them demonstrated the CHX has excellent efficacy in inactivating this virus. The results were detected after 30 s to 10 min of contact with CHX. It is strongly recommended to perform clinical trials that may confirm these results using the appropriate time of rinsing with CHX.

Patients infected with SARS-CoV-2 may unknowingly spread the virus during the subclinical period by droplets. The literature shows a higher concentration of SARS-CoV-2 may be detected in saliva, salivary gland [58] and in the gingival crevicular fluid [59]. An agreement of 100% for the detection of SARS-CoV-2 nucleic acids using real-time reverse transcription PCR (RT-PCR) between samples of saliva and nasal swabs has also been reported [60]. In this sense, it has been proposed that saliva may be a possible tool for SARS-CoV-2 detection [61]. Therefore, a mouthrinse able to reduce the viral load in the mouth or, more specifically, in saliva may be necessary during dental procedures that produce aerosol.

Among the included studies, the *in vitro* virucidal efficacy of CHX against both HCoV and SARS-CoV-2 viruses was determined as lower to none. One RCT demonstrated that the reduction of viral load is similar to the one observed in water [39]. However, one prospective clinical trial, which did not use a control group, demonstrated that a high level of viruses is detected in the saliva, but CHX was able to significantly decrease the viral load for 2 h after a single-use [62]. Similar results were also detected in the other three RCT,

of which one demonstrated a significantly higher number of patients with negative detection of SARS-CoV-2 [43].

In all clinical trials, a low number of participants were included, which decrease the clinical applicability of these findings. Other substances, such as povidone-iodine [63] and hydrogen peroxide [64], might be an additional benefit against SARS-CoV-2, but their clinical efficacy remains unproved [65, 66]. It is noteworthy that the reduced timespan for the performance of large randomized clinical trials related to SARS-CoV-2, since it has been identified recently, could explain the scarcity of the literature. In addition, this might hinder possible effects to be determined in the future.

The present study looked at the effects of CHX on viruses. The interest in such effect stands beyond the control of COVID-19. The virucidal effect of an oral antiseptic is of interest in different biosafety procedures in the dental office and also in the disinfection of material used in oral prostheses. In addition, the concentrations of CHX varied between 0.001 [27] and 1.0% [33, 47] among the included studies. The clinical efficacy of CHX has been proven with at least 0.12% [67], and these different concentrations may also explain the results detected. It is important to highlight that only a few viruses were assessed, which may be a limitation of the present study. Almost all of the included studies are *in vitro* and presented considerable heterogeneity in determining the virucidal efficacy using CHX. Therefore, the clinical applicability of results detected may be limited, and this must be considered when interpreting the results presented. However, the present study performed a broader search in the literature about the virucidal efficacy of CHX in all forms and concentrations. In addition, no restriction to language and date of publication was imposed, which are the main strengths of this systematic review. The interpretation of the findings should be performed considering the date of search and a continuous update in the literature is recommended.

In conclusion, CHX may present an interesting virucidal efficacy against HSV-1 and Influenza A viruses. However, reductions of HCoV and SARS-CoV-2 strains, when assessed *in vitro*, have not yet been demonstrated. In addition, rising with CHX may temporarily reduce the viral load of SARS-CoV-2 in patients with COVID-19.

Appendix 1

Table 3.

Table 3 Search strategy and number of studies detected in each all database

Database	Search strategy	Number of studies
PubMed	#1 Chlorhexidine[Mesh Terms] OR Chlorhexidine[Title/abstract] OR Biguanides[MeSH Terms] OR Biguanides[Title/abstract] OR chlorhexidine gluconate[Supplementary Concept] OR Mouthwashes[Mesh Terms] OR mouthwashes[Title/abstract] OR mouthrinses[Title/abstract] OR "mouthwash"[Title/abstract] OR "mouthrinse"[Title/abstract] OR Anti-Infective Agents, Local[Mesh Terms] OR Local anti-infective agents[Title/abstract] OR Tubulicid[Title/abstract] OR Novalsan[Title/abstract] OR Sebidin A[Title/abstract] #2 COVID-19[Supplementary Concept] OR COVID-19[Title/abstract] OR Coronavirus[Mesh Terms] OR Coronavirus[Title/abstract] OR Coronaviruses[Title/abstract] OR Coronaviridae[Title/abstract] OR Coronavirus Infections[Mesh Term] OR Coronavirus Infections[Title/abstract] OR Virus Inactivation[Mesh Terms] OR Virus Inactivation[Title/abstract] OR Virucidal[Title/abstract] OR Viral Inactivation[Title/abstract] OR virucide[Title/abstract] OR SARS-CoV-2[Title/abstract] OR virus ^a [Title/abstract] OR viruses[Mesh Terms] OR viruses[Title/abstract] OR viral load[Mesh Terms] OR viral load[Title/abstract] OR virus cultivation[Mesh Terms] OR virus cultivation[Title/abstract] OR viral cultivation[Title/abstract] OR viral[Title/abstract] OR virology[Title/abstract] OR Viral Burden[Title/abstract] #3—#1 AND #2	1225
Scopus	((TITLE-ABS-KEY("Hydrogen Peroxide") OR TITLE-ABS-KEY(H2O2) OR TITLE-ABS-KEY(Hydroperoxide) OR TITLE-ABS-KEY(Peroxides) OR TITLE-ABS-KEY(Superoxol) OR TITLE-ABS-KEY(Oxydol) OR TITLE-ABS-KEY(Perhydrol))) and ((TITLE-ABS-KEY(COVID-19) OR TITLE-ABS-KEY(Coronavirus) OR TITLE-ABS-KEY(Coronaviruses) OR TITLE-ABS-KEY(Coronaviridae) OR TITLE-ABS-KEY("Virus Inactivation") OR TITLE-ABS-KEY(Virucidal) OR TITLE-ABS-KEY("Viral Inactivation") OR TITLE-ABS-KEY(virucide) OR TITLE-ABS-KEY(SARS-CoV-2) OR TITLE-ABS-KEY("viral load") OR TITLE-ABS-KEY("virus cultivation") OR TITLE-ABS-KEY(viral) OR TITLE-ABS-KEY("Viral Burden"))))	1741
Web of Science	#1—TS = (chlorhexidine OR biguanides OR biguanide OR "chlorhexidine gluconate" OR mouthwashes OR mouthrinses OR mouthwash OR mouthrinse OR "Local anti-infective agents" OR tubulicid OR novalsan OR "Sebidin A") #2—TS = (covid-19 OR coronavirus OR coronaviruses OR coronaviridae OR "Coronavirus Infections" OR "Virus Inactivation" OR virucidal OR "Viral Inactivation" OR virucide OR sars-cov-2 OR virus ^a OR viruses OR "viral load" OR "virus cultivation" OR "viral cultivation" OR viral OR virology OR "Viral Burden") #3—#1 AND #2	351
Embase	#1—chlorhexidine OR biguanides OR biguanide OR "chlorhexidine gluconate" OR mouthwashes OR mouthrinses OR mouthwash OR mouthrinse OR "Local anti-infective agents" OR tubulicid OR novalsan OR "Sebidin A" #2—covid-19 OR coronavirus OR coronaviruses OR coronaviridae OR "Coronavirus Infections" OR "Virus Inactivation" OR virucidal OR "Viral Inactivation" OR virucide OR sars-cov-2 OR virus ^a OR viruses OR "viral load" OR "virus cultivation" OR "viral cultivation" OR viral OR virology OR "Viral Burden" #3—#1 AND #2	1562
Total		4879
Duplicates		2217

^aSearch strategy performed on July 9th, 2021

Author contributions Conceptualization: MIFG, CKR, JC, and FWMGM. Methodology: CKR, JC, and FWMGM. Formal analysis and investigation: MSF, MIFG, GPJL, and FWMGM. Writing—original draft preparation: MSF and FWMGM. Writing—review and editing: MSF, MIFG, GPJL, CKR, JC, and FWMGM. Supervision: FWMGM and MIFG.

Funding This study was self-funded.

Availability of data and material Data we will forward upon request.

Code availability Not applicable.

Declarations

Conflict of interest The authors report no conflict of interest related to this study.

Ethical approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

References

1. Arteagoitia I, Rodriguez Andrés C, Ramos E. Does chlorhexidine reduce bacteremia following tooth extraction? A systematic review and meta-analysis. *PLoS One*. 2018;13: e0195592. <https://doi.org/10.1371/journal.pone.0195592>.
2. Rosenberg A, Alatary SD, Peterson AF. Safety and efficacy of the antiseptic chlorhexidine gluconate. *Surg Gynecol Obstet*. 1976;143:789–92.

3. McDonnell G, Russell AD. Antiseptics and disinfectants: activity, action, and resistance. *Clin Microbiol Rev*. 1999;12:147–79.
4. Milstone AM, Passaretti CL, Perl TM. Chlorhexidine: expanding the armamentarium for infection control and prevention. *Clin Infect Dis*. 2008;46:274–81. <https://doi.org/10.1086/524736>.
5. Popovich KJ, Lyles R, Hayes R, et al. Relationship between chlorhexidine gluconate skin concentration and microbial density on the skin of critically ill patients bathed daily with chlorhexidine gluconate. *Infect Control Hosp Epidemiol*. 2012;33:889–96. <https://doi.org/10.1086/667371>.
6. James P, Worthington HV, Parnell C, et al. Chlorhexidine mouthrinse as an adjunctive treatment for gingival health. *Cochrane Database Syst Rev*. 2017;3: Cd008676. <https://doi.org/10.1002/14651858.CD008676.pub2>.
7. Afenich F, Slot DE, Hossainian N, Van der Weijden GA. The effect of hexetidine mouthwash on the prevention of plaque and gingival inflammation: a systematic review. *Int J Dent Hyg*. 2011;9:182–90. <https://doi.org/10.1111/j.1601-5037.2010.00478.x>.
8. Zhao H, Hu J, Zhao L. Adjunctive subgingival application of Chlorhexidine gel in nonsurgical periodontal treatment for chronic periodontitis: a systematic review and meta-analysis. *BMC Oral Health*. 2020;20:34. <https://doi.org/10.1186/s12903-020-1021-0>.
9. Sharma K, Acharya S, Verma E, Singhal D, Singla N. Efficacy of chlorhexidine, hydrogen peroxide and tulsi extract mouthwash in reducing halitosis using spectrophotometric analysis: a randomized controlled trial. *J Clin Exp Dent*. 2019;11:e457–63. <https://doi.org/10.4317/jced.55523>.
10. Fedorowicz Z, Aljufairi H, Nasser M, Outhouse TL, Pedrazzi V. Mouthrinses for the treatment of halitosis. *Cochrane Database Syst Rev*. 2008. <https://doi.org/10.1002/14651858.CD006701.pub2>.
11. Richards D. Chlorhexidine mouthwash plaque levels and gingival health. *Evid Based Dent*. 2017;18:37–8. <https://doi.org/10.1038/sj.ebd.6401232>.
12. Flötra L, Gjermeo P, Rölla G, Waerhaug J. Side effects of chlorhexidine mouth washes. *Scand J Dent Res*. 1971;79:119–25. <https://doi.org/10.1111/j.1600-0722.1971.tb02001.x>.
13. Bescos R, Ashworth A, Cutler C, et al. Effects of chlorhexidine mouthwash on the oral microbiome. *Sci Rep*. 2020;10:5254. <https://doi.org/10.1038/s41598-020-61912-4>.
14. Marui VC, Souto MLS, Rovai ES, Romito GA, Chambrone L, Pannuti CM. Efficacy of preprocedural mouthrinses in the reduction of microorganisms in aerosol: a systematic review. *J Am Dent Assoc*. 2019;150:1015–26.e1. <https://doi.org/10.1016/j.adaj.2019.06.024>.
15. Jones CG. Chlorhexidine: is it still the gold standard? *Periodontol*. 1997;15:55–62. <https://doi.org/10.1111/j.1600-0757.1997.tb00105.x>.
16. Page M, Moher D, Bossuyt P, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372: n160. <https://doi.org/10.1136/bmj.n160>.
17. Karpiński TM, Szkaradkiewicz AK. Chlorhexidine—pharmacological activity and application. *Eur Rev Med Pharmacol Sci*. 2015;19:1321–6.
18. Koletsi D, Belibasakis GN, Eliades T. Interventions to reduce aerosolized microbes in dental practice: a systematic review with network meta-analysis of randomized controlled trials. *J Dent Res*. 2020;99:1228–38. <https://doi.org/10.1177/0022034520943574>.
19. Assis MS, Araújo R, Lopes AMM. Safety alert for hospital environments and health professional: chlorhexidine is ineffective for coronavirus. *Rev Assoc Med Bras* (1992). 2020;66Suppl 2:124–9. <https://doi.org/10.1590/1806-9282.66.S2.124>.
20. Joanna Briggs Institute. Joanna Briggs Institute reviewers' manual: 2014 edition. Australia: The Joanna Briggs Institute; 2014.
21. Krithikadatta J, Gopikrishna V, Datta M. CRIS guidelines (checklist for reporting in-vitro studies): a concept note on the need for standardized guidelines for improving quality and transparency in reporting in-vitro studies in experimental dental research. *J Conserv Dent*. 2014;17:301–4. <https://doi.org/10.4103/0972-0707.136338>.
22. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366: 14898. <https://doi.org/10.1136/bmj.14898>.
23. Bailey A, Longson M. Virucidal activity of chlorhexidine on strains of Herpesvirus hominis, poliovirus, and adenovirus. *J Clin Pathol*. 1972;25:76.
24. Baqui AA, Kelley JI, Jabra-Rizk MA, Depaola LG, Falkler WA, Meiller TF. In vitro effect of oral antiseptics on human immunodeficiency virus-1 and herpes simplex virus type 1. *J Clin Periodontol*. 2001;28:610–6. <https://doi.org/10.1034/j.1600-051x.2001.028007610.x>.
25. Damery B, Cremieux A. Virucidal activity against Herpes and Vaccinia virus of 8 antiseptic formulations. *Int J Pharmaceut*. 1989;49:205–8.
26. Park N-H, Park JB, Min B-M, Cherrick HM. Combined synergistic antiherpetic effect of acyclovir and chlorhexidine in vitro. *Oral Sur Oral Med Oral Pathol*. 1991;71:193–6. [https://doi.org/10.1016/0030-4220\(91\)90467-q](https://doi.org/10.1016/0030-4220(91)90467-q).
27. Park JB, Park N-H. Effect of chlorhexidine on the in vitro and in vivo herpes simplex virus infection. *Oral Sur Oral Med Oral Pathol*. 1989;67:149–53. [https://doi.org/10.1016/0030-4220\(89\)90320-4](https://doi.org/10.1016/0030-4220(89)90320-4).
28. Shinkai K, Yoshino K. Different sensitivities of type 1 and type 2 Herpes simplex virus to sodium p-chloromercuribenzoate and chlorhexidine gluconate. *Proc Soc Exp Biol Med*. 1974;147:201–4. <https://doi.org/10.3181/00379727-147-38311>.
29. Tyler R, Ayliffe G. A surface test for virucidal activity of disinfectants: preliminary study with herpes virus. *J Hosp Infect*. 1987;9:22–9. [https://doi.org/10.1016/0195-6701\(87\)90090-9](https://doi.org/10.1016/0195-6701(87)90090-9).
30. Okunishi J, Nishihara Y, Maeda S, Ikeda M. In vitro evaluation of the antimicrobial activity of HM-242, a novel antiseptic compound. *J Antibiot*. 2009;62:489–93. <https://doi.org/10.1038/ja.2009.56>.
31. Bernstein D, Schiff G, Echler G, Prince A, Feller M, Briner W. In vitro virucidal effectiveness of a 0.12%-chlorhexidine gluconate mouthrinse. *J Dent Res*. 1990;69:874–6. <https://doi.org/10.1177/00220345900690030901>.
32. Kawana R, Kitamura T, Nakagomi O, et al. Inactivation of human viruses by povidone-iodine in comparison with other antiseptics. *Dermatology*. 1997;195Suppl 2:29–35. <https://doi.org/10.1159/000246027>.
33. Ansaldi F, Banfi F, Morelli P, et al. SARS-CoV, influenza A and syncytial respiratory virus resistance against common disinfectants and ultraviolet irradiation. *J Prev Med Hyg*. 2004;45:5–8.
34. Geller C, Fontanay S, Finance C, Duval REA. new Sephadex-based method for removing microbicidal and cytotoxic residues when testing antiseptics against viruses: experiments with a human coronavirus as a model. *J Virol Methods*. 2009;159:217–26. <https://doi.org/10.1016/j.jviromet.2009.03.023>.
35. Sattar SA, Springthorpe VS, Karim Y, Loro P. Chemical disinfection of non-porous inanimate surfaces experimentally contaminated with four human pathogenic viruses. *Epidemiol Infect*. 1989;102:493–505. <https://doi.org/10.1017/s095026880030211>.
36. Imai K, Hagi A, Yamagawa S, Sano D. Olanexidine gluconate formulations as environmental disinfectants for enveloped viruses infection control. *J Hosp Infect*. 2021;112:37–41. <https://doi.org/10.1016/j.jhin.2021.03.014>.

37. Meister TL, Brüggemann Y, Todt D, et al. Virucidal efficacy of different oral rinses against severe acute respiratory syndrome coronavirus 2. *J Infect Dis.* 2020;222:1289–92. <https://doi.org/10.1093/infdis/jiaa471>.
38. Xu C, Wang A, Hoskin ER, et al. Differential effects of antiseptic mouth rinses on SARS-CoV-2 infectivity in vitro. *Pathogens.* 2021;10:272. <https://doi.org/10.3390/pathogens10030272>.
39. Seneviratne CJ, Balan P, Ko KKK, et al. Efficacy of commercial mouth-rinses on SARS-CoV-2 viral load in saliva: randomized control trial in Singapore. *Infection.* 2020;2020:1–7. <https://doi.org/10.1007/s15010-020-01563-9>.
40. Davies K, Buczkowski H, Welch SR, et al. Effective in vitro inactivation of SARS-CoV-2 by commercially available mouthwashes. *J Gen Virol.* 2021;102: 001578. <https://doi.org/10.1099/jgv.0.001578>.
41. Eduardo FP, Corrêa L, Heller D, et al. Salivary SARS-CoV-2 load reduction with mouthwash use: a randomized pilot clinical trial. *Heliyon.* 2021;7: e07346. <https://doi.org/10.1016/j.heliyon.2021.e07346>.
42. Elzein R, Abdel-Sater F, Fakhreddine S, et al. In vivo evaluation of the virucidal efficacy of chlorhexidine and povidone-iodine mouthwashes against salivary SARS-CoV-2. A randomized-controlled clinical trial. *J Evid Based Dent Pract.* 2021. <https://doi.org/10.1016/j.jebdp.2021.101584>.
43. Huang YH, Huang JT. Use of chlorhexidine to eradicate oropharyngeal SARS-CoV-2 in COVID-19 patients. *J Med Virol.* 2021;93:4370–3. <https://doi.org/10.1002/jmv.26954>.
44. Jain A, Grover V, Singh C. Chlorhexidine: an effective anticovid mouth rinse. *J Indian Soc Periodontol.* 2021;25:86–8. https://doi.org/10.4103/jisp.jisp_824_20.
45. Komine A, Yamaguchi E, Okamoto N, Yamamoto K. Virucidal activity of oral care products against SARS-CoV-2 in vitro. *J Oral Maxillofac Surg Med Pathol.* 2021;33:475–7. <https://doi.org/10.1016/j.ajoms.2021.02.002>.
46. Steinhauer K, Meister TL, Todt D. Comparison of the in-vitro efficacy of different mouthwash solutions targeting SARS-CoV-2 based on the European Standard EN 14476. *J Hosp Infect.* 2021;111:180–3. <https://doi.org/10.1016/j.jhin.2021.01.031>.
47. Hirose R, Bandou R, Ikegaya H, et al. Disinfectant effectiveness against SARS-CoV-2 and influenza viruses present on human skin: model-based evaluation. *Clin Microbiol Infect.* 2021;27(1042):e1–4. <https://doi.org/10.1016/j.cmi.2021.04.009>.
48. Villar CC, Pannuti CM, Nery DM, Morillo CM, Carmona MJ, Romito GA. Effectiveness of intraoral chlorhexidine protocols in the prevention of ventilator-associated pneumonia: meta-analysis and systematic review. *Respir Care.* 2016;61:1245–59. <https://doi.org/10.4187/respcare.04610>.
49. Wade RG, Burr NE, McCauley G, Bourke G, Efthimiou O. The comparative efficacy of chlorhexidine gluconate and povidone-iodine antiseptics for the prevention of infection in clean surgery: a systematic review and network meta-analysis. *Ann Surg.* 2020. <https://doi.org/10.1097/SLA.0000000000004076>.
50. Ogah CO, Anikwe CC, Ajah LO, et al. Pre-operative vaginal cleansing with chlorhexidine solution in preventing post-caesarean section infections in a low resource setting: a randomized controlled trial. *Acta Obstet Gynecol Scand.* 2021. <https://doi.org/10.1111/aogs.14060>.
51. Looker KJ, Welton NJ, Sabin KM, et al. Global and regional estimates of the contribution of herpes simplex virus type 2 infection to HIV incidence: a population attributable fraction analysis using published epidemiological data. *Lancet Infect Dis.* 2020;20:240–9. [https://doi.org/10.1016/S1473-3099\(19\)30470-0](https://doi.org/10.1016/S1473-3099(19)30470-0).
52. Thangavelu A, Kaspar SS, Kathirvelu RP, Srinivasan B, Srinivasan S, Sundram R. Chlorhexidine: an elixir for periodontics. *J Pharm Bioallied Sci.* 2020;12:S57–9. https://doi.org/10.4103/jpbs.JPBS_162_20.
53. James C, Harfouche M, Welton NJ, et al. Herpes simplex virus: global infection prevalence and incidence estimates, 2016. *Bull World Health Organ.* 2020;98:315–29. <https://doi.org/10.2471/BLT.19.237149>.
54. Chi CC, Wang SH, Delamere FM, Wojnarowska F, Peters MC, Kanjirath PP. Interventions for prevention of herpes simplex labialis (cold sores on the lips). *Cochrane Database Syst Rev.* 2015;2015: CD010095. <https://doi.org/10.1002/14651858.CD010095.pub2>.
55. Kalonda A, Saasa N, Nkhoma P, et al. Avian influenza viruses detected in birds in sub-Saharan Africa: a systematic review. *Viruses.* 2020;12:993. <https://doi.org/10.3390/v12090993>.
56. Wong JY, Kelly H, Cheung CM, et al. Hospitalization fatality risk of influenza A(H1N1)pdm09: a systematic review and meta-analysis. *Am J Epidemiol.* 2015;182:294–301. <https://doi.org/10.1093/aje/kwv054>.
57. Bilder L, Machtei EE, Shenhar Y, Kra-Oz Z, Basis F. Salivary detection of H1N1 virus: a clinical feasibility investigation. *J Dent Res.* 2011;90:1136–9. <https://doi.org/10.1177/0022034511413283>.
58. da Silva PM, Sipert CR, Nogueira FN. Are the salivary glands the key players in spreading COVID-19 asymptomatic infection in dental practice? *J Med Virol.* 2020;93:204–5. <https://doi.org/10.1002/jmv.26316>.
59. Elisetti N. Periodontal pocket and COVID-19: could there be a possible link? *Med Hypotheses.* 2021;146: 110355. <https://doi.org/10.1016/j.mehy.2020.110355>.
60. Abasiyanik MF, Flood B, Lin J, et al. Sensitive detection and quantification of SARS-CoV-2 in saliva. *Sci Rep.* 2021;11:12425. <https://doi.org/10.1038/s41598-021-91835-7>.
61. Medeiros da Silva RC, Nogueira Marinho LC, de Araújo Silva DN, Costa de Lima K, Pirih FQ, de Aquino L, Martins AR. Saliva as a possible tool for the SARS-CoV-2 detection: a review. *Travel Med Infect Dis.* 2020;38: 101920. <https://doi.org/10.1016/j.tmaid.2020.101920>.
62. Yoon JG, Yoon J, Song JY, et al. Clinical Significance of a High SARS-CoV-2 Viral Load in the Saliva. *J Korean Med Sci.* 2020;35: e195. <https://doi.org/10.3346/jkms.2020.35.e195>.
63. Cimolai N. Efficacy of povidone-iodine to reduce viral load. *Oral Dis.* 2020;26:1832. <https://doi.org/10.1111/odi.13557>.
64. Santana LADM, Pinho JNA, de Albuquerque HIM, Souza LMA. Virucidal potential of H₂O₂-based spray against SARS-CoV-2 and biosafety in a dental environment. *Oral Dis.* 2021. <https://doi.org/10.1111/odi.13778>.
65. Muniz FWMG, Cavagni J, Langa GPJ, Stewart B, Malheiros Z, Rösing CK. A systematic review of the effect of oral rinsing with H. *Int J Dent.* 2020;2020:8841722. <https://doi.org/10.1155/2020/8841722>.
66. Burton MJ, Clarkson JE, Goulao B, et al. Antimicrobial mouthwashes (gargling) and nasal sprays administered to patients with suspected or confirmed COVID-19 infection to improve patient outcomes and to protect healthcare workers treating them. *Cochrane Database Syst Rev.* 2020;9: CD013627. <https://doi.org/10.1002/14651858.CD013627.pub2>.
67. Van Strydonc DAC, Slot DE, Van der Velden U, Van der Weijden F. Effect of a chlorhexidine mouthrinse on plaque, gingival inflammation and staining in gingivitis patients: a systematic review. *J Clin Periodontol.* 2012;39:1042–55. <https://doi.org/10.1111/j.1600-051X.2012.01883.x>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.