

Virulence contribution and  
recognition of homologs of the  
***Verticillium dahliae***  
effector Ave1

Jordi C. Boshoven

# Propositions

1. *Verticillium dahliae* effector *Ave1L* has been under selection pressure of an immune receptor.  
(this thesis)
2. Residues at both the N- and C-terminus of the *Ave1* effector are required for recognition by tomato immune receptor *Ve1*.  
(this thesis)
3. Organic agriculture alone cannot feed the future world population.
4. We should be more concerned about the impact of pesticides on the environment, than about growing genetically modified crops that can replace pesticides.
5. Competition between scientists for research grants shows parallels with competition between microorganisms in a microbial community since in both cases occupation of a unique niche increases the chances of success.
6. Open access journals should become the forum of science allowing knowledge to become available to anyone interested.
7. Raising a child during a PhD creates a healthy balance between time spent to work and private life.
8. For a Dutchman it is easier to write a PhD thesis than to learn to speak Greek.

Propositions belonging to the thesis, entitled  
Virulence contribution and recognition of homologs of the  
*Verticillium dahliae* effector *Ave1*

Jordi C. Boshoven  
Wageningen, 10 November 2017

**Virulence contribution and recognition  
of homologs of the  
*Verticillium dahliae* effector Ave1**

**Jordi C. Boshoven**

## **Thesis committee**

### **Promotors**

Prof. Dr B.P.H.J. Thomma  
Professor of Phytopathology  
Wageningen University & Research

Prof. Dr P.J.G.M de Wit  
Professor of Phytopathology  
Wageningen University & Research

### **Other members**

Prof. Dr M.G.M. Aarts, Wageningen University & Research  
Prof. Dr A. Pérez-García, University of Málaga, Spain  
Dr Y. Bai, Wageningen University & Research  
Dr M. de Vos, Keygene N.V., Wageningen

This research was conducted under the auspices of the Graduate School  
Experimental Plant Sciences.

**Virulence contribution and recognition  
of homologs of the  
*Verticillium dahliae* effector Ave1**

**Jordi C. Boshoven**

**Thesis**

submitted in fulfilment of the requirements for the degree of doctor  
at Wageningen University  
by the authority of the Rector Magnificus,  
Prof. Dr A.P.J. Mol,  
in the presence of the  
Thesis Committee appointed by the Academic Board  
to be defended in public  
on Friday 10 November 2017  
at 11 a.m. in the Aula.

Jordi C. Boshoven

Virulence contribution and recognition of homologs of the *Verticillium dahliae* effector Ave1,

184 pages.

PhD thesis, Wageningen University, Wageningen, the Netherlands (2017)

With references, with summary in English

ISBN 978-94-6343-644-1

DOI <http://dx.doi.org/10.18174/420169>



# Contents

<b>Chapter 1</b>	General introduction	<b>9</b>
<b>Chapter 2</b>	Filamentous Pathogen Effector Functions: Of Pathogens, Hosts and Microbiomes	<b>19</b>
<b>Chapter 3</b>	Homologs of <i>Verticillium dahliae</i> effector Ave1 contribute to virulence in fungal pathogens of diverse plant hosts	<b>35</b>
<b>Chapter 4</b>	A nine amino acid peptide of <i>Verticillium dahliae</i> effector Ave1 is required but not sufficient for recognition by tomato immune receptor Ve1	<b>55</b>
<b>Chapter 5</b>	Characterization of an expanded family of Ave1 homologs in the apple and pear scab fungi <i>Venturia inaequalis</i> and <i>V. pirina</i>	<b>85</b>
<b>Chapter 6</b>	An Ave1-like effector displays allelic variation within the <i>Verticillium dahliae</i> population	<b>97</b>
<b>Chapter 7</b>	Rhamnose synthase activity is required for pathogenicity of <i>Verticillium dahliae</i>	<b>121</b>
<b>Chapter 8</b>	General discussion: From effector discovery to breeding for durable pathogen resistance in crops	<b>151</b>
	<b>Summary</b>	<b>171</b>
	<b>Acknowledgements</b>	<b>175</b>
	<b>About the author</b>	<b>178</b>
	<b>List of publications</b>	<b>179</b>
	<b>Education statement</b>	<b>180</b>





*To my family*





**General introduction**

**Chapter 1**



## Introduction

Plants provide us with food, feed, fuel, fibre, shelter and drugs and are therefore an important natural resource on which we depend heavily. Since the world population has been, and still is, rapidly growing, the meat and dairy intake has been increasing and also the use of biofuel is rising. Collectively, this still results in an urgent demand for increased crop production. The green revolution that started in the 60s of the 20<sup>th</sup> century intensified global agricultural practice significantly, often resulting in vast areas of a single crop cultivar known as a monoculture (Jain 2010). Disease outbreaks caused by plant pathogens including fungi, oomycetes, bacteria and viruses can lead to severe yield losses and can even wipe out entire crops, especially when cultivated as monoculture (Oerke 2006; Strange and Scott 2005). To find novel and improved means to combat disease outbreaks, a fundamental understanding of the molecular basis of host-pathogen interactions is essential. Already in the 40s of the 20<sup>th</sup> century, Harald Flor showed that plants that carry resistance (*R*) genes are resistant to pathogens that carry corresponding avirulence (*Avr*) genes (Flor 1942). Initially, it was thought that *R* and *Avr* proteins directly interact (Keen 1990). However, most protein interaction studies failed to confirm this hypothesis, which led to the proposal of the guard hypothesis (Van der Biezen and Jones 1998). This hypothesis states that an *R* protein guards a plant protein that, when targeted by a pathogen *Avr* protein, triggers a resistance response. It subsequently became clear that *Avr* proteins not only initiate resistance responses, but actually act as pathogen virulence factors that manipulate host proteins in absence of host resistance proteins. Therefore, the term effector is more frequently used for *Avr* proteins nowadays (Boller and Felix 2009). Effectors are considered to be species-, or even race- or strain-specific (Chisholm et al. 2006). Besides sensing effectors or their activities, plants recognize conserved structural components of microorganisms such as bacterial flagellin, fungal chitin and oomycete  $\beta$ -glucans, known as microbe-associated molecular patterns (MAMPs) resulting in MAMP-triggered immunity (MTI) (Dodds and Rathjen 2010). It is generally thought that pathogens evolved effectors to overcome this MTI response. In turn, however, plants evolved receptors to recognize these effectors, resulting in effector-triggered immunity (ETI). These receptors are the earlier discussed *R* proteins. To avoid recognition by the plant, pathogens have to lose or mutate the recognized effector, or evolve new effectors that can again suppress the immune response. These processes continuously occur in the ongoing arms-race that takes place between plants and pathogens (Chisholm et al. 2006; Jones and Dangl 2006).

1

About a decade after its introduction, several misconceptions are ubiquitously professed when describing the plant immune system, such as strict separation of MTI versus ETI (Cook et al. 2015; Thomma et al. 2011). Therefore, it was recently proposed that an accurate description of inducible plant immunity should explicitly separate host perception of a given ligand from its physiological or biochemical function. This is captured in the Invasion Model in which host receptors, termed invasion pattern receptors (IPRs), detect ligands that accurately betray pathogen ingress, termed invasion patterns (IPs)(Cook et al. 2015). These IPs can either be an externally encoded or modified-self ligands that indicates invasion. Although any molecule may serve as an IP that will be detected by an IPR, the probability of any given ligand-receptor complex to develop an important role in host immunity increases with increasing ligand molecular constraint to retain function, conservation across organisms, importance in facilitating symbiosis, and accessibility (Cook et al. 2015).

## The study system

The genus *Verticillium* includes soil-borne fungal vascular pathogens that are causal agents of Verticillium wilts in a broad range of host plants, including many economically important crops (Agrios 2005), leading to billions of euros of damage annually (Pegg and Brady 2002). We use the species *V. dahliae* as a model system since it has a broad host range and, consequently, can infect model plants such as Arabidopsis, tomato and tobacco. The lifecycle of *V. dahliae* starts with the germination of resting structures, microsclerotia, in the soil after sensing of root exudates. The hyphae that grow from the germinating microsclerotia will progress towards the root and penetrate the root tips (Fitzell et al. 1980; Mol 1995). *V. dahliae* grows through the root cortex into the xylem vessels where it produces conidia that will further spread through the plant (Bishop and Cooper 1983; Fitzell et al. 1980; Garber and Houston 1966). Growth of the fungus in the xylem vessels blocks water transport, which will ultimately lead to wilting and dying of the tissue. Microsclerotia will form on dying tissue, and will eventually end up in the soil after the decomposition of plant material. *V. dahliae* is difficult to control since it has a broad host range and microsclerotia can survive up to 20 years in the soil, which makes crop rotation ineffective. Most fungicides are ineffective as well because they do not reach the fungus which grows protected inside the plant. Although soil fumigation is effective against *V. dahliae* most countries have banned this practice due to its detrimental effects on the environment (Fradin and Thomma 2006).

An effective way to control *V. dahliae* is genetic resistance conferred by tomato immune receptor Ve1 of which homologs are found amongst others in particular tobacco genotypes, mint, hop and cotton (Majer et al. 2014; Vining and Davis 2009; Zhang

et al. 2011; Zhang et al. 2013). Nevertheless, functionality of the *Ve1* homologs was only reported in *Nicotiana glutinosa* and cotton (Zhang et al. 2011; Zhang et al. 2013). Plants carrying *Ve1* are resistant to race 1 strains of *V. dahliae*, but not to race 2 strains (Fradin et al. 2009; Kawchuk et al. 2001). Recently, the effector that is recognized by *Ve1* was identified in race 1 strains and named *Ave1*, for mediating avirulence on **Ve1** plants (de Jonge et al. 2012). Besides that *Ave1* deletion strains are no longer recognized by *Ve1*, they show reduced aggressiveness on tomato plants lacking *Ve1*, indicating a role in virulence (de Jonge et al. 2012). In addition, *Ave1* deletion was shown to compromise aggressiveness on *Arabidopsis* as well (de Jonge, van Esse et al. 2012). Remarkably, homologs of *Ave1* are mainly found in plants and only in a few microorganisms that all happen to be plant pathogens, including the fungi *Fusarium oxysporum* f. sp. *lycopersici*, *Cercospora beticola*, *Colletotrichum higginsianum* and the bacterium *Xanthomonas axonopodis* (de Jonge et al. 2012). The *Ave1* homologs from plant pathogens were coexpressed with *Ve1* in *Nicotiana tabacum* to determine whether they can be recognized as well. This experiment showed that homologs from *F. oxysporum* and *C. beticola* are recognized while the homologs from *C. higginsianum* and *X. axonopodis* are not, or only poorly, recognized (de Jonge et al. 2012). Since *F. oxysporum* f. sp. *lycopersici* is a tomato pathogen the fungus was inoculated on transgenic tomato carrying *Ve1*. These *Ve1* tomato plants showed reduced disease symptoms when compared with plants that lack *Ve1*, suggesting that recognition of *FoAve1* activates host defense (de Jonge et al. 2012).

The *Ave1* homologs that were found in plants are annotated as plant natriuretic peptides or expansin-like proteins that are described to be involved in ion homeostasis, and therefore in many downstream processes including growth, net water uptake, photosynthesis, stomatal opening and gas exchange (Gehring and Irving 2003; Ludidi et al. 2004; Maryani et al. 2001; Pharmawati et al. 2001). Phylogenetic analysis showed that *Ave1* homologs found in microbes do not follow phylogeny of the species in which they occur, but rather occur among plant homologs, which is suggestive of horizontal gene transfer. Therefore, our working hypothesis is that *Ave1* homologs found in microbes are acquired from plants through horizontal gene transfer (de Jonge et al. 2012; Gottig et al. 2008). Interestingly, the *Ave1* homolog that is found in *X. axonopodis* known as *XacPNP* is a virulence factor that affects homeostasis and photosynthesis in citrus plants (Garavaglia et al. 2010; Gottig et al. 2008; Nembaware et al. 2004). Since *Ave1* homologs are found in multiple plant pathogens and are likely important for virulence, further studies are needed to gain more understanding of the function of these homologs.



## Main objective and research questions

The main objective of this PhD research is to study the *Ave1* homologs derived from *Verticillium* species and other plant pathogens. To this end we addressed the following research questions:

- How are the *Ave1* homologs distributed?
- What is the role of *Ave1* homologs in virulence in the pathogens that express them?
- What is the function of the *Ave1* homologs?
- Do all *Ave1* homologs display the same functionality?
- What determines recognition of *Ave1* homologs by *Ve1*?
- How did the *Ave1* homologs evolve?

## Thesis outline

In **Chapter 2**, a complete overview of the biological function of fungal effectors is provided. Moreover, the most recent views on various manners how fungal effectors contribute to disease establishment is provided. Furthermore we discuss potential roles of fungal effectors outside host-pathogen interactions.

In **Chapter 3**, we studied the *Ave1* homologs of three fungal pathogens: *F. oxysporum* f. sp. *lycopersici*, *C. beticola* and *C. higginsianum* that infect tomato, sugarbeet and Arabidopsis, respectively. We determine whether the *Ave1* homologs are expressed by the respective fungi during host colonization and whether these genes play a role in virulence.

In **Chapter 4**, the recognition of *V. dahliae* *Ave1* by the tomato *Ve1* immune receptor is studied. Based on comparisons of *Ave1* homologs from various species and functional assays with truncated proteins and domain swaps, attempts are made to determine the region of *Ave1* that is recognized by *Ve1*.

In **Chapter 5**, the expanded *Ave1* gene family that is found in the genomes of the apple and pear scab pathogens *Venturia inaequalis* and *V. pirina* is described. RNAseq of *V. inaequalis* revealed that some of the paralogs are induced *in planta*. It is studied whether the induced *Ave1* homologs are recognized by *Ve1* and whether they may perform the same biological function as *V. dahliae* *Ave1*.

In **Chapter 6**, we describe the occurrence of an *Ave1-like* gene that is found in many *V. dahliae* strains. In our analysis, we identified five different alleles of this gene. We determine whether the various *Ave1-like* gene alleles are expressed during host colonization, and whether *Ve1* or its close homolog *Ve2* is able to recognize these.

In **Chapter 7**, a random mutagenesis was performed in *V. dahliae* to identify novel virulence and pathogenicity factors. One of the virulence factors that was identified is further characterized.

**Chapter 8** is the general discussion and describes how the knowledge of pathogen effectors can be used as a tool to establish durable broad spectrum resistance.

## References

- Agrios, G. N. 2005. *Plant Pathology*. 5th ed. San Diego: Academic Press.
- Bishop, C., and Cooper, R. M. 1983. An ultrastructural study of root invasion in three vascular wilt diseases. *Physiol Plant Pathol* 22:15-27.
- Boller, T., and Felix, G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol* 60:379-406.
- Chisholm, S. T., Coaker, G., Day, B., and Staskawicz, B. J. 2006. Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124:803-814.
- Cook, D. E., Mesarich, C. H., and Thomma, B. P. H. J. 2015. Understanding plant immunity as a surveillance system to detect invasion. *Annu Rev Phytopathol*.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- Dodds, P. N., and Rathjen, J. P. 2010. Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* 11:539-548.
- Fitzell, R., Evans, G., and Fahy, P. 1980. Studies on the colonization of plant roots by *Verticillium dahliae* Klebahn with use of immunofluorescent staining. *Aust J Bot* 28:357-368.
- Flor, H. H. 1942. Inheritance of pathogenicity in *Melampsora lini*. *Phytopathology* 32:653-669.
- Fradin, E. F., and Thomma, B. P. H. J. 2006. Physiology and molecular aspects of *Verticillium* wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol Plant Pathol* 7:71-86.
- Fradin, E. F., Zhang, Z., Ayala, J. C. J., Castroverde, C. D., Nazar, R. N., Robb, J., Liu, C.-M., and Thomma, B. P. 2009. Genetic dissection of *Verticillium* wilt resistance mediated by tomato Ve1. *Plant Physiology* 150:320-332.
- Garavaglia, B. S., Thomas, L., Gottig, N., Dunger, G., Garofalo, C. G., Daurelio, L. D., Ndimba, B., Orellano, E. G., Gehring, C., and Ottado, J. 2010. A eukaryotic-acquired gene by a biotrophic phytopathogen allows prolonged survival on the host by counteracting the shut-down of plant photosynthesis. *PLoS One* 5:e8950.
- Garber, R., and Houston, B. 1966. Penetration and development of *Verticillium albo-atrum* in cotton plant. *Phytopathology* 56:1121.
- Gehring, C., and Irving, H. 2003. Natriuretic peptides - a class of heterologous molecules in plants. *Int J Biochem Cell Biol* 35:1318-1322.
- Gottig, N., Garavaglia, B. S., Daurelio, L. D., Valentine, A., Gehring, C., Orellano, E. G., and Ottado, J. 2008. *Xanthomonas axonopodis* pv. *citri* uses a plant natriuretic peptide-like protein to modify host homeostasis. *Proc Natl Acad Sci U S A* 105:18631-18636.
- Jain, H. 2010. *Green revolution: history, impact and future*. Studium Press LLC, Houston USA.
- Jones, J. D., and Dangl, J. L. 2006. The plant immune system. *Nature* 444:323-329.
- Kawchuk, L. M., Hachey, J., Lynch, D. R., Kulcsar, F., Van Rooijen, G., Waterer, D. R., Robertson, A., Kokko, E., Byers, R., and Howard, R. J. 2001. Tomato Ve disease resistance genes encode cell surface-like receptors. *Proceedings of the National Academy of Sciences of the United States of America* 98:6511-6515.
- Keen, N. 1990. Gene-for-gene complementarity in plant-pathogen interactions. *Annu Rev Gen* 24:447-463.
- Ludidi, N., Morse, M., Sayed, M., Wherrett, T., Shabala, S., and Gehring, C. 2004. A recombinant plant natriuretic peptide causes rapid and spatially differentiated K<sup>+</sup>, Na<sup>+</sup> and H<sup>+</sup> flux changes in *Arabidopsis thaliana* roots. *Plant Cell Physiol* 45:1093-1098.
- Majer, A., Javornik, B., Cerenak, A., and Jakse, J. 2014. Development of novel EST-derived resistance gene markers in hop (*Humulus lupulus* L.). *Mol Breed* 33:61-74.

- Maryani, M., Bradley, G., Cahill, D., and Gehring, C. 2001. Natriuretic peptides and immunoreactants modify osmoticum-dependent volume changes in *Solanum tuberosum* L. mesophyll cell protoplasts. *Plant Sci* 161:443-452.
- Mol, L. 1995. Effect of plant roots on the germination of microsclerotia of *Verticillium dahliae*. *Eur J Plant Pathol* 6:679-685.
- Nembaware, V., Seoighe, C., Sayed, M., and Gehring, C. 2004. A plant natriuretic peptide-like gene in the bacterial pathogen *Xanthomonas axonopodis* may induce hyper-hydration in the plant host: a hypothesis of molecular mimicry. *BMC Evol Biol* 4:10.
- Oerke, E. C. 2006. Crop losses to pests. *J Agric Sci* 144:31-43.
- Pegg, G. F., and Brady, B. L. 2002. *Verticillium* wilts. CABI, Wallingford.
- Pharmawati, M., Maryani, M. M., Nikolakopoulos, T., Gehring, C. A., and Irving, H. R. 2001. Cyclic GMP modulates stomatal opening induced by natriuretic peptides and immunoreactive analogues. *Plant Physiol Biochem* 39:385-394.
- Strange, R. N., and Scott, P. R. 2005. Plant disease: a threat to global food security. *Annu Rev Phytopathol* 43:83-116.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- Van der Biezen, E. A., and Jones, J. D. 1998. Plant disease-resistance proteins and the gene-for-gene concept. *Trends Biochem Sci* 23:454-456.
- Vining, K., and Davis, T. 2009. Isolation of a *Ve* homolog, *mVe1*, and its relationship to *Verticillium* wilt resistance in *Mentha longifolia* (L.) Huds. *Mol Genet Genomics* 282:173-184.
- Zhang, Y., Wang, X., Yang, S., Chi, J., Zhang, G., and Ma, Z. 2011. Cloning and characterization of a *Verticillium* wilt resistance gene from *Gossypium barbadense* and functional analysis in *Arabidopsis thaliana*. *Plant Cell Rep* 30:2085-2096.
- Zhang, Z., Fradin, E., de Jonge, R., van Esse, H. P., Smit, P., Liu, C.-M., and Thomma, B. P. H. J. 2013. Optimized agroinfiltration and virus-induced gene silencing to study *Ve1*-mediated *Verticillium* resistance in tobacco. *Mol. Plant Microbe Interact.* 26:182-190.





# **Filamentous Pathogen Effector Functions: Of Pathogens, Hosts and Microbiomes**

Hanna Rovenich#  
Jordi C. Boshoven#  
Bart P.H.J. Thomma

#These authors contributed equally

## **Chapter 2**

## Abstract

Microorganisms play essential roles in almost every environment on earth. For instance, microbes decompose organic material, or establish symbiotic relationships that range from pathogenic to mutualistic. Symbiotic relationships have been particularly well studied for microbial plant pathogens and have emphasized the role of effectors; secreted molecules that support host colonization. Most effectors characterized thus far play roles in deregulation of host immunity. Arguably, however, pathogens not only deal with immune responses during host colonization, but also encounter other microbes including competitors, (myco)parasites and even potential co-operators. Thus, part of the effector catalog may target microbiome co-inhabitants rather than host physiology.



## Introduction

During early microbial colonization stages, plant cell surface-localized pattern recognition receptors (PRRs) recognize microbe-associated molecular patterns (MAMPs), such as fungal chitin, to activate immune responses (Dodds and Rathjen 2010; Thomma et al. 2011). In order to establish themselves, adapted pathogens secrete effector molecules that deregulate immune responses and facilitate host colonization. Simultaneously, hosts evolve effector recognition by novel receptors that reinstall immunity (Dodds and Rathjen 2010; Thomma et al. 2011). Consequently, effectors are subject to various selective forces that drive their evolution, leading to diversified effector repertoires between pathogen lineages. Functional characterization of effectors and determination of their contribution to the microbial lifestyle provides insight in relevant processes for host colonization.

## Plant pathogen effectors deregulate host immunity in various subcellular compartments

Many pathogens initially enter the plant apoplast, which contains enzymes that hamper microbial colonization. For example, chitinases target fungal cell walls to release chitin fragments that activate immune receptors, leading to further chitinase accumulation to induce hyphal lysis. In turn, fungal pathogens secrete chitin-binding effectors to protect their cell walls and interfere with immune receptor activation (de Jonge et al. 2010; Kombrink et al. 2011; Marshall et al. 2011; Mentlak et al. 2012). The LysM domain-containing Ecp6 effector of the leaf mold fungus *Cladosporium fulvum* can outcompete host receptors through chitin binding with unprecedented ultrahigh (pM) affinity by intramolecular LysM domain dimerization (Sanchez-Vallet et al. 2013). Additionally, LysM effectors likely interfere with receptor dimerization that is required to activate immune signaling (Hayafune et al. 2014; Liu et al. 2012; Sanchez-Vallet et al. 2013).

Although effectors that directly target chitinases have not yet been identified, some effectors target other apoplastic hydrolytic enzymes, such as proteases. For example, sequence-unrelated effectors of *C. fulvum*, the oomycete *Phytophthora infestans*, and the parasitic nematode *Globodera rostochiensis* inhibit tomato cysteine proteases including Rcr3 (Lozano-Torres et al. 2012; Song et al. 2009; van Esse et al. 2008). The closely related oomycetes *P. infestans* and *P. mirabilis* express an orthologous pair of host protease inhibitor effectors that are subject to positive selection, which was implicated in adaptation to unique protease targets in their respective host plants (Dong et al. 2014). Besides protease inhibitors, *P. infestans* secretes the Avrblb2 effector that interferes with protease secretion (Bozkurt et al.

2011). The smut fungus *Ustilago maydis* inhibits apoplastic proteases via multiple effectors. While Pit2 directly inhibits cysteine proteases (Mueller et al. 2013), Pep1 induces the maize cystatin CC9 that inhibits apoplastic proteases in turn (van der Linde et al. 2012). Pep1 furthermore inhibits the maize peroxidase POX12 to perturb reactive oxygen species balances (Hemetsberger et al. 2012). Thus, the plant apoplast is a dynamic battlefield for plant pathogens.


In addition to apoplastic effectors, many pathogens deliver effectors that act inside host cells, although mechanisms that govern their uptake remain controversial (Petre and Kamoun 2014). The rice blast fungus *Magnaporthe oryzae* was shown to secrete various effectors that enter rice cells, and even move to non-infected neighboring cells, presumably to prepare these for infection (Khang et al. 2010). The AvrPiz-t effector targets proteasome activity through interaction with the RING E3 ubiquitin ligase APIP6, leading to their mutual degradation and suppression of PRR-mediated immunity (Park et al. 2012). Effector diffusion from infected cells into neighboring cells was similarly observed for the *U. maydis* chorismate mutase Cmu1 that targets the shikimate pathway to channel chorismate into the phenylpropanoid pathway, thus adversely affecting salicylic acid (SA) biosynthesis (Djamei et al. 2011). *U. maydis* furthermore secretes the Tin2 effector to stabilize the maize ZmTTK1 kinase that controls anthocyanin biosynthesis, possibly to suppress tissue lignification (Tanaka et al. 2014). Also the oomycete *Hyaloperonospora arabidopsidis* targets SA signaling by secreting a nuclear-localized effector that interacts with the mediator complex that controls interactions between transcriptional regulators and RNA polymerase (Caillaud et al. 2013). Host transcription is furthermore perturbed by effectors that inhibit transcription factor translocation to the nucleus (McLellan et al. 2013). Additionally, nuclear-localized effectors may affect host immunity post-transcriptionally by suppressing the biogenesis of small RNAs in the host (Qiao et al. 2013). Interestingly, *Botrytis cinerea* was recently suggested to deliver even small RNAs into host cells to affect immune responses (Weiberg et al. 2013).

Finally, several effectors target host cell death mechanisms, such as *P. infestans* Avr3a and PexRD2. While Avr3a suppresses INF1-triggered cell death by stabilizing the U-box E3 ligase CMPG1 during biotrophic growth, PexRD2 targets the kinase domain of the cell death regulator MAPKKKε (Bos et al. 2010; King et al. 2014). During later stages of infection, however, *P. infestans* relies on induction of host cell death as it switches to a necrotrophic lifestyle. Necrotrophic pathogens evolved effectors that actually induce cell death. An elegant example is provided by the *Cochliobolus victoriae* effector victorin that binds to thioredoxins including TRXh5, which is required for redox control of the transcriptional immune regulator NPR1. TRXh5 binding activates the NB-LRR-type immune receptor LOV1, facilitating necrotrophic exploitation of host cell death by *C. victoriae* (Lorang et al. 2012).

In conclusion, although information for the vast majority of pathogen effectors, particularly of filamentous pathogens, is still lacking, effector molecules are highly versatile. Clearly, recently uncovered functions revealed that virulence effectors, despite the finding that they converge onto pivotal elements of the plant immune system (Mukhtar et al. 2011), can deregulate any step of immunity in any cellular compartment (Figure 1, Supplemental table 1).

## Endophytes and mutualists use effectors to suppress host immunity too

Like pathogens, commensalistic endophytes and mutualists develop intimate host plant associations. During initiation of such symbioses, PRRs continue to perceive MAMPs. Consequently, similar to pathogens, endophytes and mutualists are recipients of immune responses. However, the precise role and fate of host immunity in the establishment of symbiosis has remained enigmatic.



**FIGURE 1 | Filamentous pathogen effectors deregulate host immunity in various host subcellular compartments.** Pathogens secrete effectors (red symbols) to deregulate plant immunity (see text for details). Whereas one group of effectors (red circles) interacts with host targets that act in immunity (black shapes), another group of effectors (red triangles) acts in self-defense to protect the pathogen from host-derived antimicrobials.

The root endophyte *Piriformospora indica* has a wide host range and induces enhanced growth and stress resistance in colonized hosts. Rather than evading host detection, the fungus actively suppresses immunity (Jacobs et al. 2011). During early biotrophic growth at the onset of symbiosis, about 10% of the transcriptome encodes putative effector proteins (Zuccaro et al. 2011). At later growth stages

the fungus requires host cell death for further colonization, thus resembling hemibiotrophic pathogens such as *Mycosphaerella graminicola* and *M. oryzae*. Like *C. fulvum*, these latter species utilize LysM effectors to suppress immune responses (de Jonge et al. 2010; Marshall et al. 2011; Mentlak et al. 2012). *P. indica* carries an expanded LysM domain-containing effector repertoire that may similarly act in immune suppression (Zuccaro et al. 2011).


Effector-like proteins are also encoded by genomes of other mutualists (Martin et al. 2010; Plett et al. 2011; Plett and Martin 2011). The ectomycorrhiza *Laccaria bicolor* genome encodes hundreds of small secreted proteins, several of which are only expressed in symbiotic tissues. Of these, MiSSP7 was shown to translocate to the nucleus of poplar host cells to stabilize the JAZ6 protein and repress jasmonate signaling (Plett et al. 2014; Plett et al. 2011). Likewise, the ectomycorrhiza *Tuber melanosporum* expresses 125 cysteine-rich small secreted proteins, including a LysM effector, which are highly upregulated during symbiosis (Martin et al. 2010).

It was recently shown that arbuscular endomycorrhizal fungi produce lipochitooligosaccharide mycorrhizal (Myc) factors that stimulate root growth and branching to initiate symbiosis (Maillet et al. 2011). Similar to endophytes and ectomycorrhiza, arbuscular endomycorrhiza secrete effector-like proteins during symbiotic interactions (Kloppholz et al. 2011; Lin et al. 2014; Tisserant et al. 2013). The genome of *Rhizophagus irregularis* encodes a family of CRN-like proteins that are abundantly found in plant pathogenic *Phytophthora* spp. (Lin et al. 2014). *R. irregularis* was furthermore found to encode an effector that interacts with the pathogenesis-related ethylene-responsive transcription factor 19 (ERF19) in the host nucleus to promote mycorrhization, potentially by counteracting MAMP-induced host defense responses that are regulated by ERF19 (Kloppholz et al. 2011).

Collectively, these findings suggest that symbiotic associations that include endophytism, mutualism and parasitism form a continuum in which effectors play essential roles (Supplemental table 1).

## Effectors act in self-defense and competition

The ability to establish symbiosis evolved multiple times in microbes, presumably from saprotrophism, and many plant pathogens still display saprotrophic life stages. Saprotrophs generally reside within the soil where they feed on decaying organic matter in the presence of a rich microbiota. In this environment, microbial competition as well as co-operation occurs (Figure 2). Threats are posed by (myco) parasites and competitors that produce antibiotics with specific or broad-spectrum activities. Consequently, microbes require molecules for self-defense and interaction with other microbiome partners.



**FIGURE 2 | How pathogens influence the local biota by exploiting effector activities.** The interaction between microbial pathogens and plant hosts occurs in environments that contain additional microbiome partners that can negatively (competition) or positively (co-operation) impact the pathogen as well as the host. Consequently, the pathogen and host may target each other directly (solid lines) as well as indirectly (dotted lines). Likely, pathogens exploit effector activities (orange lines) to not only directly modulate their hosts, but also to influence the local microbiota that can impact the outcome of the interaction with their hosts.

Similar to infected plants, many mycoparasites secrete hydrolytic enzymes including proteases, chitinases and glucanases to target fungal cell walls. Presumably, chitin-binding effectors that protect hyphal cell walls against plant-derived chitinases similarly protect against mycoparasite-derived chitinases, which may explain abundant LysM effector catalogs of non-pathogenic fungi (Kombrink and Thomma 2013; Kubicek et al. 2011). As LysM domains occur in peptidoglycan-binding proteins of various origins, LysM effector homologs that bind non-chitin substrates likely occur. Indeed, a plant pathogen LysM effector that binds bacterial cell walls was characterized (Kombrink and Thomma, unpublished data), potentially implicating this effector in bacterial competition or protection against bacterial mycoparasites. Genome analyses furthermore revealed that saprotrophic species encode abundant catalogs of small secreted proteins that resemble pathogen effector catalogs (Druzhinina et al. 2012; Kasuga et al. 2009; Kombrink and Thomma 2013; Suh et al. 2012). Although these potential effectors are poorly studied, one such effector, CipC, was implicated in competition with bacteria in *Aspergillus* spp. (Melin et al. 2002; Suh et al. 2012). The genome of the ubiquitous saprophyte and opportunistic mammalian pathogen *A. fumigatus* encodes several effector proteins (Wartenberg

et al. 2011). However, since the vast majority of fungi that cause disease in animals are soil saprophytes that opportunistically infect their hosts, to which they are not highly adapted, it has been speculated that infection does not rely on the activity of effectors (Lowe and Howlett 2012). Rather, their effectors are thought to be required for saprophytic survival (Lowe and Howlett 2012). Nevertheless, effectors that evolved to enable saprophytic survival may be co-opted for opportunistic infection as well.

Likely, competition between plant-associated microbes also occurs within hosts, although perhaps to a lesser extent than in soil due to reduced species diversity. Indeed, the second most abundantly *in planta*-expressed gene of the fungal endophyte *Epichloë festucae* encodes a secreted antifungal protein (Ambrose and Belanger 2012). Thus, effector homologs may play crucial roles in microbial competition in a broad spectrum of environments.

## Do pathogens shape local microbiomes?

For various types of multicellular organisms it is increasingly recognized that their microbiome, i.e. the community of microbes that thrives in, on, or immediately near the organism, greatly influences its performance (Ezenwa et al. 2012). For plants, it has been particularly well documented that the rhizosphere microbiota affects plant growth and stress tolerance. In addition, the importance of the phyllosphere microbiota is increasingly recognized (Vorholt 2012). These microbiota comprise members that provide direct as well as indirect pathogen protection through antibiosis and induced immunity, respectively. Whereas soil types have a major impact on root inhabiting bacterial community compositions on *Arabidopsis*, host genotypes were reported to only have a minor impact (Bulgarelli et al. 2012; Lundberg et al. 2012). In contrast, different *Arabidopsis* accessions were found to harbor different phyllosphere communities and several host genetic mutations were found to perturb the microbiota composition, demonstrating that host genetic factors shape the associated microbiota (Bodenhausen et al. 2014). It is less clear, however, whether plants evolved to actively recruit phyllosphere communities. Potentially, plants recruit founder species that further shape local microbiomes through inter-microbe interactions (Vorholt 2012). Such interactions may require effectors. Considering that plant factors control the composition of the microbiota, microbiome members may utilize effectors to modulate hosts and control competitors indirectly. Additionally, manipulation of host metabolism could even establish microbial cooperation (Figure 2). Although not immediately addressing inter-microbial interactions, an insect-transmitted phytoplasma was recently shown to utilize an effector to alter floral development of host plants, converting them

into vegetative tissues that attract leafhopper vectors (MacLean et al. 2014). This represents a striking example of the exploitation of effector activity to influence compositions of the local biome. Similarly, the rust fungus *Puccinia monoica* induces floral mimicry in the host *Boechnera stricta* to enhance its reproduction and spore dispersal by insects (Cano et al. 2013).

Considering the importance of the microbiome for the ability of plants to withstand pathogen infection, it is conceivable that pathogens evolved to affect host microbiomes, possibly through effector activities (Figure 2).

## Different mechanisms drive evolution of effector repertoires

Mechanisms underlying genome plasticity and evolution have been intensely studied, especially for plant pathogens. As genomes are structured and not just a random sequence of genes, effector genes are often found in dynamic genomic compartments, such as gene-sparse regions, subtelomeric regions or conditionally dispensable (pathogenicity) chromosomes (Raffaele and Kamoun 2012). For example, effector localization in gene-sparse regions was recorded for the endophyte *P. indica* (Zuccaro et al. 2011), while in the saprophyte *N. crassa* genes encoding small secreted proteins are found in subtelomeric regions (Kasuga et al. 2009). Genetic plasticity in such compartments is governed by diverse mechanisms including recombination and activity of transposable elements. A direct implication of genomic rearrangement in the evolution of fungal aggressiveness was shown for the vascular wilt fungus *Verticillium dahliae*, leading to the emergence of lineage-specific regions that are enriched for virulence effectors (de Jonge et al. 2013). High genetic variability in effector genes enables rapid evolutionary processes. The importance of dynamic genome compartments for accelerated gene evolution was underlined in the specialization of *P. infestans* after the host jump that separated this species from related species. Uneven evolutionary rates across the genome occur, with *in planta*-induced genes residing in fast-evolving compartments (Raffaele et al. 2010). In turn, effector specialization can lead to diversification and speciation in pathogen lineages (Dong et al. 2014). In this manner, effectors can determine microbial niches. Moreover, composition of effector catalogs can dictate microbial lifestyles. For example, the leaf epiphyte and antagonist of powdery mildews *Pseudozyma flucculosa* lost its ability to parasitize plants like its smut fungi relatives due to loss of virulence effectors (Lefebvre et al. 2013). However, the biocontrol agent has acquired other effectors that are not found in the smut relatives that may have shaped its current lifestyle (Lefebvre et al. 2013). These findings suggest that effector catalogs evolve via different mechanisms and that their composition influences a microbe's lifestyle in a given environment.



## Experimental way forward

The interaction between pathogenic (filamentous) microbes and the organisms they encounter in their niches, either while colonizing the host or during free-living stages in the environment, is poorly understood. An extensive characterization of the complex microbial communities in such niches may lead to a better understanding of the interactions that take place beyond the direct interaction between pathogen and host. Detailed transcriptome analyses may lead to the identification of particular triggers of effector gene expression derived from microbial co-inhabitants, and may hint towards functions in inter-microbial interactions (Mathioni et al. 2013; Mela et al. 2011) that can subsequently be tested for in targeted analysis to reveal components that either promote or inhibit other microbes (Kombrink and Thomma 2013).

## Conclusions

Although a paradigm in plant pathology dictates that existence of disease requires the interaction of a virulent pathogen with a susceptible host in a favorable environment, plant-microbe interactions are mostly studied as one-on-one relationships. However, in addition to host immune responses, pathogenic microbes continuously encounter other microbes that include competitors and mycoparasites that need to be dealt with simultaneously. Importantly, findings for pathogenic microbes can be extrapolated to other types of symbioses as well. After all, irrespective of the type of symbiosis, the interest of the microbial partner is merely to exploit the host for nutrition and shelter. This may also explain the thin line that is regularly observed between the different types of symbioses (Delaye et al. 2013; Malcolm et al. 2013; Plett and Martin 2011; Zuccaro et al. 2011). In all types of symbioses, the microbial partner needs to suppress host immune responses and ward off microbial antagonists. Using effectors as probes, further critical processes in host colonization will be uncovered, leading to enhanced understanding of the biology of microbes that aim to establish symbioses.

## Acknowledgements

The authors acknowledge support by the Research Council for Earth and Life sciences (ALW) of the Netherlands Organization for Scientific Research (NWO), thank Melvin Bolton, Andrea Sánchez-Vallet, and Ronnie de Jonge for critically reading the manuscript.

## References

- Ambrose, K. V., and Belanger, F. C. 2012. SOLiD-SAGE of endophyte-infected red fescue reveals numerous effects on host transcriptome and an abundance of highly expressed fungal secreted proteins. *PLoS One* 7:e53214.
- Bodenhausen, N., Bortfeld-Miller, M., Ackermann, M., and Vorholt, J. A. 2014. A synthetic community approach reveals plant genotypes affecting the phyllosphere microbiota. *PLoS Genet* 10:e1004283.
- Bos, J. I., Armstrong, M. R., Gilroy, E. M., Boevink, P. C., Hein, I., Taylor, R. M., Zhendong, T., Engelhardt, S., Vetukuri, R. R., Harrower, B., Dixelius, C., Bryan, G., Sadanandom, A., Whisson, S. C., Kamoun, S., and Birch, P. R. 2010. *Phytophthora infestans* effector AVR3a is essential for virulence and manipulates plant immunity by stabilizing host E3 ligase CMPG1. *Proc Natl Acad Sci U S A* 107:9909-9914.
- Bozkurt, T. O., Schornack, S., Win, J., Shindo, T., Ilyas, M., Oliva, R., Cano, L. M., Jones, A. M., Huitema, E., van der Hoorn, R. A., and Kamoun, S. 2011. *Phytophthora infestans* effector AVRblb2 prevents secretion of a plant immune protease at the haustorial interface. *Proc Natl Acad Sci U S A* 108:20832-20837.
- Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadijad, N., Assenza, F., Rauf, P., Huettel, B., Reinhardt, R., Schmelzer, E., Peplies, J., Gloeckner, F. O., Amann, R., Eickhorst, T., and Schulze-Lefert, P. 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. *Nature* 488:91-95.
- Caillaud, M. C., Asai, S., Rallapalli, G., Piquerez, S., Fabro, G., and Jones, J. D. 2013. A downy mildew effector attenuates salicylic acid-triggered immunity in Arabidopsis by interacting with the host mediator complex. *PLoS Biol* 11:e1001732.
- Cano, L. M., Raffaele, S., Haugen, R. H., Saunders, D. G., Leonelli, L., MacLean, D., Hogenhout, S. A., and Kamoun, S. 2013. Major transcriptome reprogramming underlies floral mimicry induced by the rust fungus *Puccinia monoica* in *Boechera stricta*. *PLoS One* 8:e75293.
- de Jonge, R., Bolton, M. D., Kombrink, A., van den Berg, G. C., Yadeta, K. A., and Thomma, B. P. H. J. 2013. Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. *Genome Res* 23:1271-1282.
- de Jonge, R., van Esse, H. P., Kombrink, A., Shinya, T., Desaki, Y., Bours, R., van der Krol, S., Shibuya, N., Joosten, M. H., and Thomma, B. P. H. J. 2010. Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science* 329:953-955.
- Delaye, L., García-Guzmán, G., and Heil, M. 2013. Endophytes versus biotrophic and necrotrophic pathogens—are fungal lifestyles evolutionarily stable traits? *Fungal Divers* 60:125-135.
- Djamei, A., Schipper, K., Rabe, F., Ghosh, A., Vincon, V., Kahnt, J., Osorio, S., Tohge, T., Fernie, A. R., Feussner, I., Feussner, K., Meinicke, P., Stierhof, Y. D., Schwarz, H., Macek, B., Mann, M., and Kahmann, R. 2011. Metabolic priming by a secreted fungal effector. *Nature* 478:395-398.
- Dodds, P. N., and Rathjen, J. P. 2010. Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* 11:539-548.
- Dong, S., Stam, R., Cano, L. M., Song, J., Sklenar, J., Yoshida, K., Bozkurt, T. O., Oliva, R., Liu, Z., Tian, M., Win, J., Banfield, M. J., Jones, A. M., van der Hoorn, R. A., and Kamoun, S. 2014. Effector specialization in a lineage of the Irish potato famine pathogen. *Science* 343:552-555.
- Druzhinina, I. S., Shelest, E., and Kubicek, C. P. 2012. Novel traits of *Trichoderma* predicted through the analysis of its secretome. *FEMS Microbiol Lett* 337:1-9.
- Ezenwa, V. O., Gerardo, N. M., Inouye, D. W., Medina, M., and Xavier, J. B. 2012. Microbiology. Animal behavior and the microbiome. *Science* 338:198-199.
- Hayafune, M., Berisio, R., Marchetti, R., Silipo, A., Kayama, M., Desaki, Y., Arima, S., Squeglia, F., Ruggiero, A., Tokuyasu, K., Molinaro, A., Kaku, H., and Shibuya, N. 2014. Chitin-induced activation of immune signaling by the rice receptor CEBiP relies on a unique sandwich-type dimerization. *Proc Natl Acad Sci U S A* 111:E404-413.

- Hemetsberger, C., Herrberger, C., Zechmann, B., Hillmer, M., and Doehlemann, G. 2012. The *Ustilago maydis* effector Pep1 suppresses plant immunity by inhibition of host peroxidase activity. *PLoS Pathog* 8:e1002684.
- Jacobs, S., Zechmann, B., Molitor, A., Trujillo, M., Petutschnig, E., Lipka, V., Kogel, K. H., and Schafer, P. 2011. Broad-spectrum suppression of innate immunity is required for colonization of Arabidopsis roots by the fungus *Piriformospora indica*. *Plant Physiol* 156:726-740.
- Kasuga, T., Mannhaupt, G., and Glass, N. L. 2009. Relationship between phylogenetic distribution and genomic features in *Neurospora crassa*. *PLoS One* 4:e5286.
- Khang, C. H., Berruyer, R., Giraldo, M. C., Kankanala, P., Park, S. Y., Czymmek, K., Kang, S., and Valent, B. 2010. Translocation of *Magnaporthe oryzae* effectors into rice cells and their subsequent cell-to-cell movement. *Plant Cell* 22:1388-1403.
- King, S. R., McLellan, H., Boevink, P. C., Armstrong, M. R., Bukharova, T., Sukarta, O., Win, J., Kamoun, S., Birch, P. R., and Banfield, M. J. 2014. *Phytophthora infestans* RXLR effector PexRD2 interacts with host MAPKKK epsilon to suppress plant immune signaling. *Plant Cell* 26:1345-1359.
- Kloppholz, S., Kuhn, H., and Requena, N. 2011. A secreted fungal effector of *Glomus intraradices* promotes symbiotic biotrophy. *Curr Biol* 21:1204-1209.
- Kombrink, A., and Thomma, B. P. H. J. 2013. LysM effectors: secreted proteins supporting fungal life. *PLoS Pathog* 9:e1003769.
- Kombrink, A., Sanchez-Vallet, A., and Thomma, B. P. H. J. 2011. The role of chitin detection in plant-pathogen interactions. *Microbes Infect* 13:1168-1176.
- Kubicek, C. P., Herrera-Estrella, A., Seidl-Seiboth, V., Martine, D. A., Druzhinina, I. S., Thon, M., Zeilinger, S., Casas-Flores, S., Horwitz, B. A., Mukherjee, P. K., Mukherjee, M., Kredics, L., Alcaraz, L. D., Aerts, A., Antal, Z., Atanasova, L., Cervantes-Badillo, M. G., Challacombe, J., Chertkov, O., McCluskey, K., Culpier, F., Deshpande, N., von Dohren, H., Ebbole, D. J., Esquivel-Naranjo, E. U., Fekete, E., Flippi, M., Glaser, F., Gomez-Rodriguez, E. Y., Gruber, S., Han, C., Henrissat, B., Hermosa, R., Hernandez-Onate, M., Karaffa, L., Kosti, I., Le Crom, S., Lindquist, E., Lucas, S., Lubeck, M., Lubeck, P. S., Margeot, A., Metz, B., Misra, M., Nevalainen, H., Omann, M., Packer, N., Perrone, G., Uresti-Rivera, E. E., Salamov, A., Schmoll, M., Seiboth, B., Shapiro, H., Sukno, S., Tamayo-Ramos, J. A., Tisch, D., Wiest, A., Wilkinson, H. H., Zhang, M., Coutinho, P. M., Kenerley, C. M., Monte, E., Baker, S. E., and Grigoriev, I. V. 2011. Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of *Trichoderma*. *Genome Biol* 12:R40.
- Lefebvre, F., Joly, D. L., Labbe, C., Teichmann, B., Linning, R., Belzile, F., Bakkeren, G., and Belanger, R. R. 2013. The transition from a phytopathogenic smut ancestor to an anamorphic biocontrol agent deciphered by comparative whole-genome analysis. *Plant Cell* 25:1946-1959.
- Lin, K., Limpens, E., Zhang, Z., Ivanov, S., Saunders, D. G., Mu, D., Pang, E., Cao, H., Cha, H., Lin, T., Zhou, Q., Shang, Y., Li, Y., Sharma, T., van Velzen, R., de Ruijter, N., Aanen, D. K., Win, J., Kamoun, S., Bisseling, T., Geurts, R., and Huang, S. 2014. Single nucleus genome sequencing reveals high similarity among nuclei of an endomycorrhizal fungus. *PLoS Genet* 10:e1004078.
- Liu, T., Liu, Z., Song, C., Hu, Y., Han, Z., She, J., Fan, F., Wang, J., Jin, C., Chang, J., Zhou, J. M., and Chai, J. 2012. Chitin-induced dimerization activates a plant immune receptor. *Science* 336:1160-1164.
- Lorang, J., Kidarsa, T., Bradford, C. S., Gilbert, B., Curtis, M., Tzeng, S. C., Maier, C. S., and Wolpert, T. J. 2012. Tricking the guard: exploiting plant defense for disease susceptibility. *Science* 338:659-662.
- Lowe, R. G., and Howlett, B. J. 2012. Indifferent, affectionate, or deceitful: lifestyles and secretomes of fungi. *PLoS Pathog* 8:e1002515.
- Lozano-Torres, J. L., Wilbers, R. H. P., Gawronski, P., Boshoven, J. C., Finkers-Tomczak, A., Cordewener, J. H. G., America, A. H. P., Overmars, H. A., Van 't Klooster, J. W., Baranowski, L., Sobczak, M., Ilyas, M., van der Hoorn, R. A. L., Schots, A., de Wit, P. J. G. M., Bakker, J., Govers, A., and Smant, G. 2012. Dual disease resistance mediated by the immune receptor Cf-2 in tomato requires a common virulence target of a fungus and a nematode. *Proc Natl Acad Sci U S A* 109:10119-10124.

- Lundberg, D. S., Lebeis, S. L., Paredes, S. H., Yourstone, S., Gehring, J., Malfatti, S., Tremblay, J., Engelbrektson, A., Kunin, V., del Rio, T. G., Edgar, R. C., Eickhorst, T., Ley, R. E., Hugenholtz, P., Tringe, S. G., and Dangl, J. L. 2012. Defining the core *Arabidopsis thaliana* root microbiome. *Nature* 488:86-90.
- MacLean, A. M., Orlovskis, Z., Kowitzanich, K., Zdziarska, A. M., Angenent, G. C., Immink, R. G., and Hogenhout, S. A. 2014. Phytoplasma effector SAP54 hijacks plant reproduction by degrading MADS-box proteins and promotes insect colonization in a RAD23-dependent manner. *PLoS Biol* 12:e1001835.
- Maillet, F., Poinso, V., Andre, O., Puech-Pages, V., Haouy, A., Gueunier, M., Cromer, L., Giraudet, D., Formey, D., Niebel, A., Martinez, E. A., Driguez, H., Becard, G., and Denarie, J. 2011. Fungal lipochitooligosaccharide symbiotic signals in arbuscular mycorrhiza. *Nature* 469:58-63.
- Malcolm, G. M., Kuldau, G. A., Gugino, B. K., and Jimenez-Gasco Mdel, M. 2013. Hidden host plant associations of soilborne fungal pathogens: an ecological perspective. *Phytopathology* 103:538-544.
- Marshall, R., Kombrink, A., Motteram, J., Loza-Reyes, E., Lucas, J., Hammond-Kosack, K. E., Thomma, B. P. H. J., and Rudd, J. J. 2011. Analysis of two in planta expressed LysM effector homologs from the fungus *Mycosphaerella graminicola* reveals novel functional properties and varying contributions to virulence on wheat. *Plant Physiol* 156:756-769.
- Martin, F., Kohler, A., Murat, C., Balestrini, R., Coutinho, P. M., Jaillon, O., Montanini, B., Morin, E., Noel, B., Percudani, R., Porcel, B., Rubini, A., Amicucci, A., Amsalem, J., Anthouard, V., Arcioni, S., Artiguenave, F., Aury, J. M., Ballario, P., Bolchi, A., Brenna, A., Brun, A., Buee, M., Cantarel, B., Chevalier, G., Couloux, A., Da Silva, C., Denoeud, F., Duplessis, S., Ghignone, S., Hilselberger, B., Iotti, M., Marçais, B., Mello, A., Miranda, M., Pacioni, G., Quesneville, H., Riccioni, C., Ruotolo, R., Splivallo, R., Stocchi, V., Tisserant, E., Viscomi, A. R., Zambonelli, A., Zampieri, E., Henrissat, B., Lebrun, M. H., Paolocci, F., Bonfante, P., Ottonello, S., and Wincker, P. 2010. Perigord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* 464:1033-1038.
- Mathioni, S. M., Patel, N., Riddick, B., Sweigard, J. A., Czymmek, K. J., Caplan, J. L., Kunjeti, S. G., Kunjeti, S., Raman, V., Hillman, B. I., Kobayashi, D. Y., and Donofrio, N. M. 2013. Transcriptomics of the rice blast fungus *Magnaporthe oryzae* in response to the bacterial antagonist *Lysobacter enzymogenes* reveals candidate fungal defense response genes. *PLoS One* 8:e76487.
- McLellan, H., Boevink, P. C., Armstrong, M. R., Pritchard, L., Gomez, S., Morales, J., Whisson, S. C., Beynon, J. L., and Birch, P. R. 2013. An RxLR effector from *Phytophthora infestans* prevents re-localisation of two plant NAC transcription factors from the endoplasmic reticulum to the nucleus. *PLoS Pathog* 9:e1003670.
- Mela, F., Fritsche, K., de Boer, W., van Veen, J. A., de Graaff, L. H., van den Berg, M., and Leveau, J. H. 2011. Dual transcriptional profiling of a bacterial/fungal confrontation: *Collimonas fungivorans* versus *Aspergillus niger*. *ISME J* 5:1494-1504.
- Melin, P., Schnurer, J., and Wagner, E. G. 2002. Proteome analysis of *Aspergillus nidulans* reveals proteins associated with the response to the antibiotic concanamycin A, produced by *Streptomyces* species. *Mol Genet Genomics* 267:695-702.
- Mentlak, T. A., Kombrink, A., Shinya, T., Ryder, L. S., Otomo, I., Saitoh, H., Terauchi, R., Nishizawa, Y., Shibuya, N., Thomma, B. P. H. J., and Talbot, N. J. 2012. Effector-mediated suppression of chitin-triggered immunity by *magnaporthe oryzae* is necessary for rice blast disease. *Plant Cell* 24:322-335.
- Mueller, A. N., Ziemann, S., Treitschke, S., Assmann, D., and Doehlemann, G. 2013. Compatibility in the *Ustilago maydis*-maize interaction requires inhibition of host cysteine proteases by the fungal effector Pit2. *PLoS Pathog* 9:e1003177.
- Mukhtar, M. S., Carvunis, A. R., Dreze, M., Eppele, P., Steinbrenner, J., Moore, J., Tasan, M., Galli, M., Hao, T., Nishimura, M. T., Pevzner, S. J., Donovan, S. E., Ghamsari, L., Santhanam, B., Romero, V., Poulin, M. M., Gebreab, F., Gutierrez, B. J., Tam, S., Monachello, D., Boxem, M., Harbort, C. J., McDonald, N., Gai, L., Chen, H., He, Y., European Union Effectoromics, C., Vandenhaute, J., Roth, F. P., Hill, D. E., Ecker, J. R., Vidal, M., Beynon, J., Braun, P., and Dangl, J. L. 2011. Independently evolved virulence effectors converge onto hubs in a plant immune system network. *Science* 333:596-601.

- Okmen, B., Etalo, D. W., Joosten, M. H., Bouwmeester, H. J., de Vos, R. C., Collemare, J., and de Wit, P. J. G. M. 2013. Detoxification of alpha-tomatine by *Cladosporium fulvum* is required for full virulence on tomato. *New Phytol* 198:1203-1214.
- Park, C. H., Chen, S., Shirsekar, G., Zhou, B., Khang, C. H., Songkumarn, P., Afzal, A. J., Ning, Y., Wang, R., Bellizzi, M., Valent, B., and Wang, G. L. 2012. The *Magnaporthe oryzae* effector AvrPiz-t targets the RING E3 ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice. *Plant Cell* 24:4748-4762.
- Petre, B., and Kamoun, S. 2014. How do filamentous pathogens deliver effector proteins into plant cells? *PLoS Biol* 12:e1001801.
- Plett, J. M., and Martin, F. 2011. Blurred boundaries: lifestyle lessons from ectomycorrhizal fungal genomes. *Trends Genet* 27:14-22.
- Plett, J. M., Kemppainen, M., Kale, S. D., Kohler, A., Legue, V., Brun, A., Tyler, B. M., Pardo, A. G., and Martin, F. 2011. A secreted effector protein of *Laccaria bicolor* is required for symbiosis development. *Curr Biol* 21:1197-1203.
- Plett, J. M., Daguere, Y., Wittulsky, S., Vayssieres, A., Deveau, A., Melton, S. J., Kohler, A., Morrell-Falvey, J. L., Brun, A., Veneault-Fourrey, C., and Martin, F. 2014. Effector MiSSP7 of the mutualistic fungus *Laccaria bicolor* stabilizes the *Populus JAZ6* protein and represses jasmonic acid (JA) responsive genes. *Proc Natl Acad Sci U S A* 111:8299-8304.
- Pretsch, K., Kemen, A., Kemen, E., Geiger, M., Mendgen, K., and Voegelé, R. 2013. The rust transferred proteins—a new family of effector proteins exhibiting protease inhibitor function. *Mol Plant Pathol* 14:96-107.
- Qiao, Y., Liu, L., Xiong, Q., Flores, C., Wong, J., Shi, J., Wang, X., Liu, X., Xiang, Q., Jiang, S., Zhang, F., Wang, Y., Judelson, H. S., Chen, X., and Ma, W. 2013. Oomycete pathogens encode RNA silencing suppressors. *Nat Genet* 45:330-333.
- Raffaële, S., and Kamoun, S. 2012. Genome evolution in filamentous plant pathogens: why bigger can be better. *Nat Rev Microbiol* 10:417-430.
- Raffaële, S., Farrer, R. A., Cano, L. M., Studholme, D. J., MacLean, D., Thines, M., Jiang, R. H., Zody, M. C., Kunjeti, S. G., Donofrio, N. M., Meyers, B. C., Nusbaum, C., and Kamoun, S. 2010. Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science* 330:1540-1543.
- Rose, J. K., Ham, K. S., Darvill, A. G., and Albersheim, P. 2002. Molecular cloning and characterization of glucanase inhibitor proteins: coevolution of a counterdefense mechanism by plant pathogens. *Plant Cell* 14:1329-1345.
- Sanchez-Vallet, A., Saleem-Batcha, R., Kombrink, A., Hansen, G., Valkenburg, D. J., Thomma, B. P. H. J., and Mesters, J. R. 2013. Fungal effector Ecp6 outcompetes host immune receptor for chitin binding through intrachain LysM dimerization. *Elife* 2:e00790.
- Schmidt, S. M., Kuhn, H., Micali, C., Liller, C., Kwaaitaal, M., and Panstruga, R. 2014. Interaction of a *Blumeria graminis* f. sp. *hordei* effector candidate with a barley ARF-GAP suggests that host vesicle trafficking is a fungal pathogenicity target. *Mol Plant Pathol* 15:535-549.
- Shabab, M., Shindo, T., Gu, C., Kaschani, F., Pansuriya, T., Chintha, R., Harzen, A., Colby, T., Kamoun, S., and van der Hoorn, R. A. 2008. Fungal effector protein AVR2 targets diversifying defense-related cysteine proteases of tomato. *Plant Cell* 20:1169-1183.
- Song, J., Win, J., Tian, M., Schornack, S., Kaschani, F., Ilyas, M., van der Hoorn, R. A., and Kamoun, S. 2009. Apoplastic effectors secreted by two unrelated eukaryotic plant pathogens target the tomato defense protease Rcr3. *Proc Natl Acad Sci U S A* 106:1654-1659.
- Stergiopoulos, I., van den Burg, H. A., Okmen, B., Beenen, H. G., van Liere, S., Kema, G. H., and de Wit, P. J. 2010. Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. *Proceedings of the National Academy of Sciences of the United States of America* 107:7610-7615.

- Suh, M. J., Fedorova, N. D., Cagas, S. E., Hastings, S., Fleischmann, R. D., Peterson, S. N., Perlin, D. S., Nierman, W. C., Pieper, R., and Momany, M. 2012. Development stage-specific proteomic profiling uncovers small, lineage specific proteins most abundant in the *Aspergillus Fumigatus* conidial proteome. *Proteome Sci* 10:30.
- Tanaka, S., Brefort, T., Neidig, N., Djamei, A., Kahnt, J., Vermerris, W., Koenig, S., Feussner, K., Feussner, I., and Kahmann, R. 2014. A secreted *Ustilago maydis* effector promotes virulence by targeting anthocyanin biosynthesis in maize. *Elife* 3:e01355.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- Tian, M., Benedetti, B., and Kamoun, S. 2005. A Second Kazal-like protease inhibitor from *Phytophthora infestans* inhibits and interacts with the apoplastic pathogenesis-related protease P69B of tomato. *Plant Physiol* 138:1785-1793.
- Tian, M., Huitema, E., Da Cunha, L., Torto-Alalibo, T., and Kamoun, S. 2004. A Kazal-like extracellular serine protease inhibitor from *Phytophthora infestans* targets the tomato pathogenesis-related protease P69B. *J Biol Chem* 279:26370-26377.
- Tian, M., Win, J., Song, J., van der Hoorn, R., van der Knaap, E., and Kamoun, S. 2007. A *Phytophthora infestans* cystatin-like protein targets a novel tomato papain-like apoplastic protease. *Plant Physiol* 143:364-377.
- Tisserant, E., Malbreil, M., Kuo, A., Kohler, A., Symeonidi, A., Balestrini, R., Charron, P., Duensing, N., Freidit Frey, N., Gianinazzi-Pearson, V., Gilbert, L. B., Handa, Y., Herr, J. R., Hijiri, M., Koul, R., Kawaguchi, M., Krajinski, F., Lammers, P. J., Masclaux, F. G., Murat, C., Morin, E., Ndikumana, S., Pagni, M., Petitpierre, D., Requena, N., Rosikiewicz, P., Riley, R., Saito, K., San Clemente, H., Shapiro, H., van Tuinen, D., Becard, G., Bonfante, P., Paszkowski, U., Shachar-Hill, Y. Y., Tuskan, G. A., Young, J. P., Sanders, I. R., Henrissat, B., Rensing, S. A., Grigoriev, I. V., Corradi, N., Roux, C., and Martin, F. 2013. Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. *Proc Natl Acad Sci U S A* 110:20117-20122.
- van den Burg, H. A., Harrison, S. J., Joosten, M. H., Vervoort, J., and de Wit, P. J. G. M. 2006. *Cladosporium fulvum* Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during infection. *Mol. Plant Microbe Interact.* 19:1420-1430.
- van der Linde, K., Hemetsberger, C., Kastner, C., Kaschani, F., van der Hoorn, R. A., Kumlehn, J., and Doehlemann, G. 2012. A maize cystatin suppresses host immunity by inhibiting apoplastic cysteine proteases. *Plant Cell* 24:1285-1300.
- van Esse, H. P., Van't Klooster, J. W., Bolton, M. D., Yadeta, K. A., van Baarlen, P., Boeren, S., Vervoort, J., de Wit, P. J. G. M., and Thomma, B. P. H. J. 2008. The *Cladosporium fulvum* virulence protein Avr2 inhibits host proteases required for basal defense. *Plant Cell* 20:1948-1963.
- Vorholt, J. A. 2012. Microbial life in the phyllosphere. *Nat Rev Microbiol* 10:828-840.
- Wartenberg, D., Lapp, K., Jacobsen, I. D., Dahse, H. M., Kniemeyer, O., Heinekamp, T., and Brakhage, A. A. 2011. Secretome analysis of *Aspergillus fumigatus* reveals Asp-hemolysin as a major secreted protein. *Int J Med Microbiol* 301:602-611.
- Weiberg, A., Wang, M., Lin, F. M., Zhao, H., Zhang, Z., Kaloshian, I., Huang, H. D., and Jin, H. 2013. Fungal small RNAs suppress plant immunity by hijacking host RNA interference pathways. *Science* 342:118-123.
- Zuccaro, A., Lahrmann, U., Guldener, U., Langen, G., Pfiffi, S., Biedenkopf, D., Wong, P., Samans, B., Grimm, C., Basiewicz, M., Murat, C., Martin, F., and Kogel, K. H. 2011. Endophytic life strategies decoded by genome and transcriptome analyses of the mutualistic root symbiont *Piriformospora indica*. *PLoS Pathog* 7:e1002290.

## Supplemental data

SUPPLEMENTAL TABLE 1 | Effectors of filamentous plant-associated microbes for which molecular virulence targets were identified

Effector	Origin	Target	Function	Reference
BEC4	<i>Blumeria graminis</i> f.sp. <i>hordei</i>	ARF-GAP proteins	Interference with host vesicle trafficking	(Schmidt et al. 2014)
Avr2	<i>Cladosporium fulvum</i>	Cysteine proteases	Cysteine protease inhibition	(Shabab et al. 2008, van Esse et al. 2008)
Avr4	<i>Cladosporium fulvum</i>	Chitin	Hyphal protection	(van den Burg et al. 2006)
Ecp6	<i>Cladosporium fulvum</i>	Chitin	Perturbation chitin-triggered immunity	(de Jonge et al. 2010)
CfTom1	<i>Cladosporium fulvum</i>	$\alpha$ -tomatine	Detoxification	(Okmen et al. 2013)
Victorin	<i>Cochliobolus victoria</i>	TRX-h5	Induction of LOV1-mediated cell death	(Lorang et al. 2012)
SP7	<i>Rhizophagus irregularis</i>	ERF19	Deregulation of host gene expression	(Kloppholz et al. 2011)
HaRxL44	<i>Hyaloperonospora arabidopsidis</i>	MED19a	Interference with SA-triggered immunity	(Caillaud et al. 2013)
MISSP7	<i>Laccaria bicolor</i>	JAZ6	Deregulation of host gene expression	(Plett et al. 2014)
AvrPiz-t	<i>Magnaporthe oryzae</i>	RING E3 ubiquitin ligase APIP6	Suppression of MAMP-triggered immunity	(Park et al. 2012)
Slp1	<i>Magnaporthe oryzae</i>	Chitin	Perturbation chitin-triggered immunity	(Mentlak et al. 2012)
MFAvr4	<i>Mycosphaerella fijiensis</i>	Chitin	Hyphal protection	(Stergiopoulos et al. 2010)
Mg1LysM	<i>Mycosphaerella graminicola</i>	Chitin	Hyphal protection	(Marshall et al. 2011)
Mg3LysM	<i>Mycosphaerella graminicola</i>	Chitin	Perturbation chitin-triggered immunity	(Marshall et al. 2011)
Avr3a	<i>Phytophthora infestans</i>	CMPG1	E3 ligase stabilization	(Bos et al. 2010)
Avrblb2	<i>Phytophthora infestans</i>	C14 protease	Suppression of protease secretion	(Bozkurt et al. 2011)
EPI1	<i>Phytophthora infestans</i>	Serine proteases	Inhibition of serine proteases	(Tian et al. 2004)
EPI10	<i>Phytophthora infestans</i>	Serine proteases	Inhibition of serine proteases	(Tian et al. 2005)
EPIc1	<i>Phytophthora infestans</i>	Cysteine proteases	Inhibition of cysteine proteases	(Song et al. 2009, Tian et al. 2007)
EPIc2B	<i>Phytophthora infestans</i>	Cysteine proteases	Inhibition of cysteine proteases	(Song et al. 2009, Tian et al. 2007)
PexRD2	<i>Phytophthora infestans</i>	MAPKKk	Suppression of host cell death	(King et al. 2014)
Pi03.192	<i>Phytophthora infestans</i>	NTP1, NTP2	Suppression of transcription factor relocation	(McLellan et al. 2013)
GIP1	<i>Phytophthora sojae</i>	$\beta$ -1,3-glucanases	Glucanase inhibition	(Rose et al. 2002)
RTP1p	<i>Uromyces fabae/U. striatus</i>	Proteases	Protease inhibition	(Pretsch et al. 2013)
Cmu1	<i>Ustilago maydis</i>	Cm2	Interference with SA biosynthesis	(Djamei et al. 2011)
Pep1	<i>Ustilago maydis</i>	POX12	Inhibition of peroxidase-mediated ROS production	(Hemetsberger et al. 2012)
Pit2	<i>Ustilago maydis</i>	CP2, CP1A/B, XCP2 proteases	Cysteine protease inhibition	(Mueller et al. 2013)
Tin2	<i>Ustilago maydis</i>	TmTTK1	Control of anthocyanin biosynthesis	(Tanaka et al. 2014)





**Homologs of *Verticillium dahliae*  
effector Ave1 contribute to  
virulence of fungal pathogens  
of diverse plant hosts**

**Chapter 3**

Jordi C. Boshoven

Malaika K. Ebert<sup>#</sup>

Yin Song<sup>#</sup>

Hanna Rovenich

Eduardo Rojas-Padilla

Melvin D. Bolton

Bart P. H. J. Thomma

<sup>#</sup>These authors contributed equally

A modified version has been submitted to *Phytopathology*



## Abstract

*Verticillium dahliae* is a soil-borne fungal pathogen with a wide host range including many crops. Tomato immune receptor Ve1 confers resistance to *V. dahliae* race 1 strains that express VdAve1. On plants that do not carry Ve1, Ave1 acts as a virulence factor of *V. dahliae*. Homologs of Ave1 are mostly found in plants and in a handful of fungal plant pathogens including *Fusarium oxysporum* (FoAve1), *Cercospora beticola* (CbAve1), and *Colletotrichum higginsianum* (ChAve1). In the bacterium *Xanthomonas axonopodis* the Ave1 homolog XacPNP was previously described as a virulence factor. The homologs FoAve1, CbAve1 are recognized by Ve1 although not as efficient as VdAve1. In this study, we analysed the role of the individual Ave1 homologs in virulence. We generated targeted gene deletions in *Fusarium oxysporum*, *Cercospora beticola* and *Colletotrichum higginsianum* and showed that both FoAve1 and CbAve1 are virulence factors in contrast to ChAve1 that does not contribute to the virulence of the pathogen. We subsequently tested whether Ave1 homologs FoAve1, CbAve1, ChAve1 and XacPNP act in a similar fashion as VdAve1 by expressing these genes in a *V. dahliae* VdAve1 deletion strain. We concluded that VdAve1 on the one hand and FoAve1, CbAve1, ChAve1 and XacPNP on the other hand act in different manners.

## Introduction

Plants are surrounded by microbes including oomycetes, fungi, bacteria and viruses. To prevent colonization by harmful microbes, plants evolved pattern recognition receptors (PRRs) that recognize microbe associated molecular patterns (MAMPs) and activate MAMP-triggered immunity (MTI) (Boller and Felix 2009; Dodds and Rathjen 2010). MAMPs generally are structural components such as bacterial elongation factor Tu (EF-Tu), flagellin, lipopolysaccharides (LPS), fungal chitin and oomycete  $\beta$ -glucans, which are typically conserved across genera of microbes. Successful pathogens secrete effector proteins that modulate host immunity to overcome PRR-mediated plant defence (de Jonge et al. 2012; Rovenich et al. 2014). Such effectors are often lineage-specific, facilitate colonization of the plant, and therefore contribute to virulence of the pathogen. In turn, plants evolved immune receptors, typically called resistance (R) proteins, which recognize particular effectors or their activities to mount an immune response known as effector-triggered immunity (ETI) (Jones and Dangl 2006). To overcome recognition by the plant, pathogens have to mutate or lose the recognized effector or, alternatively, evolve new effectors to suppress the ETI response. Nevertheless, not all pathogen-secreted molecules follow the strict MAMP-effector dichotomy but rather display traits of either class of molecules, and therefore MTI and ETI responses cannot strictly be separated (Thomma et al. 2011). Typical examples of such pathogen-secreted molecules are Nep1-like proteins, which are virulence factors that are dispersed throughout three kingdoms of life that act as a MAMP at least in *Arabidopsis* (Böhm et al. 2014; Oome et al. 2014). Although less widely distributed, phenomena have been observed for homologs of the Avr4 effector from the tomato leaf mould fungus *Cladosporium fulvum* (Stergiopoulos et al. 2010), and for homologs of the Ave1 effector from the vascular wilt fungus *Verticillium dahliae* (de Jonge et al. 2012). This has inspired the proposal of the so-called invasion model, which states that plants evolved receptors for all types of molecules, pathogen- as well as host-derived, that can betray microbial invasion to mount the appropriate immune responses (Cook et al. 2015).

*V. dahliae* is a soil-borne fungal pathogen that causes vascular wilt in over 200 plant species, including important crop species (Fradin and Thomma 2006). Resistance to *V. dahliae* is mediated by the immune receptor Ve1 (Fradin et al. 2009; Kawchuk et al. 2001), of which homologs were found in mint, hop and cotton amongst other plant species (Majer et al. 2014; Vining and Davis 2009; Zhang et al. 2011). Recently, the effector protein that is secreted by *V. dahliae* race 1 strains and that activates Ve1-mediated resistance was identified as Ave1, and all resistance-breaking race 2 strains analysed thus far lack the complete Ave1 gene (de Jonge et al. 2012; Faino et al.

2016). Importantly, *Ave1* deletion strains show reduced aggressiveness on tomato plants that lack *Ve1*, revealing that *Ave1* acts as a virulence factor on this host species. Additionally, *Ave1* was also shown to act as a virulence factor of *V. dahliae* on Arabidopsis plants (de Jonge et al. 2012). However, how *Ave1* contributes to *V. dahliae* virulence on these plant hosts remains unknown thus far.


Intriguingly, although pathogen effectors are typically lineage-specific traits, many homologs of *Ave1* can be found in public databases (de Jonge et al. 2012). Remarkably, a wealth of *Ave1* homologs was found in plants as well as in a handful of fungal plant pathogens, including *Fusarium oxysporum* (*FoAve1*), *Cercospora beticola* (*CbAve1*) and *Colletotrichum higginsianum* (*ChAve1*), and in the bacterial plant pathogen *Xanthomonas axonopodis* (*XacPNP*) (de Jonge et al. 2012; Gottig et al. 2008). Interestingly, it was shown that *Ve1* is not confined to *V. dahliae* *Ave1* (*VdAve1*), as *Ve1* is able to also recognize *FoAve1* and *CbAve1*, while *ChAve1* is not recognized (de Jonge et al. 2012). Consequently, *Ve1* was indeed found to be able to recognize *F. oxysporum* in tomato and mount a defense response (de Jonge et al. 2012). Remarkably, *Ve1* was furthermore found to recognize an endogenous *Ave1* homolog (*SlAve1*) that shares a high degree of identity with *VdAve1* (de Jonge et al. 2012).

Most plant homologs of *Ave1* have been annotated either as plant natriuretic peptides (PNPs) or as expansin-like proteins, and functionally analyzed members were implicated in the regulation of water and ion homeostasis, and consequently in many downstream processes including growth, net water uptake, photosynthesis, stomatal opening and gas exchange (Gehring and Irving 2003; Ludidi et al. 2004; Maryani et al. 2001; Pharmawati et al. 2001). The observation that the microbial-derived *Ave1* homologs do not follow the phylogeny of the species in which they occur is generally taken as evidence for horizontal acquisition, and it has thus been speculated that the microbial *Ave1* homologs were acquired from plants (de Jonge et al. 2012; Gottig et al. 2008). In this respect it is interesting that the *X. axonopodis* homolog *XacPNP* affects homeostasis and photosynthesis in citrus plants, and thus promotes bacterial proliferation as a virulence factor in citrus plants (Garavaglia et al. 2010; Gottig et al. 2008; Nembaware et al. 2004). In this study, we investigate whether the *Ave1* homologs of several fungal pathogens act as genuine virulence factors.

## Results

### Characterization of Ve1-mediated tomato defence against *Fusarium oxysporum*

We previously demonstrated that tomato Ve1 can recognize FoAve1, leading to a defence response that affects infection of Ve1-carrying tomato by *F. oxysporum* (de Jonge et al. 2012). Because recognition of FoAve1 by Ve1-carrying tomato plants does not lead to full immunity against *F. oxysporum* we investigated the occurrence of expression of FoAve1 in two *F. oxysporum* strains, Bt.01 and Fol4287, in tomato plants lacking Ve1 with reverse-transcription PCR at 14 dpi. Clear expression of FoAve1 was monitored at 14 dpi (Figure 1A).



**FIGURE 1 | FoAve1 is a virulence factor recognized by tomato Ve1.** (A) Expression of FoAve1 in *F. oxysporum* in wild type (Wt) strains Bt.01 and Fol4287 and the FoAve1 deletion strains ( $\Delta$ FoAve1) inoculated on tomato at 14 dpi. (B) (Upper) FoAve1 deletion strains ( $\Delta$ FoAve1) of *F. oxysporum* strains Bt.01 and Fol4287 escape recognition by tomato Ve1 compared with the corresponding wild type (Wt) strains evidenced by stunted Ve1 plants at 14 dpi. (Lower) FoAve1 deletion strains display reduced virulence compared with the corresponding *F. oxysporum* wild type strains Bt.01 and Fol4287 on tomato lacking Ve1 (ve1) evidenced by reduced stunting at 14 dpi. (C) Canopy area of Ve1 tomato and tomato lacking Ve1 after inoculation with FoAve1 deletion strains and the corresponding *F. oxysporum* wild type strains at 14 dpi. (D) FoAve1 deletion strains display reduced virulence compared to the corresponding *F. oxysporum* wild type strains Bt.01 and Fol4287 on tomato lacking Ve1 evidenced by reduced fungal biomass at 14 dpi. 8 plants were pooled per 2 plants (n=4). Different letter labels indicate statistically significant differences ( $P < 0.05$ ). Similar results were observed in at least 3 biological replications for all experiments.

To provide further evidence for the role of FoAve1-recognition in Ve1-mediated tomato defence against Fusarium wilt, we generated FoAve1 deletion mutants in *F. oxysporum* strains Bt.01 and FoI4287 through homologous recombination. Deletion of FoAve1 was confirmed by PCR (Supplemental Figure 1A) and the deletion strains were subsequently evaluated for their ability to colonize Ve1 tomato plants. As anticipated, inoculation of Ve1 tomato plants with FoAve1 deletion strains resulted in more extensive symptoms of Fusarium wilt disease when compared to tomato plants inoculated with the corresponding wild type strains (Figure 1B (Upper) and 1C). Altogether, these data show that tomato Ve1 is able to recognize FoAve1 and activate a defense response that inhibits colonization by *F. oxysporum*, albeit that the response does not fully arrest the fungus and immunity is not fully established. At later stages after inoculation clear symptoms of wilt disease are observed on Ve1 plants.


### **FoAve1 is a virulence factor of *F. oxysporum* during tomato colonization**

According to the paradigm that plant immune receptors recognize crucial virulence factors of microbial pathogens, it is expected that FoAve1 contributes to *F. oxysporum* virulence. To test this hypothesis, FoAve1 deletion strains were inoculated on tomato plants lacking Ve1. Indeed, FoAve1 deletion strains displayed reduced virulence on tomato plants when compared to the corresponding *F. oxysporum* wild type strain as visualized by the difference in canopy area of the inoculated tomato plants (Figure 1B (Lower) and 1C). In addition, plants inoculated with FoAve1 deletion strains resulted in less fungal colonization compared to those inoculated with wild type *F. oxysporum* (Figure 1D). These results suggest that FoAve1 is a virulence factor of *F. oxysporum*, albeit that its contribution to virulence appears to be relatively minor.

### **ChAve1 is not a virulence factor of *C. higginsianum* during Arabidopsis colonization**

Besides the vascular wilt fungi *V. dahliae* and *F. oxysporum*, that belong to the class of Sordariomycetes, an Ave1 homolog is found in the foliar Sordariomycete *C. higginsianum* (*ChAve1*). Using real-time PCR, *ChAve1* expression was monitored in wild type *C. higginsianum* during colonization of Arabidopsis plants from 1 to 4 dpi, by which time the plant tissue was completely macerated. No expression of *ChAve1* was detected at any of these time points. Nevertheless, we investigated whether *ChAve1* contributes to virulence in this foliar pathogen. For this purpose, *ChAve1* deletion strains were generated that, after confirmation with PCR (Supplemental Figure 1B), were evaluated for aggressiveness on *Arabidopsis thaliana*. As expected *ChAve1* deletion strains did not show reduced virulence when compared to the


corresponding wild type *C. higginsianum*, as infections resulted in lesions with similar sizes (Figure 2A). Moreover, similar levels of pathogen biomass were recorded in plants inoculated with the wild-type fungus and the deletion strains (Figure 2B). Together these results show that ChAve1 is not a virulence factor of *C. higginsianum* on *Arabidopsis*, likely due to the lack of *ChAve1* expression.



**FIGURE 2 | ChAve1 is not a virulence factor of *C. higginsianum*.** (A) *ChAve1* deletion strains ( $\Delta$ *ChAve1*) show a similar lesion size compared to the corresponding *C. higginsianum* wild type strain (Wt) at 3 dpi on 3-week-old *A. thaliana*. Twenty four lesions on three *A. thaliana* plants were measured per strain. No significant difference was observed ( $P < 0.05$ ). (B) No difference in fungal biomass was observed at 4 dpi on spray inoculated 3-week-old *Arabidopsis* plants ( $n = 4$ ). Different letter labels indicate statistically significant differences ( $P < 0.05$ ). Similar results were observed in at least 3 biological replications for all experiments in this figure.

### **CbAve1 is a virulence factor of *C. beticola* during sugarbeet colonization**

Besides Sordariomycetes, Ave1 homologs were also identified as we previously identified an Ave1 homolog in the Dothidiomycete *C. beticola* (*CbAve1*). Expression of *CbAve1* was investigated in *C. beticola* during colonization of sugarbeet plants between 3 and 18 dpi using real-time PCR. Clear expression was detected between these time points. To evaluate the contribution of *CbAve1* in *C. beticola* virulence, *CbAve1* deletion strains were generated and confirmed with PCR. For this pathogen, deletion of the Ave1 homolog resulted in reduced virulence compared to the corresponding wild type strain, as infection resulted in reduced numbers and size of the lesions (Figure 3A; B). As expected, *CbAve1* deletion strains showed reduced colonization on sugarbeet plants compared to the corresponding *C. beticola* wild type strain (Figure 3C). Taken together, these results show that *CbAve1* is a virulence factor of *C. beticola*.



**FIGURE 3 | *CbAve1* is a virulence factor of *C. beticola*.** (A) Expression of *CbAve1* during infection of *C. beticola* on sugar beet. Leaves of 6- to 7-week-old sugar beet plants were inoculated with wild-type *C. beticola* and collected at regular intervals from 3 to 18 days post inoculation (dpi). RT-qPCR was performed to determine the relative expression levels of *CbAve1* using the *C. beticola actin* gene as a reference, and compared with *CbAve1* expression in *C. beticola* in sugar beet plants upon the wild-type *C. beticola* inoculation at 3 dpi, which is set to 1. (B) *CbAve1* deletion strains ( $\Delta CbAve1$  #1 and  $\Delta CbAve1$  #2) show reduced virulence compared to the corresponding *C. beticola* wild type strain (WT) visualized by a reduction in lesion area on sugar beet leaves at 15 dpi. (C) Quantification of leaf lesion caused by *C. beticola* on sugar beet plants at 15 dpi ( $n > 5$ ). Bars represent the average percentage of leaf lesion area of whole leaf area with standard deviations. (D) Fungal biomass determined with qPCR in *Cercospora*-inoculated sugar beet plants at 15 dpi. The fungal biomass in sugar beet plants upon inoculation with the wild-type *C. beticola* is set to 100% (control). Different letter labels indicate statistically significant differences ( $P < 0.05$ ).




### Functional diversification among plant and fungal Ave1 homologs

VdAve1 is a virulence factor of *V. dahliae* that shows a high degree of identity with homologs from plants. It has therefore been proposed that VdAve1 has been acquired by *V. dahliae* from plants through horizontal gene transfer (de Jonge et al. 2012). However, the function of VdAve1 through which it contributes to *V. dahliae* aggressiveness remains presently enigmatic. Similarly, the function of the Ave1 homologs that are found in various pathogens also remains unclear. In order to evaluate whether all homologs share their functionality, we tested whether they can complement the virulence defect that is observed upon VdAve1 deletion in *V. dahliae*. To this end, we transformed one of the VdAve1 deletion strains of *V. dahliae* with constructs to drive expression of the plant homologs derived from tomato (*S. lycopersicum*; *SlAve1*) and grape (*V. vinifera*; *VvAve1*) by the *V. dahliae* VdAve1 promoter (de Jonge et al. 2012). To confirm that the plant homologs were expressed in *V. dahliae* upon colonization of tomato we monitored *SlAve1* and *VvAve1* expression with PCR. At 14 dpi we monitored clear expression of *SlAve1* and *VvAve1* in 3 transformants for each construct (Supplemental Figure 2a). Subsequently, we evaluated the ability of these transformants to cause disease on tomato plants lacking *Ve1*. All VdAve1 deletion strains carrying *SlAve1* and *VvAve1* showed a similar phenotype and fungal colonization as the corresponding VdAve1 deletion strain (Figure 4; Supplemental Figure 2). These results suggest that two plant Ave1 homologs that share a high identity with VdAve1 are unable to restore virulence in a VdAve1 deletion strain, suggesting that Ave1 homologs from plants act in a different manner as VdAve1.

Subsequently, we evaluated whether microbial Ave1 homologs can reinstall the compromised virulence of *V. dahliae* that results from VdAve1 deletion. To this end, we transformed one of the VdAve1 deletion strains with constructs to drive expression of the Ave1 homologs derived from *F. oxysporum* (*FoAve1*) *C. higginsianum* (*ChAve1*) and *C. beticola* (*CbAve1*) by the VdAve1 promoter. We also included the Ave1 homolog derived from the plant pathogenic bacterium *Xanthomonas axonopodis* (*XacPNP*) that was previously described as a virulence factor (Gottig et al. 2008). We confirmed in a minimum of 2 transformants that the constructs were expressed (Supplemental Figure 2a). Subsequently, we tested the *V. dahliae* transformants on tomato lacking *Ve1* and compared them with the corresponding wild type *V. dahliae* and the *V. dahliae* VdAve1 deletion strain complemented with VdAve1. VdAve1 deletion strains expressing *FoAve1*, *CbAve1*, *ChAve1* and *XacPNP* showed a similar disease phenotype and fungal colonization as VdAve1 deletion strains (Figure 4; Supplemental Figure 2). (Figure 4; Supplemental Figure 2). These results show that the Ave1 homologs *FoAve1*, *CbAve1*, *ChAve1* and *XacPNP* cannot restore virulence in a VdAve1 deletion strain which may suggest that Ave1 homologs of *V. dahliae* on the one hand, and *F. oxysporum*, *C. beticola*, *C. higginsianum* and *X. axonopodis* on the other hand, contribute to fungal virulence in a different manner.





**FIGURE 4 | VdAve1 acts in a different manner than Ave1 homologs derived from plant pathogens.** *V. dahliae* VdAve1 deletion strain ( $\Delta$ VdAve1) complemented with Ave1 homologs derived from *V. dahliae* (VdAve1), tomato (*S. lycopersicum*; SlAve1), grape (*V. vinifera*; VvAve1), *F. oxysporum* (FoAve1), *C. beticola* (CbAve1), *C. higginsianum* (ChAve1) and *Xanthomonas axonopodis* (XacPNP) inoculated on tomato plants. (A) Gene expression of Ave1 homologs by *V. dahliae* during colonization of tomato at 14 dpi. (B) Canopy area of tomato plants inoculated with *V. dahliae* strains expressing Ave1 homologs compared to *V. dahliae* wild type (Vd wt) at 14 dpi. (C) Fungal colonization of tomato plants inoculated with *V. dahliae* strains expressing Ave1 homologs compared to *V. dahliae* wild type (Vd wt) at 14 dpi. (D) Photos taken from the side and the top of tomato plants inoculated with *V. dahliae* strains expressing Ave1 homologs compared to *V. dahliae* wild type (Vd wt) at 14 dpi.

## Discussion

### The role of FoAve1 in recognition by tomato immune receptor Ve1

We have previously shown that recognition of FoAve1 by Ve1 leads to a defence response (de Jonge et al. 2012) and that Ve1 tomato plants can recognize *F. oxysporum*. To confirm that FoAve1 is involved in recognition we first analysed FoAve1 expression. A previous study could not detect FoAve1 expression in *F. oxysporum* after inoculation on tomato (Schmidt et al. 2013). However, in the conditions that we tested clear FoAve1 expression was monitored in *F. oxysporum* on tomato at 14 dpi. In addition, deletion of FoAve1 resulted in abolishment of recognition by Ve1. Our data suggest that the incomplete disease resistance cannot be attributed to a lack of FoAve1 expression, but may be the result of a less efficient detection of this Ave1 homolog by Ve1. Similar differences in recognition efficiencies were previously observed for *Cladosporium fulvum* effector Ecp2 and its homolog MfEcp2 of *Mycosphaerella fijiensis* when coexpressed with tomato immune receptor Cf-Ecp2 (Stergiopoulos et al. 2010).

### Some Ave1 homologs act as virulence factors

Previously, we have shown that Ave1 is a virulence factor of *V. dahliae* (de Jonge et al. 2012). In this study, we tested whether Ave1 homologs in other plant pathogens act as virulence factors as well. We showed that deletion of Ave1 homologs in *F. oxysporum* and *C. beticola* resulted in reduced aggressiveness on their respective plant hosts. In contrast, the Ave1 homolog ChAve1 of the foliar pathogen *C. higginsianum* does not contribute to virulence due to the lack of expression *in planta*. Possibly, the lack of expression is due to the availability of effectors with a similar function that render ChAve1 functionality redundant. Alternatively, recognition of ChAve1 by a host immune receptor posed pressure on *C. higginsianum* to avoid recognition, resulting in the lack of ChAve1 expression. Interestingly, homologs of tomato Ve1 have been identified in many other plant species, of which some within as well as outside the Solanaceae family have been shown to be functional immune receptors (Song et al. 2017a).

### Functional diversification among Ave1 homologs

Since Ave1 homologs derived from plants display high similarity to Ave1, it has been suggested that Ave1 was acquired through horizontal gene transfer from plants (de Jonge et al. 2012). As the function of the Ave1 homologs remain unknown we tested whether the Ave1 homologs share the same functionality by complementing a *V. dahliae* Ave1 deletion strain with homologs derived from plants and various plant

3

pathogens. Highly similar Ave1 homologs derived from plants are unable to restore virulence in an Ave1 deletion strain. Therefore we conclude that Ave1 homologs derived from plants act in a different manner as Ave1. Similarly, Ave1 homologs derived from *F. oxysporum*, *C. beticola*, *C. higginsianum* and *X. axonopodis* cannot restore the virulence penalty caused by the deletion of Ave1. Therefore, we conclude that Ave1 from *V. dahliae* on the one hand, and Ave1 homologs derived from *F. oxysporum*, *C. beticola*, *C. higginsianum* and *X. axonopodis* on the other hand, contribute to virulence in a different manner. Similarly, functional diversification within effector families has been observed for LysM and NLP effectors (Kombrink and Thomma 2013; Santhanam et al., 2013). Alternatively, it is important to note that *V. dahliae*, *C. beticola*, *C. higginsianum* and *X. axonopodis* have diverse plant hosts and that the reason for the inability to restore virulence in an *V. dahliae* Ave1 deletion strain could be that each homolog targets a host-specific protein. This would for example explain why the homolog of *C. beticola* that likely targets a sugarbeet protein cannot restore virulence in an *V. dahliae* Ave1 deletion strain colonizing tomato. Host specific effector adaptation has been suggested for the closely related oomycetes *Phytophthora infestans* and *P. mirabilis* that are pathogens on different hosts. Homologous effectors of both pathogens act more efficiently on their specific host target than on a “foreign” host target (Dong et al. 2014). Nevertheless, the finding that the Ave1 homolog from *F. oxysporum* f.sp. *lycopersici* cannot complement the *VdAve1* deletion strain for virulence on tomato argues against this hypothesis.

## Materials and methods

### Deletion and complementation strains

Deletion strains were generated in *F. oxysporum* and *C. higginsianum* by amplifying 1,5 kb sequences (Supplemental table 1) flanking the coding sequence of *FoAve1* and *ChAve1*, respectively, and cloned as previously described (Frandsen et al. 2008) into vector pRF-HU2 containing a nourseothricin cassette for selection. For the complementation of *V. dahliae* Ave1 deletion strains *FoAve1*, *CbAve1*, *ChAve1*, *SIPNP*, *VvPNP* and *XacPNP* were obtained by gene synthesis (Eurofins Genomics, Ebersberg, Germany). The synthesized genes were then cloned into vector pFBT005 under the *VdAve1* promoter, containing a nourseothricin cassette for selection. *F. oxysporum* (Fol4287 and Bt.01) and *C. higginsianum* (IMI349063A) conidiospores were transformed on a Hybond-N<sup>+</sup> filter by *A. tumefaciens* carrying the pRF-HU2 plasmid containing the constructs and a *V. dahliae* Ave1 deletion strain (JR2) was transformed by *A. tumefaciens* carrying the pFBT005 plasmid containing the constructs as previously described (Santhanam 2012). Transformants

were then selected on potato dextrose agar (PDA) (Thermo Fisher Scientific Inc, Breda, The Netherlands) containing hygromycin B (Duchefa Biochemie BV, Haarlem, The Netherlands) for selection of *F. oxysporum*, *C. beticola* and *C. higginsianum* transformants or nourseothricin sulphate (Sigma Aldrich Chemie BV, Zwijndrecht, The Netherlands) for selection of *V. dahliae* transformants. After five to seven days at room temperature, individual transformants were transferred from the filter to fresh PDA plates with the appropriate selection and incubated for ten days (Santhanam 2012). To verify the transformants, fungal spores were collected and genomic DNA was extracted followed by a PCR to test the presence of the hygromycin or nourseothricin cassette and the presence of the construct in the genome (Supplemental table 1).

*CbAve1* deletion mutants of *C. beticola* were generated using the split-marker approach described by Catlett *et al.* (2003). Genomic DNA of the wild type *C. beticola* strain 09-40 and pDAN vector (Friesen *et al.*, 2006) served as PCR templates to generate split-marker PCR constructs used for transformation. Primers are listed in Supplemental table 1. PEG-mediated transformation of the wild type *C. beticola* strain 09-40 was performed as previously described (Bolton *et al.*, 2016). Site-directed gene replacement was confirmed by the absence of PCR product using split-marker *CbAve1* 1F forward primer of the 5' flank of the target gene and MDB-760 reverse primer designed on the coding sequence of *CbAve1*.

### Pathogen inoculations

Plants were grown in soil in the greenhouse at 21°C/19°C during 16-h/8-h day/night periods, respectively, with 70% relative humidity and 100 W m<sup>-2</sup> supplemental light when the intensity dropped below 150 W m<sup>-2</sup>.

For *V. dahliae* and *F. oxysporum* inoculations, 10-day-old tomato (*Solanum lycopersicum*; cv. MoneyMaker or MoneyMaker 35S:Ve1) (Fradin *et al.* 2009) were uprooted, rinsed in water and dipped for 5 minutes in a suspension of 10<sup>6</sup> conidiospores per mL of water harvested from 1- to 2-week-old *V. dahliae* or *F. oxysporum* cultures on PDA as previously described (Fradin *et al.* 2009). Control plants received the same treatment, but the roots were dipped in water without conidiospores. After replanting in fresh soil, plants were incubated at standard greenhouse conditions. Disease development was monitored up to 21 days post inoculation (dpi).

For *C. higginsianum* inoculations 3-week-old Arabidopsis (*A. thaliana* ecotype Co-0 or Col-0 35S:Ve1) (Fradin *et al.* 2011) were inoculated on the leaves with either 2 µl drops or sprayed with a suspension of 10<sup>6</sup> conidia per mL of water harvested from 1- to 2-week-old *C. higginsianum* cultures on Mathurs' agar as previously

described (O'Connell et al. 2004; Sherriff et al. 1994). Control plants received the same treatment, but the leaves were sprayed with water without conidiospores. After sealing the plants inside a transparent closed box lined with wet tissue paper to provide high humidity, they were incubated at 25°C, under a 16-h/ 8-h light/dark regime (O'Connell et al. 2004). Disease development was monitored up to 4 dpi.

For *C. beticola* inoculations 6- to 7-week-old sugarbeet plants (*Beta vulgaris*) were inoculated by evenly spaying spore suspension of 10<sup>5</sup> conidia per mL of water on the lower side of the leaves harvested from *C. beticola* cultures growing on V8 solid medium as previously described (Bolton et al. 2012). Control plants received the same treatment, but the leaves were sprayed with water. Plants were incubated at standard greenhouse conditions and disease development was monitored up to 18 dpi.

### Fungal biomass and gene expression

Stem sections of tomato, cut from the base of the stem up to the cotyledons, were collected at 7 and 14 dpi from plant inoculated with *F. oxysporum* or *V. dahliae*. Arabidopsis leaves were collected at 1-4 dpi after inoculation with *C. higginsianum*. Sugarbeet leaves were collected at regular intervals between 3 and 18 dpi from plants inoculated with *C. beticola*. Collected plant tissue was flash frozen in liquid nitrogen and ground to powder, of which an aliquot of ~100 mg was used for RNA extraction with the Quick-RNA™ Miniprep kit (Zymo Research Europe GmbH, Freiburg, Germany), and cDNA was synthesized using M-MLV Reverse Transcriptase (Promega Benelux BV, Leiden, The Netherlands). Fungal biomass was determined by quantifying the expression of *VdGAPDH*, *FoTUB*, *ChTUB* and *CbAct* relative to the expression of *SIRUB* (tomato), *AtRub* (Arabidopsis), *BvAct* (sugarbeet), respectively. Expression of *VdAve1*, *FoAve1*, *CbAve1*, *ChAve1*, *SIPNP* and *VvPNP* was determined by quantifying the expression relative to the expression of *VdGAPDH*, *FoTUB*, *ChTUB* and *CbAct*, respectively (Supplemental table 1). Real-time PCR was carried out on an ABI7300 PCR machine (Applied Biosystems, Life Technologies Europe BV, Bleiswijk, The Netherlands) in combination with the qPCR SensiMix kit (BioLine, GC Biotech BV, Alphen aan den Rijn, The Netherlands). The following Real-time PCR conditions were used: an initial 95°C denaturation step for 10 minutes followed by denaturation for 15 seconds at 95°C, annealing for 60 seconds at 60°C, and extension at 72°C for 40 cycles.

### Acknowledgements

Work in the laboratory of BPHJT is supported by the Research Council Earth and Life Sciences (ALW) of the Netherlands Organization for Scientific Research (NWO). We thank Bert Essenstam and Pauline Sanderson of Unifarm for excellent plant care. We thank Jelle Postma, Dorien Oppelaar and Saskia Bosman for technical support.

## References

- Böhm, H., Albert, I., Oome, S., Raaymakers, T. M., Van den Ackerveken, G., and Nürnberger, T. 2014. A conserved peptide pattern from a widespread microbial virulence factor triggers pattern-induced immunity in *Arabidopsis*. *PLoS Pathog* 10:e1004491.
- Boller, T., and Felix, G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol* 60:379-406.
- Bolton, M. D., Birla, K., Rivera-Varas, V., Rudolph, K. D., and Secor, G. A. 2012. Characterization of *CbCyp51* from field isolates of *Cercospora beticola*. *Phytopathology* 102:298-305.
- Cook, D. E., Mesarich, C. H., and Thomma, B. P. H. J. 2015. Understanding plant immunity as a surveillance system to detect invasion. *Annu Rev Phytopathol*.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- Dodds, P. N., and Rathjen, J. P. 2010. Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* 11:539-548.
- Dong, S., Stam, R., Cano, L. M., Song, J., Sklenar, J., Yoshida, K., Bozkurt, T. O., Oliva, R., Liu, Z., Tian, M., Win, J., Banfield, M. J., Jones, A. M., van der Hoorn, R. A., and Kamoun, S. 2014. Effector specialization in a lineage of the Irish potato famine pathogen. *Science* 343:552-555.
- Faino, L., Seidl, M. F., Shi-Kunne, X., Pauper, M., van den Berg, G. C., Wittenberg, A. H., and Thomma, B. P. 2016. Transposons passively and actively contribute to evolution of the two-speed genome of a fungal pathogen. *Genome Res* 26:1091-1100.
- Fradin, E. F., and Thomma, B. P. H. J. 2006. Physiology and molecular aspects of *Verticillium* wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol Plant Pathol* 7:71-86.
- Fradin, E. F., Abd-El-Halim, A., Masini, L., van den Berg, G. C., Joosten, M. H., and Thomma, B. P. H. J. 2011. Interfamily transfer of tomato *Ve1* mediates *Verticillium* resistance in *Arabidopsis*. *Plant Physiol* 156:2255-2265.
- Fradin, E. F., Zhang, Z., Ayala, J. C. J., Castroverde, C. D., Nazar, R. N., Robb, J., Liu, C.-M., and Thomma, B. P. 2009. Genetic dissection of *Verticillium* wilt resistance mediated by tomato *Ve1*. *Plant Physiology* 150:320-332.
- Frandsen, R. J., Andersson, J. A., Kristensen, M. B., and Giese, H. 2008. Efficient four fragment cloning for the construction of vectors for targeted gene replacement in filamentous fungi. *BMC Mol Biol* 9:70.
- Garavaglia, B. S., Thomas, L., Gottig, N., Dunger, G., Garofalo, C. G., Daurelio, L. D., Ndimba, B., Orellano, E. G., Gehring, C., and Ottado, J. 2010. A eukaryotic-acquired gene by a biotrophic phytopathogen allows prolonged survival on the host by counteracting the shut-down of plant photosynthesis. *PLoS One* 5:e8950.
- Gehring, C., and Irving, H. 2003. Natriuretic peptides - a class of heterologous molecules in plants. *Int J Biochem Cell Biol* 35:1318-1322.
- Gottig, N., Garavaglia, B. S., Daurelio, L. D., Valentine, A., Gehring, C., Orellano, E. G., and Ottado, J. 2008. *Xanthomonas axonopodis* pv. *citri* uses a plant natriuretic peptide-like protein to modify host homeostasis. *Proc Natl Acad Sci U S A* 105:18631-18636.
- Jones, J. D., and Dangl, J. L. 2006. The plant immune system. *Nature* 444:323-329.
- Kawchuk, L. M., Hachey, J., Lynch, D. R., Kulcsar, F., Van Rooijen, G., Waterer, D. R., Robertson, A., Kokko, E., Byers, R., and Howard, R. J. 2001. Tomato *Ve* disease resistance genes encode cell surface-like receptors. *Proceedings of the National Academy of Sciences of the United States of America* 98:6511-6515.
- Ludidi, N., Morse, M., Sayed, M., Wherrett, T., Shabala, S., and Gehring, C. 2004. A recombinant plant natriuretic peptide causes rapid and spatially differentiated K<sup>+</sup>, Na<sup>+</sup> and H<sup>+</sup> flux changes in *Arabidopsis thaliana* roots. *Plant Cell Physiol* 45:1093-1098.

- Majer, A., Javornik, B., Cerenak, A., and Jakse, J. 2014. Development of novel EST-derived resistance gene markers in hop (*Humulus lupulus* L.). *Mol Breed* 33:61-74.
- Maryani, M., Bradley, G., Cahill, D., and Gehring, C. 2001. Natriuretic peptides and immunoreactants modify osmoticum-dependent volume changes in *Solanum tuberosum* L. mesophyll cell protoplasts. *Plant Sci* 161:443-452.
- Nembaware, V., Seoighe, C., Sayed, M., and Gehring, C. 2004. A plant natriuretic peptide-like gene in the bacterial pathogen *Xanthomonas axonopodis* may induce hyper-hydration in the plant host: a hypothesis of molecular mimicry. *BMC Evol Biol* 4:10.
- O'Connell, R., Herbert, C., Sreenivasaprasad, S., Khatib, M., Esquerré-Tugayé, M.-T., and Dumas, B. 2004. A novel Arabidopsis-Colletotrichum pathosystem for the molecular dissection of plant-fungal interactions. *Mol. Plant Microbe Interact.* 17:272-282.
- Oome, S., Raaymakers, T. M., Cabral, A., Samwel, S., Böhm, H., Albert, I., Nürnberger, T., and Van den Ackerveken, G. 2014. Nep1-like proteins from three kingdoms of life act as a microbe-associated molecular pattern in Arabidopsis. *Proc Natl Acad Sci U S A* 111:16955-16960.
- Pharmawati, M., Maryani, M. M., Nikolakopoulos, T., Gehring, C. A., and Irving, H. R. 2001. Cyclic GMP modulates stomatal opening induced by natriuretic peptides and immunoreactive analogues. *Plant Physiol Biochem* 39:385-394.
- Rovenich, H., Boshoven, J. C., and Thomma, B. P. H. J. 2014. Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. *Curr Opin Plant Biol* 20:96-103.
- Santhanam, P. 2012. Random insertional mutagenesis in fungal genomes to identify virulence factors. Pages 509-517 in: *Plant fungal pathogens. Methods in molecular biology*, vol. 835. T. B. Bolton MD, ed. Humana press, Totowa.
- Schmidt, S. M., Houterman, P. M., Schreiber, I., Ma, L., Amyotte, S., Chellappan, B., Boeren, S., Takken, F. L., and Rep, M. 2013. MITEs in the promoters of effector genes allow prediction of novel virulence genes in *Fusarium oxysporum*. *BMC Genomics* 14:119.
- Sherriff, C., Whelan, M. J., Arnold, G. M., Lafay, J.-F., Brygoo, Y., and Bailey, J. A. 1994. Ribosomal DNA sequence analysis reveals new species groupings in the genus *Colletotrichum*. *Exp Mycol* 18:121-138.
- Song, Y., Zhang, Z., Seidl, M. F., Majer, A., Jakse, J., Javornik, B., and Thomma, B. P. 2017a. Broad taxonomic characterization of *Verticillium* wilt resistance genes reveals ancient origin of the tomato Ve1 immune receptor. *Mol Plant Pathol* 18:195-209.
- Stergiopoulos, I., van den Burg, H. A., Ökmen, B., Beenen, H. G., van Liere, S., Kema, G. H., and de Wit, P. J. G. M. 2010. Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. *Proc Natl Acad Sci U S A* 107:7610-7615.
- Thomma, B. P. H. J., Nürnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- Vining, K., and Davis, T. 2009. Isolation of a Ve homolog, *mVe1*, and its relationship to *Verticillium* wilt resistance in *Mentha longifolia* (L.) Huds. *Mol Genet Genomics* 282:173-184.
- Zhang, Y., Wang, X., Yang, S., Chi, J., Zhang, G., and Ma, Z. 2011. Cloning and characterization of a *Verticillium* wilt resistance gene from *Gossypium barbadense* and functional analysis in *Arabidopsis thaliana*. *Plant Cell Rep* 30:2085-2096.




## Supplemental data

SUPPLEMENTAL TABLE 1 | Primers used in this study


Primer	Sequence (5'- 3')	Notes
SlRub_QPCR_F	GAACAGTTTCTCACTGTTGAC	<i>S. lycopersicum Rubisco</i>
SlRub_QPCR_R	CGTGAGAACCATAAAGTCACC	<i>S. lycopersicum Rubisco</i>
FolSIX1_F	GTCTCACGAGCCAAGTCTACC	<i>F. oxysporum Six1</i>
FolSIX1_R	GAACCGCAGCCTCTTGAGCAT	<i>F. oxysporum Six1</i>
FolTub_F	CTCTGGCAACAAGTATGTTCCC	<i>F. oxysporum Tubulin</i>
FolTub_R	TTGTGCGGGACGGAAGAGCTGA	<i>F. oxysporum Tubulin</i>
FolAve1_QPCR_F3	ATATCGGAAGTCAAAATATTCTCAAC	<i>F. oxysporum Ave1</i>
FolAve1_QPCR_R3	CTTATACATTTTCATCGTATACAGTCTGC	<i>F. oxysporum Ave1</i>
AtRub_QPCR_F	GCAAGTGTGGGTTCAAAGCTGGT	<i>A. thaliana Rubisco</i>
AtRub_QPCR_R	CCAGGTTGAGGAGTACTCGGAATGCTG	<i>A. thaliana Rubisco</i>
ChELF1a_F	CTGGTACAAGGGTTGGGAGA	<i>C. higginsianum Elongation factor</i>
ChELF1a_R	ACCGCCGATCTGTAGACAT	<i>C. higginsianum Elongation factor</i>
ChAve1_QPCR_F5	CAAGATGCTATGGCAACAATATGAAC	<i>C. higginsianum Ave1</i>
ChAve1_QPCR_R5	GTCCTTGAGGAAAATCTATCGTATTTCTG	<i>C. higginsianum Ave1</i>
BvAct_QPCR_F	GATTTGGCACCACACCTTCT	<i>B. vulgaris actin</i>
BvAct_QPCR_R	TCTTTTCCCTGTTTGCCTTG	<i>B. vulgaris actin</i>
CbAct_QPCR_F	ACATGGCTGGTCTGATTTG	<i>C. beticola actin</i>
CbAct_QPCR_R	TGTCGGTCAGGAAGCTCGTA	<i>C. beticola actin</i>
CbAve1_QPCR_F	ATTCCTTCAGGCAACTCT	<i>C. beticola Ave1</i>
CbAve1_QPCR_R	CGGACAAGCTTCGCAATAAT	<i>C. beticola Ave1</i>
VdGapdh_F	CGAGTCCACTGGTGTCTTCA	<i>V. dahliae GAPDH</i>
VdGapdh_R	CCCTCAACGATGGTGAACCT	<i>V. dahliae GAPDH</i>
VdAve1-Fw5	ATCCTACTATAACCCACCTACCTTC	<i>V. dahliae Ave1</i>
VdAve1-Rv5	CATCATATGAGTCTGAGATAAGATCA	<i>V. dahliae Ave1</i>
XacAve1_QPCR_F	GCAATCGGTTTGCTCTTTTC	<i>X. axonopodis XacPNP</i>
XacAve1_QPCR_R	AGCACCGTTATCCACAGAC	<i>X. axonopodis XacPNP</i>
VvAve1-QPCR-F2	CGTTATAGGTTAAGGTGCCTGAGT	<i>V. vinifera Ave1</i>
VvAve1-QPCR-R2	TTCATGCTAGGAGAGTGTGAAATG	<i>V. vinifera Ave1</i>
SlAve1_QPCR_F	CGTCGGGGAATCTATTTGTG	<i>S. lycopersicum Ave1</i>
SlAve1_QPCR_R	AAAGCATCCGTTGACAAAGC	<i>S. lycopersicum Ave1</i>
FolAve1_LB_F3	GGTCTTAAUAAACCTAACCTGTATCTAGACCAGAGTGTCT	Left border <i>FolAve1</i>
FolAve1_LB_R3	GGCATTAAUCAGCAGGTGTTTAGGTATTGTTAAGATAAG	Left border <i>FolAve1</i>
FolAve1_RB_F	GGACTTAAUGTCTTAGAGCTTAAGGTCAATGTAGC	Right border <i>FolAve1</i>
FolAve1_RB_R	GGGTTTAAUUCTGCAGCTTTCACGGGGCTA	Right border <i>FolAve1</i>
ChAve1_LB_F	GGTCTTAAUUGGCAGGAGGTCTAGGTGAGA	Left border <i>ChAve1</i>
ChAve1_LB_R	GGCATTAAUCGACTTGAGTTTTGCTGCAC	Left border <i>ChAve1</i>
ChAve1_RB_F	GGACTTAAUAAAGATTTTCGCAGTGCTTCA	Right border <i>ChAve1</i>
ChAve1_RB_R	GGGTTTAAUAGGCTGCAGAATGAGTTTCG	Right border <i>ChAve1</i>
pRF-HU2_Hyg_Fw	CTATTCCTTGGCCCTCGACGAGTGC	Hygromycin primers
pRF-HU2_Hyg_Rv	GATTGCTAGGGCGGTGGATATGTCC	Hygromycin primers
NAT_Fw	CGGGCCGATTGGTCAAGATTTCG	Nourseothricin primers
NAT_Rv	CGATTGCTCGTCCGATTCTGTCG	Nourseothricin primers



Primer	Sequence (5'- 3')	Notes
MDB-277	GACGTTGATAAACGACGGCCAGTG	Split-marker HYG-F
MDB-258	GGATGCCTCCGCTCGAAGTA	Split-marker HY-R
MDB-259	CGTTGCAAGACCTGCCTGAA	Split-marker YG-F
MDB-278	CACAGGAAACAGCTATGACCATGA	Split-marker HYG-R
MDB-1145	GGCAGGTAGATGACGACCAT	HYG R2
CbAve1 1F	CAAGATTGGGCCTTCGTATG	CbAve1 split-marker left
CbAve1 2R	CACTGGCCGTCGTTTTACACAGTCTCTTCAATGGATCC GGACTG	CbAve1 split-marker left border
CbAve1 3F	TCATGGTCATAGCTGTTTCCTGTGTGAGTGGGGTTTT GGTTTC	CbAve1 split-marker right border
CbAve1 4R	AAGCATACCTCTTCGGCAAA	CbAve1 split-marker right
MDB-760	CAACGCTGCCACAAGCTGCT	CbAve1 reverse




**SUPPLEMENTAL FIGURE 1 | Verification of deletion strains.** (A) Identification of *F. oxysporum* *FoAve1* deletion strains by amplification of *FoAve1* and selection marker nourseothricin (NAT) from DNA obtained from transformants. DNA bands were observed at the expected size. Asterisks indicate positive transformants, while transformants that lack an asterisk are ectopic transformants. (B) Identification of *C. higginsianum* *ChAve1* deletion strains by amplification of *ChAve1* and selection marker hygromycin (HYG) from DNA obtained from *C. higginsianum* transformants. DNA bands were observed at the expected size. Asterisks indicate positive transformants, while transformants that lack an asterisk are ectopic transformants. (C) Verification of *CbAve1* deletion strains (1 and 2) based on absence of *CbAve1* using gene-specific primers and presence of the hygromycin resistance cassette. Amplification on *C. beticola* Wt gDNA (3) as a control showing the presence of *CbAve1* and absence of the hygromycin resistance cassette at the expected size. Asterisks indicate positive transformants.



**SUPPLEMENTAL FIGURE 2 | VdAve1 acts in a different manner than Ave1 homologs derived from plant pathogens.** *V. dahliae* *VdAve1* deletion strain ( $\Delta VdAve1$ ) complemented with *Ave1* homologs derived from *V. dahliae* (*VdAve1*), tomato (*S. lycopersicum*; *SlAve1*), grape (*V. vinifera*; *VvAve1*), *F. oxysporum* (*FoAve1*), *C. beticola* (*CbAve1*), *C. higginsianum* (*ChAve1*) and *Xanthomonas axonopodis* (*XacPNP*) inoculated on tomato plants. (A) Expression of *VdAve1*, *SlAve1*, *VvAve1*, *FoAve1*, *CbAve1*, *ChAve1* and *XacPNP* in a minimum of 2 transformants. (B) Canopy area of tomato plants inoculated with *V. dahliae* strains expressing *Ave1* homologs in a minimum of 2 transformants compared to *V. dahliae* wild type (*Vd wt*) at 14 dpi. (C) Fungal colonization of tomato plants inoculated with *V. dahliae* strains expressing *Ave1* homologs compared to *V. dahliae* wild type (*Vd wt*) at 14 dpi. Different letter labels indicate significant differences ( $P < 0.05$ ). (D) Photos taken from the side and the top of tomato plants inoculated with *V. dahliae* strains expressing *Ave1* homologs compared to *V. dahliae* wild type (*Vd wt*) at 14 dpi.





**A nine amino acid peptide of  
*Verticillium dahliae* effector Ave1  
is required but not sufficient  
for recognition by tomato immune  
receptor Ve1**

**Chapter 4**

Zhao Zhang<sup>#</sup>

Yin Song<sup>#</sup>

Jordi C. Boshoven<sup>#</sup>

Hanna Rovenich

Michael F. Seidl

Jernej Jakše

Karunakaran Maruthachalam

Chun-Ming Liu

Krishna V. Subbarao

Branka Javornik

Bart P.H.J. Thomma

<sup>#</sup>These authors contributed equally

A modified version is available online at bioRxiv preprint server  
doi: <https://doi.org/10.1101/103473>

## Abstract

Effectors are molecules secreted by plant pathogens to facilitate infection, often through deregulation of plant immune responses. During infection, race 1 strains of the soil borne vascular wilt fungus *Verticillium dahliae* secrete the effector protein Ave1 that is recognized by tomato immune receptor Ve1. Homologs of *V. dahliae* Ave1 (VdAve1), have been identified from the bacterial pathogen *Xanthomonas axonopodis* pv. *citri* (XacPNP), and in the fungal pathogens *Fusarium oxysporum* (FoAve1), *Colletotrichum higginsianum* (ChAve1) and *Cercospora beticola* (CbAve1). In this study, we identified a novel Ave1 homolog from *V. nubilum* (VnAve1), which is mainly known as saprophyte and opportunistic pathogen. The Ave1 homologs are differentially recognized by Ve1, with the most efficient recognition of VdAve1 and VnAve1, and little to no recognition of ChAve1 and XacPNP. Interestingly, C-terminal GFP-tagging of VdAve1 resulted in loss of recognition by Ve1, suggesting that exposure of the C-terminus of VdAve1 is essential for recognition by Ve1. Truncations of VdAve1 and domain swaps between VdAve1 and XacPNP narrowed down the sequence that is required for Ve1 recognition to the C-terminal nine amino acids. Site-directed mutagenesis was performed to analyse the contribution of individual amino acid residues to recognition by Ve1, revealing the importance of five amino acids from Val127 to Phe131. Nevertheless, fusions of the VdAve1 C-terminal nine amino acids to GFP or the *Cladosporium fulvum* effector Avr9 showed that this sequence is not sufficient to activate Ve1-mediated immunity. Finally, with a complementation experiment in *V. dahliae* we confirmed that the C-terminal nine amino acids of Ave1 are required but not sufficient to activate Ve1-mediated immunity.

## Introduction

In nature, plants are threatened by diverse groups of potential microbial pathogens. However, only few of these potential pathogens actually cause disease, as plants have developed innate immunity to protect themselves against microbial attack (Dodds and Rathjen 2010; Thomma et al. 2011). In its simplest form, plant immunity against pathogen attack is governed by immune receptors that sense pathogen-derived ligands to activate defense. Originally, the interaction between plant immune receptors and pathogen ligands was described in the “gene-for-gene” model, stating that the products of plant resistance (*R*) genes induce race-specific resistance upon recognition of the products of corresponding pathogen avirulence (*Avr*) genes (Flor 1971). Decades later, an updated view of plant innate immunity has been introduced as the “zigzag” model, by incorporating pathogen-secreted effector molecules that suppress host immune responses, but that may subsequently be recognized by newly evolved immune receptors, in turn (Jones and Dangl 2006). In this model, the first line of defense is governed by plasma membrane-localized pattern recognition receptors (PRRs) that detect conserved microbe-associated molecular patterns (MAMPs) to activate MAMP-triggered immunity (MTI). In subsequent layers of defense effectors are recognized by corresponding resistance proteins (*R* proteins), resulting in effector-triggered immunity (ETI). Although initially portrayed as separate layers of defense, numerous studies on various plant-microbe interactions have revealed that the delineation between MTI and ETI is not strict, but rather a continuum (Thomma et al. 2011). Moreover, the conceptual conflict that MAMPs are defined from the perspective of the host whereas effectors are defined from the perspective of the invader creates a conceptual conflict and has recently inspired the formulation of the Invasion Model, in which host receptors (termed invasion pattern receptors; IPRs) detect either an externally encoded or modified-self ligand that betrays invasion (termed invasion patterns; IPs) (Cook et al. 2015). In this model, any molecule can serve as an IP that is detected by an IPR, but the probability of a particular ligand-receptor complex to evolve within the framework of host immunity increases with increasing ligand probability to retain function, conservation across organisms, importance in establishment of symbiosis, and accessibility (Cook et al. 2015).

*Verticillium dahliae* is a xylem invading fungal pathogen that causes *Verticillium* wilt diseases in a wide range of plant species worldwide (Fradin and Thomma 2006). *V. dahliae* persists in the soil and enters plants through their roots. Once inside the root, the fungus grows intercellularly and invades the xylem vessels, where it sporulates to spread through the vascular system. Typical symptoms of *V. dahliae* infection include stunting, wilting, chlorosis, and necrosis (Fradin and Thomma 2006). In tomato, a single dominant locus that confers *Verticillium* resistance has

been identified that controls isolates that are assigned to race 1 (Schaible et al. 1951). The locus comprises two genes, *Ve1* and *Ve2*, that both encode extracellular leucine rich repeat (eLRR) receptor-like proteins (RLPs), and of which only *Ve1* acts as a functional *Verticillium* resistance gene in tomato (Fradin et al. 2011; Fradin et al. 2009; Kawchuk et al. 2001; Wang et al. 2010). Functional homologs of *Ve1* were reported from *Nicotina glutinosa* and from cotton (Chen et al. 2016; Yang et al. 2014; Zhang et al. 2012; Zhang et al. 2011; Zhang et al. 2013).

Through comparative population genomics, the *V. dahliae* effector protein that is recognized by the *Ve1* immune receptor of tomato was identified as *Ave1* (for **A**virulence on **Ve1** tomato 1) (de Jonge et al. 2012; Fradin et al. 2009). Although the intrinsic function of *V. dahliae* *Ave1* (*VdAve1*) remains elusive, it is clear that *Ave1* contributes to fungal virulence on susceptible plant genotypes (de Jonge et al. 2012). Interestingly, *Ave1* homologs were identified from a number of fungal pathogens, including the tomato pathogen *Fusarium oxysporum* f. sp. *lycopersici* (*FoAve1*), the sugar beet pathogen *Cercospora beticola* (*CbAve1*) and the Brassicaceae pathogen *Colletotrichum higginsianum* (*ChAve1*). Strikingly, however, most *Ave1* homologs were found in plants, with the most closely related homologs derived from tomato (*Solanum lycopersicum*; *SIPNP*) and grape (*Vitis vinifera*; *VvAve1*). Finally, a more distantly related homolog was identified in the plant pathogenic bacterium *Xanthomonas axonopodis* pv. *citri* (*XacPNP*) (de Jonge et al. 2012; Nembaware et al. 2004). Co-expression of *SIPNP*, *FoAve1*, and *CbAve1* with tomato *Ve1* in tobacco triggers a hypersensitive response (HR), whereas co-expression of *ChAve1* with tomato *Ve1* did not lead to an HR (de Jonge et al. 2012). Consequently, *Ve1* was found to mediate resistance towards *F. oxysporum* in tomato, demonstrating involvement of this tomato immune receptor in resistance against multiple fungal pathogens (de Jonge et al. 2012).

It has previously been demonstrated that eLRR-containing cell-surface immune receptors recognize peptide sequences as epitopes of their pathogen ligands. For example, *flg22* is the 22 amino acid peptide derived from bacterial flagellin that is perceived by the RLK-type immune receptor *FLS2* (Zipfel et al. 2004), while the Arabidopsis *EFR* RLK-type immune receptor was shown to recognize *elf18*, an 18 amino acid peptide derived from bacterial EF-Tu (Zipfel et al. 2006). Similarly, a surface-exposed pentapeptide *TKLGE* of the 22 kDa ethylene-inducing xylanase (*EIX*) from the biocontrol fungus *Trichoderma viride* determines recognition by the tomato RLP-type receptor *LeEIX2* (Ron and Avni 2004; Rotblat et al. 2002). Furthermore, the tyrosine-sulfated 21 amino acid sequence *RaxX21-sY* present in *Xanthomonas oryzae* pv. *oryzae* *RaxX* is sufficient to activate *XA21*-mediated immunity (Pruitt et al. 2015). Finally, a conserved 20-24 amino acid sequence derived from *Nep1*-like proteins (NLPs) is sufficient to activate *RLP23*-mediated immunity in

*Arabidopsis* (Albert et al. 2015; Böhm et al. 2014; Oome et al. 2014). In the present study, we aimed to identify a minimal motif in VdAve1 that determines recognition by tomato immune receptor Ve1. Our approach was based on alignment of differentially recognized Ave1 homologs, followed by a combination of deletions, domain swaps among Ave1 homologs and mutagenesis.

## Results


### Conservation among Ave1 homologs

Previously, we reported the cloning of Ave1 from *V. dahliae* and described the absence of allelic variation among 85 Ave1 alleles of *Verticillium* strains (de Jonge et al. 2012). The Ave1 alleles were derived from *V. dahliae* and *V. albo-atrum* (presently *V. alfalfae*; (Inderbitzin et al. 2011)), whereas Ave1 alleles were not identified in any of the race 2 strains analysed, nor in *V. dahliae* and *V. albo-atrum* strains that are not pathogenic on tomato, nor in *V. longisporum* or *V. tricorpus* (de Jonge et al. 2012). To further explore Ave1 diversity, we assessed its presence in a collection of 97 *Verticillium* strains isolated from various host plants and different geographical locations, resulting in the identification of 17 novel Ave1 alleles (Supplemental Table 1). No allelic variation was found among the newly identified Ave1 alleles amplified from *V. dahliae* as well as from *V. alfalfae* and *V. nonalfalfae*, two novel species that have recently been separated from *V. albo-atrum* (Inderbitzin et al. 2011). However, an Ave1 homolog was identified in four isolates of *V. nubilum* (VnAve1), a species that is mainly known as a saprophytic and opportunistic pathogen (Isaac 1953). While the predicted VnAve1 protein sequence displays 13 amino acid polymorphisms when compared with VdAve1 (Figure 1A), the four VnAve1 alleles were found to be identical to each other.

Alignment of the amino acid sequences of VdAve1 with homologs from plants (SlAve1 and VvAve1), and plant pathogens (VnAve1, FoAve1, CbAve1, ChAve1 and XacPNP) reveals blocks of highly conserved amino acids that are alternated with more variable regions (Figure 1A). Based on prediction by SignalP 4.0 (Petersen et al. 2011), all Ave1 homologs contain N-terminal signal peptides that direct secretion of the molecules into the extracellular space (Figure 1A; D-cutoff score>0.6). Moreover, the four cysteine residues that are present in VdAve1 are conserved among all homologous proteins (Figure 1A), and *in silico* analysis using DISULFIND (Ceroni et al. 2006) suggests the formation of disulphide bridges between Cys35 and Cys63, as well as between Cys71 and Cys79. From the alignment it is apparent that XacPNP is the most divergent, while all other homologs are relatively comparable (Figure 1A).



**FIGURE 1 | Comparison of necrosis induced by Ave1 homologs through co-expression with Ve1 in *Nicotiana tabacum*.** (A) Amino acid sequence alignment of Ave1 homologs from *Verticillium dahliae* (VdAve1), *V. nubilum* (VnAve1), *Solanum lycopersicum* (SlAve1), *Vitis vinifera* (VvAve1), *Fusarium oxysporum* f. sp. *lycopersici* (FoAve1), *Cercospora beticola* (CbAve1), *Colletotrichum higginsianum* (ChAve1), and *Xanthomonas axonopodis* pv. citri (XacPNP). Purple shading indicates identical amino acids while the color intensity represents the frequency. Asterisks indicate the 13 amino acid polymorphisms between VdAve1 and VnAve1. The positions of four conserved cysteine residues are indicated with arrows in the bottom. Predicted signal peptides are framed. (B) Coexpression of tomato Ve1 and Ave1 homologs VdAve1, VnAve1, SlAve1, VvAve1, FoAve1, CbAve1, ChAve1, and XacPNP in *N. tabacum*. Expression of the Avr9 effector from the tomato leaf mold fungus *Cladosporium fulvum* in combination with Ve1 is shown as negative control. Pictures were taken at five days post infiltration. (C) Quantification of cell death resulting from recognition of Ave1 homologs by Ve1. The average percentage of necrotic leaf area of infiltration zones at five days post infiltration (n=24). Data are presented as mean with standard errors.




### Comparison of necrosis-inducing activity of Ave1 homologs

It was previously demonstrated that Ve1 recognizes not only VdAve1, but also SIPNP, FoAve1 and CbAve1 (de Jonge et al. 2012). We now also tested the HR-inducing capacity of VnAve1, VvAve1 and XacPNP that were isolated from *V. nubilum*, *V. vinifera* and *X. axonopodis*, respectively. Co-expression of the sequence-unrelated effector Avr9 from the tomato leaf mould fungus *Cladosporium fulvum* (van Kan et al. 1991) in combination with Ve1 served as a negative control. Whereas expression of VnAve1 or VvAve1 together with Ve1 in *Nicotiana tabacum* resulted in strong HR, co-expression of XacPNP or Avr9 with Ve1 triggered little to no necrosis in addition to the small wounds that were caused by the infiltration procedure (Figure 1B). To compare the HR induced by the various Ave1 homologs, they were co-expressed with Ve1 in *N. tabacum* and HR development was measured at five days post infiltration by quantification of the leaf area that developed necrosis (Zhang et al. 2013). Importantly, none of the Ave1 homologs induced necrosis in absence of Ve1 (Figure 1B). Whereas agroinfiltration of either VdAve1 or VnAve1 with Ve1 resulted in complete necrosis of the infiltrated leaf area, agroinfiltration of FoAve1 with Ve1 resulted in large necrotic spots in the infiltrated leaf area, although no complete collapse of the infiltrated area was observed. Upon agroinfiltration of CbAve1 with Ve1, spreading of smaller and larger necrotic spots was observed in all infiltrated areas, but the infiltrated leaf area did not turn completely necrotic. For ChAve1, XacPNP and Avr9, no spreading necrosis was observed beyond the wounded infiltration sites (Figure 1B). Upon agroinfiltration of the tomato and grape homologs, SlAve1 and VvAve1, most of the infiltrated leaf area developed necrosis, occasionally affecting the complete infiltrated leaf sector. To confirm that reduced levels of HR are not due to instability of the proteins in the expression assay, GFP-tagged Ave1 homologs were detected by immunoblotting. Similar to GFP-tagged VdAve1 protein or GFP-tagged VnAve1 that induce the strongest HR, all other GFP-tagged Ave1 homologs accumulated to clearly detectable protein levels (Supplemental Figure 1).

### The C-terminus of VdAve1 is required for recognition by Ve1

In order to permit functional analyses, a construct encoding C-terminal protein GFP fusion of VdAve1 was generated. However, C-terminal fusion of a GFP tag to Ave1 resulted in loss of recognition by Ve1. Considering that the GFP tag is relatively large, we engineered C-terminal fusions to VdAve1 with smaller protein tags. Nevertheless, all C-terminal tags abolished, or significantly reduced, HR development (Figure 2A; C). Importantly, the C-terminal GFP fusions could be detected by immunoblotting (Supplemental Figure 2), suggesting that accessibility of the VdAve1 C-terminus is important for recognition by Ve1.




**FIGURE 2 | C-terminal fusion of a GFP tag to VdAve1 results in loss of recognition by Ve1.** (A) The signal peptide of VdAve1 (SP) directs secretion of untagged VdAve1 (VdAve1), C-terminally tagged VdAve1 (VdAve1\_GFP, VdAve1\_HA, VdAve1\_HIS, VdAve1\_4MYC, VdAve1\_10MYC, VdAve1\_FLAG) into the extracellular space. The constructs were co-expressed with Ve1 in *N. tabacum* upon agroinfiltration and the occurrence of the hypersensitive response was monitored. (B) The signal peptide of VdAve1 (SP) directs secretion of N-terminally tagged VdAve1 (GFP\_VdAve1, HA\_VdAve1 and MYC\_VdAve1) into the extracellular space. The constructs were co-expressed with Ve1 in *N. tabacum* upon agroinfiltration and the occurrence of the hypersensitive response was monitored. All pictures were taken at five days post infiltration. (C) Quantification of cell death resulting from recognition of tagged VdAve1 proteins by Ve1. The graph shows the average percentage of necrotic leaf area of infiltration zones at five days post infiltration ( $n > 5$ ). Data are presented as mean with standard deviations.

To further investigate the role of the C-terminus in recognition of VdAve1 by Ve1, a number of C-terminal truncations was generated. Deletion of 42 amino acids of the C-terminus, from Lys93 to Ile134 (VdAve1\_M1), resulted in loss of recognition by

Ve1 (Figure 3). Subsequent analysis of step-wise smaller truncations revealed that already a C-terminal deletion of nine amino acids (VdAve1\_M4) resulted in loss of Ve1 recognition (Figure 3), despite the expression of stable protein (Supplemental Figure 2).

We subsequently performed complementation experiments in *V. dahliae* to confirm the importance of the C-terminus for immune activation. To this end we expressed the VdAve1\_M4 construct (Figure 3) in a *V. dahliae* VdAve1 deletion strain and inoculated Ve1 tomato with this transgenic fungus. Plants that were inoculated with *V. dahliae* expressing this VdAve1 construct showed a similar disease phenotype as plants inoculated with a VdAve1 deletion strain, whereas plants inoculated with wild type *V. dahliae* and the VdAve1 complementation strain resembled mock-inoculated plants (Figure 4A; B). Collectively, our results show that the C-terminal nine amino acids of VdAve1 are required to establish Ve1-mediated immunity.




**FIGURE 3 | Deletion of the C-terminal nine amino acids from VdAve1 results in loss of recognition by Ve1.** Occurrence of the hypersensitive response in *N. tabacum* upon co-expression of VdAve1 truncations with Ve1. Constructs encoding full length VdAve1 with its native signal peptide (SP) and four truncations that lack the C-terminal 42 (VdAve1\_M1), 32 (VdAve1\_M2), 18 (VdAve1\_M3) and 9 (VdAve1\_M4) amino acids of VdAve1 were tested. Pictures were taken at five days post infiltration.


### Ave1 recognition cannot be assigned to a single amino acid residue

Since the C-terminal nine amino acids appear to be essential for VdAve1 recognition, and the bacterial homolog XacPNP that is significantly divergent in this region (Figure 1A) is not recognized by Ve1 (Figure 1B and C), an expression construct for a chimeric Ave1 protein was engineered in which the C-terminal nine amino acids of VdAve1 were replaced by those of XacPNP (Vd\_Xac9AA; Figure 5A; D). As expected, co-expression of the Vd\_Xac9AA chimera with Ve1 in *N. tabacum* resulted in absence of recognition, as only small necrotic spots were observed at

most (Figure 5A). Conversely, a chimeric Xac\_Vd9AA protein was constructed, in which the last nine amino acids of XacPNP were replaced by those of VdAve1. Co-expression of Xac\_Vd9AA with Ve1 in *N. tabacum* resulted in a relatively strong HR, although full necrosis was not observed in the infiltrated leaf area (Figure 5A; D). With immunodetection the stability of the chimeric Vd\_Xac9AA and Xac\_Vd9AA proteins was confirmed (Supplemental Figure 3).



**FIGURE 4 | Complementation experiments in *V. dahliae* confirm that the C-terminal nine amino acids are required to activate Ve1 immunity in tomato.** (A) *V. dahliae* *Ave1* deletion strains expressing a construct encoding VdAve1 lacking the C-terminal 9 amino acids (VdAve1\_M4) are unable to activate Ve1 immunity in tomato when compared with *V. dahliae* wild-type (Vd wt) and genetic complementation (VdAve1). (B) Quantification of the tomato canopy area ( $n=8$ ). Data are presented as mean with standard deviations. Different letters indicate significant differences ( $P<0.05$ ; One-Way ANOVA). (C) *V. dahliae* *Ave1* deletion strains expressing a construct that encodes the C-terminal 9 amino acids fused to GFP (SP-GFP-9AA) are unable to activate Ve1 immunity in tomato compared to *V. dahliae* wild-type (Vd wt) and genetic complementation (VdAve1). (D) Quantification of the tomato canopy area ( $n=8$ ). Data are presented as mean with standard deviations. Different letters indicate significant differences ( $P<0.05$ ; One-Way ANOVA).




**FIGURE 5 | Domain swaps reveal the importance of the C-terminal nine amino acids of Ave1 homologs for recognition by Ve1.** (A) Occurrence of the hypersensitive response in *N. tabacum* upon co-expression of Ave1 chimeras with Ve1. Constructs encoding full length VdAve1 and XacPNP with their native signal peptides (SP) and two chimeras in which their C-terminal nine amino acids were exchanged were tested. In chimera Vd\_Xac9AA the C-terminal nine amino acids of VdAve1 (Vd9AA) were replaced with those of XacPNP (Xac9AA), while in chimera Xac\_Vd9AA Xac9AA was replaced by Vd9AA. (B) Construct SP\_9AA encodes the C-terminal nine amino acids of VdAve1 (Vd9AA) fused to the VdAve1 signal peptide (SP). SP\_GFP\_Vd9AA encodes GFP that is C-terminally tagged with Vd9AA, and N-terminally fused to the VdAve1 signal peptide (SP) to establish extracellular targeting, while SP\_GFP lacks the C-terminal Vd9AA fusion. Furthermore, a chemically synthesized peptide encompassing Vd9AA (AVNIEFRQI) was used. (C) Constructs encoding full length VdAve1 and ChAve1 with their native signal peptides (SP) and two chimeras in which their C-terminal 9 amino acids were exchanged were tested. In chimera Vd\_Ch9AA the C-terminal nine amino acids of VdAve1 (Vd9AA) were replaced with those of ChAve1 (Ch9AA), while in chimera Ch\_Vd9AA Ch9AA was replaced by Vd9AA. All pictures were taken at five days post infiltration. (D) Quantification of cell death resulting from recognition of chimera proteins by Ve1. The average percentage of necrotic leaf area of infiltration zones at five days post infiltration ( $n > 10$ ). Data are presented as mean with standard deviation.

To further unravel how the C-terminal nine amino acids contribute to recognition of VdAve1, we performed site-directed mutagenesis. Since the first of the nine residues (Ala126) is shared with XacPNP, we conclude that this one likely does not contribute to VdAve1 recognition. Thus, eight alanine scanning constructs were engineered in which the remaining residues were substituted by alanine (V127A, N128A, I129A, E130A, F131A, R132A, Q133A, and I134A, respectively; Figure 6A; B). The mutant constructs were tested in *N. tabacum* upon co-expression with Ve1, and HR development was measured at five days post infiltration by quantification of the leaf area that developed necrosis (Figure 6C). Similar to VdAve1, mutants R132A and Q133A activated strong Ve1-mediated HR, as >90% of the infiltrated leaf area developed necrosis (Figure 6C). In contrast, mutants N128A, I129A, E130A and F131A triggered clearly reduced necrosis, with <70% of the infiltrated leaf area developing necrosis (Figure 6C). Alanine substitution of Val127 and Ile134 (V127A and I134A, respectively) resulted in an intermediate reduction of necrosis-inducing activity, with 70-80% of the infiltrated leaf area developing necrosis (Figure 6C).

4 To further investigate the contribution of individual amino acids to recognition by Ve1, we engineered five additional mutants in which individual residues of the VdAve1 C-terminal nine amino acids were substituted by corresponding residues from XacPNP (V127R, N128T, E130P, F131W, and Q133N; Figure 6A; B). Whereas expression of F131W and Q133N induced similar necrosis as wild type VdAve1, mutants N128T and E130P exhibited significantly compromised HR upon co-expression with Ve1, with <60% of the infiltrated leaf area developing necrosis (Figure 6C). Mutant V127R did not induce HR at all in the presence of Ve1 (Figure 6C). Importantly, all site-directed mutants that showed compromised HR-inducing capacity accumulated to similar levels as VdAve1 or XacPNP (Supplemental Figure 4). These observations suggest that the region from Val127 to Phe131 is critical for recognition by Ve1. Collectively, we conclude that the C-terminal nine amino acids of VdAve1 are required for recognition by Ve1, and that recognition cannot be assigned to a single amino acid residue.





**FIGURE 6 | Contribution of individual amino acid residues to recognition of the C-terminal nine amino acids of VdAve1.** (A) Amino acid sequence alignment of the C-terminal nine amino acids of various Ave1 homologs. Identical and highly similar residues are indicated with black shading whereas weakly similar residues are indicated with grey shading. (B) Schematic representation of mutant proteins V127A, N128A, I129A, E130A, F131A, R132A, Q133A, I134A, V127R, N128T, E130P, F131W, and Q133N, respectively. Bold characters indicate the mutated residues. (C) Co-expression of tomato *Ve1* and *VdAve1* mutants in *N. tabacum*. The quantification of cell death resulting from recognition of *VdAve1* mutants by *Ve1* is shown as the average percentage of necrotic leaf area of infiltration zones at five days post infiltration (n>5). Data are presented as mean with standard errors.

### The C-terminal nine amino acids of VdAve1 are not sufficient for recognition by Ve1

To determine whether the C-terminal nine amino acids are sufficient to trigger *Ve1*-mediated recognition, we generated a construct encoding the C-terminal nine amino acids of *VdAve1* fused to the *VdAve1* signal peptide (SP\_Vd9AA) (Figure 5B; D). This construct was co-expressed with *Ve1* in *N. tabacum*, but no necrosis was observed in the infiltrated leaf (Figure 5B). Furthermore, also infiltration of a chemically synthesized peptide encompassing the C-terminal nine amino acids of *VdAve1* was not able to trigger an HR in *Ve1*-expressing tobacco up to a



concentration of 1 mg/mL (Figure 5B; D). We speculated that the nine amino acid peptide may not be stable upon secretion in the apoplast, irrespective whether it is *in planta* expressed or injected upon chemical synthesis. In an attempt to overcome such complication, we generated constructs in which the coding sequence of GFP or Avr9 was N-terminally fused to the VdAve1 signal peptide and C-terminally fused to the C-terminal nine amino acids of VdAve1 (SP\_GFP\_Vd9AA and SP-Avr9\_Vd9AA; Figure 5B; D). As negative controls, we generated constructs in which the coding sequence of GFP or Avr9 was N-terminally fused to the VdAve1 signal peptide without the C-terminal nine amino acids of VdAve1 (SP\_GFP and SP\_Avr9; Figure 5B; D). All constructs were co-expressed with *Ve1* in *N. tabacum* and HR development was monitored at five days post infiltration. However, only slight necrosis was found in the infiltrated sector despite the observation that the chimeric GFP proteins could be detected upon immunoblotting (Supplemental Figure 2).


Subsequently, a complementation assay was performed in *V. dahliae* to confirm that the C-terminal nine amino acids are not sufficient to activate immunity. To this end we expressed the SP\_GFP\_Vd9AA construct in the *Ave1* deletion strain of *V. dahliae* (Figure 4C; D) and inoculated *Ve1* tomato. Plants inoculated with strains expressing SP\_GFP\_Vd9AA showed similar symptoms as plants inoculated with the *VdAve1* deletion strain, whereas plants inoculated with wild type *V. dahliae* and the *VdAve1* complementation strain displayed similar symptoms as mock-inoculated plants (Figure 4A; D). Collectively, our results show that the C-terminal nine amino acids of *VdAve1* are required but not sufficient to establish *Ve1*-mediated immunity.

### **Evidence for contribution to recognition in addition to the C-terminal nine amino acids**

ChAve1 is closely related to VdAve1 as six of the nine C-terminal amino acids are identical. Nevertheless, only weak HR is detected upon co-expression of *ChAve1* and *Ve1* in *N. tabacum* (Figure 1B; C). To further investigate recognition of the C-terminal nine amino acids of ChAve1, an expression construct encoding a chimeric Ave1 protein was engineered in which the C-terminal nine amino acids of VdAve1 were replaced by those of ChAve1 (Vd\_Ch9AA). Intriguingly, expression of Vd\_Ch9AA resulted in complete necrosis of the infiltrated leaf area upon co-expression with *Ve1* in *N. tabacum*, whereas the reciprocal swap in which the C-terminal nine amino acids of ChAve1 were replaced by those of VdAve1 (Ch\_Vd9AA) was not recognized by *Ve1* (Figure 5C; D). With immunodetection the stability of the chimeric Vd\_Ch9AA and Ch\_Vd9AA proteins was confirmed (Supplemental Figure 5). These data provide further support for the finding that the C-terminal nine amino acids of VdAve1 are not sufficient to establish *Ve1*-mediated recognition, and confirm that additional requirements for the occurrence of the HR reside outside the stretch of the C-terminal nine amino acids.

### A structural model of Ave1 suggests a contribution of both the N- and C-terminus of Ave1 to Ve1 recognition

In an attempt to get a better insight in Ave1 recognition by Ve1 we generated a structural model of Ave1. Structural comparison with the protein databank (RCSB PDB) (Rose et al. 2013) revealed the maize protein EXPB1 (PDB ID: 2HCZ) (Yennawar et al. 2006) as a potential structural analogue (TM-Score 0.84) of VdAve1. The Ave1 structural model shows that the C-terminal nine amino acids (shown in orange in Figure 7) are exposed at the surface of the Ave1 protein. The model also predicts that the N-terminus (shown in blue in Figure 7) congregates with the C-terminus, which led us to hypothesize that Ve1 potentially recognizes a patch of the Ave1 protein that includes both the N- and C-terminal. To test this hypothesis, we fused various tags to the N-terminal of Ave1, between the signal peptide and the mature peptide. Surprisingly, while N-terminal tagging of Ave1 with GFP and HA did not affect recognition, N-terminal tagging with MYC compromised recognition (Figure 2B; C). Immuno blotting confirmed that the GFP fusions were stably expressed (Supplemental Figure 2).



**FIGURE 7 | Model for the 3D structure of the *Verticillium dahliae* effector VdAve1.** The *V. dahliae* VdAve1 structure was predicted using I-TASSER (Zhang 2008). The inferred VdAve1 structure is of high quality indicated by a confidence score (C-Score) of 1.22, and displayed as a ribbon (A), and surface (B) model from the side (left) and the top (right). The C-terminal nine amino acid sequence of VdAve1 is indicated in orange, while the N-terminal eight amino acid sequence is indicated in blue.

## Discussion

### The C-terminal nine amino acids of Ave1 are required for Ve1-mediated recognition

We have previously shown that the VdAve1 effector of *V. dahliae* is recognized by the tomato immune receptor Ve1, and we identified several homologs in other species that are differentially recognized by Ve1 (Figure 1B; C) (de Jonge et al. 2012). In this study, we demonstrate that the C-terminal nine amino acids of VdAve1 are required for the recognition by Ve1, as truncations of VdAve1 lose recognition. The sequence of the C-terminal nine amino acids of the Ave1 homologs that are recognized (VdAve1, VnAve1, SIPNP, VvAve1, FoAve1, CbAve1 and also ChAve1) only share Asn128 and Gln133 that are completely conserved (Figure 6A). However, the other amino acids in this region are characterized by a high degree of conservation. A similar sequence variation in peptide epitopes, allowing substitutions by amino acids with similar chemical properties has previously been observed, for instance in flagellin, CLV3 and AtPep1 (Kondo et al. 2008; Naito et al. 2008; Pearce et al. 2008; Song et al. 2012).

To further analyse the contribution of individual amino acids in the C-terminal nine amino acids to recognition by Ve1, we performed site-directed mutagenesis on VdAve1. The sequence alignment and mutagenesis analysis suggest that the five neighbouring residues Val127, Asn128, Ile129, Glu130 and Phe131 are the major determinants of recognition of the C-terminal nine amino acids. Whereas alanine substitution of Val127 (V127A) only slightly reduced the HR-inducing capacity of VdAve1, VdAve1 cannot be recognized when Val127 is substituted by an arginine (V127R) that occurs in XacPNP on that position. Conversely, the HR-inducing capacity of VdAve1 is significantly impaired when Phe131 was replaced by the non-similar amino acids alanine (F131A), while substitution by tryptophan (F131W), an amino acid with similar chemical properties that occurs in XacPNP on that position, still triggers strong HR.

### The C-terminal nine amino acids are not sufficient to trigger Ve1-mediated recognition

To test whether the C-terminal nine amino acids are sufficient to trigger Ve1-mediated HR a synthesized peptide and a construct in which the C-terminal nine amino acids of Ave1 were fused to GFP were co-expressed with Ve1. The C-terminal nine amino acids are not sufficient to trigger Ve1-mediated recognition as neither construct encoding the C-terminal nine amino acids peptide alone nor a GFP fusion construct can activate Ve1-mediated HR. With a *V. dahliae* complementation assay

we confirmed that the C-terminal nine amino acids of Ave1 are not sufficient to stop the fungus from colonizing Ve1 tomato. Altogether, these results show that the C-terminal nine amino acid sequence is required, but not sufficient, to trigger Ve1-mediated immunity. Thus, recognition may involve additional residues besides the nine amino acid sequence to initiate Ve1-mediated immunity. To test if recognition involves residues immediately adjacent to the nine amino acids, GFP fusion constructs should be generated with longer C-terminal sequences.

We noticed that expression of ChAve1 with Ve1 does not trigger the activation of HR although an Ave1 chimera containing the C-terminal nine amino acids of ChAve1 have the potential to be recognized. Conversely, a ChAve1 chimera containing the C-terminal nine amino acids of VdAve1 does not trigger the activation of an HR. This also supports the hypothesis that a stretch outside the nine amino acid sequence is required to activate Ve1 immunity. However, obviously it needs to be realized that Ve1 evolved in tomato to recognize attempted host colonization by *V. dahliae*, while *C. higginsianum* is not a tomato pathogen and thus Ve1 did not have the chance to co-evolve with ChAve1.

### **A structural model of Ave1**

Plasma membrane-localized PRRs are often activated upon recognition of a short peptide sequence on the surface of their PAMP ligands, such as flg22 or flgII-28 derived from flagellin, elf18 or EFa50 derived from EF-Tu, and pentapeptide TKLGE derived from EIX (Cai et al. 2011; Furukawa et al. 2014; Rotblat et al. 2002; Zipfel et al. 2006; Zipfel et al. 2004). Similarly, a 20-24 peptide in NLP effectors conserved in organisms from three kingdoms of life was shown to be sufficient to activate the PRR receptor RLP23 in Arabidopsis (Albert et al. 2015; Böhm et al. 2014; Oome et al. 2014). However, for the flax-rust effectors AvrM and AvrL567 it has been suggested that multiple contact points are required for recognition by the corresponding flax R-proteins (Ve et al. 2013; Wang et al. 2007). Similarly, multiple residues at separate locations on the surface of the ATR1 effector of *Hyaloperonospora arabidopsidis* are required for recognition by the R-protein RPP1 in Arabidopsis (Chou et al. 2011; Goritschnig et al. 2016). Fusion of various tags to the C-terminus of Ave1 resulted in loss of recognition and fusion of a MYC tag to the N-terminus of Ave1 resulted in reduced Ve1-mediated HR. We generated a structural model of Ave1 to get a better insight in the recognition by Ve1. This model shows that the N- and C-terminal congregate and are exposed at the protein surface. We therefore hypothesize that residues at both the N- and C-terminus of Ave1 are required for recognition by Ve1. This can be tested by generating N-terminal deletions of VdAve1 for co-expression with Ve1.

### Involvement of C-terminal nine amino acids in Ave1 virulence

Ave1 has been characterized as an ortholog of plant natriuretic peptides (PNPs) (de Jonge et al. 2012). Natriuretic peptides were originally identified in animals as atrial natriuretic peptides (ANPs); polypeptide hormones that are secreted by heart muscle cells that are involved in the homeostatic control of salt and water balance and blood pressure (Potter et al. 2009). It has been demonstrated that *X. axonopodis* pv. *citri* utilizes a PNP homolog, XacPNP, to modulate the regulation of host plant homeostasis to establish infection (Gottig et al. 2008). The high homology between tomato SlAve1 and *V. dahliae* VdAve1, likely due to acquisition of VdAve1 by *V. dahliae* from plants through horizontal gene transfer (HGT), in combination with the fact that VdAve1 acts as a virulence factor of *Verticillium* in susceptible tomato plants, suggests that *V. dahliae* utilizes VdAve1 to modulate host plant responses to generate conditions that favour colonization.

Absence of allelic variation in Ave1 alleles from *V. dahliae* and *V. albo-atrum* suggests that identical alleles are required for maximum virulence (de Jonge et al. 2012). However, the importance of the C-terminal amino acids for the function of VdAve1 related to virulence function is unknown. Mutational analysis within Pep-13, a surface-exposed fragment of a calcium-dependent cell wall transglutaminase (TGase) from the oomycete plant pathogen *Phytophthora sojae* that activates plant immune responses, identified the same amino acids indispensable for both TGase activity and immune activation (Brunner et al. 2002). Similarly, amino acid residues that are required for immunity-inducing activity of the flagellin-derived peptide flg22 were found to be important for the intrinsic flagellar function, as mutant flagella were unstable and impaired in flagellar motility (Naito et al. 2008). However, in contrast, the surface exposed pentapeptide TKLGE epitope of the *Trichoderma viride* ethylene-inducing xylanase (EIX) that is essential for recognition by the eLRR receptor LeEIX2 is not involved in xylanase enzymatic activity (Ron and Avni 2004; Rotblat et al. 2002). Further research, involving the complementation of VdAve1 deletion strains in *V. dahliae* with truncated versions of VdAve1, will reveal the particular involvement of the C-terminal nine amino acids in VdAve1 function.

Processing of the VdAve1 protein into smaller peptides might occur during infection. This is the case for the biologically active peptide in animal atrial natriuretic peptides (ANPs) which is a C-terminally derived 28 amino acid peptide that is cleaved from the 126 amino acid ANP precursor (Potter et al. 2009). The importance of precursor maturation has been demonstrated for the recognition of the CLV3 peptide by the eLRR-containing cell-surface receptor CLV2. Receptor CLV2 directly recognizes a 12 amino acid CLE motif of the Arabidopsis CLV3 peptide, that is also conserved in the nematode CLE-like GrCLE peptide, to modulate shoot apical meristem maintenance

during development (Guo et al. 2011; Ni et al. 2011; Wang et al. 2010). Both types of CLE peptides are cleaved from their larger precursor proteins. Studies have suggested that maturation of the CLE peptide is essential for its function (Fiers et al. 2006; Fiers et al. 2005; Guo et al. 2011; Ni and Clark 2006). In addition, elongation of chemically synthesized CLE peptides was found to decrease its activity (Kondo et al. 2008; Kondo et al. 2006).

## Materials and methods

### Plant materials

Tobacco (*Nicotiana tabacum* cv. Petite Havana SR1) and transgenic tomato (*Solanum lycopersicum* cv. MoneyMaker p35S:Ve1) (Fradin et al. 2011) plants were grown in the greenhouse at 21°C/19°C during 16/8 hours day/night periods, respectively, with 70% relative humidity and 100 W/m<sup>2</sup> supplemental light when the light intensity dropped below 150 W/m<sup>2</sup>. After agroinfiltration, tobacco plants were grown in the climate room at 22°C/19°C during 16-h/8-h day/night periods, respectively, with 70% relative humidity.

### Generation of expression constructs for Ave1 homologs

To generate constructs for constitutive expression of the Ave1 homologs, the coding sequences of the homologs were cloned into pDONR207 (Invitrogen, Carlsbad, California) through a Gateway BP reaction, and subsequently transferred into the Gateway-compatible destination vector pSol2092 (Zhang et al. 2013) to generate expression constructs driven by the constitutive CaMV35S promoter. *VdAve1*, *VnAve1* and *FoAve1* were amplified from cDNA using the primers attB-Vd-F and attB-Vd-R, attB-Vn-F and attB-Vn-R, attB-Fo-F and attB-Fo-R, respectively (Supplemental Table 2). *ChAve1*, *CbAve1*, *VvAve1*, *SlAve1* and *XacPNP* were obtained by gene synthesis (Eurofins MWG Operon, Ebersberg, Germany).

### Generation of expression constructs for GFP-VdAve1 fusion

To generate *VdAve1* fused at the C-terminus to the green fluorescent protein (GFP), the *VdAve1* coding sequence without stop codon was PCR amplified using primers containing Gateway attB sites (attB-Vd-F and attB-Vd-R-SC). The PCR fragment was cloned into pDONR207 (Invitrogen, Carlsbad, California) through a Gateway BP reaction to generate entry vector pDONR207::VdAve1-SC. Subsequently, pDONR207::VdAve1-SC was transferred into the Gateway-compatible destination vector pSol2095 (Zhang et al. 2013) to generate an expression construct driven by the constitutive CaMV35S promoter and C-terminally tagged to GFP. Similarly,

pDONR207::VdAve1-SC was transferred into the Gateway-compatible destination vector pGWB14, pGWB8, pGWB17, pGWB20 and pGBW11 (Nakagawa et al. 2007) to generate VdAve1\_HA, VdAve1\_HIS, VdAve1\_4MYC, VdAve1\_10MYC, VdAve1\_FLAG, respectively. To generate VdAve1 fused at the N-terminus to a GFP tag, the GFP coding sequence lacking the stop codon was fused at the N-terminus to the sequence encoding the signal peptide of VdAve1 to ensure extracellular targeting, and at the C-terminus to the VdAve1 coding sequence (without signal peptide). This construct was obtained by gene synthesis (Eurofins MWG Operon, Ebersberg, Germany), and subsequently cloned into Gateway destination vector pSol2092. Fusion constructs were transformed into *Agrobacterium tumefaciens* strain GV3101 by electroporation.

### Generation of deletion and domain swap constructs for VdAve1

Deletion constructs VdAve1\_M1, VdAve1\_M2, VdAve1\_M3 and VdAve1\_M4 were amplified from cDNA using the forward primers attB-Vd-F in combination with reverse primers attB-VdM1-R, attB-VdM2-R, attB-VdM3-R, and attB-VdM4-R, respectively. Vd\_Ch9AA, Ch\_Vd9AA, and SP\_GFP-Vd9AA were amplified from cDNA using the primers attB-Vd-F and attB-VdCh9AA-R, attB-Ch-F and attB-ChVd9AA-R, attB-Vd-F and attB-GFP-Vd9AA-R respectively (Supplemental Table 2). Vd\_Xac9AA, Xac\_Vd9AA and SP\_Vd9AA were obtained by gene synthesis (Eurofins MWG Operon, Ebersberg, Germany). The AVNIEFRQI peptides were synthesized by GenScript company (Piscataway, USA; Purity>75%). The deletion and domain swap constructs were cloned into pDONR207 and subsequently transferred into pSol2092 to generate expression constructs.

### Generation of site-directed mutagenesis constructs for VdAve1

Site-directed mutagenesis constructs V127A, V127R, N128A, N128T, I129A, E130A, E130P, F131A, F131W, R132A, Q133A, Q133N, I134A were amplified from pDONR207::VdAve\_M4 plasmid using the forward primers attB-Vd-F in combination with corresponding reverse primers (Supplemental Table 2). The PCR products were cloned into pDONR207 and subsequently transferred into pSol2092 to generate expression constructs. Subsequently, pDONR207 carrying mutant constructs were transferred into pSol2092 for expression *in planta*.

### *Agrobacterium tumefaciens*-mediated transient expression

Overnight cultures of *A. tumefaciens* strain GV3101 containing expression constructs were harvested at OD<sub>600</sub> of 0.8 to 1 by centrifugation and resuspended to a final OD of 2 in infiltration medium as described previously (Zhang et al. 2013).



*A. tumefaciens* cultures containing constructs to express VdAve1 and Ve1 proteins were mixed in a 1:1 ratio and infiltrated into leaves of five- to six-week-old tobacco plants. At five days post infiltration (dpi), necrosis was examined and quantified by measuring the area of necrosis as percentage of the total infiltrated area.

### **Generation of *Verticillium dahliae* strains carrying VdAve1 constructs**

pDONR207::VdAve1\_M4 and pDONR207::SP\_GFP-Vd9AA were used to clone the constructs into vector pFBT005 under the *VdAve1* promoter, containing a nourseothricin cassette for selection. *V. dahliae* was transformed by *A. tumefaciens* carrying pFBT005 as previously described (Santhanam 2012) and selected on PDA plates containing nourseothricin sulphate (Sigma Aldrich Chemie BV, Zwijndrecht, The Netherlands). After five to seven days at room temperature, individual transformants were transferred to fresh PDA plates and incubated for ten days. Genomic DNA was extracted from the transformants and PCR was performed to test presence of the inserted nourseothricin cassette and of the inserted swap constructs.

### **Verticillium inoculation**

For *V. dahliae* inoculations 10-day-old tomato plants were uprooted, rinsed in water and then dipped for 5 minutes in a suspension of  $10^6$  conidia per mL of water harvested from 1- to 2-week-old *V. dahliae* cultures on PDA as previously described (Fradin et al. 2009). Control plants got the same treatment, but the roots were dipped in water without conidiospores. After replanting in fresh soil, plants were incubated at standard greenhouse conditions. Disease development was monitored up to 21 dpi. The canopy area of 8 plants was measured with ImageJ software and a One-Way ANOVA was performed with IBM SPSS statistics software.

### **Generation of the structural model of VdAve1**

The *V. dahliae* VdAve1 structure was predicted using I-TASSER v4.3 (Zhang 2008) and rendered using UCSF Chimera v1.10.1 (Pettersen et al. 2004). Structural predictions with C-Scores  $> -1.5$  are generally considered to have a correct fold (C-Scores are typically in the range of  $[-5,2]$ ) (Roy et al. 2010). The structural analog in the protein data bank (RCSB PDB) (Rose et al. 2013) was identified using the TM-align program which is part of the I-TASSER package. Analogous structures with TM-Scores  $> 0.5$  are considered to have a similar fold (TM-Scores in the range  $[0,1]$ ) (Roy et al. 2010).



## Acknowledgements

B.P.H.J.T. is supported by a Vici grant of the Research Council for Earth and Life sciences (ALW) of the Netherlands Organization for Scientific Research (NWO). This research was funded in part by the Centre for BioSystems Genomics (CBSG) which is part of the Netherlands Genomics Initiative/NWO and by ERA-NET Plant Genomics. We thank Bert Essenstam and Pauline Sanderson at Unifarm for excellent plant care, and Sebastjan Radišek at the Institute of Hop Research and Brewing, Slovenia, for providing *Verticillium* strains.

## References

- Albert, I., Böhm, H., Albert, M., Feiler, C. E., Imkampe, J., Wallmeroth, N., Brancato, C., Raaymakers, T. M., Oome, S., Zhang, H., Krol, E., Grefen, C., Gust, A. A., Chai, J., Hedrich, R., Van den Ackerveken, G., and Nürnberger, T. 2015. An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity. *Nat Plants* 1:15140.
- Böhm, H., Albert, I., Oome, S., Raaymakers, T. M., Van den Ackerveken, G., and Nürnberger, T. 2014. A conserved peptide pattern from a widespread microbial virulence factor triggers pattern-induced immunity in *Arabidopsis*. *PLoS Pathog* 10:e1004491.
- Brunner, F., Rosahl, S., Lee, J., Rudd, J. J., Geiler, C., Kauppinen, S., Rasmussen, G., Scheel, D., and Nurnberger, T. 2002. Pep-13, a plant defense-inducing pathogen-associated pattern from *Phytophthora* transglutaminases. *EMBO J* 21:6681-6688.
- Cai, R., Lewis, J., Yan, S., Liu, H., Clarke, C. R., Campanile, F., Almeida, N. F., Studholme, D. J., Lindeberg, M., and Schneider, D. 2011. The plant pathogen *Pseudomonas syringae* pv. *tomato* is genetically monomorphic and under strong selection to evade tomato immunity. *PLoS Pathog* 7:e1002130.
- Ceroni, A., Passerini, A., Vullo, A., and Frasconi, P. 2006. DISULFIND: a disulfide bonding state and cysteine connectivity prediction server. *Nucleic Acids Res* 34:W177-181.
- Chen, T., Kan, J., Yang, Y., Ling, X., Chang, Y., and Zhang, B. 2016. A Ve homologous gene from *Gossypium barbadense*, *Gbvdr3*, enhances the defense response against *Verticillium dahliae*. *Plant Physiol Biochem* 98:101-111.
- Chou, S., Krasileva, K. V., Holton, J. M., Steinbrenner, A. D., Alber, T., and Staskawicz, B. J. 2011. *Hyaloperonospora arabidopsidis* ATR1 effector is a repeat protein with distributed recognition surfaces. *Proc Natl Acad Sci U S A* 108:13323-13328.
- Cook, D. E., Mesarich, C. H., and Thomma, B. P. H. J. 2015. Understanding plant immunity as a surveillance system to detect invasion. *Annu Rev Phytopathol*.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- Dodds, P. N., and Rathjen, J. P. 2010. Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* 11:539-548.
- Fiers, M., Golemiac, E., Xu, J., van der Geest, L., Heidstra, R., Stiekema, W., and Liu, C. M. 2005. The 14-amino acid CLV3, CLE19, and CLE40 peptides trigger consumption of the root meristem in *Arabidopsis* through a CLAVATA2-dependent pathway. *Plant Cell* 17:2542-2553.
- Fiers, M., Golemiac, E., van der Schors, R., van der Geest, L., Li, K. W., Stiekema, W. J., and Liu, C. M. 2006. The CLAVATA3/ESR motif of CLAVATA3 is functionally independent from the nonconserved flanking sequences. *Plant Physiol* 141:1284-1292.
- Flor, H. H. 1971. Current status of the gene-for-gene concept. *Annu Rev Phytopathol* 9:275-296.


- Fradin, E. F., and Thomma, B. P. H. J. 2006. Physiology and molecular aspects of *Verticillium* wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol Plant Pathol* 7:71-86.
- Fradin, E. F., Abd-El-Halim, A., Masini, L., van den Berg, G. C., Joosten, M. H., and Thomma, B. P. H. J. 2011. Interfamily transfer of tomato *Ve1* mediates *Verticillium* resistance in *Arabidopsis*. *Plant Physiol* 156:2255-2265.
- Fradin, E. F., Zhang, Z., Juarez Ayala, J. C., Castroverde, C. D., Nazar, R. N., Robb, J., Liu, C. M., and Thomma, B. P. H. J. 2009. Genetic dissection of *Verticillium* wilt resistance mediated by tomato *Ve1*. *Plant Physiol* 150:320-332.
- Furukawa, T., Inagaki, H., Takai, R., Hirai, H., and Che, F.-S. 2014. Two distinct EF-Tu epitopes induce immune responses in rice and *Arabidopsis*. *Mol. Plant Microbe Interact.* 27:113-124.
- Goritschnig, S., Steinbrenner, A. D., Grunwald, D. J., and Staskawicz, B. J. 2016. Structurally distinct *Arabidopsis thaliana* NLR immune receptors recognize tandem WY domains of an oomycete effector. *New Phytol.*
- Gottig, N., Garavaglia, B. S., Daurelio, L. D., Valentine, A., Gehring, C., Orellano, E. G., and Ottado, J. 2008. *Xanthomonas axonopodis* pv. *citri* uses a plant natriuretic peptide-like protein to modify host homeostasis. *Proc Natl Acad Sci U S A* 105:18631-18636.
- Guo, Y., Ni, J., Denver, R., Wang, X., and Clark, S. E. 2011. Mechanisms of molecular mimicry of plant CLE peptide ligands by the parasitic nematode *Globodera rostochiensis*. *Plant Physiol* 157:476-484.
- Inderbitzin, P., Bostock, R. M., Davis, R. M., Usami, T., Platt, H. W., and Subbarao, K. V. 2011. Phylogenetics and taxonomy of the fungal vascular wilt pathogen *Verticillium*, with the descriptions of five new species. *PLoS One* 6:e28341.
- Isaac, I. 1953. A further comparative study of pathogenic isolates of *Verticillium*: *V. nubilum* Pethybr. and *V. tricorpus* sp.nov. *Trans Br Mycol Soc* 36:180-IN182.
- Jones, J. D., and Dangl, J. L. 2006. The plant immune system. *Nature* 444:323-329.
- Kawchuk, L. M., Hachey, J., Lynch, D. R., Kulcsar, F., van Rooijen, G., Waterer, D. R., Robertson, A., Kokko, E., Byers, R., Howard, R. J., Fischer, R., and Pruffer, D. 2001. Tomato *Ve* disease resistance genes encode cell surface-like receptors. *Proc Natl Acad Sci U S A* 98:6511-6515.
- Kondo, T., Nakamura, T., Yokomine, K., and Sakagami, Y. 2008. Dual assay for MCLV3 activity reveals structure-activity relationship of CLE peptides. *Biochem Biophys Res Commun* 377:312-316.
- Kondo, T., Sawa, S., Kinoshita, A., Mizuno, S., Kakimoto, T., Fukuda, H., and Sakagami, Y. 2006. A plant peptide encoded by CLV3 identified by in situ MALDI-TOF MS analysis. *Science* 313:845-848.
- Naito, K., Taguchi, F., Suzuki, T., Inagaki, Y., Toyoda, K., Shiraishi, T., and Ichinose, Y. 2008. Amino acid sequence of bacterial microbe-associated molecular pattern flg22 is required for virulence. *Mol. Plant Microbe Interact.* 21:1165-1174.
- Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata, R., Kawai, T., Tanaka, K., Niwa, Y., Watanabe, Y., Nakamura, K., Kimura, T., and Ishiguro, S. 2007. Improved gateway binary vectors: High-performance vectors for creation of fusion constructs in Transgenic analysis of plants. *Biosci Biotechnol Biochem* 71:2095-2100.
- Nembaware, V., Seoighe, C., Sayed, M., and Gehring, C. 2004. A plant natriuretic peptide-like gene in the bacterial pathogen *Xanthomonas axonopodis* may induce hyper-hydration in the plant host: a hypothesis of molecular mimicry. *BMC Evol Biol* 4:10.
- Ni, J., and Clark, S. E. 2006. Evidence for functional conservation, sufficiency, and proteolytic processing of the CLAVATA3 CLE domain. *Plant Physiol* 140:726-733.
- Ni, J., Guo, Y., Jin, H., Hartsell, J., and Clark, S. E. 2011. Characterization of a CLE processing activity. *Plant Mol Biol* 75:67-75.
- Oome, S., Raaymakers, T. M., Cabral, A., Samwel, S., Böhm, H., Albert, I., Nürnberger, T., and Van den Ackerveken, G. 2014. Nep1-like proteins from three kingdoms of life act as a microbe-associated molecular pattern in *Arabidopsis*. *Proc Natl Acad Sci U S A* 111:16955-16960.
- Pearce, G., Yamaguchi, Y., Munske, G., and Ryan, C. A. 2008. Structure-activity studies of AtPep1, a plant peptide signal involved in the innate immune response. *Peptides* 29:2083-2089.

- Petersen, T. N., Brunak, S., von Heijne, G., and Nielsen, H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8:785-786.
- Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., and Ferrin, T. E. 2004. UCSF Chimera - A visualization system for exploratory research and analysis. *J Comput Chem* 25:1605-1612.
- Potter, L. R., Yoder, A. R., Flora, D. R., Antos, L. K., and Dickey, D. M. 2009. Natriuretic peptides: their structures, receptors, physiologic functions and therapeutic applications. 2008/12/18 ed.
- Pruitt, R. N., Schwessinger, B., Joe, A., Thomas, N., Liu, F., Albert, M., Robinson, M. R., Chan, L. J. G., Luu, D. D., Chen, H., Bahar, O., Daudi, A., De Vleeschauwer, D., Caddell, D., Zhang, W., Zhao, X., Li, X., Heazlewood, J. L., Ruan, D., Majumder, D., Chern, M., Kalbacher, H., Midha, S., Patil, P. B., Sonti, R. V., Petzold, C. J., Liu, C. C., Brodbelt, J. S., Felix, G., and Ronald, P. C. 2015. The rice immune receptor XA21 recognizes a tyrosine-sulfated protein from a Gram-negative bacterium. *Sci Adv* 1:e1500245.
- Ron, M., and Avni, A. 2004. The receptor for the fungal elicitor ethylene-inducing xylanase is a member of a resistance-like gene family in tomato. *Plant Cell* 16:1604-1615.
- Rose, P. W., Bi, C., Bluhm, W. F., Christie, C. H., Dimitropoulos, D., Dutta, S., Green, R. K., Goodsell, D. S., Prlić, A., Quesada, M., Quinn, G. B., Ramos, A. G., Westbrook, J. D., Young, J., Zardecki, C., Berman, H. M., and Bourne, P. E. 2013. The RCSB Protein Data Bank: new resources for research and education. *Nucleic Acids Res* 41:D475-D482.
- Rotblat, B., Enshell-Seiffers, D., Gershoni, J. M., Schuster, S., and Avni, A. 2002. Identification of an essential component of the elicitation active site of the EIX protein elicitor. *Plant J* 32:1049-1055.
- Roy, A., Kucukural, A., and Zhang, Y. 2010. I-TASSER: a unified platform for automated protein structure and function prediction. *Nat Protoc* 5:725-738.
- Santhanam, P. 2012. Random insertional mutagenesis in fungal genomes to identify virulence factors. Pages 509-517 in: *Plant fungal pathogens. Methods in molecular biology*, vol. 835. T. B. Bolton MD, ed. Humana press, Totowa.
- Schaible, L., Cannon, O. S., and Waddoups, V. 1951. Inheritance of resistance to *Verticillium* wilt in a tomato cross. *Phytopathology* 41:986-990.
- Song, X. F., Yu, D. L., Xu, T. T., Ren, S. C., Guo, P., and Liu, C. M. 2012. Contributions of individual amino acid residues to the endogenous CLV3 function in shoot apical meristem maintenance in *Arabidopsis*. *Mol Plant* 5:515-523.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- van Kan, J. A., van den Ackerveken, G., and de Wit, P. J. G. M. 1991. Cloning and characterization of cDNA of avirulence gene *avr9* of the fungal pathogen *Cladosporium fulvum*, causal agent of tomato leaf mold. *Mol. Plant Microbe Interact.* 4:52-59.
- Ve, T., Williams, S. J., Catanzariti, A.-M., Rafiqi, M., Rahman, M., Ellis, J. G., Hardham, A. R., Jones, D. A., Anderson, P. A., and Dodds, P. N. 2013. Structures of the flax-rust effector AvrM reveal insights into the molecular basis of plant-cell entry and effector-triggered immunity. *Proc Natl Acad Sci U S A* 110:17594-17599.
- Wang, C.-I. A., Gunčar, G., Forwood, J. K., Teh, T., Catanzariti, A.-M., Lawrence, G. J., Loughlin, F. E., Mackay, J. P., Schirra, H. J., and Anderson, P. A. 2007. Crystal structures of flax rust avirulence proteins AvrL567-A and-D reveal details of the structural basis for flax disease resistance specificity. *Plant Cell* 19:2898-2912.
- Wang, G., Fiers, M., Ellendorff, U., Wang, Z. Z., de Wit, P. J. G. M., Angenent, G. C., and Thomma, B. P. H. J. 2010. The diverse roles of extracellular leucine-rich repeat-containing receptor-like proteins in plants. *CRC Crit Rev Plant Sci* 29:285-299.
- Yang, Y., Ling, X., Chen, T., Cai, L., Liu, T., Wang, J., Fan, X., Ren, Y., Yuan, H., and Zhu, W. 2014. A cotton *GbvdR5* gene encoding a leucine-rich-repeat receptor-like protein confers resistance to *Verticillium dahliae* in transgenic *Arabidopsis* and upland cotton. *Plant Mol Biol Report*:1-15.

- Yennawar, N. H., Li, L. C., Dudzinski, D. M., Tabuchi, A., and Cosgrove, D. J. 2006. Crystal structure and activities of EXPB1 (Zea m 1), a beta-expansin and group-1 pollen allergen from maize. *Proc Natl Acad Sci U S A* 103:14664-14671.
- Zhang, B., Yang, Y., Chen, T., Yu, W., Liu, T., Li, H., Fan, X., Ren, Y., Shen, D., Liu, L., Dou, D., and Chang, Y. 2012. Island cotton *Gbve1* gene encoding a receptor-like protein confers resistance to both defoliating and non-defoliating isolates of *Verticillium dahliae*. *PLoS One* 7:e51091.
- Zhang, Y. 2008. I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics* 9:40.
- Zhang, Y., Wang, X., Yang, S., Chi, J., Zhang, G., and Ma, Z. 2011. Cloning and characterization of a *Verticillium* wilt resistance gene from *Gossypium barbadense* and functional analysis in *Arabidopsis thaliana*. *Plant Cell Rep* 30:2085-2096.
- Zhang, Z., Fradin, E., de Jonge, R., van Esse, H. P., Smit, P., Liu, C.-M., and Thomma, B. P. H. J. 2013. Optimized agroinfiltration and virus-induced gene silencing to study Ve1-mediated *Verticillium* resistance in tobacco. *Mol. Plant Microbe Interact.* 26:182-190.
- Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E. J., Jones, J. D. G., Felix, G., and Boller, T. 2004. Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature* 428:764-767.
- Zipfel, C., Kunze, G., Chinchilla, D., Caniard, A., Jones, J. D., Boller, T., and Felix, G. 2006. Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts *Agrobacterium*-mediated transformation. *Cell* 125:749-760.

## Supplemental data


4




**SUPPLEMENTAL FIGURE 1 | Immunoblotting of GFP-tagged Ave1 homologs.** GFP-tagged Ave1 homologs were detected by western blotting using GFP antibody ( $\alpha$ -GFP). Coomassie-stained blots (CBS) showing the 50 kDa Rubisco band present in the input samples confirm equal loading.




**SUPPLEMENTAL FIGURE 2 | Stability of truncated Ave1 proteins.** SP\_GFP, VdAve1\_GFP, GFP\_VdAve1, GFP-tagged VdAve1\_M4 and SP\_GFP\_Vd9AA proteins were detected using GFP antibody ( $\alpha$ -GFP). Coomassie-stained blots (CBS) showing the 50 kDa Rubisco band present in the input samples confirm equal loading.



**SUPPLEMENTAL FIGURE 3 | Stability of chimeric Ave proteins Vd\_Xac9AA and Xac\_Vd9AA.** GFP-tagged VdAve1, XacPNP, chimeric constructs Vd\_Xac9AA and Xac\_Vd9AA were detected by immunoblotting using GFP antibody ( $\alpha$ -GFP). Coomassie-stained blots (CBS) showing the 50 kDa Rubisco band present in the input samples confirm equal loading.



**SUPPLEMENTAL FIGURE 4 | Stability of site-directed mutagenesis constructs that showed compromised HR-inducing capacity.** GFP-tagged VdAve1, XacPNP, site-directed mutagenesis constructs V127A, V127R, N128A, N128T, I129A, E130A, F131A, F131W, R132A, Q133A, I134A were detected by immunoblotting using GFP antibody ( $\alpha$ -GFP). Coomassie-stained blots (CBS) showing the 50 kDa Rubisco band present in the input samples confirm equal loading.



**SUPPLEMENTAL FIGURE 5 | Stability of chimeric Ave proteins Vd\_Ch9AA and Ch\_Vd9AA.** GFP-tagged VdAve1, ChAve1, chimeric constructs Vd\_Ch9AA and Ch\_Vd9AA were detected by immunoblotting using GFP antibody. Coomassie-stained blots (CBS) showing the 50 kDa Rubisco band present in the input samples confirm equal loading.

SUPPLEMENTAL TABLE 1 | *Verticillium* strains analysed for presence of Ave1 homologs

Species	Isolate	Original host	Origin	Ave1
<i>V. nonalfalfae</i>	P10	hop	Germany	-
	P114/1	hop	Germany	-
	P34/1	hop	Germany	-
	P15	hop	Germany	-
	P55	hop	Germany	-
	P83	hop	Germany	-
	6/99	hop	Germany	-
	14/93	hop	Germany	-
	15/98	hop	Germany	-
	P84/2	hop	Germany	-
	16/00	hop	Germany	-
	T2	hop	Slovenia	-
	TABOR6	hop	Slovenia	-
	Ledina09-V.aa	hop	Slovenia	+
	BIZ	hop	Slovenia	-
	VranBis09	hop	Slovenia	-
	Sent4	hop	Slovenia	-
	MO3	hop	Slovenia	-
	OCer	hop	Slovenia	-
	zup	hop	Slovenia	-
	Rec91	hop	Slovenia	-
	KRES98	hop	Slovenia	-
	Gajsek	hop	Slovenia	-
	1985a	hop	U.K.	-
	11041	hop	U.K.	-
	11055	hop	U.K.	-
	11047	hop	U.K.	-
	11097	hop	U.K.	-
	11100	hop	U.K.	-
	1974	hop	U.K.	-
	298099	hop	U.K.	-
	298100	hop	U.K.	-
	298101	hop	U.K.	-
	298102	hop	U.K.	-
	11052	hop	U.K.	-
	1953	hop	U.K.	-
	298092	hop	U.K.	-
	298095	hop	U.K.	-
	Sol	hop	Poland	+
	CBS393.91	hop	Belgium	-
	kum	cucumber	Slovenia	-
	Surf	surfinias	Slovenia	-
	11077	<i>Galinsoga ciliata</i>	U.K.	-
11081	chrysanthemum	U.K.	+	
CBS102.464	cynara	Italija	-	
CBS241.82	catalpa	Italija	+	
CBS454.51	potato	U.K.	-	
CBS682.88	potato	Netherlands	-	
11066	potato	U.K.	-	
T179	tomato	U.K.	+	
CBS321.91	tomato	Netherlands	-	


Species	Isolate	Original host	Origin	Ave1
	AR01/067	tomato	U.K.	-
	AR0/140	tomato	U.K.	-
	AR01/JS1	tomato	U.K.	-
	PD83/53a	tomato	Netherlands	-
	PD2000/4186a	tomato	Netherlands	+
<i>V. alfalfae</i>	Luc	alfalfa	U.K.	+
	41	alfalfa	Canada	-
	CBS392.91	alfalfa	Netherlands	-
	107	alfalfa	USA	-
	PD693	potato	Iran	-
	314193	potato	Australia	-
	340646	potato	Spain	+
	11	alfalfa	Slovenia	-
<i>V. dahliae</i>	JKG2	catalpa	Netherlands	-
	CIG3-Vd	hop	Slovenia	-
	JKG1	potato	Netherlands	-
	JKG8	potato	Netherlands	-
	A56	potato	Slovenia	+
	DJK	chrysanthemum	Netherlands	-
	MH	chrysanthemum	Netherlands	-
	Mint	mint	USA	-
	GAJ09	hop	Slovenia	-
	PDRENU	hop	Slovenia	-
	CasD	hop	Slovenia	-
	KresD	hop	Slovenia	-
	MoD	hop	Slovenia	-
	Oset	hop	Slovenia	-
	12099	hop	U.K.	-
	12042	hop	U.K.	-
	PD335	cabbage	unknown	-
	PD584	mint	unknown	-
	V-176l	cotton	Netherlands	+
	V-138l	cotton	Germany	-
	PAPmb	pepper	Slovenia	-
	PAP	pepper	Slovenia	+
	Pap99	pepper	Slovenia	+
	Pap2008	pepper	Slovenia	+
<i>V. nubilum</i>	CBS456.51	potato	UK	+
	CBS457.51	soil	UK	+
	PD621	mushroom compost	UK	+
	PD702	potato	UK	+
<i>V. tricorpus</i>	CBS227.84	potato	Netherlands	-
	JKG20	linden	Netherlands	-
<i>V. longisporium</i>	CBS110218	<i>Brassica napus</i>	Sweden	-
	PD330	cabbage	unknown	-

**SUPPLEMENTAL TABLE 2** | Primers used in this study.

Primer name	Sequence (5'-3')
attB-Vd-F	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGAAGCTTCTACGCTT
attB-Vd-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTATATCTGTCTAAATTC
attB-Vn-F	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGAAGCTTCTACGCTT
attB-Vn-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTATATCTGTTCAAACTC
attB-Fo-F	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGAACTACTCGCACTA
attB-Fo-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTCATCTTTGTACAAAATCGATATTT
attB-Vd-R-SC	GGGGACCACTTTGTACAAGAAAGCTGGGTCTATCTGTCTAAATTC
attB-VdM1-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTACTTATGCCTCGTCCCTT
attB-VdM2-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTAAACAATGGCATCATATGAGT
attB-VdM3-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTACTTGTGTGCTGCTTTGGTAA
attB-VdM4-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTAAAGCTCTGTCAACCACCCGCA
attB-VdCh9AA-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTCAAATTTGACGTACTCAATGTTTACTTTCT-TATGCCTCGTTCCTT
attB-Ch-F	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGAAGGCCTTCTTCTAGGAG
attB-ChVd9AA-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTATATCTGTCTAAATTCGATGTTGACC-GCAGTTACGCCAAAGTCCG
attB-GFP-Vd9AA-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTATATCTGTCTAAATTCGATGTTGACCCGCTTTGTATAGTTCATCCATGC
V127A R	TTATATCTGTCTAAATTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
N128A R	TTATATCTGTCTAAATTCGATCGCGACCGCCTTATGCCTCGTTCCTTC
I129A R	TTATATCTGTCTAAATTCGCGGTTGACCGCCTTATGCCTCGTTCCTTC
E130A R	TTATATCTGTCTAAACGCGATGTTGACCGCCTTATGCCTCGTTCCTTC
F131A R	TTATATCTGTCTCGCTTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
R132A R	TTATATCTGCGCAAATTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
Q133A R	TTATATCGCTCTAAATTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
I134A R	TTACGCCTGTCTAAATTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
V127R R	TTATATCTGTCTAAATTCGATGTTTCTGCGCTTATGCCTCGTTCCTTC
N128T R	TTATATCTGTCTAAATTCGATTGTGACCGCCTTATGCCTCGTTCCTTC
E130P R	TTATATCTGTCTAAAGGGGATGTTGACCGCCTTATGCCTCGTTCCTTC
F131W R	TTATATCTGTCTCCATTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
Q133N R	TTATATATTTCTAAATTCGATGTTGACCGCCTTATGCCTCGTTCCTTC
Vd_Xac_V2 R	TTAAATATTTGCCAGGGCGCTGTGACTGCCTTATGCCTCGTTCCTTC
Vd_Xac_V5 R	TTAAATATTTGCCAATCCGCTGTTCTTGCCCTTATGCCTCGTTCCTTC
Vd_Xac_V2V5 R	TTAAATATTTGCCAATCCGCTGTGACTGCCTTATGCCTCGTTCCTTC
Vd_Xac_V2V6 R	TTAAATATTTGCAAATCCGCGTTGACTGCCTTATGCCTCGTTCCTTC







**Characterization of an expanded  
family of Ave1 homologs in the  
apple and pear scab fungi  
*Venturia inaequalis* and *V. pirina***

**Chapter 5**

## Abstract

*Venturia inaequalis* and *V. pirina* are fungal pathogens that cause scab on apple and pear, respectively. Although several gene-for-gene interactions have been identified for *V. inaequalis* and apple, no avirulence gene has been cloned thus far. Recently, a family of *Ave1* homologs was identified in *V. pirinia*. *Ave1* was first identified in race 1 strains of *Verticillium dahliae* that has a wide host range including many crops, and was shown to be recognized by the tomato *Ve1* immune receptor. In absence of *Ve1*, *Ave1* contributes to virulence of *V. dahliae*. Homologs of *Ave1* have been identified in a wide range of plants and in a handful of phylogenetically unrelated plant pathogens, including the fungi *Fusarium oxysporum*, *Cercospora beticola* and *Colletotrichum higginsianum* and the bacterium *Xanthomonas axonopodis*. In this study we found that also *V. inaequalis* contains a family of *Ave1* homologs. We show that the tomato *Ve1* immune receptor is unable to recognize a selection of *Ave1* homologs from *V. inaequalis* and *V. pirina*. Transcriptome analysis revealed that *V. inaequalis* *ViAve1-1*, *ViAve1-5* and *ViAve1-9* are expressed *in planta*. We show that *in planta* expressed *ViAve1-1*, *ViAve1-5* and *ViAve1-9* are unable to complement a *V. dahliae* *VdAve1* deletion strain, suggesting that these homologs have different functions than *V. dahliae* *VdAve1*.

## Introduction

Plants evolved an immune system that detects conserved microbe-associated molecular patterns (MAMPS) to activate defense responses against invading pathogens (Cook et al. 2015). Successful plant pathogens evolved appropriate effectors; secreted molecules of which many can deregulate immunity in the respective host (Rovenich et al. 2014) and that are typically species, strain or even isolate specific. In turn, host plants typically evolve immune receptors that can recognize such effectors and mount an immune response to prevent further invasion. However, recognition of effectors and subsequent immune responses pose a huge selection pressure on pathogens to either evolve new effectors or alter or loose recognized effectors to prevent the activation of immune responses. This continuous co-evolution between host and pathogen is elegantly visualized in the zig-zag model (Jones and Dangl 2006). However, this model is focused from the point-of-view of the plant, while the point-of-view of the pathogen is largely underrepresented. Additionally, effectors and MAMPs are depicted as two distinct and opposing features of pathogen molecules, while this distinction is often not clear cut (Thomma et al. 2011). The recently proposed invasion model states that any signal that betrays an invading organism can potentially serve as a cue to activate an appropriate host response, either to a beneficial or detrimental organism (Cook et al. 2015).

*Verticillium dahliae* is a fungal plant pathogen with a broad host range that includes many crops (Fradin and Thomma 2006). *V. dahliae* race 1 strains express the effector Ave1 that activates immunity in tomato plants carrying the Ve1 receptor (de Jonge et al. 2012). In addition, it was shown that Ave1 contributes to the virulence of *V. dahliae*. Homologs of Ave1 are mainly found in plants and only in a handful of plant pathogens, including the fungi *Fusarium oxysporum*, *Cercospora beticola* and *Colletotrichum higginsianum* and the bacterium *Xanthomonas axonopodis*. Only one homolog of Ave1 is found in the genome of each of these organisms, and these homologs share between 52-95% sequence similarity. In contrast, each *V. dahliae* strain analyzed thus far contains an additional Ave1-like (Ave1L) allele. Five alleles have been identified, denoted as Ave1L1 to Ave1L5 that share between 64 and 68% sequence similarity with *V. dahliae* VdAve1 (Chapter 6, this thesis). Co-expression of Ave1 homologs derived from *Fusarium oxysporum*, *Cercospora beticola* and *Colletotrichum higginsianum* and *Xanthomonas axonopodis* with tomato immune receptor Ve1 revealed differential recognition of the various homologs (de Jonge et al. 2012).


*Venturia inaequalis* and *V. pirina* cause devastating scab diseases on apple and pear, respectively (Bowen et al. 2011; Shabi et al. 1973). Although a number of gene-for-gene interactions have been identified for *V. inaequalis* and its host apple (Bus et al. 2011; Gessler et al. 2006), no avirulence genes have been cloned thus far (Bus et al. 2011). AvrLm6 is an effector protein of the fungal pathogen *Leptosphaeria maculans* that triggers an immune response in Brassica plants expressing the immune receptor Rlm6 (Fudal et al. 2007). Homologs of AvrLm6 have been identified in other plant pathogens, such as *Colletotrichum* and *Fusarium* species (Grandaubert et al. 2014). Interestingly, an expanded family of AvrLm6 effector homologs was identified in the genomes of *V. inaequalis* and *V. pirina* (Shiller et al. 2015). However, the *V. inaequalis* AvrLm6 homolog with the highest sequence similarity to *L. maculans* AvrLm6 was unable to trigger an immune response in Brassica plants expressing the immune receptor Rlm6 (Cooke et al. 2014). Recently it was found that the *V. pirina* genomes also contains an expanded Ave1 effector gene family, which is in contrast to the plant pathogens *Fusarium oxysporum*, *Cercospora beticola* and *Colletotrichum higginsianum* and *Xanthomonas axonopodis* that carry only one homolog of Ave1 (Cooke et al. 2014; de Jonge et al. 2012). In this study we test whether any of the homologs in the expanded Ave1 family derived from *Venturia* spp. is recognized by the tomato Ve1 immune receptor.

## Results

### **Ave1 homologs derived from *Venturia inaequalis* and *V. pirina* are not recognized by tomato Ve1**


Previously, 14 loci with homology to *V. dahliae* VdAve1 were identified in the genome of *V. pirina*, including 12 genes (VpAve1-1 to VpAve1-12) and 2 pseudogenes (VpAve1-13 and VpAve1-14) (Cooke et al. 2014) (Joanna Bowen personal communication). We similarly analysed the genome of *V. inaequalis* for the presence of Ave1 homologs. In this analysis, 17 loci were identified comprising 9 genes (ViAve1-1 to ViAve1-5 and ViAve1-8 to ViAve1-10) and 8 pseudogenes (ViAve1-6, ViAve1-7 and ViAve1-11 to ViAve1-16) (Kim Plummer personal communication). The Ave1 homologs derived from *V. inaequalis* and *V. pirina* display between 51-63% sequence similarity to *V. dahliae* Ave1 (VdAve1). Alignment of the gene products of *V. inaequalis* and *V. pirina* Ave1 homologs with *V. dahliae* VdAve1 shows a high degree of sequence conservation distributed over the whole protein sequence (Figure 1). We hypothesized that since Ave1 is recognized by tomato immune receptor Ve1, *Venturia* Ave1 homologs are potentially recognized by Ve1 as well. To test this hypothesis, we co-expressed Ve1 with Ave1 homologs from *V. inaequalis* (ViAve1-1, ViAve1-2, ViAve1-4, ViAve1-5, ViAve1-8, ViAve1-9, ViAve1-10), and Ave1

homologs from *V. pirina* (*VpAve1-1*, *VpAve1-2*, *VpAve1-3*, *VpAve1-4*, *VpAve1-5*, *VpAve1-6*, *VpAve1-7*, *VpAve1-11*). Unfortunately, we were unable to clone *Ave1* homologs *ViAve1-3*, *VpAve1-8*, *VpAve1-9*, *VpAve1-10* and *VpAve1-12*. Whereas at 5 days post inoculation (dpi) a clear hypersensitive response (HR) was observed upon co-expression of the control combination of *Ve1* and *V. dahliae* *Ave1*, no HR was observed in any of the co-inoculations of *Ve1* with the selected *V. inaequalis* and *V. pirina* *Ave1* homologs, except for some small necrotic spots in the case of *ViAve1-4* and *ViAve1-9* (Figure 2A and B). These results suggest that the tomato immune receptor *Ve1* is unable to recognize *Ave1* homologs derived from the apple and pear scab fungi *V. inaequalis* and *V. pirina*, respectively.



**FIGURE 1 | Alignment of Ave1 homologs indicates regions with high conservation** Alignment of *Ave1* homologs derived from *Venturia inaequalis* and *V. pirina* with *V. dahliae* *VdAve1*. The intensity of the blue color indicates the degree of conservation.






**FIGURE 2 | Ave1 homologs derived from *Venturia inaequalis* and *V. pirina* are not recognized by tomato Ve1.** (A) Co-expression of selected Ave1 homologs from *Venturia inaequalis* with tomato Ve1. (B) Co-expression of selected Ave1 homologs from *Venturia pirina* with tomato Ve1.

### **Ave1 homologs derived from *Venturia inaequalis* cannot complement loss of *Verticillium dahliae* Ave1**

A method to mimic leaf or fruit surface penetration and *in planta* expression is to grow *V. inaequalis* on cellophane discs. Therefore, transcriptome analysis was performed on *V. inaequalis* grown on cellophane discs (Kucheryava et al. 2008). This analysis revealed that the Ave1 homologs *ViAve1-1*, *ViAve1-5* and *ViAve1-9* are induced *in planta* (Kim Plummer personal communication). To test whether the Ave1 homologs from *V. inaequalis* share the same functionality as *V. dahliae* Ave1 to promote fungal virulence, we transformed a *V. dahliae* Ave1 deletion strain with constructs to express *ViAve1-1*, *ViAve1-5* and *ViAve1-9* driven by the *V. dahliae* Ave1 promoter. After confirmation that the strains contain the expression construct, the transformants were inoculated onto tomato plants that lack the Ve1 immune receptor and screened for virulence. Tomato plants that were inoculated with wild type *V. dahliae* and the *V. dahliae* Ave1 deletion strain that was complemented with *VdAve1* as a control showed clear symptoms of Verticillium wilt. In contrast, the Ave1 deletion strain and the transformants carrying expression constructs for *ViAve1-1*, *ViAve1-5* and *ViAve1-9* only induced reduced Verticillium wilt symptoms when compared with the symptoms induced by wild type *V. dahliae* (Figure 3A; B). To confirm that the transgenes were expressed, expression of *Ave1*, *ViAve1-1*, *ViAve1-5* and *ViAve1-9* by *V. dahliae*

*in planta* was analyzed. All genes were expressed at a similar level as *Ave1* by *V. dahliae* wild type (Figure 3D). Thus, *ViAve1-1*, *ViAve1-5* and *ViAve1-9* are unable to complement a *V. dahliae* *VdAve1* deletion strain, suggesting that these homologs display (a) different function(s) than *V. dahliae* *VdAve1*.



**FIGURE 3 | *Ave1* homologs derived from *Venturia inaequalis* cannot complement loss of *Verticillium dahliae* *Ave1*** (A) Complementation of *V. dahliae*  $\Delta Ave1$  with *ViAve1-1*, *ViAve1-5* and *ViAve1-9* on tomato plants lacking *Ve1* (top) and *Ve1* tomato (bottom) (B) Canopy area of tomato plants lacking *Ve1* inoculated with *V. dahliae*  $\Delta Ave1$  expressing *ViAve1-1*, *ViAve1-5* and *ViAve1-9* at 14 dpi (n=8). (C) Canopy area of *Ve1* tomato plants inoculated with *V. dahliae*  $\Delta Ave1$  expressing *ViAve1-1*, *ViAve1-5* and *ViAve1-9* at 14 dpi (n=8). Data are presented as the mean  $\pm$  standard deviation. Different letters indicate significant differences ( $P < 0.05$ ; One-Way ANOVA). (D) Gene expression of transgenes in a *V. dahliae* *VdAve1* deletion strain when compared with *VdAve1* expression in wild type *V. dahliae*.



### **Ave1 homologs derived from *Venturia inaequalis* are unable to initiate an immune response in Ve1 tomato plants**

From the co-expression analysis on *N. benthamiana* we concluded that the *Venturia Ave1* homologs are not recognized by tomato Ve1 due to lack of a visible HR. Nevertheless, we have previously shown that Ve1 is able to establish immunity without signs of an HR under particular conditions (Zhang et al. 2013a). We therefore tested whether the *V. dahliae* transformants expressing *ViAve1-1*, *ViAve1-5* and *ViAve1-9* are contained when inoculated on tomato plants carrying Ve1 (Figure 3A; C). Tomato plants that were inoculated with wild type *V. dahliae* and the *V. dahliae Ave1* deletion strain that was complemented with *VdAve1* as a control showed no symptoms of Verticillium wilt, indicating that they are successfully contained upon recognition by Ve1. In contrast, the tomato plants inoculated with the *Ave1* deletion strain and the transformants carrying expression constructs for *ViAve1-1*, *ViAve1-5* and *ViAve1-9* showed clear symptoms of Verticillium wilt. Together with the co-expression data this confirms that *ViAve1-1*, *ViAve1-5* and *ViAve1-9* are unable to induce a Ve1-mediated immune response.

## **Discussion**

We have previously shown that the *Verticillium dahliae* effector *VdAve1* is recognized by the tomato immune receptor Ve1, and that homologs of *Ave1* identified in particular other plant pathogens that share between 52 and 95% similarity are differentially recognized by Ve1 (de Jonge et al. 2012). Only one homolog was found in each of those plant pathogens, except for *V. dahliae* where subsequent analysis has shown that most isolates contain an additional *Ave1*-like (*Ave1L*) allele that shares between 64 and 68% sequence similarity with *Ave1* (Chapter 6 of this thesis). In the genomes of the apple scab fungus *V. inaequalis* and the pear scab fungus *V. pirinia* largely expanded families of *Ave1* homologs were identified (Cooke et al. 2014). These homologs share between 51 and 63% sequence similarity with *Ave1*. Alignment of *VdAve1* and the *Venturia Ave1* homologs shows that there is no specific region of the protein that is highly conserved. In this study we showed that *Ave1* homologs derived from *V. inaequalis* and *V. pirina* are not recognized by Ve1. A reason for the lack of recognition might be the fact that *V. inaequalis* and *V. pirina* never co-evolved with Ve1 since tomato is not a host for these pathogens. Homologs of Ve1 have been found in many plant species, including functional homologs in various plants including tobacco, potato, eggplant and hop (Song et al. 2017a). Possibly, Ve1 homologs occur in apple and pear as well, that may potentially recognize one or more of the *Ave1* homologs derived from *V. inaequalis* and *V. pirina*, respectively. Possible recognition by an immune receptor can be tested by

performing a transient expression assay with *ViAve1-1*, *ViAve1-5* and *ViAve1-9* on apple or pear (Li et al. 2004; Vleeshouwers and Oliver 2014).

We also showed that *Ave1* homologs *ViAve1-1*, *ViAve1-5* and *ViAve1-9* derived from *V. inaequalis* that are highly expressed *in planta* cannot complement the virulence function of a *V. dahliae* *Ave1* deletion strain on tomato. Possibly, to exert its function, *Ave1* targets a host specific protein. If *ViAve1-1*, *ViAve1-5* and *ViAve1-9* have a similar function as *Ave1* and target a host-specific protein, this protein may be absent from the *Verticillium*-tomato interaction and therefore *Ave1* homologs cannot exert their function in other host plants. Host specificity of effectors has been described for the closely related oomycetes *Phytophthora infestans* and *P. mirabilis* that are pathogens on different hosts (Dong et al. 2014). Effector homologs derived from both pathogens are more efficient on their specific host target than on the foreign target. To test whether one or more of the *Ave1* homologs are contributing to the virulence of *V. inaequalis*, individual deletion strains of *ViAve1-1*, *ViAve1-5* and *ViAve1-9* should be generated or gene silencing of multiple genes should be performed (Fitzgerald et al. 2004; Fitzgerald et al. 2003) and analyzed for full virulence on their host plant apple. However, unfortunately, targeted gene deletion is not straightforward in *Venturia* spp.

## Materials and methods

### Plant material

Tobacco (*Nicotiana tabacum* cv. Petite Havana SR1) and tomato (*Solanum lycopersicum* cv. MoneyMaker) plants were grown in the greenhouse at 21°C/19°C during 16/8 hours day/night periods, respectively, with 70% relative humidity and 100 W/m<sup>2</sup> supplemental light when the light intensity dropped below 150 W/m<sup>2</sup>.

### Generation of expression constructs

The coding sequences (CDSs) of *ViAve1-1*, *ViAve1-2*, *ViAve1-4*, *ViAve1-5*, *ViAve1-8*, *ViAve1-9*, *ViAve1-10* and *VpAve1-1*, *VpAve1-2*, *VpAve1-3*, *VpAve1-4*, *VpAve1-5*, *VpAve1-6*, *VpAve1-7*, *VpAve1-11*, were amplified from genomic DNA of *Venturia inaequalis* and *V. pirina*, respectively. The CDS was then cloned into pENTR-D (Invitrogen, Carlsbad, California) and subsequently transferred to the Gateway-compatible destination vector pSol2092 (Zhang et al. 2013b) to generate expression constructs driven by the constitutive CaMV35S promoter for *Agrobacterium tumefaciens*-mediated transient expression. The CDSs of *ViAve1-1*, *ViAve1-5* and *ViAve1-9* were cloned into destination vector pFBT005 to generate expression constructs driven by the *VdAve1* promoter for expression in *V. dahliae*.

### ***Agrobacterium tumefaciens*-mediated transient expression**

*A. tumefaciens* GV3101 strains containing the expression constructs in pSol2092 were grown for two nights in liquid Yeast Extract Broth (YEB) and then resuspended to an OD<sub>600</sub> of 2 in infiltration medium as previously described (Zhang et al. 2013b). *A. tumefaciens* strains that carry the expression constructs for the *V. dahliae* and *Venturia Ave1* homologs and tomato *Ve1* were mixed in a 1:1 ratio and infiltrated in the lower side of 5- to 6 weeks-old tobacco leaves. At 5 dpi the leaves were inspected for the presence or absence of a hypersensitive response.

### **Complementation of *Verticillium dahliae* *Ave1* deletion strains**

The *A. tumefaciens* Agl1 strain containing the expression construct for *ViAve1-1*, *ViAve1-5* or *ViAve1-9* in pFBT005, respectively, driven by the *VdAve1* promoter, was used to transform a *V. dahliae* strain JR2 *Ave1* deletion strain (de Jonge et al. 2012) as previously described (Santhanam 2012). Genomic DNA was extracted from transformants to verify the presence of *ViAve1-1*, *ViAve1-5* or *ViAve1-9* and the presence of the nourseothricin cassette.

### ***V. dahliae* assay on tomato**

For *V. dahliae* inoculations 10-day-old tomato seedlings were uprooted, rinsed in water and then dipped for 5 minutes in a suspension of 10<sup>6</sup> conidiospores per mL of water harvested from 1- to 2-week-old *V. dahliae* cultures on PDA as previously described (Fradin et al. 2009). Control plants received the same treatment, but the roots were dipped in water without conidiospores. After replanting in fresh soil, plants were incubated at standard greenhouse conditions. Disease development was monitored up to 21 dpi. The canopy area of 8 plants was measured with ImageJ software and a One-Way ANOVA was performed with IBM SPSS statistics software.

### **Acknowledgements**

Work in the laboratory of BPHJT is supported by the Research Council Earth and Life Sciences (ALW) of the Netherlands Organization for Scientific Research (NWO). We thank Bert Essenstam and Pauline Sanderson of Unifarm for excellent plant care.

## References

- Bowen, J. K., Mesarich, C. H., Bus, V. G., Beresford, R. M., Plummer, K. M., and Templeton, M. D. 2011. *Venturia inaequalis*: the causal agent of apple scab. *Mol Plant Pathol* 12:105-122.
- Bus, V. G., Rikkerink, E. H., Caffier, V., Durel, C.-E., and Plummer, K. M. 2011. Revision of the nomenclature of the differential host-pathogen interactions of *Venturia inaequalis* and *Malus*. *Annu Rev Phytopathol* 49:391-413.
- Cook, D. E., Mesarich, C. H., and Thomma, B. P. H. J. 2015. Understanding plant immunity as a surveillance system to detect invasion. *Annu Rev Phytopathol*.
- Cooke, I. R., Jones, D., Bowen, J. K., Deng, C., Faou, P., Hall, N. E., Jayachandran, V., Liem, M., Taranto, A. P., Plummer, K. M., and Mathivanan, S. 2014. Proteogenomic analysis of the *Venturia pirina* (Pear Scab Fungus) secretome reveals potential effectors. *J Proteome Res* 13:3635-3644.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- Dong, S., Stam, R., Cano, L. M., Song, J., Sklenar, J., Yoshida, K., Bozkurt, T. O., Oliva, R., Liu, Z., Tian, M., Win, J., Banfield, M. J., Jones, A. M., van der Hoorn, R. A., and Kamoun, S. 2014. Effector specialization in a lineage of the Irish potato famine pathogen. *Science* 343:552-555.
- Fitzgerald, A., van Kan, J. A., and Plummer, K. M. 2004. Simultaneous silencing of multiple genes in the apple scab fungus, *Venturia inaequalis*, by expression of RNA with chimeric inverted repeats. *Fungal Genet Biol* 41:963-971.
- Fitzgerald, A. M., Mudge, A. M., Gleave, A. P., and Plummer, K. M. 2003. Agrobacterium and PEG-mediated transformation of the phytopathogen *Venturia inaequalis*. *Mycological research* 107:803-810.
- Fradin, E. F., and Thomma, B. P. H. J. 2006. Physiology and molecular aspects of Verticillium wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol Plant Pathol* 7:71-86.
- Fradin, E. F., Zhang, Z., Juarez Ayala, J. C., Castroverde, C. D., Nazar, R. N., Robb, J., Liu, C. M., and Thomma, B. P. H. J. 2009. Genetic dissection of Verticillium wilt resistance mediated by tomato Ve1. *Plant Physiol* 150:320-332.
- Fudal, I., Ross, S., Gout, L., Blaise, F., Kuhn, M., Eckert, M., Cattolico, L., Bernard-Samain, S., Balesdent, M., and Rouxel, T. 2007. Heterochromatin-like regions as ecological niches for avirulence genes in the *Leptosphaeria maculans* genome: map-based cloning of *AvrLm6*. *Mol. Plant Microbe Interact.* 20:459-470.
- Gessler, C., Patocchi, A., Sansavini, S., Tartarini, S., and Gianfranceschi, L. 2006. *Venturia inaequalis* resistance in apple. *CRC Crit Rev Plant Sci* 25:473-503.
- Grandaubert, J., Lowe, R. G., Soyer, J. L., Schoch, C. L., Van de Wouw, A. P., Fudal, I., Robbertse, B., Lapalu, N., Links, M. G., and Ollivier, B. 2014. Transposable element-assisted evolution and adaptation to host plant within the *Leptosphaeria maculans*-*Leptosphaeria biglobosa* species complex of fungal pathogens. *BMC Genomics* 15:1.
- Jones, J. D., and Dangl, J. L. 2006. The plant immune system. *Nature* 444:323-329.
- Kucheryava, N., Bowen, J. K., Sutherland, P. W., Conolly, J. J., Mesarich, C. H., Rikkerink, E. H., Kemen, E., Plummer, K. M., Hahn, M., and Templeton, M. D. 2008. Two novel *Venturia inaequalis* genes induced upon morphogenetic differentiation during infection and in vitro growth on cellophane. *Fungal Genet Biol* 45:1329-1339.
- Li, C., Sasaki, N., Isogai, M., and Yoshikawa, N. 2004. Stable expression of foreign proteins in herbaceous and apple plants using Apple latent spherical virus RNA2 vectors. *Archives of virology* 149:1541-1558.
- Rovenich, H., Boshoven, J. C., and Thomma, B. P. H. J. 2014. Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. *Curr Opin Plant Biol* 20:96-103.

- Santhanam, P. 2012. Random insertional mutagenesis in fungal genomes to identify virulence factors. Pages 509-517 in: Plant fungal pathogens. Methods in molecular biology, vol. 835. T. B. Bolton MD, ed. Humana press, Totowa.
- Shabi, E., Rotem, J., and Loebenstein, G. 1973. Physiological races of *Venturia pirina* on pear. Phytopathology 63:41-43.
- Shiller, J., Van de Wouw, A. P., Taranto, A. P., Bowen, J. K., Dubois, D., Robinson, A., Deng, C. H., and Plummer, K. M. 2015. A Large Family of *AvrLm6*-like Genes in the Apple and Pear Scab Pathogens, *Venturia inaequalis* and *Venturia pirina*. Front Plant Sci 6:980.
- Song, Y., Zhang, Z., Seidl, M. F., Majer, A., Jakse, J., Javornik, B., and Thomma, B. P. 2017a. Broad taxonomic characterization of *Verticillium* wilt resistance genes reveals ancient origin of the tomato Ve1 immune receptor. Mol Plant Pathol 18:195-209.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. Plant Cell 23:4-15.
- Vleeshouwers, V. G., and Oliver, R. P. 2014. Effectors as tools in disease resistance breeding against biotrophic, hemibiotrophic, and necrotrophic plant pathogens. Mol. Plant Microbe Interact. 27:196-206.
- Zhang, Z., Esse, H. P., Damme, M., Fradin, E. F., Liu, C. M., and Thomma, B. P. H. J. 2013a. Ve1-mediated resistance against *Verticillium* does not involve a hypersensitive response in *Arabidopsis*. Mol Plant Pathol 14:719-727.
- Zhang, Z., Fradin, E., de Jonge, R., van Esse, H. P., Smit, P., Liu, C.-M., and Thomma, B. P. H. J. 2013b. Optimized agroinfiltration and virus-induced gene silencing to study Ve1-mediated *Verticillium* resistance in tobacco. Mol. Plant Microbe Interact. 26:182-190.



**An *Ave1*-like effector displays  
allelic variation within the  
*Verticillium dahliae* population**

**Chapter 6**

Jordi C. Boshoven  
Hanna Rovenich\*  
Yin Song\*  
Luigi Faino  
Michael F Seidl#  
Bart PHJ Thomma#

#These authors contributed equally

\*These authors contributed equally

To be submitted for publication

## Abstract

*Verticillium dahliae* is a soil-borne fungal pathogen that causes vascular wilt disease on a broad range of host plants. The tomato immune receptor Ve1 is able to recognize race 1 strains of *V. dahliae* that secrete the Ave1 effector. We identified a gene in *V. dahliae* strain JR2 with high similarity to the Ave1 effector gene that we called Ave1L (for Ave1-like). Subsequently, we identified in total five alleles (Ave1L1 to Ave1L5) in the *V. dahliae* population. While some of the *V. dahliae* strains do not have Ave1L, other strains have one of the five Ave1L alleles in presence or absence of Ave1. The Ave1L alleles contain multiple single nucleotide polymorphisms that result in a stop codon in some of the alleles. In addition, one of the alleles contains a retrotransposon. These observations suggest that Ave1L experienced significant selection pressure, potentially to evade recognition by an immune receptor. Thus, we attempted to predict the common progenitor sequence of the five Ave1L alleles. However, we show that neither of the Ave1L alleles, nor the progenitor, are recognized by the tomato immune receptor Ve2 that shows high homology to Ve1. We furthermore show that Ave1L2 is expressed during host colonization but does not seem to contribute to virulence on tomato. Since *V. dahliae* has a wide host range, Ave1L potentially contributes to virulence on other host plants than tomato where it may have been recognized by an immune receptor that drove its divergence.

## Introduction

Plants can sense invading microbial pathogens through the recognition of conserved microbe-associated molecular patterns (MAMPs) such as bacterial flagellin, fungal chitin and oomycete  $\beta$ -glucans (Dodds and Rathjen 2010; Thomma et al. 2011). Recognition of MAMPs leads to the activation of immune responses to stop pathogen ingress. To gain further access to the host, pathogens produce effectors; secreted molecules that support host colonization. Although effectors may contribute to host colonization in many ways, most effectors that have been functionally characterized up to now specifically target host immune responses (Rovenich et al. 2014). In response to pathogen exploitation of effectors, plants evolved immune receptors that can monitor (the activity of) specific effectors to re-initiate immune responses, typically referred to as effector-triggered immunity (ETI) (Dodds and Rathjen 2010; Thomma et al. 2011). To overcome effector recognition, pathogens in turn may lose the recognized effector, evolve mutants that are no longer recognized or, alternatively, evolve novel effectors that can suppress the immune response. As a result of all this, plant immunity evolved to a robust system to detect invasion in which immunogenic molecules form a continuum and can be host- or microbe-derived and can have various intrinsic functions. Ultimately host receptors, termed invasion pattern receptors (IPRs), detect externally encoded as well as modified-self ligands, termed invasion patterns, that betray microbial invasion (Cook et al. 2015; Thomma et al. 2011).

The *Cladosporium fulvum*-tomato interaction is a well-established fungal model system that has been used to study effector-receptor interactions (de Wit 2016; Thomma et al. 2005). The first fungal avirulence gene was cloned from *C. fulvum* and to date a dozen additional effector genes have been cloned most of which act as avirulence genes (Bolton et al. 2008; Joosten et al. 1994; Lauge et al. 2000; Luderer et al. 2002; Mesarich et al. 2014; Stergiopoulos et al. 2007; Van den Ackerveken et al. 1993; van Kan et al. 1991; Westerink et al. 2004). As a consequence, variation within a fungal effector was also first reported in *C. fulvum* for which isoforms of the Avr4 effector were identified that are no longer recognized by the corresponding tomato immune receptor Cf-4 as a result of a frameshift mutation or a single point mutation in the Avr4 gene that induces a single amino acid change (Joosten et al. 1997; Joosten et al. 1994). Other alterations were observed in *C. fulvum* effector Avr2 where, besides a frameshift mutation, the introduction of a premature stop codon or a retrotransposon in the gene results in a truncated protein that is no longer recognized by the corresponding tomato immune receptor Cf-2 (Luderer et al. 2002). In the case of *C. fulvum* effectors Avr4E and Avr9 some alleles contain point mutations while in other strains the complete gene is



deleted to avoid recognition by the corresponding tomato immune receptor Cf-4E and Cf-9, respectively (van Kan et al. 1991; Westerink et al. 2004). Importantly, while effector deletion or mutation can overcome effector recognition in plants carrying the corresponding immune receptor, such modification may compromise virulence on plants that lack this immune receptor. Although not yet observed in *C. fulvum*, pathogens may also evolve novel effectors to suppress recognition of a particular effector. This has been demonstrated for Avr1 of the tomato pathogen *Fusarium oxysporum* f. sp. *lycopersici* that can suppress immune responses that are activated by the tomato immune receptors I-2 and I-3 that recognize the Avr2 and Avr3 effectors, respectively (Houterman et al. 2008). Collectively, these examples illustrate that pathogens have evolved many ways to circumvent recognition by immune receptors of their host plants.

*Verticillium dahliae* is a soil-borne vascular pathogen and causal agent of vascular wilt disease in a broad range of plant species (Fradin and Thomma 2006). In contrast to the situation for *C. fulvum* for which many complex races have been described on tomato, the situation for *V. dahliae* is simple on that same host due to the existence of only a single dominant resistance source that divides the population in only two races. On tomato, race 1 strains of *V. dahliae* are contained by the Ve1 immune receptor whereas race 2 strains are not recognized (Fradin et al. 2009; Kawchuk et al. 2001). Recently, the race 1-specific effector that is recognized by Ve1 was identified as Ave1 (for Avirulence on Ve1 tomato) (de Jonge et al. 2012). Interestingly, it was found that all race 2 strains that were analysed lack the complete Ave1 gene. Moreover, it was shown that Ave1 contributes to the virulence of *V. dahliae* on plants that lack Ve1. Intriguingly, the Ave1 CDS is completely identical in all 85 race 1 strains that have been analyzed to date, suggesting that sequence conservation is important for virulence (de Jonge et al. 2012). Intriguingly, many homologs of Ave1 are found in plants as well as in a few plant pathogens (de Jonge et al. 2012). Most of the plant homologs are annotated as plant natriuretic peptides (PNPs) or expansin-like proteins that are thought to be involved in water and ion homeostasis and therefore in many fundamental plant processes including growth, net water uptake, photosynthesis, stomatal opening and gas exchange (Gehring and Irving 2003; Ludidi et al. 2004; Maryani et al. 2001; Pharmawati et al. 2001). Interestingly, the Ave1 homolog of the bacterial pathogen *X. axonopodis*, known as XacPNP, acts as a virulence factor by affecting homeostasis and photosynthesis in citrus to stimulate bacterial proliferation (Garavaglia et al. 2010; Gottig et al. 2008). The Ave1 homologs that occur in plant pathogens do not follow the phylogeny of the species in which they occur, which is interpreted as a sign of horizontal gene transfer, leading to believe that the microbial Ave1 homologs have been acquired from plants (de Jonge et al. 2012; Gottig et al. 2008)

In this study, a BLAST search using *Ave1* against the recently finished gapless genome of *V. dahliae* strain JR2 (Faino et al. 2015) revealed an *Ave1*-like gene. Subsequent BLAST searches identified allelic variation in a population of *V. dahliae* strains.

## Results

### Identification of an *Ave1*-like gene in *Verticillium dahliae*

Using the genome of *V. dahliae* strain JR2 (Faino et al. 2015), a BLAST analysis using *Ave1* as a query identified a gene with 67% similarity that we will further refer to as *Ave1L* (for *Ave1*-like). Intriguingly, while *Ave1* encodes a protein of 134 amino acids (de Jonge et al. 2012), the CDS of *Ave1L* likely encodes a protein of only 24 amino acids due to a predicted stop codon. To determine whether *Ave1L* can be found in other *V. dahliae* strains as well, we performed BLAST analyses using *Ave1L* as a query on twenty recently sequenced *V. dahliae* genomes (Table 1) (de Jonge et al. 2012; Faino et al. 2015; Klosterman et al. 2011). While no *Ave1L* sequence was found in four genomes, the genomes of 16 *V. dahliae* strains were each found to carry a single *Ave1L* copy that displays 93-100% similarity to the initially identified *Ave1L*, while displaying 65-68% similarity to *Ave1*. Collectively, five different *Ave1L* alleles were identified, designated *Ave1L1* to *Ave1L5*. Following the phylogeny of the *V. dahliae* strains, *Ave1L1* was found in five strains that include JR2 and VdLs17 (de Jonge et al. 2012; Faino et al. 2015; Klosterman et al. 2011), *Ave1L2* in three strains, *Ave1L3* in five strains, *Ave1L4* in one strain and *Ave1L5* in two strains (Table 1, Figure 1B). Similar to the initially identified *Ave1L1*, also *Ave1L3* and *Ave1L4* are predicted to encode proteins of 24 amino acids. In contrast, *Ave1L2* and *Ave1L5* are predicted to encode proteins of 134 amino acids similar to *Ave1* (Figure 1A). Interestingly, the presence of *Ave1L* alleles did not correlate with presence of *Ave1*, as *Ave1L* alleles were found in race 1 as well as in race 2 strains (Table 1, Figure 1B).

Like *Ave1*, *Ave1L* shows similarity to plant natriuretic peptides (PNPs) or expansin-like proteins (de Jonge et al. 2012). To analyse whether *Ave1L* carries a conserved PNP domain as has been characterized in *Arabidopsis thaliana* PNP (AtPNP-A) (Ludidi et al. 2004; Morse et al. 2004) the amino acid sequences of *Ave1*, *Ave1L* and various PNPs were aligned to AtPNP-A and similarity was determined (Fig. 1C). Intriguingly, there does not seem to be a particularly conserved region among the proteins, and also the region that aligns with the PNP domain of AtPNP-A is not particularly conserved (Fig. 1C). The bacterial pathogen *X. axonopodis* secretes a PNP-like effector known as *XacPNP* that modulates host responses to facilitate bacterial proliferation on citrus. *XacPNP* comprises a 12 amino acid peptide that

displays a high degree of similarity to the PNP domain of AtPNP-A, and that exerts the biological activity of XacPNP (Gottig et al. 2008). However, Ave1L does not carry this 12 amino acid peptide, in contrast to Ave1 and the other PNPs included in the alignment (Figure 1C) (Gottig et al. 2008). This finding suggests that Ave1L does not function as a natriuretic peptide.

**TABLE 1 | Ave1L alleles identified in sequenced *V. dahliae* genomes**

Strain	Origin	Isolated from	Race**	Ave1	Ave1L allele	Number and position of premature stop codon(s)
JR2 <sup>\$</sup>	Canada	<i>Solanum lycopersicum</i>	1	Yes		
VdLs17 <sup>Δ</sup>	USA	<i>Lactuca sativa</i>	2	No		
St16.01*	Syria	<i>Gossypium spp</i>	2	No	Ave1L1	1 (25W)
Vd57*	Germany	<i>Fragaria x ananassa</i>	2	No		
V <sup>#</sup>	Netherlands	<i>Chrysanthemum spp.</i>	2	No		
CBS38166 <sup>\$</sup>	Canada	<i>Solanum lycopersicum</i>	1	Yes		
St14.01 <sup>\$</sup>	USA	<i>Pistacia vera</i>	1	Yes	Ave1L2	No
DVD-S26 <sup>\$</sup>	Canada	Soil	2	No		
DVD-3 <sup>\$</sup>	Canada	<i>Solanum tuberosum</i>	2	No		
DVD-31 <sup>\$</sup>	Canada	<i>Solanum lycopersicum</i>	2	No		
DVD-S94 <sup>\$</sup>	Canada	Soil	2	No	Ave1L3	4 (25W, 57Q, 120Q, 128Q)
DVD-161 <sup>\$</sup>	Canada	<i>Solanum tuberosum</i>	2	No		
JKG8 <sup>#</sup>	Netherlands	<i>Solanum tuberosum</i>	2	No		
DVD-S29 <sup>\$</sup>	Canada	Soil	2	No	Ave1L4	1 (25W)
St.100 <sup>μ</sup>	Belgium	Soil	2	No	Ave1L5	No
463*	Mexico	<i>Gossypium spp.</i>	2	No		
2009-605*	Ukraine	<i>Capsicum annuum</i>	1	Yes		
Vd152*	Hungary	<i>Quercus spp</i>	2	No	No	x
Vd52*	Austria	<i>Capsicum annuum</i>	1	Yes		
Vd39*	Germany	<i>Helianthus annuus</i>	2	No		

\* newly sequenced *V. dahliae* genomes


\*\* The race nomination in *V. dahliae* is restricted to strains that are pathogenic on Ve1 tomato in which race 1 strains are contained and race 2 strains are not.

\$ de Jonge et al., 2012


Δ Klosterman et al., 2011

# Ispahani et al., 2008

μ de Jonge et al., 2013



**FIGURE 1 | The identification of five *Ave1L* alleles.** (A) Alignment of predicted *Ave1L* proteins *Ave1L1* to *Ave1L5*. The black box indicates the predicted signal peptide, while asterisks (\*) indicate predicted stop codons and the hashtag (#) indicates discontiguity. (B) Phylogenetic tree based on the genomes of *V. dahliae* strains and the presence of *Ave1* and *Ave1L* alleles in the respective genomes. (C) continues at next page.



**FIGURE 1 | The identification of five *Ave1L* alleles.** (C) Alignment of AtPNP-A, XacPNP, PNPs (*Nicotiana tomentosiformis* Nt; *N. sylvestris* Ns; *Solanum lycopersicum* Sl; *S. tuberosum* St; *Vitis vinifera* Vv), Ave1 and Ave1L. The purple colour indicates amino acid conservation. The black box indicates the active PNP domain of XacPNP (Gottig et al. 2008).


### ***Ave1L* alleles occur in a highly dynamic, lineage-specific, genomic region**

It was previously shown that extensive chromosomal rearrangements occur among strains of *V. dahliae* (de Jonge et al. 2013). The occurrence of these rearrangements coincides with the occurrence of lineage-specific (LS) regions that are only present in sub-sets of *V. dahliae* strains and that are significantly enriched for *in planta*-expressed genes that also include effectors (de Jonge et al. 2013). Consequently, it has been suggested that chromosomal rearrangements facilitate adaptive evolution in *V. dahliae*, giving rise to lineage-specific segmental duplications that may evolve new traits (de Jonge et al. 2013; Faino et al. 2015; Seidl and Thomma 2014). Since *Ave1L* does not occur in all *V. dahliae* strains, the gene likely occurs in an LS region. To confirm this, we aligned reads from the sixteen *Ave1L*-carrying *V. dahliae* strains to the gapless genome of *V. dahliae* strain JR2 (Faino et al. 2015). We indeed found that the five *Ave1L* alleles are positioned in the same genomic environment in a repeat-rich LS region (Figure 2A).

### ***Ave1L3* is disrupted by a transposable element**

We noticed that each *Ave1L3* allele was consistently located on two contigs, which suggests discontinuity of this particular allele (Figure 2B). To confirm this *in silico* observation, we designed PCR primers on conserved sequences that are present in all *Ave1L* alleles, spanning the predicted site where the discontinuity occurs, as well as PCR primers that amplify a conserved region up- and downstream of this site. We used these primers on genomic DNA of a panel of *V. dahliae* strains that represent the five different *Ave1L* alleles. The primers spanning the site of the observed discontinuity in *Ave1L3* amplified a 200 bp PCR product from strains VdLs17 (*Ave1L1*), Dvd-S26 (*Ave1L2*), Dvd-S29 (*Ave1L4*) and St.100 (*Ave1L5*), while no PCR product was amplified from the *Ave1L3*-carrying strains DVD-3, DVD-31, DVD-S94, DVD-161. However, primers that were targeted to a region upstream and downstream of the predicted discontinuity amplified a product from all strains, including those carrying *Ave1L3* (Figure 2C). These results confirm that *Ave1L3* is discontinuous in the genome of all strains that carry this allele.


Discontinuity of *Ave1L3* may be caused by the insertion of a transposable element or by a gross chromosomal rearrangement that occurred within the allele. To test whether *Ave1L3* is disrupted by an insertion we repeated the PCR with primers spanning the site of the discontinuity using a significantly enhanced elongation time to allow for amplification of considerably larger PCR fragments. We used an *Ave1L1*-carrying *V. dahliae* strain as a control for absence of discontinuity. A fragment of around 7 kb was amplified from *V. dahliae* strain DVD-3 that carries *Ave1L3* (Figure 2D). To determine the sequence of the PCR fragment we used an Oxford Nanopore MinION sequencer. Subsequently, we obtained a consensus sequence from the reads and performed a BLAST analysis. This analysis revealed that *Ave1L3* is interrupted by a long terminal repeat retrotransposon that is classified as VdLTRE3 (Faino et al. 2016).



**FIGURE 2 | The genomic context of the *Ave1L* alleles.** (A) Each coloured bar represents the genomic context of *Ave1L* in different *V. dahliae* strains. The red arrow indicates the position of the *Ave1L* gene while the black bars indicate other genes and repeats in the region. (B) Genomic region flanking *Ave1L* and the read coverage in *V. dahliae* strain VdLs17 and DVD-161. The horizontal red and blue arrows indicate the position of *Ave1L*. The black arrows indicate discontinuity of *Ave1L3* in DVD-161. (C) Amplification products on gel using primers spanning, are upstream, and downstream of (\*) the predicted side. The absence of an amplification product on gel when primers were used spanning the predicted side where discontinuity occurs confirms discontinuity in *V. dahliae* strains that carry *Ave1L3*. (D) The presence of an amplification product (left lane) confirms that *Ave1L3* contains an insert of approximately 7 kb deduced from the 10 kb ladder that was run alongside (right lane).

### Ave1L2 is expressed during host colonization

To assay whether any of the *Ave1L* alleles is expressed during host colonization, we inoculated Arabidopsis plants with *V. dahliae* strains VdLs17 (*Ave1L1*), DVD-S26 (*Ave1L2*), DVD-S94 (*Ave1L3*), DVD-S29 (*Ave1L4*) and St.100 (*Ave1L5*) and expression of the *Ave1L* alleles was monitored with real-time PCR at 14 and 21 days post inoculation, showing that only *Ave1L2* is expressed *in planta* (Figure 3). To further confirm this observation, we monitored the expression of *Ave1L2* in another strain as well. To this end, we used the race 1 strain St14.01 that carries *Ave1L2* in addition to *Ave1*, in contrast to the race 2 strain DVD-S26 that carries *Ave1L2* in absence of *Ave1*. Interestingly, also in this strain *Ave1L2* was found to be expressed (Figure 3).




**FIGURE 3 | *Ave1L2* is expressed by *V. dahliae* on Arabidopsis.** *Ave1L* expression in *V. dahliae* on Arabidopsis at 14 and 21 dpi after mock treatment or inoculation with *V. dahliae* strain VdLs17, St14.01 DVD-S26, DVD-S94, DVD-S29 or St.100.

### *Ave1L2* is not a virulence factor on tomato plants

Since *Ave1L2* is expressed *in planta* we tested whether *Ave1L2* is a virulence factor of *V. dahliae*. To this end, we deleted *Ave1L2* from the genomes of *V. dahliae* strains St14.01 and CBS38166 (Supplemental Figure 1). We then inoculated three transformants of St14.01 and four transformants of CBS38166 on tomato plants and tested their aggressiveness in comparison to the corresponding wild type *V. dahliae* strain. However, tomato plants inoculated with the *Ave1L2* deletion strains showed a similar level of stunting as the corresponding wild type strains (Figure 4A;B). Moreover, a similar amount of fungal biomass was detected in tomato upon inoculation with the *Ave1L2* deletion strains as well as with the corresponding wild type strains (Figure 4C). Therefore, we conclude that *Ave1L2* does not contribute to the virulence of *V. dahliae* on tomato.





**FIGURE 4 | *Ave1L2* does not markedly contribute to virulence on tomato.** (A) Tomato plants inoculated with *Ave1L2* deletion strains (#) and the corresponding wild strain (St14.01 or CBS38166). (B) The average canopy area of inoculated tomato plants at 14 dpi (n=10). Error bars represent standard errors. (C) Data points indicate the accumulated fungal biomass in tomato plants at 14 dpi. The horizontal line indicates the average fungal biomass.

### ***Ave1L* proteins are not recognized by *Ve2***

*Ve1* is the only plant immune receptor known to confer resistance to *V. dahliae* (Fradin et al. 2009; Kawchuk et al. 2001). In tomato, the *Ve1* gene is located at the *Ve* locus, together with *Ve2*, a gene that similarly encodes a receptor-like protein and that has high (~85%) sequence similarity to *Ve1* (Fradin et al. 2009). *Ve2* is expressed similarly as *Ve1*, but the encoded protein does not recognize *Ave1* and thus far its function remains unknown (de Jonge et al. 2012; Fradin et al. 2009). The occurrence of *Ave1L* alleles that encode proteins that display considerable sequence divergence, or that are severely truncated and thus likely not functional due to premature stop codons in the coding sequences, suggests that the progenitor of these alleles has experienced significant selection pressure, potentially to avoid recognition by a plant immune system. Although such selection pressure may have occurred in any plant species, considering the broad host range

of *V. dahliae*, we tested the possibility that Ve2 was able to recognize ancestral forms of Ave1L. To this end, a library of potentially ancestral Ave1L alleles was constructed for co-expression with Ve2 in *N. tabacum*. The predicted non-functional Ave1L alleles Ave1L1, Ave1L3 and Ave1L4 were chosen to reverse the disruptive mutations using site-directed mutagenesis to yield the derivatives Ave1L1\*, Ave1L3\* and Ave1L4 by replacing the nucleotides that are responsible for generating the stop codons by nucleotides that are present at the same positions in Ave1L2 and Ave1L5, respectively (Figure 5A and B). As Ave1L3 contains multiple mutations and a retrotransposon, we synthesized Ave1L3\* in which the stop codons were replaced and the retrotransposon was removed (Figure 5A and B). While each of the artificial derivatives (Ave1L1\*, Ave1L3\* and Ave1L4\*) is derived from only one of the current Ave1L alleles in which only stop codons and the retrotransposon were replaced, we aimed to determine their most likely common ancestor based on the observed substitutions in the current Ave1L alleles. Therefore, a consensus sequence was generated from the alignment of Ave1L alleles, leading to replacement of the various point mutations by the most frequently occurring nucleotide at the same position in the various Ave1L alleles, that we synthesized as Ave1L\*\* (Figure 5A and B). To test the hypothesis that Ve2 potentially recognized any of these potentially ancestral forms of Ave1L, we co-expressed the artificially generated alleles with Ve2 and screened for the occurrence of a hypersensitive response. However, at five days post infiltration, no necrosis was observed in leaf tissue where Ve2 was co-expressed with either Ave1L\*\*, Ave1L1\*, Ave1L2, Ave1L3\*, Ave1L4\*, Ave1L5, as well as with the negative control Ave1 (Figure 5C). As expected, in leaf tissue where Ve1 was co-expressed with Ave1 as a control, clear necrosis was observed (Figure 5C). Moreover, none of the Ave1Ls is recognized by Ve1 (Figure 5C). Therefore, we conclude that it is unlikely that Ve2 recognized an Ave1L progenitor allele, and was thus responsible for its diversification.



## Discussion

In this study we identified a gene in *V. dahliae* with high similarity to the previously characterized *V. dahliae* effector *Ave1* (de Jonge et al. 2012) that we named *Ave1L*. BLAST analyses of this gene in twenty sequenced *V. dahliae* strains revealed four additional alleles. These alleles follow the phylogenetic distribution of *V. dahliae* strains as previously described (de Jonge et al. 2013). Some strains do not have an *Ave1L* allele while other strains have a single *Ave1L* allele. Intriguingly, presence or absence of *Ave1L* does not correspond to presence or absence of *Ave1*. Particular strains contain *Ave1* but no *Ave1L* allele. Moreover, several of the alleles contain point mutations, including mutations that cause stop codons, suggesting they experienced selection pressure to inactivate the gene product. It is tempting to speculate that this pressure arose to avoid recognition by an immune receptor. However, unequivocal evidence for this hypothesis, including knowledge of the nature of the immune receptor or the host plant that carried it, remains absent today.

Since we found that *Ave1L2* is expressed during host colonization we hypothesized that *Ave1L2* is a virulence factor. However, *Ave1L2* deletion strains of *V. dahliae* did not display attenuated virulence on tomato. The *Ave1L2* deletion strains were generated in two *V. dahliae* race 1 strains that also contain *Ave1*. Since *Ave1* makes a major contribution to virulence it cannot be excluded that redundancy in virulence functions obscures the identification of virulence defects upon *Ave1L2* deletion. Unknowingly, the virulence function of *Ave1* was unravelled in the JR2 strain that does not carry a functional *Ave1L* homolog, as the *Ave1L1* homolog that is found in this strain is inactivated by an early stop codon. It can thus be explained that a potential redundancy in virulence function between *Ave1* and *Ave1L* remained unnoticed thus far. Therefore, deletion strains should be generated in *V. dahliae* race 2 strain DVD-S26 to test the contribution to virulence in absence of *Ave1*.

Some virulence factors have previously been shown to be host-specific, which is particularly relevant for broad host-range pathogens such as *V. dahliae*. For example, the *V. dahliae* effector NLP2 acts as a virulence factor on *Arabidopsis* but not on *Nicotiana benthamiana* (Santhanam et al. 2013). Similarly, the EPIC effectors of *Phytophthora infestans* and *P. mirabilis* are only functional on their respective plant hosts (Dong et al. 2014). Arguably, *Ave1L2* may act as a virulence factor on other host plants of *V. dahliae*. Potent virulence factors pose a strong selection pressure on host plants to evolve recognition (Cook et al. 2015; Thomma et al. 2011). Therefore, *Ave1L2* is likely to act as a virulence factor on the host where recognition occurs.

Ave1 is recognized by the tomato immune receptor Ve1 that is encoded by a gene at a locus that also encodes the highly identical orphan receptor Ve2. Considering the high degree of identity between the Ave1 homologs, we tested the possibility that Ve2 recognized the progenitor of the current Ave1L alleles or any of the current alleles. However, the current Ave1L alleles nor the predicted progenitor or any of the intermediate alleles appears to be recognized by Ve2. We therefore conclude that Ve2 does not recognize Ave1L. The fact that Ve2 does not recognize Ave1L is not surprising since we do not know whether Ave1L was recognized in tomato in the first place. Moreover, if recognition occurred in tomato this may have been mediated by any of the hundreds of potential immune receptors that are encoded in the genome. However, it is not unlikely that recognition occurred in another host than tomato since *V. dahliae* has a very wide host range. The observation that Ave1L2 does not contribute to virulence of *V. dahliae* on tomato supports the latter hypothesis.

Ave1L and Ave1 are both located in highly dynamic lineage-specific regions of the genome, which could have contributed to the duplication of one of the two genes resulting in the other one. It has been suggested that Ave1 was acquired from plants through horizontal gene transfer (de Jonge et al. 2012) and then was lost multiple times to result in the current population of race 2 strains (Faino et al. 2016). Ave1 shows higher sequence similarity to PNPs than Ave1L and the active PNP site appears to be degenerated and no longer functional in Ave1L (Figure 1C), whereas the PNP site in Ave1 is conserved. The sequence of Ave1 is conserved in the population and either present or absent, whereas the Ave1L alleles display variation that includes various mutations in addition to presence-absence polymorphisms. Thus, for Ave1 it has been suggested that sequence conservation is essential for its function (de Jonge et al. 2012). In addition, Ave1L alleles are present in both *V. dahliae* race 1 and race 2 strains. Collectively, these observations suggest that selection pressure on Ave1 is more recent than selection pressure on Ave1L and argues that not Ave1 but Ave1L was acquired from plants first.

## Materials and methods

### *Verticillium dahliae* genomics

Genome sequence data in this study was partly obtained from previous studies (de Jonge et al. 2013; de Jonge et al. 2012; Faino et al. 2015), and additionally contains nine newly sequenced *V. dahliae* genomes (Table 1). Library preparation (500 bp insert size) and genomic sequencing (100 bp paired-end reads) were conducted at the Beijing Genome Institute (BGI, Hong Kong, China). The genome of each individual *V. dahliae* strain was assembled using the A5 pipeline (Tritt et al. 2012). Discontiguity in a single Ave1L allele was analyzed by mapping the genomic

paired-end sequences to the *V. dahliae* strain JR2 reference genome using BWA (Faino et al. 2015; Li and Durbin 2009).

The genetic diversity and the population structure of the sequenced *V. dahliae* strains was assessed using the reference sequence alignment based phylogeny builder (REALPHY) v. 1.11 (Bertels et al. 2014). Briefly, short paired-end genomic reads were mapped against the reference genome (*V. dahliae* strain JR2 (Faino et al. 2015)) using Bowtie2 (Langmead and Salzberg 2012). Polymorphic as well as non-polymorphic sites were extracted (per base quality [20], coverage [10] and polymorphism frequency [0.95]) and sites that were present in all species were retained. With this pseudo-molecule, a maximum likelihood phylogenetic tree was inferred using PhyML (Guindon et al. 2009) with the generalized time reversible (GTR) nucleotide substitution model, and the robustness of the phylogeny was assessed by 500 bootstrap replicates.

### **Ave1L sequence analysis**

To identify *Ave1L* sequences in the twenty *V. dahliae* genome assemblies, we used the CDS of *Ave1* as a BLASTn query (Altschul et al. 1990). Sequence alignments of the five unique *Ave1L* alleles were performed online by using Clustal Omega (EMBL-EBI, Hinxton, UK), and locally using the software package MEGA6 (Tamura et al. 2013).

Phylogeny analyses of *Ave1*-like homologs in plant pathogenic microbes and in plants was conducted using MrBayes (Huelsenbeck and Ronquist 2001). *Ave1*-like homologs in plant pathogenic microbes were previously identified, and plant homologs were identified by sequence similarity search (BLASTp) against the predicted proteomes deposited in Phytosome v9.1 (Goodstein et al. 2012), supplemented by the proteome from *Amborella trichopoda*, and spurious sequence hits were subsequently manually filtered. Protein sequences were subsequently aligned using MAFFT (Kato et al. 2002), and the alignment was curated using trimAl by removing alignment positions that contained >5 % gaps (Capella-Gutiérrez et al. 2009). Phylogenetic analysis was performed using MrBayes (Huelsenbeck and Ronquist 2001) with the Whelan and Goldman amino acid substitution matrix, and the gamma-shaped rate categories with a proportion of invariable sites were estimated using four rate categories. Markov Chain Monte Carlo (Marjoram et al. 2003) was run with two independent runs of four chains ( $T = 0.1$ ) and a sample frequency of 500 for 40,000,000 generations. At this point, the average standard deviation of split frequency, a measure of convergence, was well below 0.02. The first 25% of samples of each run were discarded as burn-in, and the remaining samples were used to infer tree and branch topology.

### **Ave1L gene expression and gene deletions**

*Arabidopsis* (*Arabidopsis thaliana*) ecotype Col-0 and tomato (*Solanum lycopersicum*) cultivar MoneyMaker were inoculated with *V. dahliae* strains VdLs17 (containing *Ave1L1*), DVD-S26 (*Ave1L2*), St14.01 (*Ave1L2*), DVD-S94 (*Ave1L3*), DVD-S29 (*Ave1L4*) and St.100 (*Ave1L5*) as previously described (Fradin et al. 2011; Fradin et al. 2009). The stems of tomato plants were cut from the base to the first cotyledons and flash frozen in liquid nitrogen. The base of *Arabidopsis* plants of which the leaves were removed were harvested and flash frozen in liquid nitrogen. Plant material was ground to a powder and RNA was extracted using Quick-RNA™ Miniprep (Zymo Research Europe GmbH, Freiburg, Germany) and cDNA was synthesized using M-MLV Reverse Transcriptase (Promega Benelux BV, Leiden, The Netherlands). Real-time PCR was conducted using an ABI7300 PCR machine (Applied Biosystems, Life Technologies Europe BV, Bleiswijk, The Netherlands) in combination with the qPCR SensiMix kit (BioLine, GC Biotech BV, Alphen aan den Rijn, The Netherlands). Primers that were used for gene expression and fungal biomass are described in supplemental table 2. Real-time PCR conditions were as follows: an initial 95°C denaturation step for 10 minutes followed by denaturation for 15 seconds at 95°C, annealing for 60 seconds at 60°C, and extension at 72°C for 40 cycles. Gene expression and fungal biomass was analyzed using the software package 7300 System Software (Applied Biosystems, Life Technologies Europe BV, Bleiswijk, The Netherlands). To generate *Ave1L2* deletion strains, primers were designed to amplify approximately 1500 bp up- and downstream of the *Ave1L2* CDS (Supplemental table 2). Both amplicons were used to generate a USER friendly cloning construct to replace *Ave1L2* by a hygromycin cassette through homologous recombination (Frandsen et al. 2008).

### **Production of derived Ave1L alleles**

The potential progenitor of the *Ave1L* alleles was obtained by generating a consensus sequence from an alignment of the CDS of *Ave1L* alleles and was synthesized as *Ave1L\*\**. The predicted derivatives of *Ave1L1* and *Ave1L4* were obtained by replacing the predicted premature stop codons. To this end primers were designed (Supplemental table 2) to replace 1 nucleotide in the stop codon with a nucleotide at the same position in *Ave1L2* and *Ave1L5* that do not contain a predicted premature stop codon. Overlap PCR was performed to introduce the mutation in *Ave1L1* and *Ave1L4*. The *Ave1L1* and *Ave1L4* sequences without a predicted premature stop codon were designated as *Ave1L1\** and *Ave1L4\**, respectively (Figure 4A). The derivative of *Ave1L3* was obtained by synthesizing the gene without stop codons and without the retrotransposon and was designated as *Ave1L3\** (Eurofins Genomics, Ebersberg, Germany) (Figure 4A).

### Functional analysis of *Ave1L* alleles

*V. dahliae* strains were grown on potato dextrose agar (PDA) (Thermo Fisher Scientific Inc, Breda, The Netherlands) for 1-2 weeks and conidia were harvested for DNA extraction with high-salt CTAB buffer (100 mM TrisHCl, 20 mM EDTA, 2 M NaCl, 3% CTAB). PCR primers were designed to amplify *Ave1*, *Ave1L* or to detect discontinuity (Supplemental table 2). For constitutive expression the CDS of *Ave1L* alleles was amplified from genomic DNA and cloned in the adjusted, Gateway compatible, pBIN vector variant pSol2092 (Zhang et al. 2013) and transformed into *Agrobacterium tumefaciens* strain GV3101. *Nicotiana tabacum* cv. Petite Havana SR1 was infiltrated in a 1:1 ratio with *A. tumefaciens* that express either *Ve1* or *Ve2* and an *Ave1(-like)* construct, as previously described (Zhang et al. 2013). After infiltration, plants were transferred to a climate chamber and incubated at 22°C and 19°C during 16-h day and 8-h night periods, respectively, with 70% relative humidity. At 5 dpi leaves were inspected for necrosis development.

### Oxford Nanopore MinION sequencing

Library preparation of the PCR fragment was performed according to the protocol of Oxford Nanopore (<https://wiki.nanoporetech.com/display/BP/Nanopore+Sequencing+Kit+protocol+SQK-MAP005>) skipping the DNA fragmentation step. The library was loaded on the Nanopore flowcell and the sequencing was stopped after ~4 hr. The run generated about 870 high quality long reads with an average length of 3,869 bp and the longest read of about 14 kb. Using Nanocorrect software (<https://github.com/jts/nanocorrect>), we used all reads to correct the longest 50 reads. Out of the 50 longest reads, only 28 were corrected to generate a consensus. The average protected read length was of 5,666.8 bp with a median of 6,483 bp. All the reads were used in BLAST analysis to confirm the presence of *Ave1L3* fragments at both ends. Out of the 28 corrected fragments, eight showed high similarity (~99%) to *Ave1L3* at both ends.

### Acknowledgements

We thank Bert Essenstam and Pauline Sanderson at Unifarm for excellent plant care. We thank Jose Espejo Valle-Inclan for assistance with the Oxford Nanopore MinION sequencing. We thank Dorien Oppelaar for technical assistance in the lab.



## References

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. 1990. Basic local alignment search tool. *J Mol Biol* 215:403-410.
- Bertels, F., Silander, O. K., Pachkov, M., Rainey, P. B., and van Nimwegen, E. 2014. Automated reconstruction of whole-genome phylogenies from short-sequence reads. *Mol Biol Evol* 31:1077-1088.
- Bolton, M. D., van Esse, H. P., Vossen, J. H., de Jonge, R., Stergiopoulos, I., Stulemeijer, I. J., van den Berg, G. C., Borrás-Hidalgo, O., Dekker, H. L., de Koster, C. G., de Wit, P. J. G. M., Joosten, M. H., and Thomma, B. P. H. J. 2008. The novel *Cladosporium fulvum* lysin motif effector Ecp6 is a virulence factor with orthologues in other fungal species. *Mol Microbiol* 69:119-136.
- Capella-Gutiérrez, S., Silla-Martínez, J. M., and Gabaldón, T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972-1973.
- Cook, D. E., Mesarich, C. H., and Thomma, B. P. H. J. 2015. Understanding plant immunity as a surveillance system to detect invasion. *Annu Rev Phytopathol*.
- de Jonge, R., Bolton, M. D., Kombrink, A., van den Berg, G. C., Yadeta, K. A., and Thomma, B. P. H. J. 2013. Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. *Genome Res* 23:1271-1282.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- de Wit, P. J. 2016. *Cladosporium fulvum* Effectors: Weapons in the Arms Race with Tomato. *Annu Rev Phytopathol* 54:1-23.
- Dodds, P. N., and Rathjen, J. P. 2010. Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* 11:539-548.
- Dong, S., Stam, R., Cano, L. M., Song, J., Sklenar, J., Yoshida, K., Bozkurt, T. O., Oliva, R., Liu, Z., Tian, M., Win, J., Banfield, M. J., Jones, A. M., van der Hoorn, R. A., and Kamoun, S. 2014. Effector specialization in a lineage of the Irish potato famine pathogen. *Science* 343:552-555.
- Faino, L., Seidl, M. F., Datema, E., van den Berg, G. C., Janssen, A., Wittenberg, A. H., and Thomma, B. P. H. J. 2015. Single-Molecule Real-Time Sequencing Combined with Optical Mapping Yields Completely Finished Fungal Genome. *MBio* 6.
- Faino, L., Seidl, M. F., Shi-Kunne, X., Pauper, M., van den Berg, G. C., Wittenberg, A. H., and Thomma, B. P. 2016. Transposons passively and actively contribute to evolution of the two-speed genome of a fungal pathogen. *Genome Res* 26:1091-1100.
- Fradin, E. F., and Thomma, B. P. H. J. 2006. Physiology and molecular aspects of Verticillium wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol Plant Pathol* 7:71-86.
- Fradin, E. F., Abd-El-Haliem, A., Masini, L., van den Berg, G. C., Joosten, M. H., and Thomma, B. P. H. J. 2011. Interfamily transfer of tomato Ve1 mediates Verticillium resistance in Arabidopsis. *Plant Physiol* 156:2255-2265.
- Fradin, E. F., Zhang, Z., Juarez Ayala, J. C., Castroverde, C. D., Nazar, R. N., Robb, J., Liu, C. M., and Thomma, B. P. H. J. 2009. Genetic dissection of Verticillium wilt resistance mediated by tomato Ve1. *Plant Physiol* 150:320-332.
- Frandsen, R. J., Andersson, J. A., Kristensen, M. B., and Giese, H. 2008. Efficient four fragment cloning for the construction of vectors for targeted gene replacement in filamentous fungi. *BMC Mol Biol* 9:70.
- Garavaglia, B. S., Thomas, L., Gottig, N., Dunger, G., Garofalo, C. G., Daurelio, L. D., Ndimba, B., Orellano, E. G., Gehring, C., and Ottado, J. 2010. A eukaryotic-acquired gene by a biotrophic phytopathogen allows prolonged survival on the host by counteracting the shut-down of plant photosynthesis. *PLoS One* 5:e8950.


- Gehring, C., and Irving, H. 2003. Natriuretic peptides - a class of heterologous molecules in plants. *Int J Biochem Cell Biol* 35:1318-1322.
- Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., and Putnam, N. 2012. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res* 40:D1178-D1186.
- Gottig, N., Garavaglia, B. S., Daurelio, L. D., Valentine, A., Gehring, C., Orellano, E. G., and Ottado, J. 2008. *Xanthomonas axonopodis* pv. *citri* uses a plant natriuretic peptide-like protein to modify host homeostasis. *Proc Natl Acad Sci U S A* 105:18631-18636.
- Guindon, S., Delsuc, F., Dufayard, J.-F., and Gascuel, O. 2009. Estimating maximum likelihood phylogenies with PhyML. Pages 113-137 in: *Bioinformatics for DNA Sequence Analysis*. Springer.
- Houterman, P. M., Cornelissen, B. J., and Rep, M. 2008. Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* 4:e1000061.
- Huelsenbeck, J. P., and Ronquist, F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754-755.
- Joosten, M., Vogelsang, R., Cozijnsen, T. J., Verberne, M. C., and de Wit, P. J. G. M. 1997. The biotrophic fungus *Cladosporium fulvum* circumvents Cf-4-mediated resistance by producing unstable AVR4 elicitors. *Plant Cell* 9:367-379.
- Joosten, M. H., Cozijnsen, T. J., and de Wit, P. J. G. M. 1994. Host resistance to a fungal tomato pathogen lost by a single base-pair change in an avirulence gene. *Nature* 367:384-386.
- Katoh, K., Misawa, K., Kuma, K. i., and Miyata, T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30:3059-3066.
- Kawchuk, L. M., Hachey, J., Lynch, D. R., Kulcsar, F., van Rooijen, G., Waterer, D. R., Robertson, A., Kokko, E., Byers, R., Howard, R. J., Fischer, R., and Pruffer, D. 2001. Tomato Ve disease resistance genes encode cell surface-like receptors. *Proc Natl Acad Sci U S A* 98:6511-6515.
- Klosterman, S. J., Subbarao, K. V., Kang, S., Veronese, P., Gold, S. E., Thomma, B. P. H. J., Chen, Z., Henrissat, B., Lee, Y. H., Park, J., Garcia-Pedrajas, M. D., Barbara, D. J., Anchieta, A., de Jonge, R., Santhanam, P., Maruthachalam, K., Atallah, Z., Amyotte, S. G., Paz, Z., Inderbitzin, P., Hayes, R. J., Heiman, D. I., Young, S., Zeng, Q., Engels, R., Galagan, J., Cuomo, C. A., Dobinson, K. F., and Ma, L. J. 2011. Comparative genomics yields insights into niche adaptation of plant vascular wilt pathogens. *PLoS Pathog* 7:e1002137.
- Langmead, B., and Salzberg, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357-359.
- Lauge, R., Goodwin, P. H., de Wit, P. J. G. M., and Joosten, M. H. 2000. Specific HR-associated recognition of secreted proteins from *Cladosporium fulvum* occurs in both host and non-host plants. *Plant J* 23:735-745.
- Li, H., and Durbin, R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754-1760.
- Luderer, R., Takken, F. L., de Wit, P. J. G. M., and Joosten, M. H. 2002. *Cladosporium fulvum* overcomes Cf-2-mediated resistance by producing truncated AVR2 elicitor proteins. *Mol Microbiol* 45:875-884.
- Ludidi, N., Morse, M., Sayed, M., Wherrett, T., Shabala, S., and Gehring, C. 2004. A recombinant plant natriuretic peptide causes rapid and spatially differentiated K<sup>+</sup>, Na<sup>+</sup> and H<sup>+</sup> flux changes in *Arabidopsis thaliana* roots. *Plant Cell Physiol* 45:1093-1098.
- Marjoram, P., Molitor, J., Plagnol, V., and Tavaré, S. 2003. Markov chain Monte Carlo without likelihoods. *Proc Natl Acad Sci U S A* 100:15324-15328.
- Maryani, M., Bradley, G., Cahill, D., and Gehring, C. 2001. Natriuretic peptides and immunoreactants modify osmoticum-dependent volume changes in *Solanum tuberosum* L. mesophyll cell protoplasts. *Plant Sci* 161:443-452.
- Mesarich, C. H., Griffiths, S. A., van der Burgt, A., Okmen, B., Beenen, H. G., Etalo, D. W., Joosten, M. H., and de Wit, P. J. G. M. 2014. Transcriptome sequencing uncovers the Avr5 avirulence gene of the tomato leaf mold pathogen *Cladosporium fulvum*. *Mol. Plant Microbe Interact.* 27:846-857.

- Morse, M., Pironcheva, G., and Gehring, C. 2004. AtPNP-A is a systemically mobile natriuretic peptide immunoanalogue with a role in *Arabidopsis thaliana* cell volume regulation. *FEBS Lett* 556:99-103.
- Pharmawati, M., Maryani, M. M., Nikolakopoulos, T., Gehring, C. A., and Irving, H. R. 2001. Cyclic GMP modulates stomatal opening induced by natriuretic peptides and immunoreactive analogues. *Plant Physiol Biochem* 39:385-394.
- Rovenich, H., Boshoven, J. C., and Thomma, B. P. H. J. 2014. Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. *Curr Opin Plant Biol* 20:96-103.
- Santhanam, P., van Esse, H. P., Albert, I., Faino, L., Nurnberger, T., and Thomma, B. P. H. J. 2013. Evidence for functional diversification within a fungal NEP1-like protein family. *Mol. Plant Microbe Interact.* 26:278-286.
- Seidl, M. F., and Thomma, B. P. H. J. 2014. Sex or no sex: Evolutionary adaptation occurs regardless. *BioEssays* 36:335-345.
- Stergiopoulos, I., De Kock, M. J., Lindhout, P., and de Wit, P. J. G. M. 2007. Allelic variation in the effector genes of the tomato pathogen *Cladosporium fulvum* reveals different modes of adaptive evolution. *Mol. Plant Microbe Interact.* 20:1271-1283.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725-2729.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- Thomma, B. P. H. J., van Esse, H. P., Crous, P. W., and de Wit, P. J. G. M. 2005. *Cladosporium fulvum* (syn. *Passalora fulva*), a highly specialized plant pathogen as a model for functional studies on plant pathogenic Mycosphaerellaceae. *Mol Plant Pathol* 6:379-393.
- Tritt, A., Eisen, J. A., Facciotti, M. T., and Darling, A. E. 2012. An integrated pipeline for de novo assembly of microbial genomes. *PLoS One* 7:e42304.
- Van den Ackerveken, G. F., Van Kan, J. A., Joosten, M. H., Muisers, J. M., Verbakel, H. M., and de Wit, P. J. G. M. 1993. Characterization of two putative pathogenicity genes of the fungal tomato pathogen *Cladosporium fulvum*. *Mol. Plant Microbe Interact.* 6:210-215.
- van Kan, J. A., van den Ackerveken, G., and de Wit, P. J. G. M. 1991. Cloning and characterization of cDNA of avirulence gene *avr9* of the fungal pathogen *Cladosporium fulvum*, causal agent of tomato leaf mold. *Mol. Plant Microbe Interact.* 4:52-59.
- Westerink, N., Brandwagt, B. F., de Wit, P. J. G. M., and Joosten, M. H. A. J. 2004. *Cladosporium fulvum* circumvents the second functional resistance gene homologue at the Cf-4 locus (*Hcr9-4E*) by secretion of a stable *avr4E* isoform. *Mol Microbiol* 54:533-545.
- Zhang, Z., Fradin, E., de Jonge, R., van Esse, H. P., Smit, P., Liu, C.-M., and Thomma, B. P. H. J. 2013. Optimized agroinfiltration and virus-induced gene silencing to study Ve1-mediated *Verticillium* resistance in tobacco. *Mol. Plant Microbe Interact.* 26:182-190.

## Supplemental data

SUPPLEMENTAL TABLE 2 | Primers used in this study


Primer	Sequence (5'-3')	Notes
Ave1L-Break-Fw	TCGATGCGAGCAGATTCC	Primers spanning discontinuity
Ave1L-Break-Rv	GAGTAGTACCTTCCTTGCCCTTGA	Primers spanning discontinuity
Ave1L-Up-Fw	CTACGCTTGGAGCCCTCAT	Primers upstream of discontinuity
Ave1L-Up-Rv	AGGAGTGTAGGGTGGAGGAT	Primers upstream of discontinuity
Ave1L-Down-Fw	GGCATACTGAACCGATTGAAG	Primers downstream of discontinuity
Ave1L-Down-Rv	CTCCTTAGAAAGAACCATTTCG	Primers downstream of discontinuity
VdLs17del.stop_Fw	CGCATCCTGGTATCCTCCACC	Replace stop codon in Ave1L1 and Ave1L4
VdLs17del.stop_Rv	TGGAGGATACAGGATGCGGT	Replace stop codon in Ave1L1 and Ave1L4
Topo_Ave1L_Fw	CACCATGAAGCTTTCTACGCTTGGAG	Combined with primer VdLs17del.stop_Rv
Topo_St14.01_Rv	TTATTTTTTCTACTCAACTTGAATGAAC	Combined with primer VdLs17del.stop_Fw
Topo_St.100_Rv	TTATTTTATTCTACTCAACTTGAATGAAC	Combined with primer VdLs17del.stop_Fw
Ave1L2-LB-F3	GGTCTTAAUTTTTATCTCTCCCTTCCTCTTATCT	1500bp fragment upstream of Ave1L2
Ave1L2-LB-R3	GGCATTAAUTTTTAAGCCTTTCTAGCTTATTCTT	1500bp fragment upstream of Ave1L2
Ave1L2-RB-F3	GGACTTAAUGCTATCTTCACGAGAGCAGAGT	1500bp fragment downstream of Ave1L2
Ave1L2-RB-R3	GGGTTTAAU TTCGCGTTTATATATTCTTATCTT	1500bp fragment downstream of Ave1L2
QPCR-Ave1L-Fw3	ATCCTCCACCCTACACTCCT	Ave1L expression
QPCR-Ave1L-Rv3	CCTTGATGAGTAGTACCTTCCTTG	Ave1L expression
Vd-ITS-Fw	AAAGTTTAAATGGTTCGCTAAGA	<i>V. dahliae</i> biomass
Vd-ITS-Rv	CTTGGTCAATTAGAGGAAGTAA	<i>V. dahliae</i> biomass
SI-RUB-Fw	GAACAGTTTCTCACTGTTGAC	Tomato rubisco gene
SI-RUB-Rv	CGTGAGAACCATAAGTCACC	Tomato rubisco gene



SUPPLEMENTAL FIGURE 1 |

Verification of Ave1L2 deletion strains by amplification of VdITS and Ave1L2 from *V. dahliae* strains CBS38166 and St14.01.





**Rhamnose synthase activity  
is required for pathogenicity  
of the vascular wilt fungus  
*Verticillium dahliae***

**Chapter 7**

Parthasarathy Santhanam  
Jordi C. Boshoven#  
Omar Salas#  
Kyle Bowler#  
Md Tohidul Islam  
Mojtaba Keykha Saber  
Grady C.M. van den Berg  
Maor Bar-Peled  
Bart P.H.J. Thomma  
#These authors contributed equally

## Abstract

The initial interaction of a pathogenic fungus with its host is complex and involves numerous metabolic pathways and regulatory proteins. Here we present a random genetic screen that enabled us to identify 58 novel candidate genes that are involved in the pathogenic potential of the fungal pathogen *Verticillium dahliae* that causes vascular wilt diseases in over 200 dicotyledonous plant species, including economically important crops. One of the candidate genes that was identified concerns a putative biosynthetic gene involved in nucleotide-sugar precursor formation as it encodes a putative nucleotide-rhamnose synthase/epimerase-reductase (NRS/ER). This enzyme has homology to bacterial enzymes involved in biosynthesis of the nucleotide sugar dTDP-rhamnose, a precursor of L-rhamnose that was shown to be required for virulence in several human pathogenic bacteria. Rhamnose is known as a minor cell wall glycan in fungi and has therefore not been suspected as a crucial molecule in fungal-host interactions. Nevertheless, our study shows that deletion of the *VdNRS/ER* gene from the *V. dahliae* genome results in complete loss of pathogenicity on tomato and *N. benthamiana* plants, whereas vegetative growth and sporulation are not affected. We demonstrate that *VdNRS/ER* is a functional enzyme in the biosynthesis of UDP-rhamnose, and further analysis revealed that *VdNRS/ER* deletion strains are impaired in colonization of tomato roots. Collectively, our results demonstrate that rhamnose, although only being a minor cell wall component, is essential for pathogenicity of *V. dahliae*.

## Introduction

Vascular wilt diseases caused by soil-borne pathogens are among the most devastating plant diseases worldwide. The fungus *Verticillium dahliae* causes vascular wilt diseases in over 200 dicotyledonous plant species, including important crops such as eggplant, lettuce, olive, spinach and tomato (Fradin and Thomma 2006; Klosterman et al. 2009). Triggered by root exudates the melanised resting structures, microsclerotia, germinate and penetrate roots through the root tips, wounds, or sites of lateral root formation (Fradin and Thomma 2006). After crossing the cortex, hyphae of the fungus grow into the xylem vessels. The mycelium remains exclusively in these vessels and produces conidia which are transported acropetally with the water flow in the xylem throughout the plant. Once senescing, microsclerotia are produced that are released into the soil upon tissue decomposition (Wilhelm 1955). Typical symptoms of plants infected with *V. dahliae* comprise stunting, wilting, chlorosis and vascular browning (Pegg and Brady 2002).

*Verticillium* wilt diseases are difficult to control due to the longevity of the microsclerotia and inability of fungicides to eliminate the fungus once it has entered the xylem tissues of the host plant (Fradin and Thomma 2006; Wilhelm 1955). Moreover, crop rotation is ineffective due to the broad host range of *V. dahliae*. The only effective control measure, soil fumigation, is expensive and has harmful environmental effects (Rowe et al. 1987). As a consequence, genetic resistance is presently preferred to control *Verticillium* wilt diseases. Importantly, *Verticillium* resistance has been described in several plant species, including crops (Bolek et al. 2005; Hayes et al. 2011; Schaible et al. 1951; Simko et al. 2004). From tomato (*Solanum lycopersicum*) the *Ve1* *Verticillium* resistance gene has been cloned that mediates resistance to race 1 strains of *V. dahliae* that express the secreted Ave1 effector (de Jonge et al. 2012; Fradin et al. 2009). Putative *Ve1* orthologs have been identified within and outside the Solanaceae family, suggesting that *Ve1*-mediated *Verticillium* resistance may be widespread in plants (Thomma et al. 2011). Interestingly, Ave1 contributes to *V. dahliae* virulence on susceptible tomato plants, and homologs occur in a handful of taxonomically unrelated plant pathogenic fungi (de Jonge et al. 2012).

Various other virulence genes have been identified in *V. dahliae* including members of the expanded LysM effector and NEP-like (NLP) protein family (de Jonge et al. 2013; Klimes et al. 2015; Klosterman et al. 2011; Santhanam et al. 2013). Genome analysis furthermore revealed the relative abundance of particular families of cell-wall degrading enzymes (CWDE) when compared with other fungi (Klosterman et al. 2011). The importance of CWDE during pathogenicity was demonstrated by the disruption of the CWDE regulator VdSNF1 which reduced *V. dahliae* virulence (Tzima



et al. 2011). In addition, genes involved in nitrogen and amino acid metabolism were found to be important for establishment in the nutrient-poor vascular system (Singh et al. 2010; Timpner et al. 2013). Finally, virulence genes were described that are also involved in stress tolerance (Klimes et al. 2015).

The identification and subsequent characterization of novel genes involved in disease establishment is of fundamental importance to understand the interaction between *V. dahliae* and its hosts, and to potentially design novel strategies for disease control. An effective way to identify novel pathogenicity genes in plant pathogenic fungi is the use of random mutagenesis coupled with a quick manner to screen the mutants for phenotypes. *Agrobacterium tumefaciens* is a Gram-negative soil bacterium that can infect hosts as diverse as fungi, oomycetes and plants, and can be used to randomly insert a transfer-DNA (T-DNA) into the host genome. Over recent years, *A. tumefaciens*-mediated transformation (ATMT) was used to identify genes involved in disease development in various plant pathogens (Giesbert et al. 2012; Hüser et al. 2009; Jeon et al. 2007; Korn et al. 2015; Maruthachalam et al. 2011; Michielse et al. 2009; Munch et al. 2011; Ramos et al. 2013; Schumacher et al. 2014; Xu and Chen 2013). In this study, we report on the identification of pathogenicity and virulence genes in *V. dahliae* through ATMT. A collection of 900 transformants of *V. dahliae* was generated and tested for reduced disease development on susceptible tomato plants. We subsequently studied one of the candidates in detail, a gene that putatively encodes a biosynthetic gene involved in nucleotide-sugar precursor formation, to reveal its role in pathogenicity.

## Results

### Identification of random *V. dahliae* transformants affected in aggressiveness

In this study, we generated 900 random T-DNA insertion mutants of *V. dahliae* by transforming conidiospores through ATMT. All transformants were assessed for reduced potential to cause disease on tomato seedlings by root-dip inoculation and scored for typical disease symptoms, including stunting, wilting, chlorosis and necrosis, for up to 14 days post inoculation (dpi). Seedlings that exhibited reduced *Verticillium* wilt symptoms when compared with plants inoculated with wild type *V. dahliae* were selected, and the corresponding *V. dahliae* mutants were retained for rescreening, leading to a set of 200 transformants. Subsequently, after calibration of the conidial concentration in the inoculum to  $10^6$  conidia/mL, all 200 transformants were reassessed for compromised ability to cause wilt disease on tomato plants for up to 21 dpi. Eventually, 80 transformants showed a reproducible defect in disease development when compared with the wild-type progenitor strain (Figure 1).



**FIGURE 1 | Typical assay to identify transformants with reduced pathogenicity or virulence.** Ten-day-old tomato seedlings were mock-inoculated (M) or inoculated with conidiospores of wild-type *V. dahliae* (WT) or random T-DNA insertion mutants. At 21 days post inoculation the plants were scored for disease development by comparing plants inoculated with wild-type *V. dahliae* with those inoculated with the T-DNA insertion mutants. Side (upper panel) and top (bottom panel) of plants inoculated with five T-DNA insertion mutants that are impaired in aggressiveness (upper panel) and the top (bottom panel).

### Isolation of T-DNA flanking regions in selected *V. dahliae* transformants

Inverse PCR (iPCR) was carried out on the 80 selected transformants to determine the T-DNA insertion sites. The genomic borders flanking the T-DNA were amplified, sequenced, and the resulting sequences were queried against the *V. dahliae* genome (Faino et al. 2015). In this manner, sequences flanking the T-DNA were determined for 65 mutants, while T-DNA flanking regions repeatedly could not be amplified from any of the borders of the remaining 15 mutants. Of the 65 flanking sequences, 12 sequences corresponded to the backbone of the vector that was used for ATMT while 10 sequences could not unambiguously be assigned to a single genomic location. The remaining 43 sequences resulted in single genomic hits, allowing determination of T-DNA integration sites. In total, 14 insertions were found within predicted open reading frames (ORFs; Table 1) of which two transformants carried an insertion in the same ORF, while the remaining 29 insertions occurred in non-coding regions. For these 29 insertions the predicted genes flanking the T-DNA were identified. In one case we found that for two transformants the T-DNA was integrated in the same non-coding region, albeit at a slightly different location, resulting in the identification of the same flanking genes. In another case the same candidate gene was identified twice in two transformants due to an insertion upstream and downstream of the gene. One flanking gene was only identified at 55,077 bp from the insertion site, for which we disqualified it as a candidate gene. In total, this resulted in the identification of 55 candidate genes from the 29 mutants with insertions in non-coding regions.

In order to try to determine which of the two genes flanking a particular insertion site in the non-coding region is likely causal to the reduced disease development, *in planta* expression of the flanking genes was assessed making use of previously generated RNAseq data of *V. dahliae*-inoculated *Nicotiana benthamiana* plants (de Jonge et al. 2012; Faino et al. 2012). For 10 insertions, this revealed that only one of the two flanking genes is expressed during infection, disqualifying the 10 candidate genes that are not expressed. Thus, 45 candidate genes were considered from mutants with an insertion in the non-coding region and 13 candidate genes were identified from mutants with an insertion in the ORF (Table 1). Subsequently, the 58 candidate genes were queried against the pathogen-host interaction (PHI) database that contains experimentally verified pathogenicity and virulence factors of plant and animal pathogens (Winnenburg et al. 2006; Winnenburg et al. 2008), which resulted in the identification of homologs ( $E < 10^{-6}$ ) for only 19 of the 58 candidate genes (Table 2).

**TABLE 1 | List of candidate genes identified based on the T-DNA insertion sites in the random mutants that showed compromised virulence on tomato.**

T-DNA insertion within the CDS				
Mutant	Old locus <sup>b</sup>	New locus <sup>c</sup>	Annotation	Expr. <sup>d</sup> CDS <sup>e</sup>
78	Not available	VDAG_JR2_Chr1g08240	glycoside hydrolase	+ -
73	Not available	VDAG_JR2_Chr1g08240	glycoside hydrolase	+ -
407	VDAG_00500	VDAG_JR2_Chr1g19180	carboxypeptidase d	+ -
659	VDAG_02200	VDAG_JR2_Chr2g03300	alpha-beta hydrolase fold family	- -
SD-133	VDAG_04418	VDAG_JR2_Chr3g02140	ran gtp-ase activating protein 1	+ -
444-B	VDAG_08830	VDAG_JR2_Chr3g07650	acetate kinase	+ -
211	VDAG_06985	VDAG_JR2_Chr4g06020	leucine rich-repeat protein	+ -
375	VDAG_07314	VDAG_JR2_Chr4g10770	copper amine oxidase 1	+ -
416	VDAG_10292	VDAG_JR2_Chr5g06230	conserved hypothetical protein	- -
174	VDAG_05684	VDAG_JR2_Chr5g09640	hypothetical protein	+ -
657-2	VDAG_04769	VDAG_JR2_Chr6g09500	phospho-2-dehydro-3-deoxyheptonate aldolase	+ -
367	VDAG_06253	VDAG_JR2_Chr7g00300	cyclic nucleotide-binding protein	+ -
618	VDAG_08151	VDAG_JR2_Chr8g01560	hypothetical protein	+ -
48	VDAG_08150	VDAG_JR2_Chr8g01570	caib baif family enzyme	- -

T-DNA insertion in intergenic region					
Upstream direction					
Mutant	Dist. <sup>a</sup>	Old locus <sup>b</sup>	New locus <sup>c</sup>	Annotation	Expr. <sup>d</sup> CDS <sup>e</sup>
459-1	2485	VDAG_00838	VDAG_JR2_Chr1g01770	predicted protein	+ -
143	374	VDAG_00843	VDAG_JR2_Chr1g01820	not annotated	+ -
SD-102	270	VDAG_00599	VDAG_JR2_Chr1g20260	histidinol-phosphate aminotransferase	+ -
44	545	VDAG_08030	VDAG_JR2_Chr1g22770	deuterolysin metalloprotease	- -
11	100	VDAG_08003	VDAG_JR2_Chr1g23050	f-box protein	- -
SD-107	165	VDAG_07796	VDAG_JR2_Chr1g25310	hydrolase	+ -
182	470	VDAG_09439	VDAG_JR2_Chr2g01480	smc n-terminal containing protein	+ -
525	895	VDAG_02343	VDAG_JR2_Chr2g01760	cell wall glucanase	+ 100
SD-7	215	VDAG_02014	VDAG_JR2_Chr2g05250	myb-like dna-binding protein	+ -
540	622	VDAG_01960	VDAG_JR2_Chr2g05800	kh protein	+ -
402	1788	VDAG_01891	VDAG_JR2_Chr2g06520	domain-containing protein	+ 400
417	88	VDAG_01651	VDAG_JR2_Chr2g09280	f-box protein	+ -
622	700	Not available	VDAG_JR2_Chr3g02050	unknown	+ -
SD-89	335	VDAG_07703	VDAG_JR2_Chr4g00600	transcriptional regulatory protein pro-1	- -
SD-78	1214	VDAG_07011	VDAG_JR2_Chr4g05750	subunit of golgi mannosyltransferase complex	+ -
414	4	Not available	VDAG_JR2_Chr4g08330	Myomesin	- -
584	51	VDAG_09007	VDAG_JR2_Chr4g08540	family protein	+ -
534	737	VDAG_09007	VDAG_JR2_Chr4g08540	family protein	+ -
199	64	VDAG_05141	VDAG_JR2_Chr5g01900	low-affinity potassium transport protein	- -
412	1647	VDAG_09301	VDAG_JR2_Chr5g03950	potassium sodium efflux p-type fungal type	+ -
550	78	VDAG_04591	VDAG_JR2_Chr6g07500	calmodulin	+ -
SD-94	716	VDAG_04810	VDAG_JR2_Chr6g09950	nadph dehydrogenase	+ -
75	1060	VDAG_09054	VDAG_JR2_Chr6g10940	ubiquitin fusion degradation protein	+ -
374	1023	VDAG_06275	VDAG_JR2_Chr7g00090	maintenance of telomere capping 1	+ -
389	56	VDAG_06010	VDAG_JR2_Chr7g02960	nad dependent epimerase dehydratase	+ -
438	977	VDAG_10534	VDAG_JR2_Chr7g06650	ctp synthase	+ -
59	47	VDAG_10192	VDAG_JR2_Chr7g10460	benzoate 4-monooxygenase cytochrome P450	- -
646	246	VDAG_10191	VDAG_JR2_Chr7g10470	methyltransferase	+ -
575	466	VDAG_03431	VDAG_JR2_Chr8g09910	vegetative cell wall protein gp1-like	- -

T-DNA insertion in intergenic region					
Downstream direction					
Dist. <sup>a</sup>	Old locus <sup>b</sup>	New locus <sup>c</sup>	Annotation	Expr. <sup>d</sup> CDS <sup>e</sup>	
1784	VDAG_00839	VDAG_JR2_Chr1g01780	transcription initiation factor tfiid subunit 12	+	-
100	VDAG_00844	VDAG_JR2_Chr1g01830	exosome complex exonuclease	+	-
746	VDAG_00600	VDAG_JR2_Chr1g20270	acetyl esterase	+	-
669	VDAG_08031	VDAG_JR2_Chr1g22760	saff protein	+	-
3330	VDAG_08988	VDAG_JR2_Chr1g23040	umta methyltransferase	+	-
1586	VDAG_07797	VDAG_JR2_Chr1g25300	briggsae cbr-smu-2 protein	+	-
257	VDAG_09440	VDAG_JR2_Chr2g01470	sgt1-like protein	+	-
71	VDAG_02344	VDAG_JR2_Chr2g01750	phosphoribosylglycinamide formyltransferase	+	-
1004	VDAG_02015	VDAG_JR2_Chr2g05240	zinc nfx1-type containing protein	-	-
336	VDAG_01961	VDAG_JR2_Chr2g05790	duf974 protein	+	-
1318	VDAG_01892	VDAG_JR2_Chr2g06510	hypothetical protein	+	100
731	VDAG_01652	VDAG_JR2_Chr2g09270	cytochrome p450 protein	-	-
101	VDAG_04427	VDAG_JR2_Chr3g02040	agc akt protein kinase	+	-
423	VDAG_07704	VDAG_JR2_Chr4g00580	pathway-specific nitrogen regulator	+	-
1914	VDAG_07012	VDAG_JR2_Chr4g05740	fungal specific transcription factor	+	-
1466	Not available	VDAG_JR2_Chr4g08340	umta methyltransferase	-	-
1221	VDAG_09008	VDAG_JR2_Chr4g08550	zinc-binding dehydrogenase	+	-
535	VDAG_09008	VDAG_JR2_Chr4g08550	zinc-binding dehydrogenase	+	-
703	VDAG_05142	VDAG_JR2_Chr5g01910	sodium ion /h ion antiporter	+	-
1459	VDAG_09302	VDAG_JR2_Chr5g03940	glycoside hydrolase family 43 protein	+	100
667	VDAG_04592	VDAG_JR2_Chr6g07510	ap-1 complex subunit gamma-1	+	-
1014	VDAG_04811	VDAG_JR2_Chr6g09960	hypothetical protein	-	100
196	VDAG_09055	VDAG_JR2_Chr6g10930	ubiquitin fusion degradation protein	+	-
279	VDAG_06276	VDAG_JR2_Chr7g00080	major facilitator superfamily transporter	+	-
2363	VDAG_06011	VDAG_JR2_Chr7g02950	squalene synthetase	+	-
55077	VDAG_06595	VDAG_JR2_Chr7g06640	sad1 unc protein	+	-
1565	VDAG_10193	VDAG_JR2_Chr7g10450	cellulose-binding family II	+	-
422	VDAG_10192	VDAG_JR2_Chr7g10460	benzoate 4-monooxygenase cytochrome p450	-	-
1040	VDAG_03432	VDAG_JR2_Chr8g09920	duf221 protein	+	-

<sup>a</sup> Distance from the T-DNA to the coding sequence (CDS).

<sup>b</sup> Locus corresponds to the *V. dahliae* strain VdLs.17 assembly deposited on <http://genome.jgi.oe.gov/programs/fungi/> (Klosterman et al. 2011).

<sup>c</sup> Locus corresponds to the *V. dahliae* strain JR2 assembly deposited on <http://fungi.ensembl.org/> (Faino et al. 2015).

<sup>d</sup> Evidence for *in planta* expression.

<sup>e</sup> Evidence for expression in the region between T-DNA and the closest ORF. Distances from T-DNA are indicated in base pairs (bp).

**TABLE 2 | Homologs identified in the PHI database among the 58 potential *V. dahliae* pathogenicity genes.**

Old locus <sup>a</sup>	New locus <sup>b</sup>	Annotation	E-value	Accession <sup>c</sup>
VDAG_00839	VDAG_JR2_Chr1g01780	transcription initiation factor tfiid subunit 12	7E-19	PHI:2168
VDAG_00844	VDAG_JR2_Chr1g01830	exosome complex exonuclease	5E-12	PHI:3472
VDAG_01461	VDAG_JR2_Chr1g08260	ring finger protein	2E-09	PHI:1393
VDAG_00500	VDAG_JR2_Chr1g19180	carboxypeptidase d	4E-55	PHI:901
VDAG_09440	VDAG_JR2_Chr2g01470	sgt-1-like protein	1E-156	PHI:1615
VDAG_09439	VDAG_JR2_Chr2g01480	smc n-terminal containing protein	2E-10	PHI:359
VDAG_02343	VDAG_JR2_Chr2g01760	cell wall glucanase	9E-30	PHI:816
VDAG_02014	VDAG_JR2_Chr2g05250	myb-like dna-binding protein	1E-105	PHI:1540
VDAG_01960	VDAG_JR2_Chr2g05800	kh protein	3E-22	PHI:1662
VDAG_04427	VDAG_JR2_Chr3g02040	agc akt protein kinase	0E+00	PHI:3131
VDAG_07704	VDAG_JR2_Chr4g00580	pathway-specific nitrogen regulator	0E+00	PHI:1960
VDAG_07012	VDAG_JR2_Chr4g05740	fungal specific transcription factor	0E+00	PHI:1954
VDAG_09008	VDAG_JR2_Chr4g08550	zinc-binding dehydrogenase	7E-13	PHI:2895
VDAG_07314	VDAG_JR2_Chr4g10770	copper amine oxidase 1	1E-106	PHI:2171
VDAG_09301	VDAG_JR2_Chr5g03950	potassium sodium efflux p-type fungal type	0E+00	PHI:2098
VDAG_04591	VDAG_JR2_Chr6g07500	calmodulin	4E-20	PHI:2640
VDAG_06253	VDAG_JR2_Chr7g00300	cyclic nucleotide-binding protein	2E-10	PHI:372
VDAG_10193	VDAG_JR2_Chr7g10450	cellulose-binding family II	6E-13	PHI:2388
VDAG_10191	VDAG_JR2_Chr7g10470	methyltransferase	9E-40	PHI:2315

<sup>a</sup> Locus corresponds to the *V. dahliae* strain VdLs.17 assembly deposited on <http://genome.jgi.doe.gov/programs/fungi/> (Klosterman et al. 2011).

<sup>b</sup> Locus corresponds to the *V. dahliae* strain JR2 assembly deposited on <http://fungi.ensembl.org/> (Faino et al. 2015).

<sup>c</sup> Pathogen-host interaction database accession.

### Analysis of a T-DNA insertion site links *VdNRS/ER* to virulence


In random mutant 389, the T-DNA was integrated 56 bp upstream of the coding region of a nucleotide-rhamnose synthase/epimerase-reductase (NRS/ER) homolog VDAG\_06010 (Table 1; (Klosterman et al. 2011)), which corresponds to VDAG\_JR2\_Chr7g02960 in the *V. dahliae* JR2 assembly deposited at <http://fungi.ensembl.org/> (Faino et al. 2015). The first CDS downstream of the T-DNA insertion site is found only at 2.4 kb, encoding a squalene synthetase (VDAG\_06011 or VDAG\_JR2\_Chr7g02950), an enzyme that has been implicated in sterol and triterpene biosynthesis. However, considering the distance between the T-DNA insertion and the CDS of this gene it is unlikely that it is affected and causal to the reduced pathogenicity phenotype of the RM-389 mutant. To assess whether any genes have been overlooked by the automated gene prediction in the region between the

T-DNA insertion and the squalene synthetase CDS, RNAseq reads from *V. dahliae*-inoculated *N. benthamiana* (de Jonge et al. 2012; Faino et al. 2012) were queried. However, no reads were found to map to this region suggesting that no *in planta* transcribed genes reside in this region. Finally, with real-time PCR on genomic DNA, using the single copy *Ave1* gene as a reference (de Jonge et al. 2012), it was determined that only a single T-DNA insertion was present in the genome of the RM-389 mutant.

*V. dahliae* NRS/ER (VdNRS/ER) shows weak (17%) amino-acid sequence similarity to the *Salmonella enterica* dTDP-rhamnose biosynthesis component *rmlD*, a bacterial protein involved in the last step in the conversion of dTDP-glucose to dTDP-rhamnose. In bacteria the nucleotide sugar dTDP-rhamnose is the precursor that facilitates the incorporation of L-rhamnose into glycan, which has been implicated in virulence of pathogenic bacteria such as *Salmonella enterica*, *Vibrio cholerae* and *Streptococcus mutans* (Giraud et al. 2000). Thus, considering all of the above, functional analysis of the *VdNRS/ER* gene was pursued.

### **VdNRS/ER is required for *V. dahliae* pathogenicity**


Targeted replacement of *VdNRS/ER* by a hygromycin resistance cassette through homologous recombination was pursued in wild-type *V. dahliae*. Several independent *VdNRS/ER* deletion strains were obtained, of which two ( $\Delta 6010-1$  and  $\Delta 6010-2$ ) were used for further analysis (Supplemental Figure 1). To assess the role of *VdNRS/ER* in *V. dahliae* vegetative growth and conidiospore production, radial growth and sporulation of the *VdNRS/ER* deletion strains on potato dextrose agar was assessed together with the random mutant (RM-389) and an ectopic transformant with a random T-DNA insertion outside of the *VdNRS/ER* gene. This analysis showed that growth and conidiation was not markedly affected upon *VdNRS/ER* deletion (Figure 2).



**FIGURE 2 | Targeted deletion of *VdNRS/ER* does not impair growth and conidiogenesis.** (A) Radial growth and colony morphology of wild type *V. dahliae* (WT), random transformant RM-389, an ectopic transformant (EC) and two *VdNRS/ER* deletion strains ( $\Delta 6010-1$  and  $\Delta 6010-2$ ) after 7 days of incubation on PDA medium at 22°C. (B) Average number of conidia produced after 7 days of growth on PDA medium based on two independent experiments. Letters indicate significant differences ( $P < 0.05$ ) calculated with the Student's *t* test.




Subsequently, the role of *VdNRS/ER* in *V. dahliae* pathogenicity was addressed. To this end, the expression of *VdNRS/ER* was assessed in wild-type *V. dahliae* during infection of tomato plants in a time course harvested at regular intervals after inoculation, showing that *VdNRS/ER* is induced during *in planta* growth with a peak in expression at around 12 days post inoculation (Figure 3). Subsequently, the *VdNRS/ER* deletion strains were inoculated on tomato plants alongside the random mutant (RM-389) and an ectopic transformant to investigate the role of *VdNRS/ER* in fungal virulence. While inoculation of the plants with the wild-type *V. dahliae* strain and the ectopic transformant resulted in the development of symptoms of *Verticillium* wilt disease that include stunting of the plants and wilting, inoculation with RM-398 or the *VdNRS/ER* deletion strains did not result in disease development (Figure 4A). The absence of disease symptoms was confirmed by measuring the surface area of the foliage of the plants that were inoculated with the various fungal strains. Plants that were inoculated with *VdNRS/ER* deletion strains developed similar foliage (canopy) areas as mock-inoculated plants and plants inoculated with RM-389 (Figure 5A). In contrast, less foliage developed for plants that were inoculated with the wild-type *V. dahliae* strain and the ectopic transformant (Figure 5A). Thus, we conclude that *VdNRS/ER* is crucial for disease development by *V. dahliae*. To assess whether *VdNRS/ER* deletion strains retained the capability to colonize tomato plants, fungal outgrowth assays were performed by plating of stem sections harvested from the hypocotyls of the inoculated plants. Interestingly, while the ectopic transformants and the wild-type strain had extensively colonized the stems of the tomato plants, no outgrowth was observed for the *VdNRS/ER* deletion strains and RM-389, demonstrating that these genotypes were not able to colonize the xylem vessels of the inoculated tomato plants (Figure 4A). The absence of tomato colonization by *VdNRS/ER* deletion strains was confirmed with real-time PCR quantification of fungal biomass (Figure 5B). Importantly, pathogenicity was restored upon complementation of the *VdNRS/ER* deletion strains with a genomic construct containing the wild-type *VdNRS/ER* locus (Supplemental Figure 2). Thus, it can be concluded that *VdNRS/ER* is required for pathogenicity on tomato.



**FIGURE 3 | Expression of *VdNRS/ER* during infection of *Verticillium dahliae* on tomato.** Ten-day-old tomato (*S. lycopersicum*) cultivar MoneyMaker plants were root-inoculated with *V. dahliae* and plants were harvested at regular intervals from 4 to 16 days post inoculation (dpi). After RNA isolation and cDNA synthesis, real-time PCR was performed to determine the relative expression levels of *VdNRS/ER* using the *V. dahliae* elongation factor 1-alpha gene as a reference. Expression at 4 dpi is set to one for all panels.




To investigate whether the observed role of *VdNRS/ER* in pathogenicity is confined to tomato only, or also extends to other host species, we tested the pathogenicity of the *VdNRS/ER* deletion strains on the Solanaceous model plant *Nicotiana benthamiana*. Similar to tomato, targeted deletion of *VdNRS/ER* resulted in compromised pathogenicity, as also these plants remained devoid of *Verticillium* wilt symptoms (Figure 4B). Again, loss of pathogenicity was confirmed by measurement of the canopy surface area (Figure 5C), and loss of pathogenicity was confirmed by real-time PCR quantification of fungal biomass (Figure 5D).



**FIGURE 4 | *VdNRS/ER* is required for pathogenicity of *Verticillium dahliae* on tomato and *Nicotiana benthamiana*.** (A) Top and side view of tomato cultivar MoneyMaker plants that were mock-inoculated (M), or inoculated with wild type *V. dahliae* (WT), random transformant RM-389, an ectopic transformant (EC), and two *VdNRS/ER* deletion strains ( $\Delta 6010-1$  and  $\Delta 6010-2$ ) at 14 dpi. Fungal outgrowth at 7 days after plating of stem sections harvested at 14 days post inoculation is shown at the bottom of the panel. (B) Top and side view of *N. benthamiana* plants inoculated as specified for panel A.

### Absence of *VdNRS/ER* variation in a collection of *V. dahliae* strains


Allelic variation can be a sign of selection pressure and adaptation. To analyze potential *VdNRS/ER* diversity, we mined the genomes of 39 sequenced *V. dahliae* strains for the *VdNRS/ER* gene sequence (de Jonge et al. 2012) (Faino and Thomma, unpublished data). Intriguingly, only two single nucleotide polymorphisms (SNPs) were identified in two strains, each at a different position and both resulting in a synonymous substitution that does not affect the sequence of the encoded protein. We thus conclude that these substitutions do not affect the function of *VdNRS/ER*.



**FIGURE 5 | *VdNRS/ER* is required for pathogenicity of *Verticillium dahliae* on tomato and *Nicotiana benthamiana*.** (A) Average canopy area of 6 tomato plants, and (B) real-time PCR quantification of fungal biomass at 14 days after mock-inoculation (mock), or inoculation with wild type *Verticillium dahliae* (WT), random transformant (RM-389), an ectopic transformant (EC), and two *VdNRS/ER* deletion strains ( $\Delta 6010-1$  and  $\Delta 6010-2$ ). Different letter labels indicate significant differences ( $P < 0.05$ ). (C) Average canopy area of 6 *N. benthamiana* plants, and (D) real-time PCR quantification of fungal biomass upon inoculation as specified for panels A and B.

### ***VdNRS/ER* is not required for cell wall integrity and osmotic stress resistance**


To examine the role of *VdNRS/ER* in cell wall integrity and osmotic stress resistance, the random mutant (RM-389), the *VdNRS/ER* deletion strains and the ectopic transformant were grown on minimal agar medium supplemented with mannitol, sodium chloride, sorbitol or congo red. After seven days of growth the colony diameter was measured, showing that mutant and wild type strains were equally sensitive to the cell wall stress reagents (Figure 6). These findings suggest that *VdNRS/ER* is not involved in cell wall integrity or osmotic stress resistance during mycelium development.



**FIGURE 6 | Targeted deletion of *VdNRS/ER* does not affect cell wall integrity and osmotic stress resistance.** Stress sensitivity assays were performed by placing a 5  $\mu$ L droplet ( $10^6$  conidia/mL) of wild type *V. dahliae* (WT), an ectopic transformant (EC) and two *VdNRS/ER* deletion strains ( $\Delta 6010-1$  and  $\Delta 6010-2$ ) on Czapek-dox medium (C), or Czapek-dox supplemented with congo red (250 mM, 500 mM, 750 mM and 1 M), NaCl (250 mM, 500 mM, 750 mM and 1 M), mannitol (250 mM, 500 mM, 750 mM and 1 M), or sorbitol (300 mM, 600 mM, 900 mM, 1.2 M). The colony diameter was measured after 7 days of incubation at 22°C.

### ***VdNRS/ER* is required for tomato root colonization**


To examine the role of *VdNRS/ER* in the initial stages of *V. dahliae* infection, roots of ten-day-old tomato plants grown in a hydroponics solution were inoculated with conidiospores of the random mutant (RM-389), the *VdNRS/ER* deletion strains and the ectopic transformant. After five days of incubation, the tomato roots were inspected using a binocular microscope. Our data show that wild type spores efficiently attached to the root surface, resulting in extensive fungal growth on the tomato roots (Figure 7A). In contrast, little fungal growth was observed on roots inoculated with the random mutant (RM-389) or the *VdNRS/ER* deletion strains (Figure 7A). The reduced fungal biomass of the random mutant (RM-389) and the *VdNRS/ER* deletion strains on the tomato roots was confirmed by quantitative PCR (Figure 7B). Collectively, these data show that *VdNRS/ER* functions to contribute to successful colonization of roots by *V. dahliae*, potentially through root attachment.



**FIGURE 7 | *VdNRS/ER* is required for colonization of tomato roots.** Roots of ten-day-old tomato cultivar MoneyMaker seedlings were immersed in one-fifth PDB containing  $10^6$  conidia/mL of wild type *V. dahliae* (WT), random transformant (RM-389), an ectopic transformant (EC) and two *VdNRS/ER* deletion strains ( $\Delta 6010-1$  and  $\Delta 6010-2$ ) for 72-96 h. A) Roots were rinsed with water and photographed under a microscope. B) Real-time PCR quantification of fungal biomass on the roots. Different letter labels indicate significant differences ( $P < 0.05$ ).


### ***VdNRS/ER* is a functional enzyme that converts UDP-KDG to UDP-rhamnose**

To examine whether *VdNRS/ER* is involved in the biosynthesis of UDP-rhamnose and determine whether this activated sugar is the precursor for the synthesis of rhamnose-containing glycan, we examined the polysaccharide composition of *V. dahliae*. Polysaccharides were extracted from conidiospores of wild type *V. dahliae*, the *VdNRS/ER* deletion strain and the complementation strain, and analysed by GC-MS. In the samples derived from wild type *V. dahliae* and the complementation strain a peak was detected that corresponds to rhamnose, which was absent in the *VdNRS/ER* deletion strain (Figure 8). This finding confirms that *VdNRS/ER* is involved in rhamnose biosynthesis.



**FIGURE 8 | *VdNRS/ER* deletion strains are depleted in UDP-rhamnose.** GC-MS analysis of alditol acetate derivatives of the glycosyl residues from conidiospore polysaccharides (A) Total ion count chromatogram of alditol-acetate derived standard rhamnose (Rha), and mass spectra (insert) showing fragmentation pattern fingerprints with  $m/z$  129, 171, 231, 303. (B) Polysaccharides from wild type *V. dahliae* (WT) conidiospores display a peak eluting at 16.10 minutes that has the same retention time as rhamnose standard. (C) *VdNRS/ER* deletion strain ( $\Delta 6010-1$ ) lacks a peak eluting at 16.10 minutes. (D) *VdNRS/ER* deletion strain complemented with native *VdNRS/ER* gene (Comp-2), displays a peak eluting at 16.10 minutes that has the same retention time as the rhamnose standard.

If rhamnose-containing glycans play a role during early stages of host colonization, such as attachment of conidiospores to the root, we anticipated that such glycans are produced during conidiospore germination. Therefore, we determined the sugar composition of germinating conidiospores. Indeed, the sample derived from wild type *V. dahliae* comprised a clear rhamnose peak (Figure 9A;B) that was absent in samples from the *VdNRS/ER* deletion strain (Figure 9C). Taken together, *VdNRS/ER* plays a role in the formation of rhamnose-containing glycan in germinating conidiospores during early stages of host colonization.




**FIGURE 9 | UDP-rhamnose is produced in germinating conidiospores.** GC-MS analysis of alditol acetate derivatives of the glycosyl residues from germinating conidiospore polysaccharides (A) Total ion count chromatogram of alditol-acetate derived standard rhamnose (Rha) (B) Polysaccharides of conidiospores from wild type *V. dahliae* (WT) display a peak with the same retention time as the rhamnose standard. (C) *VdNRS/ER* deletion strain ( $\Delta 6010-1$ ) does not display a rhamnose peak. Other peaks in the chromatogram correspond to ribose (rib), arabinose (ara) and xylose (xyl). Asterisks indicate unidentified compounds that are not sugars.

In contrast to bacteria that produce dTDP-rhamnose in a three-step pathway, plants follow a two-step pathway to synthesize UDP-rhamnose (Oka et al. 2007; Watt et al. 2004). Genes involved in UDP-rhamnose were recently also identified in other fungi, including the rice pathogen *Magnaporthe oryzae* and the broad host range pathogen *Botrytis cinerea* (Martinez et al. 2012). In *B. cinerea*, UDP-glucose is converted by a 4,6-dehydratase (BfDh) into an intermediate UDP-KDG (UDP-4-keto-6-deoxy-glucose) that is subsequently converted by a bifunctional 3,5-epimerase/4-reductase (BfER, i.e. NRS/ER) into UDP-rhamnose (Martinez et al. 2012). Thus, we examined whether *VdNRS/ER*, like BfER, is a functional enzyme that can convert UDP-KDG to UDP-rhamnose. To this end, the dual enzyme activities of recombinant *VdNRS/ER* and BfDh was analyzed by LC-MS using UDP-glucose as a substrate (Martinez et al. 2012). The first enzyme in the pathway is capable to convert UDP-Glc to UDP-KDG (Figure 10B, and D) as expected, and with the addition of the NRS/ER activity (*VdNRS/ER*) UDP-KDG was converted to UDP-rhamnose (Figure 10C). The authenticity of UDP-rhamnose was determined by its elution and by its mass and ms/ms ion fragmentation (see insert in Fig 10C). The enzyme activity of recombinant *VdNRS/ER* unambiguously shows that *VdNRS/ER* is a bifunctional enzyme with a 3,5-epimerase and 4-reductase



activities (Figure 10E). Thus, we conclude that deletion of *VdNRS/ER* contributed to the loss of UDP-rhamnose formation and the lack of rhamnose-containing glycans during spore germination and early stages of infection.



**FIGURE 10 | VdNRS/ER converts UDP-KDG to UDP-rhamnose.** (A) UDP-glucose (UDP-Glc) standard elutes from HPLC column at 12.8 minutes and is detected by MS with m/z 565 [M-H]. The ms/ms of the parent ion (insert) gave m/z 323 and 385 diagnostic ion fragments [UMP-H] and [UDP-H<sub>2</sub>O-H], respectively. (B) HPLC separation of products from the enzymatic reaction with BfDh displays a peak with the same retention time as UDP-4-keto-6-deoxyglucose (UDP-KDG) with diagnostic [M-H] m/z 547 and ms/ms ion fragment with m/z 323. (C) HPLC separation of products from the dual enzymatic reaction with BfDh and VdNRS/ER displays a peak with the same retention time as UDP-rhamnose (UDP-Rha) with diagnostic [M-H] m/z 549, and ms/ms of 403, 323, 210. (D) Dual enzymatic reaction with BfDh and an empty vector control displays a peak with the same retention time as UDP-4-keto-6-deoxyglucose (UDP-KDG) with diagnostic [M-H] m/z 547 and ms/ms ion fragment with m/z 323. (E) UDP-rhamnose metabolic pathway in fungi showing the enzymes involved in the sequential conversion of UDP-Glc to UDP-Rha.

## Discussion

In our study on a random mutant library that was generated in the vascular wilt fungus *V. dahliae*, we show for the first time that rhamnose containing macromolecules are required for fungal pathogenicity through their contribution to colonization of tomato roots. In random mutant 389, the T-DNA was integrated 56 bp upstream of the coding region of a putative nucleotide-rhamnose synthase/epimerase-reductase (NRS/ER). Our biochemical analyses of cell wall polysaccharides (Figures 8 and 9) and enzyme activity (Figure 10C) confirmed that VdNRS/ER is involved in UDP-rhamnose biosynthesis and shows that this activated sugar is the precursor for rhamnose-containing glycan.

VdNRS/ER shows similarity to the bacterial rmlD substrate binding domain, which is involved in biosynthesis of dTDP-rhamnose. In Gram-negative bacteria such as *S. enterica*, *V. cholerae* or *Escherichia coli*, L-rhamnose is an important residue in the O-antigen of lipopolysaccharides, which are essential for resistance to serum killing and intestinal colonization (Chiang and Mekalanos 1999). In mycobacteria such as *Mycobacterium tuberculosis*, L-rhamnose maintains the structural integrity of the cell wall through connecting the inner peptidoglycan layer to the arabinogalactan polysaccharides (Giraud et al. 2000). Moreover, disruption of dTDP-rhamnose biosynthesis in the plant growth-promoting rhizobacteria *Azospirillum brasilense* and *Azorhizobium caulinodans* resulted in reduced root attachment and colonization (Gao et al. 2001; Jofré et al. 2004). Work on *Colletotrichum graminicola*, causal agent of stalk rot of cereals and anthracnose of maize revealed the secretion of mucilaginous matrix by the fungus to promote survival of fungal spores, the major component of which appeared to be high molecular weight glycoproteins containing rhamnose and mannose oligomers (Ramadoss et al. 1985). Subsequently, this matrix was proposed to allow for the rapid adhesion of even ungerminated conidia to the hydrophobic plant surface to ensure germination and appressorium formation and establishment of an infection court (Nicholson and Kunoh 1995). Interestingly, further rhamnose-containing extracellular matrix is released upon germ tube elongation to surround and extend between hyphae (Sugui et al. 1999). Finally, a recent study identified two genes encoding UG4,6-Dh and U4k6dG-ER enzymes from *Magnaporthe oryzae* and *Botrytis cinerea* involved in UDP-rhamnose formation, although their contribution to pathogenicity was not investigated (Martinez et al. 2012). Based on our study, however, strong evidence is presented that fungal UG4,6-Dh and U4k6dG-ER enzymes are involved in the formation of UDP-rhamnose that is incorporated into rhamnose-containing polysaccharides that are crucial for fungal pathogenicity. Here, we have shown that targeted deletion of VdNRS/ER contributes to loss of *V. dahliae* pathogenicity on various host plants.



The role of rhamnose-containing macromolecules in cell wall integrity and oxidative stress tolerance in fungi is largely unknown. *VdNRS/ER* deletion strains were able to grow and sporulate without any visible defects, suggesting that rhamnose-containing glycans such as glycoproteins, polysaccharides or exopolysaccharides, are not essential for vegetative growth or sporulation. In addition, the growth rate of *VdNRS/ER* knock-out mutants was not affected in the presence of inducers of cell wall stress and osmotic stress, suggesting that the rhamnose containing glycans are not involved in functional cell wall structure and osmotic stress tolerance.

Based on the findings reported in this study, we propose a model to explain the contribution of UDP-rhamnose to the virulence of *V. dahliae* during infection on host plants. In the cytoplasm UDP-glucose is converted into UDP-rhamnose in two steps using *VdUG4,6-Dh* and *VdNRS/ER*. The rhamnose of UDP-rhamnose is then transferred to glycan(s). The nature of these glycans remains unknown, but they consist of NaOH-solubilized cell wall polysaccharides and potentially glycoproteins. Presently, we cannot exclude the possibility that the rhamnose-containing polysaccharide is not an extracellular polysaccharide (EPS) as shown to be the case in *Botrytis cinerea*. We propose that the rhamnose-containing glycan, produced during spore germination, plays a role in the attachment of spores to the root surface which results in successful colonization of host plants. As *VdNRS/ER* deletion strains are unable to synthesize UDP-rhamnose and unable to cause disease, inhibitors of UDP-rhamnose synthesis may result in impaired pathogenicity and could present a novel strategy for disease control.

## Materials And Methods

### ***Agrobacterium tumefaciens* mediated transformation (ATMT)**

The binary vector (pBHt2) harboring the hygromycin B resistance gene (*hph*) under control of the *Aspergillus nidulans trpC* promoter was used for ATMT (Mullins et al. 2001). This vector was introduced to *Agrobacterium tumefaciens* strain SK1044 to transform conidia of race 1 *V. dahliae* strain JR2 as described previously (Faino et al. 2015; Santhanam 2012). Briefly, *A. tumefaciens* was grown at 28°C for 2 days in minimal medium supplemented with kanamycin (25 µg/mL). The *A. tumefaciens* cells were diluted to an OD<sub>600</sub> of 0.15 in induction medium (IM), supplemented with 200 µM acetosyringone (AS). The cells were grown for an additional six hours before mixing them with an equal volume of *V. dahliae* conidiospore suspension (10<sup>6</sup> conidia/mL). 200 µl of this mixture was plated onto a Hybond-N<sup>+</sup> filter placed on induction medium supplemented with 200 µM acetosyringone. The plates were incubated in the dark at room temperature for 48 h after which the filter was

transferred onto a selection plate (PDA supplemented with 50 µg/mL of hygromycin B and 200 µM of cefotaxime). After 10 to 14 days, individual transformants were transferred to 24-well culture plates containing one mL of selection medium and incubated for 7 to 10 days. Spores from these cultures were stored in 30% glycerol at -80°C until further analysis.

### Plant inoculations

Pathogenicity assays were performed as described previously (Santhanam and Thomma 2013). Briefly, individual transformants were sub-cultured in six-welled culture plates for 7 to 10 days. Subsequently, 10 glass beads (~3 mm) and 3 mL of tap water were added to each of the wells and the plates were sealed. The conidiospores were released by shaking the plates for 15 min at 200 rpm on a reciprocal shaker after which the sealing was removed. The roots of 10-day-old tomato (*Solanum lycopersicum* cv. MoneyMaker) seedlings were rinsed in water and dipped into the conidiospore suspension for 5 minutes. Seedlings were replanted in soil and scored for symptom development (wilting and stunting) up to 14 days. Seedlings that exhibited reduced *Verticillium* wilt symptoms when compared to inoculation with wild type *V. dahliae* inoculated plants were identified and the corresponding mutants were retained for rescreening.

During rescreening, the infection assay was carried out essentially as described above, with the modification that the conidiospore concentration of the transformants was calibrated to 10<sup>6</sup> conidiospores/mL. The rescreening was repeated at least two times for each of the mutants that were retained in the initial screen.

### Identification of T-DNA insertion sites

Genomic sequences flanking the T-DNA were isolated from the selected transformants with inverse PCR (iPCR) as described previously (Meng et al. 2007; Santhanam 2012). Essentially, 500 ng of genomic DNA was digested overnight with *MspI* or *NcoI* and heat-inactivated at 65°C for 20 min. Subsequently, 50 µL of ligation mix (10 µL of 10X T4 DNA ligase buffer, 5 units of T4 DNA ligase and 38 µL of H<sub>2</sub>O) was added and incubated overnight at 15°C. Next, the DNA was precipitated and dissolved in 50 µL of demineralized water. The genomic region flanking the T-DNA was amplified in 50 µL reaction mix using 2 µL of ligation product, 1 µL of each primer (Supplemental Table 1), 1X PCR buffer, 0.6 µL dNTP mix (10 mM), 0.8 units GO Taq DNA polymerase and 35.25 µL water. Cycling conditions consisted of 1 cycle for 2 min at 94°C, 35 cycles of 30 sec at 94°C, 30 sec at 55°C and 3 min at 72°C, and a final extension step for 10 min at 72°C. PCR products were purified

and sequenced. The obtained sequences were used as query to blast against the *V. dahliae* genome ([http://fungi.ensembl.org/Verticillium\\_dahliaejr2/Info/Index](http://fungi.ensembl.org/Verticillium_dahliaejr2/Info/Index)) (Faino et al. 2015).

### Targeted mutagenesis and complementation

To generate a *VdNRS/ER* deletion construct, sequences flanking the *VdNRS/ER* coding sequence were amplified from genomic DNA of *V. dahliae* strain JR2 using the primers KO-6010-LF with KO-6010-LR to amplify the left border, and KO-6010-RF with KO-6010-RR to amplify the right border, respectively (Supplemental Table 1). The resulting amplicons were cloned into pRF-HU2 as described previously (Frandsen et al. 2008). *A. tumefaciens*-mediated transformation of *V. dahliae* was performed as described previously (Santhanam 2012), and transformants were selected on PDA supplemented with 200 µg/mL of cefotaxime (Duchefa, Haarlem, The Netherlands), 50 µg/mL of hygromycin (Duchefa, Haarlem, The Netherlands). Homologous recombination was verified by PCR.

To generate a *VdNRS/ER* complementation construct, a 2404 bp *EcoRI/PacI* fragment containing the VDAG\_06010 (VDAG\_JR2\_Chr7g02960) coding sequence with 1000 bp upstream and 450 bp downstream sequence was amplified from *V. dahliae* strain JR2 genomic DNA, and cloned into binary vector pBT081 (Houterman et al. 2008). Complementation transformants were selected on PDA supplemented with 200 µg/mL of cefotaxime and 100 µg/mL phleomycin (InvivoGen, San Diego, USA).

### Growth, conidiogenesis and stress assays

Radial growth was monitored by placing a 2 µL droplet of a conidial suspension of 10<sup>6</sup> conidiospores/mL in the centre of PDA or Czapek-dox medium, incubated at 22°C and the colony diameter was measured after 10 days. For quantification of conidiospore production, 5 mL of water was added to the culture and a conidial suspension was prepared by gently rubbing the mycelium. A ten-fold dilution of the conidial suspension was counted using a haemocytometer. Stress sensitivity assays were performed by placing a 2 µL droplet with 10<sup>6</sup> conidiospores/mL of wild type *V. dahliae*, two *VdNRS/ER* deletion strains and an ectopic transformant in the centre of a Czapek-dox plate supplemented with congo red (250 mM, 500 mM, 750 mM and 1 M), NaCl (250 mM, 500 mM, 750 mM and 1 M), mannitol (250 mM, 500 mM, 750 mM and 1 M), or sorbitol (300 mM, 600 mM, 900 mM, 1.2 M), and incubated at 22°C. Plates were photographed at 7 dpi and the colony diameter was measured using ImageJ software.

### Root colonization assay

The root colonization assay was performed as previously described (Di Pietro et al. 2001). Briefly, the roots of 10-day-old tomato seedlings were rinsed in water and placed in Erlenmeyer flasks containing a suspension of  $10^6$  conidia/mL in one-fifth PDB and incubated at 22°C and 100 r.p.m. Fungal colonization of the root surface was observed macroscopically from 3 up to 5 days post inoculation. The experiments were performed three times with similar results.

### Expression of Recombinant VdNRS/ER

The following expression constructs were expressed in *E. coli* strain BL21(DE3): pET-6hSUMO-VDAG\_06010, contains a 6-histidine-SUMO tag fused to the N-terminal of *V. dahliae* VdNRS/ER (VDAG\_06010); pET-6hSUMO was used as a empty vector control (pET-SUMO); pET28-BfDH4.1 (Martinez et al. 2012) was used to express and purify the 6His tagged *Botrytis fuckeliana* UDP-Glc 4,6-dehydratase (BfDH). The *E. coli* strains carrying the different constructs were cultured for 16 h at 37°C in LB medium supplemented with 50 µg/mL kanamycin and 35 µg/mL chloramphenicol. The following day, 5 mL cultures were transferred into 245 mL of fresh LB liquid medium supplemented with the same antibiotics and cell growth was resumed at 37°C at 220 rpm until the cell density reached an OD<sub>600</sub> of ~0.6. Gene expression was then induced by the addition of isopropyl β-D-thiogalactoside (0.5 mM), and cultures were grown for additional 20 h at 18°C at 220 rpm. Cells were collected by centrifugation (6,000 × g for 10 min at 4°C) and suspended in 10 mL lysis buffer [50 mM Tris-HCl, pH 7.5, 10% glycerol, 1 mM EDTA, 2 mM DTT, 0.5 mM PMSF, with or without 50 mM NaCl]. Cell were lysed in an ice bath with a S-4000 sonicator (Misonix Inc, Farmingdale, NY, USA) equipped with 1/8-inch microtip probe using 12 sonication cycles each (10-s pulse; 20-s off). The lysed cells were centrifuged (6,000 × g for 10 min at 4°C); the supernatant was supplemented with 1 mM DTT and centrifuged again (20,000 × g for 30 min at 4°C). The resulting supernatant was kept at -20°C. Protein purification was carried out as previously described (Martinez et al. 2012).

### Enzyme Assays

The dual activity of recombinant enzymes BfDH and VdNRS/ER was analyzed by LC-MS, in a total volume of 50 µl, containing 50 mM Tris-HCl pH 6.5, 0.25 mM UDP-Glc, 0.5 mM NAD<sup>+</sup>, 0.5 mM NADPH, 5 µl of purified *B. fuckeliana* BfDH with or without the addition of 2 µl of VdNRS/ER. The reaction was incubated at 37°C for 1 h and terminated by heating for 5 min at 95°C. After the addition of chloroform and centrifugation (10,000 × g 4 min) an aliquot (20 µl) of the upper layer phase was

removed and mixed with 38  $\mu\text{L}$  acetonitrile and 2  $\mu\text{L}$  0.5 M ammonium-acetate pH 4.35. A portion (25  $\mu\text{L}$ ) of this mixture was analyzed using HILIC chromatography and mass spectrometry analyses. This was done using a LC-MS/MS IT-TOF MS system (Shimadzu Scientific Instruments, Columbia, MD, USA) operating in the negative ion mode with a Nexera UFPLC LC-30AD pump, autosampler (Sil30), and column heater (set to 37°C). Nucleotide sugars were separated on an Accucore 150-amid HILIC column (150  $\times$  4.6 mm, 2.6  $\mu\text{m}$  particle size, ThermoScientific) using 40 mM ammonium acetate, pH 4.3 (solvent A), and acetonitrile (solvent B) with the gradient conditions as in (Li et al. 2015).

### Isolation of polysaccharide and sugar analyses by GC-MS

Conidiospores were harvested from 7-day-old *V. dahliae* cultures grown on potato dextrose agar (PDA) (Thermo Fisher Scientific Inc, Breda, The Netherlands). Germinating spores were prepared by inoculating 30  $\mu\text{L}$  of conidiospores into 12.5 mL of Gamborg's Glc (2%) in a medium sized plate (9 cm). The plates were incubated for 66 hours at room temperature under regular light to allow the spores to germinate. Polysaccharides were either directly extracted from conidiospore samples or from germinating conidiospores (< 100  $\mu\text{g}$ ) starting by incubation for 3h at 80°C while stirring with 5 mL 1 M NaOH (5 mL). After centrifugation (10,000  $\times g$ , 4°C, 10 min) the alkaline extract was mixed with an equal volume of 96% ethanol. Following centrifugation (10,000  $\times g$ , 4°C, 15 min), the pellet was resuspended in 10 mL water, and dialyzed (3,000 molecular weight cutoff) against two 10-liter changes of deionized water. The dialyzed solution was centrifuged (10,000  $\times g$ , 4°C, 15 min) and the water-soluble supernatant (aWS) was collected and freeze-dried. The glycosyl residue compositions of these extracts were then determined.

An aliquot of each extract (1 mg) was supplemented with 10  $\mu\text{g}$  inositol and hydrolyzed with 1 mL of 2 M trifluoroacetic acid at 120°C. The released monosaccharides were reduced to their alditols (York et al. 1986), and acetylated. The resulting alditol acetate derivatives were analyzed by GC-MS system consisting of GC (Agilent 7890a) (Agilent Technologies, Santa Clara, CA, USA) equipped with an EI-MS detector (5975c). A 1  $\mu\text{L}$  sample or std was injected via Agilent 7693 autosampler injector port (Agilent Technologies, Santa Clara, CA, USA) (set at 250°C; 3 mL/min helium) in a split (1:50) mode into a non-polar Equity-1 capillary column (30 m  $\times$  0.25 mm i.d.; 0.25  $\mu\text{m}$  film thickness). The chromatography gas carrier was helium (1 mL/min) and the column oven temperature program used to separate sugar derivatives was as follows: initial injection temperature was 60°C for 1 min, followed by an increase of 27.5°C  $\text{min}^{-1}$  to 170°C, then at 4°C  $\text{min}^{-1}$  to 235°C, and a hold at 235°C for 2 min. The column was then kept at 260°C for 12 min, cooled to 60°C and kept

at 60°C for 1 min prior to the next sample injection. The temperature of the transfer line between the column ends to the MS was 250°C. Detection was achieved using EI-MS operating with an electron impact ionization of energy 70 V with the temperature of MS ion source set at 230°C and that of the quad at 150°C. MS data was collected in full scan monitoring mode at  $m/z$  range 50-550 from 5-50 min. The spectra were analyzed using Software MSD ChemStation D.02.00.275 (Agilent Technologies, Santa Clara, CA, USA). Alditol-acetate derivatives of monosaccharide standards (50 µg each of rhamnose, fucose, xylose, mannose, glucose, and 30 µg each ribose, arabinose, galactose) supplemented with 10 µg inositol (internal std.) were prepared under the same conditions as the samples. For data analysis, sugar residue peaks were identified based on retention times of standard monosaccharides and their characteristic mass spectrum.

## Acknowledgements

The authors thank Hugo de Vries and Emilie Fradin for technical assistance, and Bert Essenstam and Henk Smid for excellent plant care. B.P.H.J.T. is supported by a Vici grant of the Research Council for Earth and Life sciences (ALW) of the Netherlands Organization for Scientific Research (NWO). This research was funded in part by BARD grant # IS 4501-12 to M.B.P.

## References

- Bolek, Y., El-Zik, K. M., Pepper, A. E., Bell, A. A., Magill, C. W., Thaxton, P. M., and Reddy, O. U. K. 2005. Mapping of *Verticillium* wilt resistance genes in cotton. *Plant Sci* 168:1581-1590.
- Chiang, S. L., and Mekalanos, J. J. 1999. *rfb* mutations in *Vibrio cholerae* do not affect surface production of toxin-coregulated pili but still inhibit intestinal colonization. *Infect Immun* 67:976-980.
- de Jonge, R., Bolton, M. D., Kombrink, A., van den Berg, G. C., Yadeta, K. A., and Thomma, B. P. H. J. 2013. Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. *Genome Res* 23:1271-1282.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- Di Pietro, A., Garcia-MacEira, F. I., Meglec, E., and Roncero, M. I. 2001. A MAP kinase of the vascular wilt fungus *Fusarium oxysporum* is essential for root penetration and pathogenesis. *Mol Microbiol* 39:1140-1152.
- Faino, L., de Jonge, R., and Thomma, B. P. H. J. 2012. The transcriptome of *Verticillium dahliae*-infected *Nicotiana benthamiana* determined by deep RNA sequencing. *Plant Signal Behav* 7:1065-1069.
- Faino, L., Seidl, M. F., Datema, E., van den Berg, G. C., Janssen, A., Wittenberg, A. H., and Thomma, B. P. H. J. 2015. Single-Molecule Real-Time Sequencing Combined with Optical Mapping Yields Completely Finished Fungal Genome. *MBio* 6.
- Fradin, E. F., and Thomma, B. P. H. J. 2006. Physiology and molecular aspects of *Verticillium* wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol Plant Pathol* 7:71-86.
- Fradin, E. F., Zhang, Z., Juarez Ayala, J. C., Casterverde, C. D., Nazar, R. N., Robb, J., Liu, C. M., and Thomma, B. P. H. J. 2009. Genetic dissection of *Verticillium* wilt resistance mediated by tomato Ve1. *Plant Physiol* 150:320-332.
- Frandsen, R. J., Andersson, J. A., Kristensen, M. B., and Giese, H. 2008. Efficient four fragment cloning for the construction of vectors for targeted gene replacement in filamentous fungi. *BMC Mol Biol* 9:70.
- Gao, M., D'Haese, W., De Rycke, R., Wolucka, B., and Holsters, M. 2001. Knockout of an azorhizobial dTDP-L-rhamnose synthase affects lipopolysaccharide and extracellular polysaccharide production and disables symbiosis with *Sesbania rostrata*. *Mol. Plant Microbe Interact.* 14:857-866.
- Giesbert, S., Schumacher, J., Kupas, V., Espino, J., Segmuller, N., Haeuser-Hahn, I., Schreier, P. H., and Tudzynski, P. 2012. Identification of pathogenesis-associated genes by T-DNA-mediated insertional mutagenesis in *Botrytis cinerea*: a type 2A phosphoprotein phosphatase and an SPT3 transcription factor have significant impact on virulence. *Mol. Plant Microbe Interact.* 25:481-495.
- Giraud, M. F., Leonard, G. A., Field, R. A., Berlind, C., and Naismith, J. H. 2000. RmlC, the third enzyme of dTDP-L-rhamnose pathway, is a new class of epimerase. *Nat Struct Mol Biol* 7:398-402.
- Hayes, R. J., McHale, L. K., Vallad, G. E., Truco, M. J., Michelmore, R. W., Klosterman, S. J., Maruthachalam, K., and Subbarao, K. V. 2011. The inheritance of resistance to *Verticillium* wilt caused by race 1 isolates of *Verticillium dahliae* in the lettuce cultivar La Brillante. *Theor Appl Genet* 123:509-517.
- Houterman, P. M., Cornelissen, B. J., and Rep, M. 2008. Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* 4:e1000061.
- Hüser, A., Takahara, H., Schmalenbach, W., and O'Connell, R. 2009. Discovery of pathogenicity genes in the crucifer anthracnose fungus *Colletotrichum higginsianum*, using random insertional mutagenesis. *Mol. Plant Microbe Interact.* 22:143-156.
- Jeon, J., Park, S. Y., Chi, M. H., Choi, J., Park, J., Rho, H. S., Kim, S., Goh, J., Yoo, S., Choi, J., Park, J. Y., Yi, M., Yang, S., Kwon, M. J., Han, S. S., Kim, B. R., Khang, C. H., Park, B., Lim, S. E., Jung, K., Kong, S., Karunakaran, M., Oh, H. S., Kim, H., Kim, S., Park, J., Kang, S., Choi, W. B., Kang, S., and Lee, Y. H. 2007. Genome-wide functional analysis of pathogenicity genes in the rice blast fungus. *Nat Genet* 39:561-565.

- Jofré, E., Lagares, A., and Mori, G. 2004. Disruption of dTDP-rhamnose biosynthesis modifies lipopolysaccharide core, exopolysaccharide production, and root colonization in *Azospirillum brasilense*. *FEMS Microbiol Lett* 231:267-275.
- Klimes, A., Dobinson, K. F., Klosterman, S. J., and Thomma, B. P. H. J. 2015. Genomics spurs rapid advances in our understanding of the basic biology of vascular wilt pathogens in the genus *Verticillium*. *Annu Rev Phytopathol* 53:181-198.
- Klosterman, S. J., Atallah, Z. K., Vallad, G. E., and Subbarao, K. V. 2009. Diversity, pathogenicity, and management of *Verticillium* species. *Annu Rev Phytopathol* 47:39-62.
- Klosterman, S. J., Subbarao, K. V., Kang, S., Veronese, P., Gold, S. E., Thomma, B. P. H. J., Chen, Z., Henrissat, B., Lee, Y. H., Park, J., Garcia-Pedrajas, M. D., Barbara, D. J., Anchieta, A., de Jonge, R., Santhanam, P., Maruthachalam, K., Atallah, Z., Amyotte, S. G., Paz, Z., Inderbitzin, P., Hayes, R. J., Heiman, D. I., Young, S., Zeng, Q., Engels, R., Galagan, J., Cuomo, C. A., Dobinson, K. F., and Ma, L. J. 2011. Comparative genomics yields insights into niche adaptation of plant vascular wilt pathogens. *PLoS Pathog* 7:e1002137.
- Korn, M., Schmidpeter, J., Dahl, M., Muller, S., Voll, L. M., and Koch, C. 2015. A genetic screen for pathogenicity genes in the hemibiotrophic fungus *Colletotrichum higginsianum* identifies the plasma membrane proton pump *pma2* required for host penetration. *PLoS One* 10:e0125960.
- Li, Z., Hwang, S., Ericson, J., Bowler, K., and Bar-Peled, M. 2015. Pen and Pal are nucleotide-sugar dehydratases that convert UDP-GlcNac to UDP-6-deoxy-D-GlcNac-5, 6-ene and then to UDP-4-keto-6-deoxy-L-AltNac for CMP-Pseudaminic acid synthesis in *Bacillus thuringiensis*. *J Biol Chem* 290:691-704.
- Martinez, V., Ingwers, M., Smith, J., Glushka, J., Yang, T., and Bar-Peled, M. 2012. Biosynthesis of UDP-4-keto-6-deoxyglucose and UDP-rhamnose in pathogenic fungi *Magnaporthe grisea* and *Botryotinia fuckeliana*. *J Biol Chem* 287:879-892.
- Maruthachalam, K., Klosterman, S. J., Kang, S., Hayes, R. J., and Subbarao, K. V. 2011. Identification of pathogenicity-related genes in the vascular wilt fungus *Verticillium dahliae* by *Agrobacterium tumefaciens*-mediated T-DNA insertional mutagenesis. *Mol Biotechnol* 49:209-221.
- Meng, Y., Patel, G., Heist, M., Betts, M. F., Tucker, S. L., Galadima, N., Donofrio, N. M., Brown, D., Mitchell, T. K., Li, L., Xu, J. R., Orbach, M., Thon, M., Dean, R. A., and Farman, M. L. 2007. A systematic analysis of T-DNA insertion events in *Magnaporthe oryzae*. *Fungal Genet Biol* 44:1050-1064.
- Michiels, C. B., van Wijk, R., Reijnen, L., Cornelissen, B. J., and Rep, M. 2009. Insight into the molecular requirements for pathogenicity of *Fusarium oxysporum* f. sp. *lycopersici* through large-scale insertional mutagenesis. *Genome Biol* 10:R4.
- Mullins, E., Chen, X., Romaine, P., Raina, R., Geiser, D., and Kang, S. 2001. *Agrobacterium*-mediated transformation of *Fusarium oxysporum*: an efficient tool for insertional mutagenesis and gene transfer. *Phytopathology* 91:173-180.
- Munch, S., Ludwig, N., Floss, D. S., Sugui, J. A., Koszucka, A. M., Voll, L. M., Sonnewald, U., and Deising, H. B. 2011. Identification of virulence genes in the corn pathogen *Colletotrichum graminicola* by *Agrobacterium tumefaciens*-mediated transformation. *Mol Plant Pathol* 12:43-55.
- Nicholson, R. L., and Kunoh, H. 1995. Early interactions, adhesion, and establishment of the infection court by *Erysiphe graminis*. *Canadian Journal of Botany* 73:609-615.
- Oka, T., Nemoto, T., and Jigami, Y. 2007. Functional analysis of *Arabidopsis thaliana* RHM2/MUM4, a multidomain protein involved in UDP-D-glucose to UDP-L-rhamnose conversion. *J Biol Chem* 282:5389-5403.
- Pegg, G. F., and Brady, B. L. 2002. *Verticillium* wilts. CABI, Wallingford.
- Ramadoss, C. S., Uhlig, J., Carlson, D. M., Butler, L. G., and Nicholson, R. L. 1985. Composition of the mucilaginous spore matrix of *Colletotrichum graminicola*, a pathogen of corn, sorghum, and other grasses. *J Agric Food Chem* 33:728-732.




- Ramos, B., Gonzalez-Melendi, P., Sanchez-Vallet, A., Sanchez-Rodriguez, C., Lopez, G., and Molina, A. 2013. Functional genomics tools to decipher the pathogenicity mechanisms of the necrotrophic fungus *Plectosphaerella cucumerina* in *Arabidopsis thaliana*. *Mol Plant Pathol* 14:44-57.
- Rowe, R. C., Davis, J. R., Powelson, M. L., and Rouse, D. I. 1987. Potato early dying: causal agents and management strategies. *Plant Dis* 71:482-489.
- Santhanam, P. 2012. Random insertional mutagenesis in fungal genomes to identify virulence factors. Pages 509-517 in: *Plant fungal pathogens. Methods in molecular biology*, vol. 835. T. B. Bolton MD, ed. Humana press, Totowa.
- Santhanam, P., and Thomma, B. P. H. J. 2013. *Verticillium dahliae* Sge1 differentially regulates expression of candidate effector genes. *Mol. Plant Microbe Interact.* 26:249-256.
- Santhanam, P., van Esse, H. P., Albert, I., Faino, L., Nurnberger, T., and Thomma, B. P. H. J. 2013. Evidence for functional diversification within a fungal NEP1-like protein family. *Mol. Plant Microbe Interact.* 26:278-286.
- Schaible, L., Cannon, O. S., and Waddoups, V. 1951. Inheritance of resistance to *Verticillium* wilt in a tomato cross. *Phytopathology* 41:986-990.
- Schumacher, J., Simon, A., Cohrs, K. C., Viaud, M., and Tudzynski, P. 2014. The transcription factor BcLTF1 regulates virulence and light responses in the necrotrophic plant pathogen *Botrytis cinerea*. *PLoS Genet* 10:e1004040.
- Simko, I., Haynes, K. G., Ewing, E. E., Costanzo, S., Christ, B. J., and Jones, R. W. 2004. Mapping genes for resistance to *Verticillium albo-atrum* in tetraploid and diploid potato populations using haplotype association tests and genetic linkage analysis. *Mol Genet Genomics* 271:522-531.
- Singh, S., Braus-Stromeier, S. A., Timpner, C., Tran, V. T., Lohaus, G., Reusche, M., Knufer, J., Teichmann, T., von Tiedemann, A., and Braus, G. H. 2010. Silencing of Vlaro2 for chorismate synthase revealed that the phytopathogen *Verticillium longisporum* induces the cross-pathway control in the xylem. *Appl Microbiol Biotechnol* 85:1961-1976.
- Sugui, J. A., Kunoh, H., and Nicholson, R. L. 1999. Detection of protein and carbohydrate in the extracellular matrix released by *Cochliobolus heterostrophus* during germination. *Mycoscience* 40:11-19.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- Timpner, C., Braus-Stromeier, S. A., Tran, V. T., and Braus, G. H. 2013. The Cpc1 regulator of the cross-pathway control of amino acid biosynthesis is required for pathogenicity of the vascular pathogen *Verticillium longisporum*. *Mol. Plant Microbe Interact.* 26:1312-1324.
- Tzima, A. K., Paplomatas, E. J., Rauyaree, P., Ospina-Giraldo, M. D., and Kang, S. 2011. VdSNF1, the sucrose nonfermenting protein kinase gene of *Verticillium dahliae*, is required for virulence and expression of genes involved in cell-wall degradation. *Mol. Plant Microbe Interact.* 24:129-142.
- Watt, G., Leoff, C., Harper, A. D., and Bar-Peled, M. 2004. A bifunctional 3, 5-epimerase/4-keto reductase for nucleotide-rhamnose synthesis in *Arabidopsis*. *Plant Physiol* 134:1337-1346.
- Wilhelm, S. 1955. Longevity of the *Verticillium* wilt fungus in the laboratory and field. *Phytopathology* 45:180-181.
- Winnenburg, R., Baldwin, T. K., Urban, M., Rawlings, C., Köhler, J., and Hammond-Kosack, K. E. 2006. PHI-base: a new database for pathogen host interactions. *Nucleic Acids Res* 34:D459-D464.
- Winnenburg, R., Urban, M., Beacham, A., Baldwin, T. K., Holland, S., Lindeberg, M., Hansen, H., Rawlings, C., Hammond-Kosack, K. E., and Kohler, J. 2008. PHI-base update: additions to the pathogen host interaction database. *Nucleic Acids Res* 36:D572-576.
- Xu, L., and Chen, W. 2013. Random T-DNA mutagenesis identifies a Cu/Zn superoxide dismutase gene as a virulence factor of *Sclerotinia sclerotiorum*. *Mol. Plant Microbe Interact.* 26:431-441.
- York, W. S., Darvill, A. G., McNeil, M., Stevenson, T. T., and Albersheim, P. 1986. Isolation and characterization of plant cell walls and cell wall components. *Meth Enzymol* 118:3-40.

## Supplemental data


SUPPLEMENTAL TABLE 1 | Primers used in this study

Primer	Sequence (5' - 3')	Remarks
MLBF	GGATTTTGGTTTTAGGAATTAGA	<i>MspI</i> left border, forward
MLBR	AATTCGGCGTTAATTCAGTACA	<i>MspI</i> left border, reverse
MLSeq	TCAGTACATTAAAAACGTCCGCAA	<i>MspI</i> left border, sequencing
MRBF	CAACTGTTGGGAAGGGCGATC	<i>MspI</i> right border, forward
MRBR	CAGCCTGAATGGCGAATGCTA	<i>MspI</i> right border, reverse
MRSeq	GAATGCTAGAGCAGCTTGAGCT	<i>MspI</i> right border, sequencing
NLBF	AGTGATTGACCGATTCTTGC	<i>NcoI</i> left border, forward
NLBR	AGGGTTCCTATAGGTTTTCGCTCATG	<i>NcoI</i> left border, reverse
NLSeq	GAATTAATTCGGCGTTAATTCAGT	<i>NcoI</i> left border, sequencing
NRBF	CGTTATGTTTATCGGCACCTTG	<i>NcoI</i> right border, forward
NRBR	GGCACTGGCCGCTGTTTTACAAC	<i>NcoI</i> right border, reverse
NRSeq	CCCTCCCAACAGTTGCGCA	<i>NcoI</i> right border, sequencing
KO-6010-LF	GGTCTTAAUACTGCGCTGATGACCTCAC	Left border, forward
KO-6010-LR	GGCATTAAUCAGTGGATGCTGTTGTCGAT	Left border, reverse
KO-6010-RF	GGACTTAAUAATGCGAAATGCGTAAGGAG	Right border, forward
KO-6010-RR	GGGTTTAAUGCGGTAGGCCTTCTTGATCT	Right border, reverse
6010-F	GCGGCCGCATGTCAGTTACGAACGGTGC	Full length, forward
6010-R	CCATGGTCAATTAACACCAGCAGCCTT	Full length, reverse
6010 -comp_F	GAATTCTGCGCTGATGACCTCACC	Complementation, forward
6010 -comp_R	TTAATTAACCGCCTTGGATCACCAC	Complementation, reverse
qPCR-6010-F	ACACGATCCTCCACG ATCTC	Real-time PCR, forward
qPCR-6010-R	GCGAGAAGTCTTCCAGGTG	Real-time PCR, reverse
VdELF1-a-F	CCATTGATATCGCACTGTGG	Real-time PCR, forward
VdELF1-a-F	TGGAGATACCAGCCTCGAAC	Real-time PCR, reverse
qPCR-Hyg-F	ATAGGTCAGGCTCTCGCTGA	Real-time PCR, forward
qPCR-Hyg-R	GATGTAGGAGGGCGTGGATA	Real-time PCR, reverse
6010-BamH1-F	GGATCCTCAGTTACGAACGGTGCCA	pET-SUMO cloning, forward
6010-Hind3-R	AAGCTTCTATTATCAATTAACACCAGCAGCCTT	pET-SUMO cloning, reverse




**SUPPLEMENTAL FIGURE 1 | Verification of deletion and complementation strains by PCR.**

(A) Amplification of the T-DNA flanking sequences from wild type *V. dahliae* (w), 10 independent random transformants (lanes 1 to 10), and the marker (m) is indicated. (B) Amplification of the hygromycin resistance gene (*hph*) and *VdNRS/ER* from 10 independent transformants (lanes 1 to 10) and wild type *V. dahliae* (w). The marker (m) is indicated. True deletion strains are marked with asterisks, while the ectopic transformant is marked with an arrow. (C) Amplification of the zeocin resistance gene (*ble*) and *VdNRS/ER* from 5 independent complementation strains (lanes 1 to 5), a *VdNRS/ER* deletion strain (\*) and wild type *V. dahliae* (w). The marker (m), and true complementation strains (+) are indicated.



**SUPPLEMENTAL FIGURE 2 | Complementation of *VdNRS/ER* deletion strain restores pathogenicity on tomato.**

(A) Side and top view of tomato cultivar MoneyMaker plants inoculated with wild type *V. dahliae* (WT), a *VdNRS/ER* deletion strain ( $\Delta 6010-1$ ) and two complementation strains (comp-1 and comp-2), or mock-inoculated at 14 days post inoculation. (B) Fungal outgrowth at 7 days after plating of stem sections harvested at 14 days post inoculation. (C) Average canopy area of 6 plants at 14 days after inoculation as described for panel A. Different letter labels indicate significant differences ( $P < 0.05$ ).



**General discussion:  
From effector discovery to  
breeding for durable pathogen  
resistance in crops**

# **Chapter 8**

## Abstract

One of the major challenges in agriculture is to prevent yield loss caused by plant pathogens. Arguably, the most sustainable way to control disease is the deployment of genetic resistance. Pathogen resistance in plants is often conferred by immune receptors that, directly or indirectly, recognize pathogens or their activity. Effectors are pathogen-secreted molecules that support host colonization and therefore pose a strong selection pressure on plant hosts to recognize them. Effector identification and characterization has shed light on our understanding of plant-pathogen interactions, and on the various molecular mechanisms of pathogens and hosts that are involved in shaping those interactions. Here, I discuss the current understanding of pathogen effectors and how they can be exploited in breeding programs to establish durable pathogen resistance in crops.

## Introduction

Crop production is continuously threatened by plant pathogens, and genetic resistance is the most important source to prevent disease development. Already in the 40's of the 20<sup>th</sup> century, Harold Flor proposed the "gene-for-gene" theory, stating that for any given plant that carries a resistance (*R*) gene, a corresponding avirulence (*Avr*) gene is present in the pathogen to which it confers resistance (Flor 1942). This led to the assumption that *R* proteins directly interact with the corresponding *Avr* proteins (Keen 1990). However, most studies on *R*-*Avr* pairs failed to demonstrate such direct interaction, which led to the formulation of "the guard hypothesis" (Van der Biezen and Jones 1998). This hypothesis states that plant *R* proteins monitor, or "guard", the status of a plant protein that may be targeted by pathogen effectors. Once this indeed happens and the "guardee" is modulated by a pathogen effector, the guard becomes activated and triggers an immune response. This hypothesis was formulated because meanwhile it had become evident that *Avr* proteins often have functions to contribute to the virulence of the pathogen, for instance by targeting host proteins that are involved in defence responses, and were therefore referred to as effectors (Boller and Felix 2009). Currently, pathogen effectors are best described as secreted molecules that support host colonization (Rovenich et al. 2014). Besides pathogen effectors, plant immune receptors may also recognize conserved microbial signatures such as fungal chitin and bacterial flagellin, also known as microbe-associated molecular patterns (MAMPs). Whereas recognition of effectors is referred to as effector-triggered immunity (ETI), recognition of MAMPs is referred to as MAMP-triggered immunity (MTI) in the so-called "zigzag model" (Chisholm et al. 2006; Jones and Dangl 2006). It is generally accepted that successful plant pathogens use effectors to overcome MTI. In turn, to overcome ETI upon recognition of an effector by a host receptor, the pathogen needs to mutate or purge this effector, or evolve a novel effector to suppress the immune response and successfully colonize the plant.

### Effectors and MAMPs cannot be strictly separated

The discovery and characterization of various effectors and MAMPs from plant pathogens, including fungi, oomycetes and bacteria, have resulted in molecules that cannot be strictly separated as was proposed in "the zigzag model" (Thomma et al. 2011). In this model, effectors are described as species-, race- or even isolate-specific and contribute to pathogen virulence, whereas MAMPs are described as molecular patterns that are conserved in a broad range of species and contribute to microbial fitness. Arguably, these two groups represent the ends of a spectrum of molecules that should be seen as a continuum. Indeed, these definitions do not apply to pathogen molecules such as the NEP1-like proteins (NLPs) that are

found in a multitude of species belonging even to multiple kingdoms, and act as effectors that stimulate virulence on the one hand, and as MAMP on the other hand (Oome et al. 2014; Pemberton and Salmond 2004). Similarly, lysin-motif containing (LysM) effectors are conserved and widespread in the fungal kingdom (de Jonge and Thomma 2009; Kombrink and Thomma 2013). Also *Verticillium dahliae* effector Ave1 has homologs in other plant pathogens, including the fungi *Fusarium oxysporum*, *Colletotrichum higginsianum*, *Cercospora beticola* and the bacterium *Xanthomonas axonopodis* (de Jonge et al. 2012). In these organisms, the Ave1 homologs act as virulence factor as well (Ficarra et al. 2017; Gottig et al. 2008), with the exception of *C. higginsianum* (Chapter 3, this thesis). In addition, multiple Ave1 homologs were identified in the fungi *Venturia inaequalis* and *V. pirina* that are pathogens of apple and pear, respectively (Chapter 5, this thesis) (Cooke et al. 2014; Deng et al. 2017). However, whether these homologs act as virulence factors too remains to be demonstrated. Tomato immune receptor Ve1 differentially recognizes Ave1 effector homologs from various plant pathogens (Chapter, 4 this thesis) (de Jonge et al. 2012; Song et al. 2017). Finally, the effector proteins Pep1 and Pee1 of the maize smut fungus *Ustilago maydis*, which are essential effectors at the biotrophic stage, are conserved in other *Ustilago* species (Hemetsberger et al. 2012; Redkar et al. 2015). These examples illustrate that pathogen effectors are not always lineage-specific but can be conserved even across multiple kingdoms of life.

Bacterial flagellin is a MAMP that is broadly conserved between species. However, within species polymorphisms in flagellin have been identified in which some versions are recognized and others are not (Cai et al. 2011; Clarke et al. 2013; Sun et al. 2006). This shows that MAMPs are not always conserved and can also mutate to avoid recognition. Therefore, MAMPs and effectors and, consequently, MTI and ETI cannot be strictly separated, but rather form a continuum (Thomma et al. 2011).

The zigzag model adequately describes the interaction of biotrophic pathogens with their hosts, as they are generally affected by immune responses that involve host cell death. However, necrotrophic pathogens that actively kill host cells to access nutrients actually benefit from cell death activated by immune receptors (Glazebrook 2005). This has been shown for the fungal pathogen *Cochliobolus victoriae* effector victorin that binds to TRXh5 which activates immune receptor LOV1, resulting in cell death that benefits the pathogen instead of stopping the pathogen (Lorang et al. 2012). Thus, although recognition of effectors of biotrophic and necrotrophic pathogens similarly leads to cell death, the outcome for the interaction is significantly different. Thus, the interaction between necrotrophic pathogens and their hosts is generally not well described by the zigzag model.



Similarly, the zigzag model does not account for the interaction between plants and mutualistic microbes that use effectors as well (Plett and Martin 2015). Cell death does not benefit the 'desired' outcome of the interaction between plants and mutualists, and therefore the zigzag model does not accurately describe these interactions.

Examples such as those above have been the inspiration to propose an alternative model to describe plant-microbe interactions, in which a continuum between MAMPs and effectors is appreciated on the one hand, and similarly between MTI and ETI on the other hand. This alternative model is named the "the invasion model" (Cook et al. 2015), which states that any signal that reveals invasion is a so-called invasion pattern (IP) that can potentially be detected by the plant through invasion pattern receptors (IPRs), leading to an appropriate response (IP-triggered response; IPTR) that either supports or obstructs microbial colonization of the host.

### **Population genomics to identify genomic regions that are associated with pathogenicity**

The ease and low cost of present-day genome sequencing (Faino and Thomma 2014) has made it possible to sequence pathogen populations instead of a single genome per species. This has led to new insights in genome and effector evolution. Certain regions of fungal pathogen genomes are enriched in effector genes. For example, in the genome of the asexual pathogen *V. dahliae* chromosomal rearrangements are associated with highly dynamic lineage-specific regions that are enriched for *in planta*-induced effector genes and active transposons (de Jonge et al. 2013; Faino et al. 2016). Also in the fungal Brassica pathogen *Leptosphaeria maculans* and the potato famine oomycete *Phytophthora infestans* it was shown that gene-poor and repeat-rich regions of the genome are highly dynamic and enriched for *in planta* expressed effector genes, in contrast to gene-rich and repeat-poor regions of the genome (Dong et al. 2015). These observations illustrate the "the two-speed genome" model, in which effectors reside in fast evolving regions of the genome (Croll and McDonald 2012; Stukenbrock et al. 2010).

Various pathogens carry conditionally dispensable chromosomes (CDCs) that are enriched in effector genes. In the genome of the tomato pathogen *Alternaria alternate* a CDC contains all the genes responsible for AAL-toxin production (Akagi et al. 2009). More strikingly, in the genome of the tomato pathogen *F. oxysporum* f. sp. *lycopersici* a CDC contains all but one SIX (for secreted in xylem) effector genes (Ma et al. 2010; Schmidt et al. 2013; Van Dam et al. 2016), including *FoAve1* (Chapter 3, this thesis)(de Jonge et al. 2012). When this chromosome, which is also referred to as a pathogenicity chromosome, is transferred to a non-pathogenic isolate of *F. oxysporum*, this isolate gains the capacity to infect tomato (Ma et al. 2010).



### Effector discovery based on motifs and similarity

Many effectors have been identified based on their characteristics as small secreted cysteine-rich proteins. Depending on the (type of) organism, additional motifs have been associated with effectors, such as the RXLR or CRN (Crinkler) motif in *Phytophthora* species (Haas et al. 2009), MIMP (miniature impala) motif presence in the promotor of *F. oxysporum* effectors (Schmidt et al. 2013), lysin motif (LysM) in fungal effectors (de Jonge and Thomma 2009), the type-III secretion signal in effectors of gram-negative bacteria and the TAL (transcription activator-like) effector DNA binding motif in *Xanthomonas* species (Boch and Bonas 2010). However, for most pathogens effector motifs have not yet been identified and effectors often do not display sequence homology to other effectors, which makes it challenging to identify novel effectors. Comparison of the 3D-structure of two sequence-unrelated effector proteins of the fungal rice pathogen *Magnaporthe oryzae* revealed structural similarity (de Guillen et al. 2015). Subsequent structure-informed pattern searches identified novel effectors in *M. oryzae* and *M. grisea* that were named MAX (**M**agnaporthe **A**vrs and **T**oxB like) effectors (de Guillen et al. 2015). A similar approach to identify novel effectors can likely be applied to other pathogens as well (Franceschetti et al. 2017).

### Identification of effectors through random mutagenesis

Since the majority of effector genes that have been identified thus far are unique and do not have homologs in other pathogens, an unbiased approach should be applied to identify them. Indeed, using random mutagenesis through *Agrobacterium tumefaciens*-mediated transformation (ATMT) followed by a screen for reduced virulence on the host plant has proven to be successful in identifying novel effectors and pathogenicity factors in fungal pathogens including for example *F. oxysporum* (Michielse et al. 2009), *Colletotrichum graminicola* (Munch et al. 2011) and *Botrytis cinerea* (Giesbert et al. 2012). In a random mutagenesis screen on *V. dahliae* using ATMT 900 mutants were generated that, after thorough virulence testing and sequence analysis, led to the identification of 58 candidate pathogenicity and effector genes. One of the candidate genes, *VdNRS/ER*, was functionally analyzed and led to the finding that rhamnose synthase activity is required for pathogenicity of *V. dahliae* (Chapter 7, this thesis) (Santhanam et al. 2017).

### Population genomics as a tool for effector identification

Population genomics has contributed to the discovery of effectors that are recognized by host receptors. For example, by sequencing the genomes of 11 race 1 and race 2 strains of *V. dahliae* and subsequent RNA sequencing, the race

1-specific Ave1 effector that is recognized by tomato immune receptor Ve1 was identified (de Jonge et al. 2012). A similar approach was used to identify the Avr5 effector of the tomato leaf mould fungus *Cladosporium fulvum* that is recognized by the tomato immune receptor Cf-5 (Mesarich et al. 2014). Finally, the *F. oxysporum* f. sp. *melonis* effector AvrFom2 that is recognized by melon immune receptor Fom-2 was identified using a comparable approach (Schmidt et al. 2016). Potentially, this powerful strategy may be applied to identify potent virulence factors that are differentially present in other pathogen populations.

### Effectors define host specificity

Population genomics has revealed that effectors may be responsible for host specificity of a pathogen. Within the species complex of *F. oxysporum*, the *formae speciales* indicates on which host plant an isolate is pathogenic. *Formae speciales* are generally polyphyletic and therefore hard to identify based on conserved genes. However, effectors identified in *F. oxysporum* isolates from various host plants group together based on the same *formae speciales*, providing an identification tool based on effector presence or absence (Lievens et al. 2009; Van Dam et al. 2016). However, it should be taken into account that effector presence alone is not always proof that an isolate is pathogenic as non-pathogenic *Fusarium* isolates were found that carry effector genes too (Jelinski et al. 2017). Also, comparative genomics between strains of the bacterium *Ralstonia solanacearum* that are pathogenic on different host plants revealed effector repertoires associated with a specific host (Ailloud et al. 2015). Finally, *P. infestans* and its close relative *P. mirabilis* that is pathogenic on *Mirabilis jalapa* share a homologous effector that is only effective on their corresponding plant host (Dong et al. 2014).

Gene and effector expansions and contractions are linked to the host range of a pathogen. In *V. dahliae*, NLP effector expansion has been associated with functional diversification (Santhanam et al. 2013; Zhou et al. 2012). Only a subset of the NLP effectors in *V. dahliae* is cytotoxic (typical for NLPs) while others are non-cytotoxic. Besides a role in virulence, NLP1 was also shown to be involved in asexual reproduction and vegetative growth. NLP1 and NLP2 are virulence factors on tomato and *Arabidopsis*, but not on cotton, and only NLP1 is required for virulence on *Nicotiana benthamiana* (Santhanam et al. 2013; Zhou et al. 2012). This suggests that NLP diversification might have contributed to host range expansion. A comparison of species in the genus *Colletotrichum* that have a broad host range with species that have a limited host range revealed that species with broad host ranges rely on secreted enzymes and NLPs, whereas narrow host range species have lost proteases and are more reliant on lineage-specific effectors (Baroncelli et al. 2016). The examples above illustrate that effector repertoires define the host

(range) of a pathogen. Moreover, as effectors are important virulence factors of pathogens that are potentially recognized by related or unrelated plant species of the host, they can be exploited as tools to identify novel resistance sources.

### Effectors as tools for pathogen resistance breeding

Traditional plant breeding is a tedious and time-consuming practice in which marker-assisted selection has significantly increased the speed and efficiency of the selection process. However, pathogen assays are still an essential aspect for the development and verification of markers. Pathogen assays have limitations since they require extensive knowledge of the pathogen's life cycle and can often only be performed with a single species or race of a given pathogen as resistance is usually species- or even race- specific. Once effectors have been identified for a given pathogen, these effectors can be used to screen germplasm for novel recognition specificities, and thus new resistance sources (Vleeshouwers and Oliver 2014). This can be done by transient expression of effector genes in germplasm by *Agrobacterium*-mediated expression or potato virus X (PVX) (Chapman et al. 1992; Janssen and Gardner 1990). In this manner, many effectors can be tested in parallel, circumventing the limitations of disease assays, while simultaneously providing the possibility of screening germplasm that is a non-host to the pathogen or the possibility of screening effectors of pathogens with a complex lifecycle.

Since many effectors are race- or even strain- specific it is not very efficient to use these in a germplasm screen for resistance. Recognition specificity towards effectors that are conserved in a population of pathogens, also referred to as core effectors, is far more valuable. In a population genomics study on 65 isolates of the cassava bacterial blight pathogen *Xanthomonas axonopodis* pv. *manihotis* 9 effectors were identified that are conserved in the population (Bart et al. 2012). In a similar approach, 13 conserved effectors were identified in 67 isolates that belong to three *Xanthomonas* species that are pathogenic on tomato and pepper (Potnis et al. 2011; Schwartz et al. 2015). Population genomics on strains of *V. dahliae* also revealed effectors that are conserved in the population (de Jonge et al. 2013). Identification and exploitation of *R* genes that recognize these conserved effectors might potentially lead to durable pathogen resistance.

Knowledge of the effectors that are present in a given population can also aid in developing recognition specificity towards (a) certain effector(s). A single amino acid substitution can already broaden the specificity of an *R* gene. This was shown for potato NB-LRR immune receptor R3a by performing a gain-of-recognition screen after random mutagenesis in which the recognition specificity was extended to isoforms of the recognized *P. infestans* effector Avr3A that are predominant in *Phytophthora* populations world-wide (Segretin et al. 2014). I2 is a NB-LRR immune receptor of

tomato that confers resistance to *F. oxysporum* and partial resistance to *P. infestans* and shows high similarity to R3a (Giannakopoulou et al. 2015). Mutations that extended recognition in potato R3a were also mutated in tomato I2 which resulted in a broader and enhanced specificity to both *P. infestans* effector isoforms of Avr3a and *F. oxysporum* effector isoforms of Avr2 (Giannakopoulou et al. 2015). These examples illustrate that with relatively small adjustments the recognition specificity of an immune receptor can be extended. Potentially, this strategy can be exploited to increase the recognition specificity of Ve1 towards pathogens that secrete homologs of Ave1 that are poorly recognized.

Like I2, other *R* genes or combinations of genes have been described to provide resistance to two or more pathogens. Tomato immune receptor Cf-2 guards Rcr3 that is targeted by effectors of the fungus *C. fulvum* and the nematode *Globodera rostochiensis* resulting in a defence response (Lozano-Torres et al. 2012). Thus, naturally occurring *R* genes can be exploited that confer resistance to multiple pathogens.

In a map-based cloning approach combined with natural variation analysis in *Arabidopsis* a dual *R* gene system consisting of the nucleotide binding, leucine-rich repeat (NLR) domain receptors *RPS4-Ws* and *RRS1-Ws* was identified that provides resistance to the fungus *C. higginsianum*, the bacteria *Ralstonia solanacearum* and *Pseudomonas syringae* pv. *tomato* (Deslandes et al. 2002; Narusaka et al. 2009). Other dual *R* gene systems have been characterized as well, such as the rice NLRs RGA4 and RGA5 that interact to provide resistance to *M. oryzae* (Césari et al. 2014a). In these NLR pairs integrated domains (NLR-IDs) have been identified that serve as a bait for effectors to activate immunity as has been described in the integrated decoy hypothesis (Césari et al. 2014b). NLR-ID integration in plant immune receptors is broadly distributed in the plant kingdom (Kroj et al. 2016; Sarris et al. 2016) and analyses of grass genomes suggests the presence of NLR ‘acceptor’ genes that likely acquire these domains from elsewhere in the genome (Bailey et al. 2017). Potentially, modification or transfer of NLR-IDs to other NLRs can contribute to synthesizing novel *R* genes (Kim et al. 2016; Rodriguez-Moreno et al. 2017).

Often *R* genes are identified or characterized in wild species or model organisms and cannot easily be introduced into crops by crossing. However, interfamily transfer of receptor-like proteins (RLPs) and receptor kinases (RKs) has been quite successful. For example, the *Arabidopsis thaliana* receptor kinase EFR that recognizes bacterial elongation factor Tu (EF-Tu) was transferred to tomato (*Solanum lycopersicum*) providing broad bacterial resistance (Lacombe et al. 2010). Similarly, the *A. thaliana* receptor kinase *LecRK-1.9* which provides resistance to *P. infestans* was functionally transferred to potato (*S. tuberosum*) (Bouwmeester et al. 2014). Interfamily transfer of NB-LRRs seems to be less efficient compared to transfer of RLPs and RKs. Perhaps, for NLRs it is more efficient to transfer genes from wild relatives into crops (Bent 2016).

### Susceptibility genes as a source of resistance

Resistance is often overcome within a couple of years after the introduction of an *R* gene (Rouxel et al. 2003; Sprague et al. 2006a; Sprague et al. 2006b). It has been proposed that mutation of host components that are required for a pathogen to successfully colonize its host, also known as susceptibility (*S*) genes (Eckardt 2002), is a more durable strategy to provide pathogen resistance (McDonald and Linde 2002). An example of an *S* gene is the *Mlo* gene that provides resistance to powdery mildew in many crops including monocots and dicots (Kusch and Panstruga 2017). In barley, a recessive mutant of this gene was introduced 70 years ago and is still providing resistance to all powdery mildews in the field (Jørgensen 1992). Effector targets can be mutated with the consequence that the corresponding effector becomes useless. However, *S* genes have other functions in the plant as well. Therefore it should be taken into account that mutating an *S* gene can have detrimental effects on the plant (van Schie and Takken 2014).


### Genome editing in crops

Genome editing in crops has made considerable steps forward over the last few years. With synthetic nucleases such as zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs) and Cas9 of the clustered regularly interspaced short palindromic repeats (CRISPR) system it is possible to knockout genes, delete or insert sequences without selection markers (Puchta 2017; Schiml and Puchta 2016). Arguably, CRISPR/Cas9 is the most efficient and flexible system to work with (Puchta 2017). Concerns about off-target mutations seem to be a problem of the past and efficiencies of 4-9% have been reached. The strength of the CRISPR/Cas9 system is that it is possible to make precise changes in a sequence without the introduction of foreign DNA and marker genes. This was elegantly shown by editing the genomes of maize and bread wheat (Liang et al. 2017; Svitashv et al. 2016). In tomato, CRISPR/Cas9 was used to inactivate the *S* gene orthologue *DMR6* (for downy mildew resistance 6) that was characterized in *A. thaliana* (Van Damme et al. 2008), resulting in resistance to *Pseudomonas syringae*, *Phytophthora capsici* and *Xanthomonas* species (de Toledo Thomazella et al. 2016). Similarly, a deletion generated in an *Mlo* orthologue by CRISPR/Cas9 in tomato resulted in the variety Tomelo which is resistant to the powdery mildew fungal pathogen *Oidium neolycopersici* (Nekrasov et al. 2017). For this variety it was shown with whole-genome sequencing that it does not carry any foreign DNA, does not have off-target mutations and carries a single deletion that is indistinguishable from naturally occurring mutations. Whether genome editing using CRISPR/Cas9 will skip regulation as a genetically modified organism (GMO) in the EU should become clear in the near future (Waltz 2016; Wolter and Puchta 2017).

### Management strategies to prevent resistance breakdown

Unfortunately, current agricultural practice is still based on planting vast areas of monocultures, often carrying only a single *R* gene against a given pathogen, season-after-season. This puts a huge selection pressure on pathogen populations to overcome resistance. Resistance breakdown can occur within a few years after the introduction of an *R* gene (Rouxel et al. 2003; Sprague et al. 2006a; Sprague et al. 2006b).

*R* gene stacking, the introduction of multiple *R* genes in a single variety, significantly increases the hurdle for a pathogen to overcome resistance (McDonald and Linde 2002; Michelmore et al. 2013). *R* gene stacking can be achieved by gene transfer or marker-assisted breeding (Dangl et al. 2013). Using a cisgenic approach, broad spectrum *Rpi* (for resistance against *P. infestans*) genes were stacked and introduced in the susceptible potato cultivar Désirée to provide durable resistance to *P. infestans* and were subsequently monitored in field trials in the Netherlands (Haverkort et al. 2016; Tan et al. 2010; Zhu et al. 2013; Zhu et al. 2012). Several combinations of *Rpi* genes provided resistance to *P. infestans* and spray applications were significantly reduced (Haverkort et al. 2016). With marker-assisted breeding multiple *R* genes were stacked in soybean to provide durable resistance to *Soybean mosaic virus* (SMV) (Maroof et al. 2008) and in wheat to provide resistance to powdery mildew (Liu et al. 2000). Criteria for stacking genes should be taken into account to prevent rapid resistance breakdown. These include: the stacked *R* genes should target the same pathogen species, they should be introduced as a tandem construct to prevent segregation, have similar levels of expression in the same tissue and should target species that are susceptible to at least two of the stacked *R* genes (Gressel et al. 2017). Obviously, extensive knowledge on *R* genes and their corresponding effectors is essential to make educated decisions on which genes to stack. *R* gene catalogues are available for multiple crop-pathogen interactions including, for example, potato and *P. infestans*, and lettuce and *Bremia lactucae* (Parra et al. 2016; Vleeshouwers et al. 2011). Monitoring pathogen populations in the field is essential to know which effectors are present in the pathogen population and to know at an early time point whether resistance breakdown occurs (Howlett et al. 2015; Michelmore et al. 2013). Subsequently, *R* genes should be selected that provide resistance to conserved core effectors present in a pathogen population to increase the durability of resistance (Dangl et al. 2013). In addition, stacks of different *R* genes should be used either within a variety or between different varieties to provide distinct selection pressure and therefore increased durability (Chin and Wolfe 1984; Michelmore et al. 2013). Finally, different *R* genes or *R* gene stacks should be used in time and space to break the continuous selection pressure that is put on pathogens. The criteria described above will ultimately lead to durable broad spectrum resistance in crops (Figure 1).



**FIGURE 1** | Summarizing figure to visualize the many strategies that potentially lead to durable broad spectrum resistance in crops.

## References

- Ailloud, F., Lowe, T., Cellier, G., Roche, D., Allen, C., and Prior, P. 2015. Comparative genomic analysis of *Ralstonia solanacearum* reveals candidate genes for host specificity. *BMC Genomics* 16:1.
- Akagi, Y., Akamatsu, H., Otani, H., and Kodama, M. 2009. Horizontal chromosome transfer, a mechanism for the evolution and differentiation of a plant-pathogenic fungus. *Eukaryotic cell* 8:1732-1738.
- Bailey, P. C., Dagdas, G., Baggs, E., Haerty, W., and Krasileva, K. V. 2017. Immune receptors with exogenous domain fusions form evolutionary hotspots in grass genomes. *BioRxiv*:100834.
- Baroncelli, R., Amby, D. B., Zapparata, A., Sarrocco, S., Vannacci, G., Le Floch, G., Harrison, R. J., Holub, E., Sukno, S. A., and Sreenivasaprasad, S. 2016. Gene family expansions and contractions are associated with host range in plant pathogens of the genus *Colletotrichum*. *BMC Genomics* 17:555.
- Bart, R., Cohn, M., Kassen, A., McCallum, E. J., Shybut, M., Petriello, A., Krasileva, K., Dahlbeck, D., Medina, C., and Alicai, T. 2012. High-throughput genomic sequencing of cassava bacterial blight strains identifies conserved effectors to target for durable resistance. *Proceedings of the National Academy of Sciences* 109:E1972-E1979.
- Bent, A. 2016. Resistance from relatives. *Nat Biotechnol* 34:620-621.
- Boch, J., and Bonas, U. 2010. *Xanthomonas* AvrBs3 family-type III effectors: discovery and function. *Phytopathology* 48:419.
- Boller, T., and Felix, G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol* 60:379-406.
- Bouwmeester, K., Han, M., Blanco-Portales, R., Song, W., Weide, R., Guo, L.-Y., van der Vossen, E. A., and Govers, F. 2014. The Arabidopsis lectin receptor kinase LecRK-I. 9 enhances resistance to *Phytophthora infestans* in Solanaceous plants. *Plant Biotechnology Journal* 12:10-16.
- Cai, R., Lewis, J., Yan, S., Liu, H., Clarke, C. R., Campanile, F., Almeida, N. F., Studholme, D. J., Lindeberg, M., and Schneider, D. 2011. The plant pathogen *Pseudomonas syringae* pv. *tomato* is genetically monomorphic and under strong selection to evade tomato immunity. *PLoS Pathog* 7:e1002130.
- Césari, S., Bernoux, M., Moncuquet, P., Kroj, T., and Dodds, P. 2014b. A novel conserved mechanism for plant NLR protein pairs: the 'integrated decoy' hypothesis. *Front Plant Sci* 5.
- Césari, S., Kanzaki, H., Fujiwara, T., Bernoux, M., Chalvon, V., Kawano, Y., Shimamoto, K., Dodds, P., Terauchi, R., and Kroj, T. 2014a. The NB-LRR proteins RGA4 and RGA5 interact functionally and physically to confer disease resistance. *The EMBO journal* 33:1941-1959.
- Chapman, S., Kavanagh, T., and Baulcombe, D. 1992. Potato virus X as a vector for gene expression in plants. *Plant J* 2:549-557.
- Chin, K., and Wolfe, M. 1984. The spread of *Erysiphe graminis* f. sp. *hordei* in mixtures of barley varieties. *Plant Pathol* 33:89-100.
- Chisholm, S. T., Coaker, G., Day, B., and Staskawicz, B. J. 2006. Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124:803-814.
- Clarke, C. R., Chinchilla, D., Hind, S. R., Taguchi, F., Miki, R., Ichinose, Y., Martin, G. B., Felix, G., and Vinatzer, B. A. 2013. Allelic variation in two distinct *Pseudomonas syringae* flagellin epitopes modulates the strength of plant immune responses but not bacterial motility. *New Phytol* 200:847-860.
- Cook, D. E., Mesarich, C. H., and Thomma, B. P. H. J. 2015. Understanding plant immunity as a surveillance system to detect invasion. *Annu Rev Phytopathol*.
- Cooke, I. R., Jones, D., Bowen, J. K., Deng, C., Faou, P., Hall, N. E., Jayachandran, V., Liem, M., Taranto, A. P., Plummer, K. M., and Mathivanan, S. 2014. Proteogenomic analysis of the *Venturia pirina* (Pear Scab Fungus) secretome reveals potential effectors. *J Proteome Res* 13:3635-3644.
- Croll, D., and McDonald, B. A. 2012. The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS Pathog* 8:e1002608.
- Dangl, J. L., Horvath, D. M., and Staskawicz, B. J. 2013. Pivoting the plant immune system from dissection to deployment. *Science* 341:746-751.



- de Guillen, K., Ortiz-Vallejo, D., Gracy, J., Fournier, E., Kroj, T., and Padilla, A. 2015. Structure analysis uncovers a highly diverse but structurally conserved effector family in phytopathogenic fungi. *PLoS Pathog* 11:e1005228.
- de Jonge, R., and Thomma, B. P. 2009. Fungal LysM effectors: extinguishers of host immunity? *Trends Microbiol* 17:151-157.
- de Jonge, R., Bolton, M. D., Kombrink, A., van den Berg, G. C., Yadeta, K. A., and Thomma, B. P. H. J. 2013. Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. *Genome Res* 23:1271-1282.
- de Jonge, R., van Esse, H. P., Maruthachalam, K., Bolton, M. D., Santhanam, P., Saber, M. K., Zhang, Z., Usami, T., Lievens, B., Subbarao, K. V., and Thomma, B. P. H. J. 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proc Natl Acad Sci U S A* 109:5110-5115.
- de Toledo Thomazella, D. P., Brail, Q., Dahlbeck, D., and Staskawicz, B. J. 2016. CRISPR-Cas9 mediated mutagenesis of a DMR6 ortholog in tomato confers broad-spectrum disease resistance. [BioRxiv:064824](https://doi.org/10.1101/064824).
- Deng, C. H., Plummer, K. M., Jones, D. A. B., Mesarich, C. H., Shiller, J., Taranto, A. P., Robinson, A. J., Kastner, P., Hall, N. E., Templeton, M. D., and Bowen, J. K. 2017. Comparative analysis of the predicted secretomes of Rosaceae scab pathogens *Venturia inaequalis* and *V. pirina* reveals expanded effector families and putative determinants of host range. *BMC Genomics* 18:339.
- Deslandes, L., Olivier, J., Theulières, F., Hirsch, J., Feng, D. X., Bittner-Eddy, P., Beynon, J., and Marco, Y. 2002. Resistance to *Ralstonia solanacearum* in *Arabidopsis thaliana* is conferred by the recessive *RRS1-R* gene, a member of a novel family of resistance genes. *Proceedings of the National Academy of Sciences* 99:2404-2409.
- Dong, S., Raffaele, S., and Kamoun, S. 2015. The two-speed genomes of filamentous pathogens: waltz with plants. *Current Opinion in Genetics & Development* 35:57-65.
- Dong, S., Stam, R., Cano, L. M., Song, J., Sklenar, J., Yoshida, K., Bozkurt, T. O., Oliva, R., Liu, Z., Tian, M., Win, J., Banfield, M. J., Jones, A. M., van der Hoorn, R. A., and Kamoun, S. 2014. Effector specialization in a lineage of the Irish potato famine pathogen. *Science* 343:552-555.
- Eckardt, N. A. 2002. Plant disease susceptibility genes? *Plant Cell* 14:1983-1986.
- Faino, L., and Thomma, B. P. 2014. Get your high-quality low-cost genome sequence. *Trends Plant Sci* 19:288-291.
- Faino, L., Seidl, M. F., Shi-Kunne, X., Pauper, M., van den Berg, G. C., Wittenberg, A. H., and Thomma, B. P. 2016. Transposons passively and actively contribute to evolution of the two-speed genome of a fungal pathogen. *Genome Res* 26:1091-1100.
- Ficarra, F. A., Grandellis, C., Garavaglia, B. S., Gottig, N., and Ottado, J. 2017. Bacterial and plant natriuretic peptides improve plant defense responses against pathogens. *Mol Plant Pathol*.
- Flor, H. H. 1942. Inheritance of pathogenicity in *Melampsora lini*. *Phytopathology* 32:653-669.
- Franceschetti, M., Maqbool, A., Jiménez-Dalmaroni, M. J., Pennington, H. G., Kamoun, S., and Banfield, M. J. 2017. Effectors of filamentous plant pathogens: commonalities amid diversity. *Microbiology and Molecular Biology Reviews* 81:e00066-00016.
- Giannakopoulou, A., Steele, J. F., Segretin, M. E., Bozkurt, T. O., Zhou, J., Robatzek, S., Banfield, M. J., Pais, M., and Kamoun, S. 2015. Tomato I2 immune receptor can be engineered to confer partial resistance to the oomycete *Phytophthora infestans* in addition to the fungus *Fusarium oxysporum*. *Mol. Plant Microbe Interact.* 28:1316-1329.
- Giesbert, S., Schumacher, J., Kupas, V., Espino, J., Segmuller, N., Haeuser-Hahn, I., Schreier, P. H., and Tudzynski, P. 2012. Identification of pathogenesis-associated genes by T-DNA-mediated insertional mutagenesis in *Botrytis cinerea*: a type 2A phosphoprotein phosphatase and an SPT3 transcription factor have significant impact on virulence. *Mol. Plant Microbe Interact.* 25:481-495.
- Glazebrook, J. 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol* 43:205-227.

- Gottig, N., Garavaglia, B. S., Daurelio, L. D., Valentine, A., Gehring, C., Orellano, E. G., and Ottado, J. 2008. *Xanthomonas axonopodis* pv. *citri* uses a plant natriuretic peptide-like protein to modify host homeostasis. *Proc Natl Acad Sci U S A* 105:18631-18636.
- Gressel, J., Gassmann, A. J., and Owen, M. D. 2017. How well will stacked transgenic pest/herbicide resistances delay pests from evolving resistance? *Pest Management Science* 73:22-34.
- Haas, B. J., Kamoun, S., Zody, M. C., Jiang, R. H., Handsaker, R. E., Cano, L. M., Grabherr, M., Kodira, C. D., Raffaele, S., and Torto-Alalibo, T. 2009. Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature* 461:393-398.
- Haverkort, A. J., Boonekamp, P. M., Hutten, R., Jacobsen, E., Lotz, L. A. P., Kessel, G. J. T., Vossen, J. H., and Visser, R. G. F. 2016. Durable Late Blight Resistance in Potato Through Dynamic Varieties Obtained by Cisgenesis: Scientific and Societal Advances in the DuRPh Project. *Potato Research* 59:35-66.
- Hemetsberger, C., Herrberger, C., Zechmann, B., Hillmer, M., and Doeblemann, G. 2012. The *Ustilago maydis* effector Pep1 suppresses plant immunity by inhibition of host peroxidase activity. *PLoS Pathog* 8:e1002684.
- Howlett, B. J., Lowe, R. G., Marcroft, S. J., and van de Wouw, A. P. 2015. Evolution of virulence in fungal plant pathogens: exploiting fungal genomics to control plant disease. *Mycologia* 107:441-451.
- Janssen, B.-J., and Gardner, R. C. 1990. Localized transient expression of GUS in leaf discs following cocultivation with *Agrobacterium*. *Plant Mol Biol* 14:61-72.
- Jelinski, N., Broz, K., Jonkers, W., Ma, L.-J., and Kistler, H. C. 2017. Effector Gene Suites in Some Soil Isolates of *Fusarium oxysporum* Are not Sufficient Predictors of Vascular Wilt in Tomato. *Phytopathology*.
- Jones, J. D., and Dangl, J. L. 2006. The plant immune system. *Nature* 444:323-329.
- Jørgensen, J. H. 1992. Discovery, characterization and exploitation of Mlo powdery mildew resistance in barley. Pages 141-152 in: *Breeding for disease resistance*. Springer.
- Keen, N. 1990. Gene-for-gene complementarity in plant-pathogen interactions. *Annu Rev Gen* 24:447-463.
- Kim, S. H., Qi, D., Ashfield, T., Helm, M., and Innes, R. W. 2016. Using decoys to expand the recognition specificity of a plant disease resistance protein. *Science* 351:684-687.
- Kombrink, A., and Thomma, B. P. H. J. 2013. LysM effectors: secreted proteins supporting fungal life. *PLoS Pathog* 9:e1003769.
- Kroj, T., Chanclud, E., Michel-Romiti, C., Grand, X., and Morel, J. B. 2016. Integration of decoy domains derived from protein targets of pathogen effectors into plant immune receptors is widespread. *New Phytol* 210:618-626.
- Kusch, S., and Panstruga, R. 2017. *mlo*-Based Resistance: An Apparently Universal "Weapon" to Defeat Powdery Mildew Disease. *Mol. Plant Microbe Interact.* 30:179-189.
- Lacombe, S., Rougon-Cardoso, A., Sherwood, E., Peeters, N., Dahlbeck, D., van Esse, H. P., Smoker, M., Rallapalli, G., Thomma, B. P. H. J., Staskawicz, B. J., Jones, J. D., and Zipfel, C. 2010. Interfamily transfer of a plant pattern-recognition receptor confers broad-spectrum bacterial resistance. *Nat Biotechnol* 28:365-369.
- Liang, Z., Chen, K., Li, T., Zhang, Y., Wang, Y., Zhao, Q., Liu, J., Zhang, H., Liu, C., and Ran, Y. 2017. Efficient DNA-free genome editing of bread wheat using CRISPR/Cas9 ribonucleoprotein complexes. *Nat Commun* 8.
- Lievens, B., Houterman, P. M., and Rep, M. 2009. Effector gene screening allows unambiguous identification of *Fusarium oxysporum* f. sp. *lycopersici* races and discrimination from other formae speciales. *FEMS Microbiol Lett* 300:201-215.
- Liu, J., Liu, D., Tao, W., Li, W., Wang, S., Chen, P., Cheng, S., and Gao, D. 2000. Molecular marker-facilitated pyramiding of different genes for powdery mildew resistance in wheat. *Plant Breeding* 119:21-24.

- Lorang, J., Kidarsa, T., Bradford, C. S., Gilbert, B., Curtis, M., Tzeng, S. C., Maier, C. S., and Wolpert, T. J. 2012. Tricking the guard: exploiting plant defense for disease susceptibility. *Science* 338:659-662.
- Lozano-Torres, J. L., Wilbers, R. H. P., Gawronski, P., Boshoven, J. C., Finkers-Tomczak, A., Cordewener, J. H. G., America, A. H. P., Overmars, H. A., Van 't Klooster, J. W., Baranowski, L., Sobczak, M., Ilyas, M., van der Hoorn, R. A. L., Schots, A., de Wit, P. J. G. M., Bakker, J., Goverse, A., and Smant, G. 2012. Dual disease resistance mediated by the immune receptor Cf-2 in tomato requires a common virulence target of a fungus and a nematode. *Proc Natl Acad Sci U S A* 109:10119-10124.
- Ma, L. J., van der Does, H. C., Borkovich, K. A., Coleman, J. J., Daboussi, M. J., Di Pietro, A., Dufresne, M., Freitag, M., Grabherr, M., Henrissat, B., Houterman, P. M., Kang, S., Shim, W. B., Woloshuk, C., Xie, X., Xu, J. R., Antoniw, J., Baker, S. E., Bluhm, B. H., Breakspear, A., Brown, D. W., Butchko, R. A., Chapman, S., Coulson, R., Coutinho, P. M., Danchin, E. G., Diener, A., Gale, L. R., Gardiner, D. M., Goff, S., Hammond-Kosack, K. E., Hilburn, K., Hua-Van, A., Jonkers, W., Kazan, K., Kodira, C. D., Koehrsen, M., Kumar, L., Lee, Y. H., Li, L., Manners, J. M., Miranda-Saavedra, D., Mukherjee, M., Park, G., Park, J., Park, S. Y., Proctor, R. H., Regev, A., Ruiz-Roldan, M. C., Sain, D., Sakthikumar, S., Sykes, S., Schwartz, D. C., Turgeon, B. G., Wapinski, I., Yoder, O., Young, S., Zeng, Q., Zhou, S., Galagan, J., Cuomo, C. A., Kistler, H. C., and Rep, M. 2010. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464:367-373.
- Maroof, S., Jeong, S., Gunduz, I., Tucker, D., Buss, G., and Tolin, S. 2008. Pyramiding of soybean mosaic virus resistance genes by marker-assisted selection. *Crop Science* 48:517-526.
- McDonald, B. A., and Linde, C. 2002. Pathogen population genetics, evolutionary potential, and durable resistance. *Annu Rev Phytopathol* 40:349-379.
- Mesarich, C. H., Griffiths, S. A., van der Burgt, A., Okmen, B., Beenen, H. G., Etalo, D. W., Joosten, M. H., and de Wit, P. J. G. M. 2014. Transcriptome sequencing uncovers the Avr5 avirulence gene of the tomato leaf mold pathogen *Cladosporium fulvum*. *Mol. Plant Microbe Interact.* 27:846-857.
- Michelmore, R. W., Christopoulou, M., and Caldwell, K. S. 2013. Impacts of resistance gene genetics, function, and evolution on a durable future. *Annu Rev Phytopathol* 51:291-319.
- Michielse, C. B., van Wijk, R., Reijnen, L., Cornelissen, B. J., and Rep, M. 2009. Insight into the molecular requirements for pathogenicity of *Fusarium oxysporum* f. sp. *lycopersici* through large-scale insertional mutagenesis. *Genome Biol* 10:R4.
- Munch, S., Ludwig, N., Floss, D. S., Sugui, J. A., Koszucka, A. M., Voll, L. M., Sonnewald, U., and Deising, H. B. 2011. Identification of virulence genes in the corn pathogen *Colletotrichum graminicola* by *Agrobacterium tumefaciens*-mediated transformation. *Mol Plant Pathol* 12:43-55.
- Narusaka, M., Shirasu, K., Noutoshi, Y., Kubo, Y., Shiraishi, T., Iwabuchi, M., and Narusaka, Y. 2009. RRS1 and RPS4 provide a dual Resistance-gene system against fungal and bacterial pathogens. *Plant J* 60:218-226.
- Nekrasov, V., Wang, C., Win, J., Lanz, C., Weigel, D., and Kamoun, S. 2017. Rapid generation of a transgene-free powdery mildew resistant tomato by genome deletion. *Scientific Reports* 7:482.
- Oome, S., Raaymakers, T. M., Cabral, A., Samwel, S., Böhm, H., Albert, I., Nürnberger, T., and Van den Ackerveken, G. 2014. Nep1-like proteins from three kingdoms of life act as a microbe-associated molecular pattern in Arabidopsis. *Proc Natl Acad Sci U S A* 111:16955-16960.
- Parra, L., Maisonneuve, B., Lebeda, A., Schut, J., Christopoulou, M., Jeuken, M., McHale, L., Truco, M.-J., Crute, I., and Michelmore, R. 2016. Rationalization of genes for resistance to *Bremia lactucae*. *Euphytica* 210:309-326.
- Pemberton, C. L., and Salmond, G. P. 2004. The Nep1-like proteins—a growing family of microbial elicitors of plant necrosis. *Mol Plant Pathol* 5:353-359.
- Plett, J. M., and Martin, F. 2015. Reconsidering mutualistic plant-fungal interactions through the lens of effector biology. *Curr Opin Plant Biol* 26:45-50.
- Potnis, N., Minsavage, G., Smith, J. K., Hurlbert, J. C., Norman, D., Rodrigues, R., Stall, R. E., and Jones, J. B. 2011. Avirulence Proteins AvrBs7 from *Xanthomonas gardneri* and AvrBs1.1 from *Xanthomonas euvesicatoria* Contribute to a Novel Gene-for-Gene Interaction in Pepper. *Mol. Plant Microbe Interact.* 25:307-320.

- Puchta, H. 2017. Applying CRISPR/Cas for genome engineering in plants: the best is yet to come. *Curr Opin Plant Biol* 36:1-8.
- Redkar, A., Villajuana-Bonequi, M., and Doehlemann, G. 2015. Conservation of the *Ustilago maydis* Effector See1 in related smuts. *Plant Signal Behav* 10:e1086855.
- Rodriguez-Moreno, L., Song, Y., and Thomma, B. P. H. J. 2017. Transfer and engineering of immune receptors to improve recognition capacities in crops. *Curr Opin Plant Biol* 38:42-49.
- Rouxel, T., Penaud, A., Pinochet, X., Brun, H., Gout, L., Delourme, R., Schmit, J., and Balesdent, M.-H. 2003. A 10-year survey of populations of *Leptosphaeria maculans* in France indicates a rapid adaptation towards the *Rlm1* resistance gene of oilseed rape. *Eur J Plant Pathol* 109:871-881.
- Rovenich, H., Boshoven, J. C., and Thomma, B. P. H. J. 2014. Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. *Curr Opin Plant Biol* 20:96-103.
- Santhanam, P., van Esse, H. P., Albert, I., Faino, L., Nurnberger, T., and Thomma, B. P. H. J. 2013. Evidence for functional diversification within a fungal NEP1-like protein family. *Mol. Plant Microbe Interact.* 26:278-286.
- Santhanam, P., Boshoven, J. C., Salas, O., Bowler, K., Islam, M. T., Saber, M. K., van den Berg, G., Barpeled, M., and Thomma, B. P. 2017. Rhamnose synthase activity is required for pathogenicity of the vascular wilt fungus *Verticillium dahliae*. *Mol Plant Pathol* 18:347-362.
- Sarris, P. F., Cevik, V., Dagdas, G., Jones, J. D., and Krasileva, K. V. 2016. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. *BMC biology* 14:8.
- Schimpl, S., and Puchta, H. 2016. Revolutionizing plant biology: multiple ways of genome engineering by CRISPR/Cas. *Plant methods* 12:8.
- Schmidt, S. M., Lukasiewicz, J., Farrer, R., Dam, P., Bertoldo, C., and Rep, M. 2016. Comparative genomics of *Fusarium oxysporum* f. sp. *melonis* reveals the secreted protein recognized by the Fom2 resistance gene in melon. *New Phytol* 209:307-318.
- Schmidt, S. M., Houterman, P. M., Schreiber, I., Ma, L., Amyotte, S., Chellappan, B., Boeren, S., Takken, F. L., and Rep, M. 2013. MITEs in the promoters of effector genes allow prediction of novel virulence genes in *Fusarium oxysporum*. *BMC Genomics* 14:119.
- Schwartz, A. R., Potnis, N., Timilsina, S., Wilson, M., Patané, J., Martins, J., Minsavage, G. V., Dahlbeck, D., Akhunova, A., and Almeida, N. 2015. Phylogenomics of *Xanthomonas* field strains infecting pepper and tomato reveals diversity in effector repertoires and identifies determinants of host specificity. *Frontiers in Microbiology* 6:535.
- Segretin, M. E., Pais, M., Franceschetti, M., Chaparro-Garcia, A., Bos, J. I. B., Banfield, M. J., and Kamoun, S. 2014. Single Amino Acid Mutations in the Potato Immune Receptor R3a Expand Response to *Phytophthora* Effectors. *Mol. Plant Microbe Interact.* 27:624-637.
- Song, Y., Zhang, Z., Boshoven, J., Rovenich, H., Seidl, M., Jakse, J., Maruthachalam, K., Liu, C.-M., Subbarao, K., and Javornik, B. 2017. Tomato immune receptor Ve1 recognizes surface-exposed co-localized N- and C-termini of *Verticillium dahliae* effector Ave1. *BioRxiv*:103473.
- Sprague, S., Marcroft, S., Hayden, H., and Howlett, B. 2006a. Major gene resistance to blackleg in *Brassica napus* overcome within three years of commercial production in southeastern Australia. *Plant Dis* 90:190-198.
- Sprague, S. J., Balesdent, M.-H., Brun, H., Hayden, H. L., Marcroft, S. J., Pinochet, X., Rouxel, T., and Howlett, B. J. 2006b. Major gene resistance in *Brassica napus* (oilseed rape) is overcome by changes in virulence of populations of *Leptosphaeria maculans* in France and Australia. *Eur J Plant Pathol* 114:33-40.
- Stukenbrock, E. H., Jorgensen, F. G., Zala, M., Hansen, T. T., McDonald, B. A., and Schierup, M. H. 2010. Whole-genome and chromosome evolution associated with host adaptation and speciation of the wheat pathogen *Mycosphaerella graminicola*. *PLoS Genet* 6:e1001189.

- Sun, W., Dunning, F. M., Pfund, C., Weingarten, R., and Bent, A. F. 2006. Within-species flagellin polymorphism in *Xanthomonas campestris* pv *campestris* and its impact on elicitation of Arabidopsis FLAGELLIN SENSING2-dependent defenses. *Plant Cell* 18:764-779.
- Svitashev, S., Schwartz, C., Lenderts, B., Young, J. K., and Cigan, A. M. 2016. Genome editing in maize directed by CRISPR-Cas9 ribonucleoprotein complexes. *Nat Commun* 7:13274.
- Tan, M. Y. A., Hutten, R. C. B., Visser, R. G. F., and van Eck, H. J. 2010. The effect of pyramiding *Phytophthora infestans* resistance genes R Pi-mcd1 and R Pi-ber in potato. *Theor Appl Genet* 121:117-125.
- Thomma, B. P. H. J., Nurnberger, T., and Joosten, M. H. A. J. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* 23:4-15.
- Van Dam, P., Fokkens, L., Schmidt, S. M., Linmans, J. H., Kistler, H. C., Ma, L. J., and Rep, M. 2016. Effector profiles distinguish formae speciales of *Fusarium oxysporum*. *Environmental Microbiology*.
- Van Damme, M., Huibers, R. P., Elberse, J., and Van den Ackerveken, G. 2008. Arabidopsis DMR6 encodes a putative 2OG-Fe (II) oxygenase that is defense-associated but required for susceptibility to downy mildew. *Plant J* 54:785-793.
- Van der Biezen, E. A., and Jones, J. D. 1998. Plant disease-resistance proteins and the gene-for-gene concept. *Trends Biochem Sci* 23:454-456.
- van Schie, C. C., and Takken, F. L. 2014. Susceptibility genes 101: how to be a good host. *Annu Rev Phytopathol* 52:551-581.
- Vleeshouwers, V. G., and Oliver, R. P. 2014. Effectors as tools in disease resistance breeding against biotrophic, hemibiotrophic, and necrotrophic plant pathogens. *Mol. Plant Microbe Interact.* 27:196-206.
- Vleeshouwers, V. G., Raffaele, S., Vossen, J. H., Champouret, N., Oliva, R., Segretin, M. E., Rietman, H., Cano, L. M., Lokossou, A., and Kessel, G. 2011. Understanding and exploiting late blight resistance in the age of effectors. *Annu Rev Phytopathol* 49:507-531.
- Waltz, E. 2016. CRISPR-edited crops free to enter market, skip regulation. *Nature Publishing Group*.
- Wolter, F., and Puchta, H. 2017. Knocking out consumer concerns and regulator's rules: efficient use of CRISPR/Cas ribonucleoprotein complexes for genome editing in cereals. *Genome Biol* 18:43.
- Zhou, B. J., Jia, P. S., Gao, F., and Guo, H. S. 2012. Molecular characterization and functional analysis of a necrosis- and ethylene-inducing, protein-encoding gene family from *Verticillium dahliae*. *Mol. Plant Microbe Interact.* 25:964-975.
- Zhu, S., Li, Y., Vossen, J. H., Visser, R. G., and Jacobsen, E. 2012. Functional stacking of three resistance genes against *Phytophthora infestans* in potato. *Transgenic research* 21:89-99.
- Zhu, S., Duwal, A., Su, Q., Vossen, J. H., Visser, R. G., and Jacobsen, E. 2013. Vector integration in triple *R* gene transformants and the clustered inheritance of resistance against potato late blight. *Transgenic research* 22:315-325.



**Summary**

**Acknowledgements**

**About the author**

**List of publications**

**Overview of training activities**



## Summary

Disease resistance in crops is an important aspect of securing global food security. Resistant plants carry immune receptors that sense pathogen invasion often through the recognition of important pathogen virulence factors, known as effectors. Thus, identification and characterization of effectors is important for the fundamental understanding of virulence mechanisms and to aid in resistance breeding. In **Chapter 1**, the motivation for selecting the plant pathogen *Verticillium dahliae* as a study object is explained. The main objectives and research questions are described and an outline of the thesis is provided. **Chapter 2** reviews the various roles of effectors of filamentous plant pathogens and explains that effectors can basically deregulate any step of plant immunity in any cellular compartment. Besides in plant pathogens, endophytes and mutualistic organisms carry effectors that suppress host immunity as well. In addition, effectors have been identified in saprophytes and other non-pathogenic plant-associated microbes and are proposed to act in microbial competition and cooperation. We propose a role for effectors in shaping the plant microbiome and argue that it is important for a microbe to rapidly diversify its effector repertoire to establish a microbial niche.

The genus *Verticillium* contains soil-borne fungal pathogens causing *Verticillium* wilts in a wide range of host plants. The most effective means of controlling these pathogens is genetic disease resistance. The tomato immune receptor Ve1 provides resistance to race 1 *V. dahliae* strains that carry *Ave1* (*VdAve1*). Previously, it was shown that *VdAve1* contributes to the virulence of *V. dahliae* and that *VdAve1* is the effector that is recognized by Ve1. Homologs of *VdAve1* were mostly found in plants and in some plant pathogens, including the fungi *Fusarium oxysporum* f. sp. *lycopersici* (*FoAve1*), *Colletotrichum higginsianum* (*ChAve1*), *Cercospora beticola* (*CbAve1*) and the bacterium *Xanthomonas axonopodis* (*XacPNP*). In **Chapter 3** we analysed the role of the various *Ave1* homologs in fungal virulence. Targeted *Ave1* deletions were generated in *F. oxysporum*, *C. beticola* and *C. higginsianum* and we showed that *FoAve1* and *CbAve1* are virulence factors. This is in contrast to *ChAve1*, which does not contribute to fungal virulence on *Arabidopsis thaliana* plants, likely because this gene is not expressed during infection. In a complementation experiment we found that none of the fungal *Ave1* homologs can restore virulence in a *V. dahliae* *Ave1* deletion strain. Therefore we conclude that *VdAve1* on the one hand and *FoAve1*, *CbAve1*, *ChAve1* and *XacPNP* on the other hand, act in different manners. These homologs are differentially recognized by Ve1, with the most efficient recognition of *VdAve1*, and little to no recognition of *ChAve1* and *XacPNP*. In **Chapter 4** we show that C-terminal GFP-tagging of *VdAve1* resulted in loss of recognition by Ve1, suggesting that exposure of the C-terminus of *VdAve1* is essential for recognition by Ve1. Truncations of *VdAve1*,



and domain swaps between VdAve1 and XacPNP, narrowed down the sequence that is required for Ve1 recognition to the C-terminal nine amino acids. Site-directed mutagenesis was performed to analyse the contribution of individual amino acid residues to recognition by Ve1, revealing the importance of five amino acids from Val127 to Phe131. Nevertheless, fusions of the VdAve1 C-terminal nine amino acids to GFP or to the *Cladosporium fulvum* effector Avr9 showed that this sequence is not sufficient to activate Ve1-mediated immunity. Finally, with a complementation experiment in *V. dahliae* we confirmed that the C-terminal nine amino acids of Ave1 are required, but not sufficient, to activate Ve1-mediated immunity. **Chapter 5** describes the identification of an expanded family of Ave1 homologs in the apple and pear scab pathogens *Venturia inaequalis* and *V. pirina*, respectively. We show that tomato immune receptor Ve1 is unable to recognize Ave1 homologs derived from *V. inaequalis* and *V. pirina*. Transcriptome analysis revealed that *V. inaequalis* ViAve1-1, ViAve1-5 and ViAve1-9 are expressed *in planta* and that these homologs are unable to complement a *V. dahliae* VdAve1 deletion strain, suggesting that these homologs have different functions than *V. dahliae* VdAve1.

In **Chapter 6**, a gene with high similarity to VdAve1 was identified in *V. dahliae* strain JR2 as an Ave1-like (Ave1L). Subsequently, we identified a total of five alleles (Ave1L1 to Ave1L5) in the *V. dahliae* population. While some of the *V. dahliae* strains do not have Ave1L, other strains have one of the five Ave1L alleles in presence or absence of Ave1 itself. Some of the Ave1L alleles contain multiple single nucleotide polymorphisms, some of which result in a premature stop codon. In addition, one of the alleles contains a retrotransposon insertion. These observations suggest that Ave1L experienced significant selection pressure, potentially to evade recognition by an immune receptor. Thus, we attempted to predict the common progenitor sequence of the five Ave1L alleles. However, we show that neither of the Ave1L alleles, nor the predicted progenitor, are recognized by tomato immune receptor Ve2 that shows high homology to Ve1. We furthermore show that Ave1L2 is expressed during host colonization, but does not seem to contribute to virulence on tomato. Since *V. dahliae* has a wide host range, Ave1L potentially contributes to virulence on other host plants than tomato where it may have been recognized by an immune receptor that drove its divergence.

In **Chapter 7** random mutagenesis was performed on the *V. dahliae* genome, followed by a pathogenicity screen to identify genes that are important for host invasion. This resulted in the identification of 58 novel candidate genes. We focused on one of these candidates and showed that deletion of the VdNRS/ER gene from the *V. dahliae* genome results in complete loss of pathogenicity on tomato and *N. benthamiana* plants, whereas vegetative growth and sporulation are not affected.

We demonstrate that VdNRS/ER is a functional enzyme in the biosynthesis of UDP-rhamnose, and further analysis revealed that *VdNRS/ER* deletion strains are impaired in colonization of tomato roots. Collectively, our results demonstrate that rhamnose, although only being a minor cell wall component, is essential for pathogenicity of *V. dahliae*.

**Chapter 8** is the general discussion and describes how the knowledge of pathogen effectors can be used as a tool to establish durable broad spectrum resistance.



## Acknowledgements

Here I am finishing the last part of my PhD thesis. The last part that is written, but often the first thing that people read when they open a PhD thesis. I guess it is because the acknowledgements show the personal side of the story. It is not possible to finish a PhD thesis on your own but it involves many people including your supervisor, colleagues, friends and family. Here I would like to take the opportunity to thank everybody involved.

I have mixed feelings on finishing the book but I feel mostly proud and relieved that it is done. Without my promotor Bart it would not have happened. I met Bart already in 2008, for my Hbo internship in the Verticillium group. Your style of supervision, your enthusiasm and the nice atmosphere in the group convinced me already then to keep my eyes open for a PhD position in your group. After obtaining my MSc degree I joined the Verticillium group as a PhD candidate. Bart, I am very happy you gave me the opportunity to start a project under your supervision. I enjoyed the freedom you gave me to find my own way in the project. Your office door was always open. Whenever I got stuck or lost in a situation you were able to guide me through it without giving me the answers. I am grateful for the many discussions we had about the project and the fast corrections and feedback I got on my manuscripts.

Pierre, my second promotor, thank you for the discussions in the beginning of the project and for helping me with the formalities at the end of the project. I feel honored to be your last PhD candidate.

I would like to thank the people in the Verticillium group who directly or indirectly helped me to get the experiments done in this thesis. Ronnie, for providing the data and material to initiate the project. Peter, for providing plasmids and constructs and showing me how to make good photos for publications. Partha, for teaching me how to transform *Verticillium dahliae*. Zhao and Yin, for providing strains and showing me how to do Agrobacterium infiltrations. Malaika, for helping me with the *Cercospora beticola* work. Eduardo, for all the qPCRs you run. Grady, for teaching me how the Verticillium lab is operating and for all the help. Dirk Jan, for the help you gave me with plasmids and constructs. Michael, Luigi and Xiaoqian, for helping me with the many questions I had regarding evolution, genomes, rearrangements, phylogenetic trees, alignments, blasts and other bioinformatics related questions. David and Andrea, for the many discussions we had in the lab. Jinling, Luis and Jasper, and all other Verticillium lab members thank you for the nice atmosphere in the lab. I also thank you all for the nice discussions during the coffee breaks and the time we spend together.

Then, I want to thank Jan, Francine and Matthieu for your input in the project. Klaas, Thomas, Daniela, Carl, Scott and Mansoor, thank you for all the useful suggestions and ideas but also the nice talks and times together. Thank you Ali, Rob, Grardy, Ester, Laurens and Natalie for making life easier in the lab, You made sure that the lab was up and running and that materials were available.

I would like to thank all the students that helped me with my projects. Jelle, Israel, Dorien, Saskia and Pascal without you it would have been a lot more work. Thank you Bert Essenstam and Pauline Sanderson at Unifarm for taking such good care of the plants in the greenhouse.

Dear Hanna, it was great to work with you on the same project. We had many work-related and also work-unrelated discussions and you made me feel that I was not struggling alone ☺. Our companionship gave me the positive energy to continue when things got difficult. Without you, these years would not have been the same. Apart from the PhD, I am happy we got the chance to become so close to and share beautiful moments together with you and Ale. I cherish all of them and I know that Iliana, Christof and Milo also do. I wish you all the best for the coming weeks and I am looking forward to meet the little one.

Mireille, I think we have managed to discuss almost everything, from experimental setup to family life. You were always in for a chat in the lab or at the coffee table. The talks were a nice distraction from all the lab work and writing. I am convinced that without you my PhD time would have been a lot more boring and much more challenging. Thank you for being my paronymph.

Anjaatje, as I used to reply when you called me either Jordietje or Gordi, we have managed to built up a lot of memories in the past few years; skiing together with the Verticillium group in Winterberg after our meeting in Marbourg; visiting San Francisco before the Fungal Genetics conference in California. I really missed your presence in the lab when you moved to Zurich for your Postdoc. Thank you for the talks, dinners, drinks during my PhD. I am happy we keep contact and we always have things to share and talk about. Being my paronymph will be one more thing to remember ☺

Thanks to my Bejo colleagues for the support and talks during the breaks and lunch walks.

I would like to thank all my friends in Wageningen (few of you still there) during my studies which I am proud to still call friends. Without you it would not have been fun. Thank you Bart, Alejandro, Hanna, Aniek, Melchior, Andrius, Vaida, Abel, Maaïke, Pavlo, Celine, Luigi, Alice, Bram, Anne, Xiaoqian, Tim, Bram O, Sandra, Leandre, Johanna, Victoria, Tulipan, Alexander, Jule, Peter, Lorena, Jeroen, Wouter and my friends in Athens Gianni, Ioanna and Eleni.

Thanks to my Greek family Dimitri, Maria, Stefano and Christina for supporting me. You always make me feel welcome and loved.

Many thanks to my parents Bert en Margreet for supporting me in all the years. I could not have done this without the confidence you had in me and the financial support you provided. I will always be very grateful for the opportunity you gave me to develop myself into the young professional I am now. Margreet, thank you for all the days you came to help in the house the last two years. Without your help I could not have finished this book.

Then last, but certainly not least, I want to thank my wife Iliana. I know it has been a tough time for you too. Thank you so much for your support and patience when I spent hours writing in the evenings and weekends. Thank you for taking care of everything else so that I wouldn't have to stress. Thank you for listening to all my frustration and desperation and for the confidence you show in me. Thank you for your love and care and for the lovely family we have together. Christof and Philip are beautiful boys that have put things in perspective for me. Without the three of you I would not have been able to write this thesis.

## About the author

Jordi C. Boshoven was born in Alphen aan den Rijn, The Netherlands, on April 1<sup>st</sup> 1986. At the age of 5 he moved to Maasdijk where he went to the PC Basisschool Het Kompas. His secondary education was performed at the Interconfessionele Scholengroep Westland (ISW) in Naaldwijk. During his holidays Jordi was working in the company of his uncle that is growing freesias in the greenhouse. In 2002, he went to the ROC Mondriaan in Delft where he was educated at Mbo level with a specialization in Botany to work in a laboratory. His first internship was performed at Rijk Zwaan in De Lier under the supervision of Theresia van Luijk where he studied seed pathology. His second internship was performed at Groen Agro Control in Delfgauw under the supervision of Dr. Adriaan Vermunt and Laurens Hanemaaijer where he studied the molecular detection of plant pathogens in diagnostic samples.



In 2006, he continued his studies at Hbo level at the Hogeschool van Arnhem en Nijmegen (HAN) in Nijmegen where he studied Plant Biotechnology. During his education in Nijmegen he performed a minor, an internship and a thesis at Wageningen University. His internship was performed at the Laboratory of Phytopathology under the supervision of Prof. Dr. Bart Thomma where he studied plant recognition of the human bacterial pathogen *Staphylococcus aureus*. His thesis was performed in the Laboratory of Nematology under the supervision of Dr. Jose Lozano-Torres where he studied the recognition of the nematode effector Gr-VAP-1 by the tomato immune receptor Cf-2 that resulted in a publication in the scientific journal Proceedings of the National Academy of Sciences of the United States of America.

After his studies in Nijmegen he moved to Wageningen in 2009 to do a MSc in Plant Biotechnology with the specialization Breeding and Pathology at Wageningen University. He performed his MSc internship in the group of Prof. Dr. Brian Staskawicz at the University of California in Berkley, USA under the supervision of Dr. Sandra Goritschnig where he studied the recognition of the oomycete *Hyaloperonospora arabidopsidis* effector ATR1 by the Arabidopsis immune receptor RPP1. His MSc thesis was performed at the Plant-Microbe Interactions group at Utrecht University under the supervision of Dr. Dieuwertje van der Does where he studied the jasmonic acid and salicylic acid plant defense signaling pathways in Arabidopsis.

After obtaining his MSc degree he started in 2012 as a PhD candidate in the Laboratory of Phytopathology under the supervision of Prof. Dr. Bart Thomma. He studied the virulence and recognition of *Verticillium dahliae* effector Ave1 homologs that resulted in the publication of this PhD thesis. During his PhD he organized several activities as a member of the PhD council of the graduate school of Experimental Plant Sciences (EPS). Since 2016 Jordi is working as a researcher phytopathology at Bejo Zaden in Warmenhuizen and he lives with his wife and two children in Koedijk.

## List of publications


- Lozano-Torres, J. L., Wilbers, R. H. P., Gawronski, P., **Boshoven, J. C.**, Finkers-Tomczak, A., Cordewener, J. H. G., America, A. H. P., Overmars, H. A., Van 't Klooster, J. W., Baranowski, L., Sobczak, M., Ilyas, M., van der Hoorn, R. A. L., Schots, A., de Wit, P. J. G. M., Bakker, J., Goverse, A., and Smant, G. 2012. Dual disease resistance mediated by the immune receptor Cf-2 in tomato requires a common virulence target of a fungus and a nematode. *Proc Natl Acad Sci U S A* 109:10119-10124.
- Rovenich, H., **Boshoven, J. C.**, and Thomma, B. P. H. J. 2014. Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. *Curr Opin Plant Biol* 20:96-103.
- Santhanam, P., **Boshoven, J. C.**, Salas, O., Bowler, K., Islam, M. T., Saber, M. K., van den Berg, G., Bar-Peled, M., and Thomma, B. P. 2017. Rhamnose synthase activity is required for pathogenicity of the vascular wilt fungus *Verticillium dahliae*. *Mol Plant Pathol* 18:347-362.
- Song, Y., Zhang, Z., **Boshoven, J. C.**, Rovenich, H., Seidl, M., Jakse, J., Maruthachalam, K., Liu, C.-M., Subbarao, K., and Javornik, B. 2017b. Tomato immune receptor Ve1 recognizes surface-exposed co-localized N-and C-termini of *Verticillium dahliae* effector Ave1. *BioRxiv*:103473.



## Education Statement of the Graduate School

### Experimental Plant Sciences

Issued to: Jordi C. Boshoven  
 Date: 10 November 2017  
 Group: Laboratory of Phytopathology  
 University: Wageningen University & Research



1) Start-up phase	<u>date</u>
▶ <b>First presentation of your project</b>	
Title: The role of <i>Vorticillium dahliae</i> Ave1	16 Jan 2012
▶ <b>Writing or rewriting a project proposal</b>	
Title: The role of <i>Vorticillium dahliae</i> Ave1 in fungal virulence and its recognition by the tomato Ve1 immune receptor	25 jan 2012
▶ <b>Writing a review or book chapter</b>	
Title: Filamentous Pathogen Effector Functions: Of Pathogens, Hosts and the Microbiome, <i>Current Opinion in Plant Biology</i> 20, 2014, 96-103; DOI: 10.1016/j.pbi.2014.05.001	31 Jan 2014
▶ <b>MSc courses</b>	
<b>Laboratory use of isotopes</b>	
<i>Subtotal Start-up Phase</i> 8.5 credits*	

2) Scientific Exposure	<u>date</u>
▶ <b>EPS PhD student days</b>	
EPS PhD student day, Amsterdam, NL	30 Nov 2012
5th European Plant Science Retreat, Ghent University, Ghent, Belgium	23-26 Jul 2013
EPS PhD student day, Leiden, NL	29 Nov 2013
6th European Plant Science Retreat, UvA, Amsterdam, NL	01-04 Jul 2014
EPS PhD student day (GET2GETHER) Soest, NL	29-30 Jan 2015
▶ <b>EPS theme symposia</b>	
EPS Theme 2: Interactions between Plants and Biotic Agents & Willie Commelin Scholten Day, Wageningen, NL	10 Feb 2012
EPS Theme 2: Interactions between Plants and Biotic Agents & Willie Commelin Scholten Day, Utrecht, NL	24 Jan 2013
EPS Theme 2: Interactions between Plants and Biotic Agents & Willie Commelin Scholten Day, Amsterdam, NL	25 Feb 2014
EPS Theme 2: Interactions between Plants and Biotic Agents & Willie Commelin Scholten Day, Utrecht, NL	20 Feb 2015
▶ <b>National meetings (e.g. Lunteren days) and other National Platforms</b>	
Annual meeting 'Experimental Plant Sciences', Lunteren, NL	02-03 Apr 2012
Annual meeting 'Experimental Plant Sciences', Lunteren, NL	22-23 Apr 2013
Annual meeting 'Experimental Plant Sciences', Lunteren, NL	14-15 Apr 2014
Annual meeting 'Experimental Plant Sciences', Lunteren, NL	13 Apr 2015

► Seminars (series), workshops and symposia	
<i>Workshops:</i>	
KLV speedreading course	08 Feb 2012
Lunchlezing "Initiatief nemen en assertiviteit - Hoe assertief kun je zijn?"	24 Sep 2014
Lunchlezing "Van spanning naar ontspanning, spelen met de dialoog"	18 Nov 2015
<i>Seminars:</i>	
Monique van Oers 'Insect viruses and their applications'	23 Feb 2012
Ralph Panstruga 'Comparative pathogenomics of powdery mildew fungi'.	04 Dec 2012
Andrew Sugden (Editor of Science magazine) 'Demystifying Science'	08 Feb 2013
Detleif Weigel 'Origin and consequence of genetic and epigenetic variation'	27 Feb 2013
Kazuki Saito 'Metabolomics-based functional genomics - from Arabidopsis to crops and medicinal plants'	08 Apr 2013
Brian Staskawicz "Effector-Targeted Breeding for Durable Disease Control of Xanthomonas diseases in Tomato and Cassava"	21 May 2013
Pieter Dorrestein "A "GoogleMAP"-type molecular view of microbes - from culture to people"	22 Aug 2013
Hanhui Kuang "Using the Nicotiana-TMV system to study resistance gene evolution and plant genome stability"	11 Sep 2013
David Weller 'Soilborne Pathogens and their Natural Biocontrol Agents in Cereal-Based Production Systems'	25 Sep 2013
Ross Mann "Endophytes in agriculture - evaluating their application via metabolomics and genomics"	04 Oct 2013
Jos Raaijmakers 'Back to the roots: exploring and exploiting the plant microbiome'	07 Jan 2014
Plant Sciences Seminar: Mutualism in Action (Toby Kiers and René Geurts)	11 Mar 2014
Jeroen Mesters 'From protein solution to single crystal X-ray diffraction: Chitin binding by LysM domains at atomic resolution'	31 Mar 2014
Jane Parker 'Reprogramming cells for defence in plant innate immunity'	09 Apr 2014
Frank van Breusegem 'Plant Metacaspases'	09 Apr 2014
Yuanchao Wang "Dissecting the interaction between Phytophthora sojae and soybean: making sense out of signalling and effectors"	16 Jul 2014
Joy Bergelson WEES seminar "Maintaining an ancient balanced polymorphism for resistance amidst diffuse intereactions"	26 Sep 2014
Michael Freitag "Chromatin structure controls centromeres and secondary metabolism in filamentous fungi"	21 Oct 2014
Ortrun Mittelsten Scheid "Genetics and epigenetics: a complex relationship"	19 Nov 2014
Kevin Foster WEES seminar "The evolution of cooperation and competition in microbes"	22 Jan 2015
Yves van de Peer 'The evolutionary significance of gene and genome duplications'	03 Feb 2015
Monica Höfte 'Towards understanding rice brown spot, a disease induced by physiological stress'	06 Feb 2015
Chih-Hang Wu 'Helper NLR proteins of the NRC family in solanaceous plants'	05 Mar 2015
Gero Steinberg 'Long-distance endosome trafficking drives fungal effector production during plant infection'	05 Jun 2015

Gert Kema application as Special Professor in Tropical Phytopathology	17 Nov 2015
<i>Symposia:</i>	
Plantum 'Intraspecific Pathogen Variation - Implications and Opportunities'	22 Jan 2013
<i>Minisymposium: How to write a world-class paper</i>	17 Oct 2013
KNPV fall meeting 'The process to progress'	27 Nov 2013
Minisymposium: Responses of plants to pathogenic micro-organisms and their secreted proteins	02 Dec 2013
Farewell Address Prof. dr. ir. Pierre de Wit - Fungal plant pathogens and the plant immune system	05 Jun 2014
All-inclusive Breeding: Integrating high-throughput science	16 Oct 2014
Minisymposium: Phytopathology	24 Nov 2014
Meeting BU Biointeractions & Plant Health – Phytopathology WU	10 Feb 2015
▶ <b>Seminar plus</b>	
Michael Freitag "Chromatin structure controls centromeres and secondary metabolism in filamentous fungi"	21 Oct 2014
Gero Steinberg 'Long-distance endosome trafficking drives fungal effector production during plant infection'	05 Jun 2015
▶ <b>International symposia and congresses</b>	
2nd joined Wageningen - Marburg meeting on plant-fungal interactions	30-31 Jan 2012
The 27th Fungal Genetics Conference 2013, Asilomar, California, USA	12-17 Mar 2013
The 28th Fungal Genetics Conference 2015, Asilomar, California, USA	17-22 Mar 2015
▶ <b>Presentations</b>	
<i>Poster:</i> 27th Fungal Genetics Conference 2013, Asilomar, California, USA	15 Mar 2013
<i>Poster:</i> Symposium "All-inclusive Breeding: Integrating high-throughput science", Wageningen, NL	16 Oct 2014
<i>Poster:</i> 28th Fungal Genetics Conference 2015, Asilomar, California, USA	17-22 Mar 2015
<i>Plant Science Slam:</i> Annual meeting 'Experimental Plant Sciences', Lunteren, NL	22 Apr 2013
<i>Talk:</i> 7th International Utrecht PhD Summer School, Utrecht, NL	27 Aug 2013
<i>Talk:</i> KNPV fall meeting 'The process to progress', Wageningen, NL	26 Nov 2013
<i>Talk:</i> 6th European Plant Science Retreat, UvA, Amsterdam, NL	03 Jul 2014
<i>Flash:</i> Symposium 'All-inclusive Breeding: Integrating high-throughput science', Wageningen, NL	16 Oct 2014
<i>Talk:</i> EPS Theme 2: Interactions between Plants and Biotic Agents & Willie Commelin Scholten Day Utrecht, NL	20 Feb 2015
▶ <b>IAB interview</b>	
▶ <b>Excursions</b>	
Company visit: Rijk Zwaan de Lier	27 Sep 2013

*Subtotal Scientific Exposure* 24.9 credits\*

<b>3) In-Depth Studies</b>	<i>date</i>
▶ <b>EPS courses or other PhD courses</b>	
7th International Utrecht PhD Summer School on Environmental Signaling, Utrecht, NL	26-28 Aug 2013
▶ <b>Journal club</b>	
Participated in the literature discussion of the Verticillium group	2012-2015
▶ <b>Individual research training</b>	
<i>Subtotal In-Depth Studies</i> 3.9 credits*	

<b>4) Personal development</b>	<i>date</i>
▶ <b>Skill training courses</b>	
Course: PhD competence assessment	20 Mar 2012
Course: Scientific publishing	19 Jun 2012
Course: Project- and Time Management	30 Oct-11 Dec 2012
Course: Interpersonal Communication for PhD Students	12-13 Dec -2012
WGS PhD Workshop Carousel 2015	17 Apr 2015
Course: Last Stretch of the PhD Programme	25 Sep 2015
▶ <b>Organisation of PhD students day, course or conference</b>	
EPS Expectations day 2013	01 Feb 2013
Company visit: Rijk Zwaan de Lier	27 Sep 2013
EPS Expectations day 2014	28 Mar 2014
Company visit: Genetwister and In2Care Wageningen	19 Sep 2014
EPS PhD student day (GET2GETHER)	29-30 Jan 2015
▶ <b>Membership of Board, Committee or PhD council</b>	
EPS PhD council member	2012-2015
<i>Subtotal Personal Development</i> 7.7 credits*	

<b>TOTAL NUMBER OF CREDIT POINTS*</b>	<b>45.0</b>
---------------------------------------	-------------

Herewith the Graduate School declares that the PhD candidate has complied with the educational requirements set by the Educational Committee of EPS which comprises of a minimum total of 30 ECTS credits

\* A credit represents a normative study load of 28 hours of study.

This work was supported by a VICI grant from the Research Council for Earth and Life sciences (ALW) of the Netherlands Organization for Scientific Research (NWO). The work was carried out in the Laboratory of Phytopathology, Wageningen University, the Netherlands.

**Cover design:** Iliana Boshoven-Gkini | AgileColor.com

**Thesis layout:** Iliana Boshoven-Gkini | AgileColor.com

**Print:** GVO drukkers & vormgevers | gvo.nl

