

ANNALES DE L'INSTITUT FOURIER

Andrei A. AGRACHEV, DAVIDE BARILARI & ELISA PAOLI

**Volume geodesic distortion and Ricci curvature for
Hamiltonian dynamics**

Tome 69, n° 3 (2019), p. 1187-1228.

http://aif.centre-mersenne.org/item/AIF_2019__69_3_1187_0

© Association des Annales de l'institut Fourier, 2019,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>



VOLUME GEODESIC DISTORTION AND RICCI CURVATURE FOR HAMILTONIAN DYNAMICS

by **Andrei A. AGRACHEV**,
Davide BARILARI & Elisa PAOLI (*)

ABSTRACT. — We study the variation of a smooth volume form along extremals of a variational problem with nonholonomic constraints and an action-like Lagrangian. We introduce a new invariant, called volume geodesic derivative, describing the interaction of the volume with the dynamics and we study its basic properties. We then show how this invariant, together with curvature-like invariants of the dynamics, appear in the asymptotic expansion of the volume. This generalizes the well-known expansion of the Riemannian volume in terms of Ricci curvature to a wide class of Hamiltonian flows, including all sub-Riemannian geodesic flows.

RÉSUMÉ. — On considère la variation d'un volume lisse le long des extrema d'un problème variationnel avec contraintes non holonomes et lagrangien de type action. On introduit un nouvel invariant, appelé dérivée canonique du volume, qui décrit l'interaction entre la forme volume et la dynamique. On montre comment cet invariant, avec des invariants de type courbure associés à la dynamique, apparaissent dans le développement asymptotique du volume. Cela généralise le développement classique du volume riemannien le long du flot géodésique en termes de la courbure de Ricci à une vaste classe de flots hamiltoniens, notamment tous les flots géodésiques sous-riemanniens.

1. Introduction

One of the possible ways of introducing Ricci curvature in Riemannian geometry is by computing the variation of the Riemannian volume under the geodesic flow.

Keywords: volume, geodesics, Ricci curvature, Hamiltonian systems, sub-Riemannian geometry.

2010 *Mathematics Subject Classification:* 53C17, 53B21, 53B15.

(*) This research has been supported by the European Research Council, ERC StG 2009 “GeCoMethods”, contract number 239748 and by the ANR project SRGI “Sub-Riemannian Geometry and Interactions”, contract number ANR-15-CE40-0018.

Given a point x on a Riemannian manifold (M, g) and a tangent unit vector $v \in T_x M$, it is well-known that the asymptotic expansion of the Riemannian volume vol_g in the direction of v depends on the Ricci curvature at x . More precisely, let us consider a geodesic $\gamma(t) = \exp_x(tv)$ starting at x with initial tangent vector v . In normal coordinates, the volume element is written as $\text{vol}_g = \sqrt{\det g_{ij}} dx_1 \dots dx_n$ and satisfies the following expansion for $t \rightarrow 0$

$$(1.1) \quad \sqrt{\det g_{ij}(\exp_x(tv))} = 1 - \frac{1}{6} \text{Ric}^g(v, v)t^2 + O(t^3),$$

where Ric^g is the Ricci curvature tensor associated with g (see for instance [13, Chapter 3] or [21, Chapter 14]).

The left hand side of (1.1) has a clear geometric interpretation. Indeed, fix an orthonormal basis e_1, \dots, e_n in $T_x M$ and let

$$\partial_i|_{\gamma(t)} := (d_{tv} \exp_x)(e_i) = \left. \frac{\partial}{\partial s} \right|_{s=0} \exp_x(tv + se_i), \quad 1 \leq i \leq n,$$

be the image of e_i through the differential of the Riemannian exponential map $\exp_x : T_x M \rightarrow M$ at tv . Once we take a set of normal coordinates centered at x , the vector fields ∂_i are the coordinate vector fields at $\gamma(t)$. Then the left hand side of (1.1) measures the Riemannian volume of the parallelotope with edges ∂_i at the point $\gamma(t)$, more explicitly

$$\sqrt{\det g_{ij}(\gamma(t))} = \text{vol}_g \left(\bigwedge_{i=1}^n \partial_i|_{\gamma(t)} \right).$$

The purpose of this paper is to study the variation of a smooth volume form along extremals of a variational problem with nonholonomic constraints and an action-like Lagrangian. To this aim, let us first consider the case of a weighted Riemannian manifold (M, g, μ) endowed with a smooth volume $\mu = e^\psi \text{vol}_g$, where ψ is a smooth function on M . Let $\exp_x(t, v)$ denote the exponential map defined at time t starting from x , i.e., set $\exp_x(t, v) := \exp_x(tv)$. Then

$$(1.2) \quad (d_v \exp_x(t, \cdot))(e_i) = \left. \frac{\partial}{\partial s} \right|_{s=0} \exp_x(t(v + se_i)) = t \partial_i|_{\gamma(t)}.$$

The volume of the parallelotope Q_t with edges $t \partial_i|_{\gamma(t)}$ has the following expansion for $t \rightarrow 0$,

$$(1.3) \quad \mu(Q_t) = t^n e^{\psi(\gamma(t))} \left(1 - \frac{1}{6} \text{Ric}^g(v, v)t^2 + O(t^3) \right),$$




Figure 1.1. Volume distortion on a weighted Riemannian manifold with volume $\mu = e^\psi \text{vol}_g$

as a direct consequence of (1.1) (see also Figure 1.1). By writing

$$\psi(\gamma(t)) = \psi(x) + \int_0^t g(\nabla\psi(\gamma(s)), \dot{\gamma}(s)) \, ds$$

we reduce the previous identity to tensorial quantities as follows

$$(1.4) \quad \mu(Q_t) = c_0 t^n e^{\int_0^t \rho(\dot{\gamma}(s)) \, ds} \left(1 - \frac{1}{6} \text{Ric}^g(v, v)t^2 + O(t^3) \right),$$

where we defined $\rho(w) = g(\nabla\psi(x), w)$ for every $w \in T_xM$ and $c_0 = e^{\psi(x_0)}$.

To understand the general case, it is convenient to reinterpret the last variation of volume from an Hamiltonian viewpoint. Indeed the Riemannian exponential map on M can be written in terms of the Hamiltonian flow associated with the smooth function $H : T^*M \rightarrow \mathbb{R}$ given in coordinates by

$$H(p, x) = \frac{1}{2} \sum_{i,j=1}^n g^{ij}(x) p_i p_j, \quad (p, x) \in T^*M,$$

where g^{ij} is the inverse matrix of the metric g . More geometrically, the Riemannian metric g induces a canonical linear isomorphism $\mathfrak{i} : T_xM \rightarrow T_x^*M$ between each tangent space T_xM and its dual T_x^*M . The function H is then (one half of the square of) the cometric, i.e., the metric g read as a function on covectors. If $\lambda = \mathfrak{i}(v)$ denotes the element in T_x^*M corresponding to $v \in T_xM$ under the above isomorphism, the exponential map satisfies

$$(1.5) \quad \gamma(t) = \exp_x(t, v) = \pi(e^{t\vec{H}}(\lambda)),$$

where $\pi : T^*M \rightarrow M$ is the canonical projection and \vec{H} is the Hamiltonian vector field on T^*M associated with H , whose coordinate expression is

$$(1.6) \quad \vec{H} = \sum_{j=1}^n \frac{\partial H}{\partial p_j} \frac{\partial}{\partial x_j} - \frac{\partial H}{\partial x_j} \frac{\partial}{\partial p_j}.$$




Figure 1.2. Volume distortion under the Hamiltonian flow

Denote now $E_i := \mathbf{i}(e_i)$ the frame of cotangent vectors in T_x^*M associated with the orthonormal frame $\{e_i\}_{i=1}^n$ of T_xM . Then, combining (1.2) and (1.5), we have $t\partial_i = (\pi \circ e^{t\bar{H}})_* E_i$ and the left hand side of (1.3) can be written as

$$\begin{aligned}
 \mu(Q_t) &= \langle \mu_{\gamma(t)}, (t\partial_1, \dots, t\partial_n) \rangle \\
 (1.7) \quad &= \left\langle \mu_{\pi(e^{t\bar{H}}(\lambda))}, \left((\pi \circ e^{t\bar{H}})_* E_1, \dots, (\pi \circ e^{t\bar{H}})_* E_n \right) \right\rangle \\
 &= \langle (\pi \circ e^{t\bar{H}})^* \mu|_\lambda, (E_1, \dots, E_n) \rangle.
 \end{aligned}$$

We stress that in the last formula E_i , which is an element of T_x^*M is treated as a tangent vector to the fiber, i.e., an element of $T_\lambda(T_x^*M)$ (see Figure 1.2). Indeed the pull-back $(\pi \circ e^{t\bar{H}})^* \mu$ defines an n -form on T^*M , that has dimension $2n$, and the quantity that we compute is the restriction of this form to the n -dimensional vertical space $\mathcal{V}_\lambda \simeq T_\lambda(T_x^*M)$.

To write a coordinate-independent formula, we compare this volume with the volume μ_λ^* defined naturally on the fiber \mathcal{V}_λ by the restriction of μ at x . Recall that given a smooth volume form μ on M its value μ_x at a point is a nonzero element of the one-dimensional vector space $\Lambda^n(T_xM)$. We can associate with it the unique element μ_x^* in $\Lambda^n(T_xM)^* = \Lambda^n(T_x^*M)$ satisfying $\mu_x^*(\mu_x) = 1$. This defines a volume form on the fiber T_x^*M . By the canonical identification $T_x^*M \simeq T_\lambda(T_x^*M)$ of a vector space with its tangent space to a point, this induces a volume form μ_λ^* on \mathcal{V}_λ .

With this interpretation, the Riemannian asymptotics (1.4) computes the asymptotics in t of $(\pi \circ e^{t\bar{H}})^* \mu$ restricted to the fiber \mathcal{V}_λ , with respect

to the volume μ_λ^* , i.e.,

$$(1.8) \quad (\pi \circ e^{t\vec{H}})^* \mu \Big|_{\mathcal{V}_\lambda} = t^n e^{\int_0^t \rho(\dot{\gamma}(s)) ds} \left(1 - \frac{1}{6} \text{Ric}^g(v, v)t^2 + O(t^3) \right) \mu_\lambda^*.$$

The constant c_0 appearing in (1.4) is reabsorbed in the volume μ_λ^* .

Remark 1.1. — As it follows from (1.8), the quantity ρ can be equivalently characterized as follows

$$(1.9) \quad \rho(v)\mu_\lambda^* = \frac{d}{dt} \Big|_{t=0} \log \left(t^{-n} (\pi \circ e^{t\vec{H}})^* \mu \Big|_{\mathcal{V}_\lambda} \right).$$

where $\lambda = i(v)$. The last formula inspires indeed the definition of ρ in the general case.

Identity (1.8) can be generalized to every Hamiltonian that is quadratic and convex on fibers, giving a suitable meaning to the terms in the right hand side. More precisely, we consider Hamiltonians $H : T^*M \rightarrow \mathbb{R}$ of the following form

$$(1.10) \quad H(p, x) = \frac{1}{2} \sum_{i=1}^k \langle p, X_i(x) \rangle^2 + \langle p, X_0(x) \rangle + \frac{1}{2} Q(x), \quad (p, x) \in T^*M.$$

where X_0, X_1, \dots, X_k are smooth vector fields on M and Q is a smooth function. We assume that

- (H0) X_1, \dots, X_k are everywhere linearly independent,
- (H1) $\text{Lie}\{(\text{ad } X_0)^j X_i \mid i = 1, \dots, k, j \geq 0\} \Big|_x = T_x M$ for every $x \in M$.

where $(\text{ad } Y)X = [Y, X]$ and $\text{Lie } \mathcal{F}$ denotes the smallest Lie algebra containing a family of vector fields \mathcal{F} . The Hamiltonian (1.10) is naturally associated with the optimal control problem where the dynamics is affine in the control

$$(1.11) \quad \dot{x}(t) = X_0(x(t)) + \sum_{i=1}^k u_i(t) X_i(x(t)), \quad x \in M,$$

and one wants to minimize a quadratic cost with potential (here x_u denotes the solution of (1.11) associated with $u \in L^\infty([0, T], \mathbb{R}^k)$)

$$(1.12) \quad J_T(u) := \frac{1}{2} \int_0^T \|u(s)\|^2 - Q(x_u(s)) ds.$$

We stress that when $X_0 = 0, Q = 0$ and $k = n$, the optimal control problem described above is the geodesic problem associated with the Riemannian metric defined by the orthonormal frame X_1, \dots, X_n on M and H is the corresponding Hamiltonian. The case $X_0 = 0, Q = 0$ and $k < n$ includes the geodesic problem in sub-Riemannian geometry.

Consider the projections on M of integral curves of the Hamiltonian vector field \vec{H} in T^*M . Under our assumptions, short pieces of these curves are minimizers for the optimal control problem (i.e., geodesics in the case of Riemannian or sub-Riemannian geometry). However, in general, not all minimizers can be obtained in this way. This is due to the possible presence of the so-called abnormal minimizers [16]. The projected trajectories, as solutions of an Hamiltonian system in T^*M , are smooth and parametrized by the initial covector in the cotangent bundle.

If the initial covector λ corresponds to an *ample* and *equiregular* trajectory (cf. Section 3 for precise definitions) then the exponential map $\pi \circ e^{t\vec{H}}$ is a local diffeomorphism at λ and it is possible to compute the variation of a smooth volume μ under the exponential map, as in the Riemannian case.

Let us stress that in the Riemannian case all $\lambda \in T^*M$ satisfy these assumptions, while in the sub-Riemannian case one can prove that there exists a non-empty open and dense set of covectors $\mathcal{A} \subset T^*M$ such that the corresponding geodesic is ample and equiregular (see Proposition 7.1). If the initial point x is fixed, then there exists a non-empty Zariski open set of covectors in T_x^*M such that the corresponding geodesic is ample, but the existence of equiregular geodesics is not guaranteed. On the other hand, on any ample geodesic, there exists an open dense set of t such that the germ of the geodesic at $\gamma(t)$ is equiregular (cf. Section 7).

The main result of this paper is the generalization of the asymptotics (1.8) to any flow arising from an Hamiltonian that is quadratic and convex on fibers, along any trajectory satisfying our regularity assumptions. In particular we give a geometric characterization of the terms appearing in the asymptotic expansion of a smooth volume μ under the Hamiltonian flow $\pi \circ e^{t\vec{H}}$ and we interpret every coefficient as the generalization of the corresponding Riemannian element.

Fix $x \in M$ and $\lambda \in T_x^*M$. Let $\gamma(t) = \pi(e^{t\vec{H}}(\lambda))$ be the associated geodesic on M and assume that it is ample and equiregular. With such a geodesic it is possible to associate an integer $\mathcal{N}(\lambda)$ which is defined through the structure of the Lie brackets of the controlled vector fields X_1, \dots, X_k along γ (cf. Definition 3.7). This is an invariant that depends only on the Lie algebraic structure of the drift field X_0 and the distribution $\mathcal{D} = \text{span}\{X_1, \dots, X_k\}$ along the trajectory and not on the particular frame (that induces the metric) on \mathcal{D} . The notation stresses that this integer can a priori depend on the trajectory.

The results obtained in [4, Section 6.5] imply that there exists $C_\lambda > 0$ such that, for $t \rightarrow 0$

$$(1.13) \quad (\pi \circ e^{t\vec{H}})^* \mu \Big|_{\mathcal{V}_\lambda} = t^{\mathcal{N}(\lambda)} (C_\lambda + O(t)) \mu_\lambda^*.$$

Once the order of the asymptotics is determined, one can introduce the *volume geodesic derivative* as follows

$$(1.14) \quad \rho(\lambda) \mu_\lambda^* := \frac{d}{dt} \Big|_{t=0} \log \left(t^{-\mathcal{N}(\lambda)} (\pi \circ e^{t\vec{H}})^* \mu \Big|_{\mathcal{V}_\lambda} \right),$$

This is the natural generalization of the quantity introduced in (1.9). In Section 6 we show an explicit formula to compute ρ , in terms of the *symbol of the structure along the geodesic* (cf. Definition 3.8) and we compute it explicitly for contact manifolds endowed with Popp’s volume.

The Riemannian Ricci tensor appearing in (1.8) is replaced by the trace of a curvature operator in the direction of λ . This curvature operator $\mathcal{R}_\lambda : \mathcal{D}_x \rightarrow \mathcal{D}_x$, is a generalization of the sectional curvature and is defined in [4] for the wide class of geometric structures arising from affine control systems. In the Riemannian case $\mathcal{R}_\lambda(w) = R^g(w, v)v$, where R^g is the Riemann tensor associated with g , $\lambda = i(v)$ and $w \in \mathcal{D}_x = T_x M$. Notice that in this case $\text{tr } \mathcal{R}_\lambda = \text{Ricci}^g(v, v)$.

All the geometric invariants are rational functions in the initial covector λ . The precise statement of our theorem reads as follows.

THEOREM 1.2. — *Let μ be a smooth volume form on M and $\gamma(t) = \pi(\lambda(t)) = \pi(e^{t\vec{H}}(\lambda))$ be an ample and equiregular trajectory, with $\lambda \in T_x^* M$. Then we have the following asymptotic expansion*

$$(1.15) \quad (\pi \circ e^{t\vec{H}})^* \mu \Big|_{\mathcal{V}_\lambda} = C_\lambda t^{\mathcal{N}(\lambda)} e^{\int_0^t \rho(\lambda(s)) ds} \left(1 - t^2 \frac{\text{tr } \mathcal{R}_\lambda}{6} + o(t^2) \right) \mu_\lambda^*.$$

where μ_λ^* is the canonical volume induced by μ on $\mathcal{V}_\lambda \simeq T_\lambda(T_x^* M)$.

Again, in the asymptotics (1.15), the choice of the volume form μ affects only the volume geodesic derivative ρ . Notice also that the constant C_λ and the main order $t^{\mathcal{N}(\lambda)}$ depend only on the Young diagram associated with the curve γ , while the term \mathcal{R}_λ (and actually the whole asymptotic expansion contained in the parentheses) depends only on the curvature like-invariants of the cost of the optimal control problem (1.11)–(1.12), i.e., on the Hamiltonian (1.10).

We stress that in the Riemannian case every trajectory is ample and equiregular and the constant C_λ is always equal to one and $\mathcal{N}(\lambda) = n = \dim M$ (cf. formula (1.8)).

In other words, the asymptotics (1.15) “isolates” the contribution given by the volume form with respect to the contribution given by the dynamics/geometry.

Remark 1.3. — Another geometric interpretation of the variation of the volume is given in Figure 1.3. Let $\Omega \subset T_x^*M$ be an infinitesimal neighborhood of λ and let $\Omega_{x,t} := \pi \circ e^{t\vec{H}}(\Omega)$ be its image on M with respect to the Hamiltonian flow. For every t the set $\Omega_{x,t} \subset M$ is a neighborhood of $\gamma(t)$. By construction

$$\mu(\Omega_{x,t}) = \int_{\Omega} (\pi \circ e^{t\vec{H}})^* \mu,$$

and (1.15) represents exactly the variation of the volume element along γ .




Figure 1.3. Infinitesimal variation of the volume along a geodesic

1.1. The invariant ρ in the Riemannian case

In the Riemannian case, for $\mu = e^\psi \text{vol}_g$, one has $\rho(v) = g(\nabla\psi, v)$ for every tangent vector v . If one writes explicitly the expansion of the exponential term in (1.8) at order 2 one finds, for $\gamma(t) = \exp_x(tv)$

$$(1.16) \quad (\pi \circ e^{t\vec{H}})^* \mu \Big|_{\mathcal{V}_\lambda} = t^n \left(1 + \rho(v)t - \left(\frac{1}{6} \text{Ric}^g(v, v) - \frac{1}{2} \rho(v)^2 - \frac{1}{2} \nabla \rho(v, v) \right) t^2 + O(t^3) \right) \mu_\lambda^*.$$

In particular for X, Y vector fields on M

$$(1.17) \quad \rho^2 = \nabla\psi \otimes \nabla\psi, \quad \rho(X)^2 = g(\nabla\psi, X)^2,$$

$$(1.18) \quad \nabla\rho = \text{Hess } \psi, \quad \nabla\rho(X, Y) = Y(X\psi) - (\nabla_Y X)\psi,$$

and summing over a local orthonormal basis X_1, \dots, X_n for g one can compute the traces

$$(1.19) \quad \text{tr } \rho^2 = \|\nabla\psi\|^2, \quad \text{tr } \nabla\rho = \Delta_g\psi.$$

1.2. On the relation with the small time heat kernel asymptotics

The volume geodesic derivative ρ introduced in this paper, together with the curvature-like invariants of the dynamics, characterize in the Riemannian case the small time heat kernel expansion on the diagonal associated with a weighted Laplacian $\Delta_\mu = \text{div}_\mu \nabla$.

Indeed let us consider a weighted Riemannian manifold (M, g, μ) with $\mu = e^\psi \text{vol}_g$, and denote by $p_\mu(t, x, y)$ the fundamental solution of the heat equation $\partial_t - \Delta_\mu = 0$ associated with Δ_μ . Recall that $\Delta_\mu = \Delta_g + g(\nabla\psi, \nabla \cdot)$, where Δ_g is the Laplace–Beltrami operator of (M, g) . One has the following small time asymptotics (see for instance [12])

$$p_\mu(t, x, x) = \frac{c}{t^{n/2}} \left[1 + t \left(\frac{S(x)}{12} - \frac{\|\nabla\psi(x)\|^2}{8} - \frac{\Delta_g(\psi)}{4} \right) + o(t) \right]$$

for some $c > 0$.

Hence the terms appearing in the heat kernel expansion are exactly the trace of the invariants that determine the expansion of the exponential in (1.8) at order 2.

A natural conjecture is that the same three coefficients describe the heat kernel small time asymptotics expansion also in the sub-Riemannian case. This conjecture is true in the 3D contact case as it follows by the results obtained in [6], since on a 3D contact manifold if one chooses μ equal to Popp volume one has $\rho = 0$ (cf. Remark 7.8). See also [7, 8, 18] for some results about small time heat kernel expansion for Hörmander operators with drift.

1.3. Structure of the paper

In Section 2 we describe the general setting, while in Section 3 and 4 we introduce some preliminaries needed for the proof of the main result.

Section 5 is devoted to the definition of the volume geodesic derivative ρ and the analysis of its properties, while in Section 6 we give a formula that permits us to compute it through an adapted frame. Section 7 specifies the whole construction to sub-Riemannian case. Finally, Section 8 contains the proof of the main result.

2. General setting

Let M be an n -dimensional connected manifold and let $X_0, X_1, \dots, X_k \in \text{Vec}(M)$ be smooth vector fields, with $k \leq n$. We consider the following *affine control system* on M

$$(2.1) \quad \dot{x}(t) = X_0(x(t)) + \sum_{i=1}^k u_i(t)X_i(x(t)), \quad x \in M,$$

where $u \in L^\infty([0, T], \mathbb{R}^k)$ is a measurable and essentially bounded function called *control*. In what follows we assume that

- (H0) X_1, \dots, X_k are everywhere linearly independent,
- (H1) $\text{Lie}\{(\text{ad } X_0)^j X_i \mid i = 1, \dots, k, j \geq 0\}|_x = T_x M$ for every $x \in M$.

where $(\text{ad } Y)X = [Y, X]$ and $\text{Lie } \mathcal{F}$ denotes the smallest Lie algebra containing a family of vector fields \mathcal{F} .

A Lipschitz curve $\gamma : [0, T] \rightarrow M$ is said to be *admissible* for the system (2.1) if there exists a control $u \in L^\infty([0, T], \mathbb{R}^k)$ such that γ satisfies (2.1) for a.e. $t \in [0, T]$. The pair (γ, u) of an admissible curve γ and its control u is called *admissible pair*.

Remark 2.1. — The affine control system can be defined more generally as a pair (\mathbb{U}, f) , where \mathbb{U} is a smooth vector bundle of rank k with base M and fiber \mathbb{U}_x , and $f : \mathbb{U} \rightarrow TM$ is a smooth affine morphism of vector bundles such that $\pi \circ f(u) = x$, for every $u \in \mathbb{U}_x$. Locally, by taking a local trivialization of \mathbb{U} , we can write $f(u) = X_0 + \sum_{i=1}^k u_i X_i$ for $u \in \mathbb{U}$. For more details about this approach see [2, 4].

We denote by $\mathcal{D} \subset TM$ the *distribution*, that is the family of subspaces spanned by the linear part of the control problem at a point, i.e.,

$$\mathcal{D} = \{\mathcal{D}_x\}_{x \in M}, \quad \text{where } \mathcal{D}_x := \text{span}\{X_1, \dots, X_k\}|_x \subset T_x M.$$

The distribution \mathcal{D} has constant rank by assumption (H0), and we endow \mathcal{D} with the inner product such that the fields X_1, \dots, X_k are orthonormal. We denote by $\Gamma(\mathcal{D})$ the set of smooth sections of \mathcal{D} , also called *horizontal*

vector fields. Among all admissible trajectories that join two fixed points in time $T > 0$, we want to minimize the *quadratic cost functional*

$$J_T(u) := \frac{1}{2} \int_0^T \|u(s)\|^2 - Q(x_u(s)) \, ds,$$

where Q is a smooth function on M , playing the role of a potential. Here x_u denotes the solution of (2.1) associated with u .

DEFINITION 2.2. — For $x_0, x_1 \in M$ and $T > 0$, we define the value function

$$(2.2) \quad S_T(x_0, x_1) := \inf \{J_T(u) \mid (\gamma, u) \text{ admissible pair, } \gamma(0) = x_0, \gamma(T) = x_1\}.$$

The assumption (H1) implies, by Krener’s theorem (see [14, Theorem 3.10] or [15, Chapter 3]), that the attainable set in time $T > 0$ from a fixed point $x_0 \in M$, that is the set

$$A_{x_0, T} = \{x_1 \in M : S_T(x_0, x_1) < +\infty\}$$

has non-empty interior for all $T > 0$. This is a necessary assumption to the existence of ample geodesics.

Important examples of affine control problems are sub-Riemannian structures. These are triples (M, \mathcal{D}, g) , where M is a smooth manifold, \mathcal{D} is a smooth, completely non-integrable vector sub-bundle of TM and g is a smooth inner product on \mathcal{D} . In our context, these are included in the case $X_0 = 0$ and $Q = 0$. The value function in this case coincides with (one half of the square of) the sub-Riemannian distance, i.e., the infimum of the length of absolutely continuous admissible curves joining two points. In this case the assumption (H1) on \mathcal{D} implies, by the Rashevskii–Chow theorem, that the sub-Riemannian distance is finite on M . Moreover the metric topology coincides with the one of M . A more detailed introduction on sub-Riemannian geometry can be found in [2, 17].

For an affine optimal control system, the associated Hamiltonian is defined as follows

$$(2.3) \quad H(p, x) = \frac{1}{2} \sum_{i=1}^k \langle p, X_i(x) \rangle^2 + \langle p, X_0(x) \rangle + \frac{1}{2} Q(x), \quad (p, x) \in T^*M.$$

Hamilton’s equations are written as follows

$$(2.4) \quad \dot{x} = \frac{\partial H}{\partial p}, \quad \dot{p} = -\frac{\partial H}{\partial x} \quad (p, x) \in T^*M,$$

THEOREM 2.3 (PMP [5, 19]). — *Consider a solution $\lambda(t) = (p(t), \gamma(t))$ defined on $[0, T]$ of the Hamilton equations (2.4) on T^*M . Then short pieces of the trajectory $\gamma(t) = \pi(\lambda(t))$ minimize the cost between their endpoints.*

From now on, using a slight abuse of notation, we call *geodesic* any projection $\gamma : [0, T] \rightarrow M$ of an integral line of the Hamiltonian vector field. In the general case, some minimizers of the cost might not satisfy this equation. These are the so-called strictly abnormal minimizers [16], and they are related with hard open problems in control theory and in sub-Riemannian geometry. In what follows we will focus on those minimizers that come from the Hamilton equations (also called normal) and that satisfy a suitable regularity assumption. Notice that normal geodesics are smooth.

3. Geodesic flag and symbol

In this section we define the flag and the symbol of a geodesic, that are elements carrying information about the germ of the distribution and the drift along the trajectory. The symbol is the graded vector space associated with the flag and is endowed with an inner product induced by the metric on the distribution.

3.1. The class of symbols

We start by defining the class of objects we deal with.

DEFINITION 3.1. — *A symbol S is a pair (V, L) where*

- (1) *V is a graded vector space $V = \oplus_{i=1}^m V_i$, endowed with an inner product $\langle \cdot, \cdot \rangle$ on its first layer V_1 ,*
- (2) *$L = \{L_i\}_{i=1}^m$ is a family of surjective linear maps $L_i : V_1 \rightarrow V_i$.*

Remark 3.2. — Through the surjective linear maps $L_i : V_1 \rightarrow V_i$, the inner product on V_1 naturally induces a norm of $v \in V_i$ as follows

$$\|v\|_{V_i} := \min \{ \|u\|_{V_1} \text{ s.t. } L_i(u) = v \}.$$

It is easy to check that, since $\| \cdot \|_{V_1}$ is induced by an inner product, then $\| \cdot \|_{V_i}$ is induced by an inner product as well. Hence the family of surjective maps endows V with a global inner product by defining the subspaces $\{V_i\}_{i=1}^m$ to be mutually orthogonal.

DEFINITION 3.3. — We say that the symbols $S = (V, L)$ and $S' = (V', L')$ are isomorphic if there exists an invertible linear map $\phi : V \rightarrow V'$ such that $\phi|_{V_1} : V_1 \rightarrow V'_1$ is an isometry and $L'_i \circ \phi = \phi \circ L_i$ for $i \geq 1$.

LEMMA 3.4. — If two symbols S and S' are isomorphic, then they are isometric as inner product spaces.

Proof. — Let $V = \bigoplus_{i=1}^m V_i$ and $V' = \bigoplus_{i=1}^{m'} V'_i$ and let ϕ be the map given in Definition 3.3. Let $v \in V_1$ and $v' = \phi(v) \in V'_1$. By the commutation property satisfied by ϕ one has

$$L'_i(v') = L'_i(\phi(v)) = \phi(L_i(v)),$$

therefore $V'_i = \phi(V_i)$ for every $i \geq 1$ and in particular $m = m'$. As a consequence the map ϕ descends to a family of maps between every layer of the stratification as follows: for $v_i \in V_i$ write $v_i = L_i(v)$ for some $v \in V_1$ and define $\phi_i : V_i \rightarrow V'_i$ by $\phi_i(v_i) := L'_i(\phi(v))$. Since ϕ is an isometry on V_1 , then the map $\phi|_{V_i} : V_i \rightarrow V'_i$ is an isometry on each layer. □

3.2. The symbol of a geodesic

Let $\gamma : [0, T] \rightarrow M$ be a geodesic and consider a smooth admissible extension of its tangent vector, namely a vector field $\mathbb{T} = X_0 + X$, with $X \in \Gamma(\mathcal{D})$, such that $\mathbb{T}(\gamma(t)) = \dot{\gamma}(t)$ for every $t \in [0, T]$.

DEFINITION 3.5. — The flag of the geodesic $\gamma : [0, T] \rightarrow M$ is the one-parameter family of subspaces

$$(3.1) \quad \mathcal{F}_{\gamma(t)}^i := \text{span}\{\mathcal{L}_{\mathbb{T}}^j(X)|_{\gamma(t)} \mid X \in \Gamma(\mathcal{D}), j \leq i-1\} \subseteq T_{\gamma(t)}M, \quad \forall i \geq 1,$$

for any fixed $t \in [0, T]$, where $\mathcal{L}_{\mathbb{T}}$ denotes the Lie derivative in the direction of \mathbb{T} .

Definition 3.5 is well posed, namely it does not depend on the choice of the admissible extension \mathbb{T} (see [4, Section 3.4]). By construction, the flag is a filtration of $T_{\gamma(t)}M$, i.e., $\mathcal{F}_{\gamma(t)}^i \subseteq \mathcal{F}_{\gamma(t)}^{i+1}$, for all $i \geq 1$. Moreover, $\mathcal{F}_{\gamma(t)}^1 = \mathcal{D}_{\gamma(t)}$. The growth vector of the geodesic $\gamma(t)$ is the one-parameter family of sequences of integers

$$(3.2) \quad \mathcal{G}_{\gamma(t)} := \{\dim \mathcal{F}_{\gamma(t)}^1, \dim \mathcal{F}_{\gamma(t)}^2, \dots\}.$$

A geodesic $\gamma(t)$, with growth vector $\mathcal{G}_{\gamma(t)}$, is said to be

- (i) *equiregular* if, for every $i \geq 1$, the integer $\dim \mathcal{F}_{\gamma(t)}^i$ does not depend on t ,

- (ii) *ample* if for every t there exists $m = m(t) \geq 1$ such that $\dim \mathcal{F}_{\gamma(t)}^m = \dim T_{\gamma(t)}M$.

We define the integers $d_i(t) := \dim \mathcal{F}_{\gamma(t)}^i - \dim \mathcal{F}_{\gamma(t)}^{i-1}$, for $i \geq 0$, with the convention $\dim \mathcal{F}_{\gamma(t)}^0 = 0$.

Ample (resp. equiregular) geodesics are the microlocal counterpart of bracket-generating (resp. equiregular) distributions. For an equiregular and ample geodesic the integers d_i are independent on t and represent the increment of dimension of the flag of the geodesic at each step. The following result is proved in [4, Lemma 3.5].

LEMMA 3.6. — *For any ample, equiregular geodesic, it holds $d_1 \geq d_2 \geq \dots \geq d_m$.*

DEFINITION 3.7. — *Given an ample and equiregular geodesic with initial covector $\lambda \in T_x^*M$ we define*

$$(3.3) \quad \mathcal{N}(\lambda) := \sum_{i=1}^m (2i - 1)d_i.$$

Fix an ample and equiregular geodesic $\gamma : [0, T] \rightarrow M$ and let \mathbb{T} be an admissible extension of its tangent vector. For every vector $v \in \mathcal{F}_{\gamma(t)}^i$, consider a smooth vector field X such that $X|_{\gamma(t)} = v$ and $X|_{\gamma(s)} \in \mathcal{F}_{\gamma(s)}^i$ for every $s \in [0, T]$. The Lie derivative $\mathcal{L}_{\mathbb{T}}$ in the direction of \mathbb{T} induces a well defined linear map

$$(3.4) \quad \mathcal{L}_{\mathbb{T}} : \mathcal{F}_{\gamma(s)}^i \rightarrow \mathcal{F}_{\gamma(t)}^{i+1}/\mathcal{F}_{\gamma(t)}^i, \quad v \mapsto [\mathbb{T}, X]|_{\gamma(t)} \pmod{\mathcal{F}_{\gamma(t)}^i}.$$

Indeed a direct computation shows that this map does not depend on the admissible extension \mathbb{T} and on the extension of X , under the equiregularity assumption on γ . Moreover the kernel of (3.4) is given by $\mathcal{F}_{\gamma(t)}^{i-1}$. So one obtains well-defined linear surjective maps (denoted by the same symbol)

$$(3.5) \quad \mathcal{L}_{\mathbb{T}} : \mathcal{F}_{\gamma(t)}^i/\mathcal{F}_{\gamma(t)}^{i-1} \rightarrow \mathcal{F}_{\gamma(t)}^{i+1}/\mathcal{F}_{\gamma(t)}^i, \quad i \geq 1.$$

In particular $\mathcal{L}_{\mathbb{T}}^i : \mathcal{D}_{\gamma(t)} \rightarrow \mathcal{F}_{\gamma(t)}^{i+1}/\mathcal{F}_{\gamma(t)}^i$ are surjective linear maps defined on the distribution $\mathcal{D}_{\gamma(t)} = \mathcal{F}_{\gamma(t)}^1$.

DEFINITION 3.8. — *Given an ample and equiregular geodesic $\gamma : [0, T] \rightarrow M$ we define its symbol at $\gamma(t)$, denoted by $S_{\gamma(t)}$, as the pair*

- (1) *the graded vector space: $\text{gr}_{\gamma(t)}(\mathcal{F}) := \bigoplus_{i=0}^{m-1} \mathcal{F}_{\gamma(t)}^{i+1}/\mathcal{F}_{\gamma(t)}^i$,*
- (2) *the family of operators: $\mathcal{L}_{\mathbb{T}}^i : \mathcal{D}_{\gamma(t)} \rightarrow \mathcal{F}_{\gamma(t)}^{i+1}/\mathcal{F}_{\gamma(t)}^i$ for $i \geq 1$,*


where \mathbb{T} denotes any admissible extension of $\dot{\gamma}$.

Remark 3.9. — Notice that, for the symbol (V, L) associated with an ample and equiregular geodesic, the family of maps $L = \{L_i\}_{i=1}^m$ where $L_i = \mathcal{L}_T^i$ satisfies the factorization property $\ker L_i \subset \ker L_{i+1}$ for all $i \geq 1$.

4. Young diagram, canonical frame and Jacobi fields

In this section we briefly recall how to define the canonical frame that can be associated with any ample and equiregular geodesic, introduced in [22]. We follow the approach contained in [4, 10], where the interested reader can find more details.

For an ample, equiregular geodesic we can build a tableau D with m columns of length d_i , for $i = 1, \dots, m$, as follows:



The total number of boxes in D is $n = \dim M = \sum_{i=1}^m d_i$.

Consider an ample, equiregular geodesic, with Young diagram D , with k rows, and denote the length of the rows by n_1, \dots, n_k . Indeed $n_1 + \dots + n_k = n$. We are going to introduce a moving frame on $T_{\lambda(t)}(T^*M)$ indexed by the boxes of the Young diagram. The notation $ai \in D$ denotes the generic box of the diagram, where $a = 1, \dots, k$ is the row index, and $i = 1, \dots, n_a$ is the progressive box number, starting from the left, in the specified row. We employ letters a, b, c, \dots for rows, and i, j, h, \dots for the position of the box in the row.




Figure 4.1. Levels (shaded regions) and superboxes (delimited by bold lines) for different Young diagrams: (a) Riemannian, (b) contact, (c) a more general structure.

We collect the rows with the same length in D , and we call them levels of the Young diagram. In particular, a level is the union of r rows D_1, \dots, D_r ,

and r is called the size of the level. The set of all the boxes $ai \in D$ that belong to the same column and the same level of D is called *superbox*. We use Greek letters α, β, \dots to denote superboxes. Notice that two boxes ai, bj are in the same superbox if and only if ai and bj are in the same column of D and in possibly distinct rows but with same length, i.e. if and only if $i = j$ and $n_a = n_b$ (see Figure 4.1).

In what follows, for $V(t)$ a vector field along an integral line $\lambda(t)$ of the Hamiltonian flow, we denote by

$$(4.1) \quad \dot{V}(t) := \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} e_*^{-\varepsilon \vec{H}} V(t + \varepsilon).$$

the Lie derivative of V in the direction of \vec{H} . The following theorem is proved in [22].

THEOREM 4.1. — *Assume $\lambda(t)$ is the lift of an ample and equiregular geodesic $\gamma(t)$ with Young diagram D . Then there exists a smooth moving frame $\{E_{ai}, F_{ai}\}_{ai \in D}$ along $\lambda(t)$ such that*

(1) $\pi_* E_{ai}|_{\lambda(t)} = 0.$

(2) *It is a Darboux basis, namely*

$$\sigma(E_{ai}, E_{bj}) = \sigma(F_{ai}, F_{bj}) = \sigma(E_{ai}, F_{bj}) - \delta_{ab} \delta_{ij} = 0, \quad ai, bj \in D.$$

(3) *The frame satisfies the structural equations*

$$(4.2) \quad \begin{cases} \dot{E}_{ai} = E_{a(i-1)} & a = 1, \dots, k, \quad i = 2, \dots, n_a, \\ \dot{E}_{a1} = -F_{a1} & a = 1, \dots, k, \\ \dot{F}_{ai} = \sum_{bj \in D} R_{ai,bj}(t) E_{bj} - F_{a(i+1)} & a = 1, \dots, k, \quad i = 1, \dots, n_a - 1, \\ \dot{F}_{ana} = \sum_{bj \in D} R_{ana,bj}(t) E_{bj} & a = 1, \dots, k, \end{cases}$$

for some smooth family of $n \times n$ symmetric matrices $R(t)$, with components $R_{ai,bj}(t) = R_{bj,ai}(t)$, indexed by the boxes of the Young diagram D . The matrix $R(t)$ is normal in the sense of [22].

If $\{\tilde{E}_{ai}, \tilde{F}_{ai}\}_{ai \in D}$ is another smooth moving frame along $\lambda(t)$ satisfying (1)–(3), with some normal matrix $\tilde{R}(t)$, then for any superbox α of size r there exists an orthogonal constant $r \times r$ matrix O^α such that

$$(4.3) \quad \tilde{E}_{ai} = \sum_{bj \in \alpha} O_{ai,bj}^\alpha E_{bj}, \quad \tilde{F}_{ai} = \sum_{bj \in \alpha} O_{ai,bj}^\alpha F_{bj}, \quad ai \in \alpha.$$

The explicit condition for the matrix $R(t)$ to be normal can be found in [4, Appendix F] (cf. also the original definition in [22]).

Remark 4.2. — For $a = 1, \dots, k$, we denote by E_a the n_a -dimensional row vector $E_a = (E_{a1}, \dots, E_{an_a})$, with analogous notation for F_a . Denote then by E the n -dimensional row vector $E = (E_1, \dots, E_k)$, and similarly for F . Then, we rewrite the system (4.2) as follows

$$(4.4) \quad \begin{pmatrix} \dot{E}^* \\ \dot{F}^* \end{pmatrix} = \begin{pmatrix} C_1^* & -C_2 \\ R(t) & -C_1 \end{pmatrix} \begin{pmatrix} E^* \\ F^* \end{pmatrix},$$

where $C_1 = C_1(D)$, $C_2 = C_2(D)$ are $n \times n$ matrices, depending on the Young diagram D , defined as follows: for $a, b = 1, \dots, k$, $i = 1, \dots, n_a$, $j = 1, \dots, n_b$:

$$(4.5) \quad [C_1]_{ai,bj} := \delta_{ab}\delta_{i,j-1}, \quad [C_2]_{ai,bj} := \delta_{ab}\delta_{i1}\delta_{j1}.$$

It is convenient to see C_1 and C_2 as block diagonal matrices:

$$(4.6) \quad C_i(D) := \begin{pmatrix} C_i(D_1) & & \\ & \ddots & \\ & & C_i(D_k) \end{pmatrix}, \quad i = 1, 2,$$

the a -th block being the $n_a \times n_a$ matrices

$$(4.7) \quad C_1(D_a) := \begin{pmatrix} 0 & \mathbb{I}_{n_a-1} \\ 0 & 0 \end{pmatrix}, \quad C_2(D_a) := \begin{pmatrix} 1 & 0 \\ 0 & 0_{n_a-1} \end{pmatrix},$$

where \mathbb{I}_m is the $m \times m$ identity matrix and 0_m is the $m \times m$ zero matrix.

4.1. The Jacobi equation

A vector field $\mathcal{J}(t)$ along $\lambda(t)$ is called a *Jacobi field* if it satisfies

$$(4.8) \quad \dot{\mathcal{J}} = 0.$$

The space of solutions of (4.8) is a $2n$ -dimensional vector space. The projections $J = \pi_*\mathcal{J}$ are vector fields on M corresponding to one-parameter variations of $\gamma(t) = \pi(\lambda(t))$ through geodesics; in the Riemannian case (without drift field) they coincide with the classical Jacobi fields.

We intend to write (4.8) using the natural symplectic structure σ of T^*M and the canonical frame. First, observe that on T^*M there is a natural smooth sub-bundle of Lagrangian⁽¹⁾ spaces:

$$(4.9) \quad \mathcal{V}_\lambda := \ker \pi_*|_\lambda = T_\lambda(T_{\pi(\lambda)}^*M).$$

⁽¹⁾A Lagrangian subspace $L \subset \Sigma$ of a symplectic vector space (Σ, σ) is a subspace with $2 \dim L = \dim \Sigma$ and $\sigma|_L = 0$.




Figure 4.2. The geodesic cost function

We call this the *vertical subspace*. Then, let $\{E_i(t), F_i(t)\}_{i=1}^n$ be a canonical frame along $\lambda(t)$. The fields E_1, \dots, E_n belong to the vertical subspace. In terms of this frame, $\mathcal{J}(t)$ has components $(p(t), x(t)) \in \mathbb{R}^{2n}$:

$$(4.10) \quad \mathcal{J}(t) = \sum_{i=1}^n p_i(t)E_i(t) + x_i(t)F_i(t).$$

In turn, the Jacobi equation, written in terms of the components $(p(t), x(t))$, becomes

$$(4.11) \quad \begin{pmatrix} \dot{p} \\ \dot{x} \end{pmatrix} = \begin{pmatrix} -C_1 & -R(t) \\ C_2 & C_1^* \end{pmatrix} \begin{pmatrix} p \\ x \end{pmatrix}.$$

This is a generalization of the classical Jacobi equation seen as first-order equation for fields on the cotangent bundle. Its structure depends on the Young diagram of the geodesic through the matrices $C_i(D)$, while the remaining invariants are contained in the curvature matrix $R(t)$. Notice that this includes the Riemannian case, where D is the same for every geodesic, with $C_1 = 0$ and $C_2 = \mathbb{I}$.

4.2. Geodesic cost and curvature operator

In this section we define the geodesic cost and the curvature operator associated with a geodesic γ . This operator generalizes the Riemannian sectional curvature operator.

DEFINITION 4.3. — *Let $x_0 \in M$ and consider an ample geodesic γ such that $\gamma(0) = x_0$. The geodesic cost associated with γ is the family of functions*

$$c_t(x) := -S_t(x, \gamma(t)), \quad x \in M, t > 0,$$

where S_t is the value function defined in (2.2).

By [4, Theorem 4.2], given an ample curve $\gamma(t) = \pi(e^{t\tilde{H}}(\lambda))$ starting at x_0 , the geodesic cost function $c_t(x)$ is smooth in a neighborhood of x_0 and for $t > 0$ sufficiently small. Moreover the differential $d_{x_0}c_t = \lambda$ for every t small.

Let $\dot{c}_t = \frac{\partial}{\partial t}c_t$ denote the derivative with respect to t of the geodesic cost. Then \dot{c}_t has a critical point at x_0 and its second differential $d_{x_0}^2\dot{c}_t : T_{x_0}M \rightarrow \mathbb{R}$ is defined as

$$d_{x_0}^2\dot{c}_t(v) = \left. \frac{d^2}{ds^2} \right|_{s=0} \dot{c}_t(\gamma(s)), \quad \gamma(0) = x_0, \quad \dot{\gamma}(0) = v.$$

We restrict the second differential of \dot{c}_t to the distribution \mathcal{D}_{x_0} and we define the following family of symmetric operators $\mathcal{Q}_\lambda(t) : \mathcal{D}_{x_0} \rightarrow \mathcal{D}_{x_0}$, for small t , associated with $d_{x_0}^2\dot{c}_t$ through the inner product defined on \mathcal{D}_{x_0} :

$$(4.12) \quad d_{x_0}^2\dot{c}_t(v) := \langle \mathcal{Q}_\lambda(t)v | v \rangle_{x_0}, \quad t > 0, v \in \mathcal{D}_{x_0}.$$

The following result is contained in [4, Theorem A].

THEOREM 4.4. — *Let $\gamma : [0, T] \rightarrow M$ be an ample geodesic with initial covector $\lambda \in T_{x_0}^*M$ and let $\mathcal{Q}_\lambda(t) : \mathcal{D}_{x_0} \rightarrow \mathcal{D}_{x_0}$ be defined by (4.12). Then $t \mapsto t^2\mathcal{Q}_\lambda(t)$ can be extended to a smooth family of symmetric operators on \mathcal{D}_{x_0} for small $t \geq 0$. Moreover*

$$\mathcal{I}_\lambda := \lim_{t \searrow 0} t^2\mathcal{Q}_\lambda(t) \geq \mathbb{I} > 0, \quad \left. \frac{d}{dt} \right|_{t=0} t^2\mathcal{Q}_\lambda(t) = 0,$$

where \mathbb{I} is the identity operator. In particular, there exists a symmetric operator $\mathcal{R}_\lambda : \mathcal{D}_{x_0} \rightarrow \mathcal{D}_{x_0}$ such that

$$(4.13) \quad \mathcal{Q}_\lambda(t) = \frac{1}{t^2}\mathcal{I}_\lambda + \frac{1}{3}\mathcal{R}_\lambda + O(t), \quad t > 0.$$

DEFINITION 4.5. — *We call the symmetric operator $\mathcal{R}_\lambda : \mathcal{D}_{x_0} \rightarrow \mathcal{D}_{x_0}$ in (4.13) the curvature at λ . Its trace $\text{tr}\mathcal{R}_\lambda$ is the Ricci curvature at λ .*

When γ is also equiregular, the curvature operator \mathcal{R}_λ can be written in terms of the smooth n -dimensional symmetric matrix $R(t)$ introduced in the canonical equations (4.2), as we now describe.

Let $\gamma(t) = \pi(e^{t\tilde{H}}(\lambda))$ be ample and equiregular, and let $\{E_{ai}(t), F_{ai}(t)\}_{ai \in D}$ be a canonical frame along the curve $\lambda(t)$.

LEMMA 4.6 ([4, Lemma 8.3]). — *The set of vector fields along $\gamma(t)$*

$$X_{ai}(t) := \pi_*F_{ai}(t), \quad ai \in D$$

is a basis for $T_{\gamma(t)}M$ adapted to the flag $\{\mathcal{F}_{\gamma(t)}^i\}_{i=1}^m$ and $\{X_{a1}(t)\}_{a=1}^k$ is an orthonormal basis for $\mathcal{D}_{\gamma(t)}$ along the geodesic.

The following proposition is proved in [4, Section 7.4].

PROPOSITION 4.7. — *Let γ be an ample and equiregular geodesic with initial covector λ . The matrix representing \mathcal{R}_λ in terms of the orthonormal basis $\{X_{a1}(t)\}_{a=1}^k$ depends only on the elements of $R_{a1,b1}(0)$ corresponding to the first column of the associated Young diagram. More precisely we have*

$$(4.14) \quad (\mathcal{R}_\lambda)_{ab} = 3\Omega(n_a, n_b)R_{a1,b1}(0), \quad a, b = 1, \dots, k,$$

where for $i, j \in \mathbb{N}$ we set

$$\Omega(i, j) = \begin{cases} 0, & |i - j| \geq 2, \\ \frac{1}{4(i + j)}, & |i - j| = 1, \\ \frac{i}{4i^2 - 1}, & i = j. \end{cases}$$

5. Volume geodesic derivative

In this section we introduce the volume geodesic derivative ρ describing the interaction between the dynamics and the volume μ on the manifold, and we study its basic properties.

Recall that, given a smooth volume form μ on M , its value μ_x at a point is a nonzero element of the space $\Lambda^n(T_x M)$. We can associate with it the unique element μ_x^* in $\Lambda^n(T_x M)^* = \Lambda^n(T_x^* M)$ satisfying $\mu_x^*(\mu_x) = 1$. This defines a volume form on the fiber $T_x^* M$. By the canonical identification $T_x^* M \simeq T_\lambda(T_x^* M)$ of a vector space with its tangent space to a point, this induces a volume form μ_λ^* on the vertical space $\mathcal{V}_\lambda := T_\lambda(T_x^* M)$ for each $\lambda \in T_x^* M$.

Let $\gamma(t) = \pi(\lambda(t))$ be an ample and equiregular geodesic defined on $[0, T]$, with $\lambda(t) = e^{t\tilde{H}}(\lambda)$ and $\lambda \in T_x^* M$. Denote by \mathcal{A} the set of $\lambda \in T^* M$ such that the corresponding trajectory is ample and equiregular. For a fixed $x \in M$, we set $\mathcal{A}_x := \mathcal{A} \cap T_x^* M$.

Notice that, if $\lambda \in \mathcal{A}_x$, then the exponential map $\pi \circ e^{t\tilde{H}} : T_x^* M \rightarrow M$ is a local diffeomorphism at λ , for small $t \geq 0$. Then it makes sense to consider the pull-back measure $(\pi \circ e^{t\tilde{H}})^* \mu$ and compare its restriction to the vertical space \mathcal{V}_λ with μ_λ^* .

DEFINITION 5.1. — *For every $\lambda \in \mathcal{A}_x$ we define the volume geodesic derivative by the identity*

$$(5.1) \quad \rho(\lambda)\mu_\lambda^* = \frac{d}{dt} \Big|_{t=0} \log \left(t^{-\mathcal{N}(\lambda)} (\pi \circ e^{t\tilde{H}})^* \mu \Big|_{\mathcal{V}_\lambda} \right),$$

where $\mathcal{N}(\lambda)$ is defined in (3.3).

Let $\{\theta_{ai}(t)\}_{ai \in D} \in T^*_{\gamma(t)}M$ be the coframe dual to $X_{ai}(t) = \pi_* F_{ai}(t)$ and define a volume form ω along γ as

$$(5.2) \quad \omega_{\gamma(t)} := \theta_{11}(t) \wedge \theta_{12}(t) \wedge \cdots \wedge \theta_{kn_k}(t).$$

Given a fixed smooth volume μ on M , let $g_\lambda : [0, T] \rightarrow \mathbb{R}$ be the smooth function such that

$$(5.3) \quad \mu_{\gamma(t)} = e^{g_\lambda(t)} \omega_{\gamma(t)}.$$

The first main result of this section is the relation between the invariant ρ and the function $g_\lambda(t)$ just introduced.

PROPOSITION 5.2. — For every $\lambda \in \mathcal{A}_x$ one has $\rho(\lambda) = \dot{g}_\lambda(0)$.

The proof of this Proposition is a corollary of the proof of the main theorem, that is proved in Section 8. We exploit the previous identity to prove some useful properties of the invariant ρ . We start by the following lemma.

LEMMA 5.3. — Let $\gamma(t) = \pi(e^{t\bar{H}}(\lambda))$ be an ample and equiregular geodesic. Then we have

$$\dot{g}_\lambda(t) = \dot{g}_{\lambda(t)}(0), \quad \forall t \in [0, T].$$

Proof. — Let $\lambda(t) = e^{t\bar{H}}(\lambda) \in T^*M$ be the lifted extremal and denote by $\gamma_t(s) := \gamma(t + s)$. Then $\gamma_t(s) = \pi(e^{s\bar{H}}(\lambda(t)))$ and we have the sequence of identities

$$e^{g_\lambda(t+s)} \omega_{\gamma(t+s)} = \mu_{\gamma(t+s)} = \mu_{\gamma_t(s)} = e^{g_{\lambda(t)}(s)} \omega_{\gamma_t(s)}$$

Moreover $\omega_{\gamma(t+s)} = \omega_{\gamma_t(s)}$ since, if $(E_{\lambda(t+s)}, F_{\lambda(t+s)})$ is a canonical frame along $\lambda(t + s)$, it is a canonical frame also for $e^{s\bar{H}}\lambda(t)$. It follows that $g_\lambda(t + s) = g_{\lambda(t)}(s)$ for every s and differentiating with respect to s at $s = 0$ one gets the result. □

Lemma 5.3 allows us to write g as a function of ρ , as follows

$$(5.4) \quad g_\lambda(t) = g_\lambda(0) + \int_0^t \dot{g}_\lambda(s) ds = g_\lambda(0) + \int_0^t \rho(\lambda(s)) ds.$$

PROPOSITION 5.4. — Let \mathbb{T} be any admissible extension of $\dot{\gamma}$ and ω the n -form defined in (5.2). Then for every $\lambda \in T^*_xM$

$$(5.5) \quad \rho(\lambda) = (\operatorname{div}_\mu \mathbb{T} - \operatorname{div}_\omega \mathbb{T})|_x.$$

Proof. — It is a direct consequence of the classical identity

$$\operatorname{div}_{f\omega} X - \operatorname{div}_\omega X = X(\log f)$$

which holds for every smooth volume form ω , smooth function f and smooth vector field X . □

Remark 5.5 (On the volume form ω , I). — In the Riemannian case $\{X_{ai}(t)\}_{ai \in D}$ is an orthonormal frame for the Riemannian metric by Lemma 4.6 and the form ω coincides with the restriction of the canonical Riemannian volume vol_g on the curve γ . Hence

$$(5.6) \quad \rho(v) = (\operatorname{div}_\mu \mathbb{T})|_x - (\operatorname{div}_{\operatorname{vol}_g} \mathbb{T})|_x.$$

In the general case ρ can still be represented as the difference of two divergences but the volume form ω depends on the curve γ and is not the restriction to the curve of a global volume form.

Next we recall a refinement of Lemma 4.6.

LEMMA 5.6 ([4], Lemma 8.5). — For $t \in [0, T]$, the projections $X_{ai}(t) = \pi_* F_{ai}(t)$ satisfy

$$X_{ai}(t) = (-1)^{i-1} \mathcal{L}_T^{i-1}(X_{a1}(t)) \pmod{\mathcal{F}_{\gamma(t)}^{i-1}}, \quad a = 1, \dots, k, \quad i = 1, \dots, n_a.$$

PROPOSITION 5.7 (ρ depends only on μ and the symbol along γ). — Let γ, γ' be two geodesics associated with initial covectors $\lambda \in \mathcal{A}_{\gamma(0)}$ and $\lambda' \in \mathcal{A}_{\gamma'(0)}$ respectively. Assume that there exists a diffeomorphism ϕ on M such that for $t \geq 0$ small enough

- (1) $\phi(\gamma(t)) = \gamma'(t)$,
- (2) $\phi_*|_{\gamma(t)}$ induces an isomorphism of symbols between $S_{\gamma(t)}$ and $S_{\gamma'(t)}$
- (3) $\phi^* \mu_{\gamma'(t)} = \mu_{\gamma(t)}$,

Then $\rho(\lambda) = \rho(\lambda')$.

Proof. — Let $\{E_{ai}(t), F_{ai}(t)\}_{ai \in D}$ and $\{E'_{ai}(t), F'_{ai}(t)\}_{ai \in D}$ be canonical frames with respect to λ and λ' respectively, and $X_{ai}(t) = \pi_*(F_{ai}(t))$, $X'_{ai}(t) = \pi_*(F'_{ai}(t))$ be the associated basis of $T_{\gamma(t)}M$ and $T_{\gamma'(t)}M$. Since ω evaluated on the projection of the canonical frame gives 1 by construction, we have

$$(5.7) \quad e^{g_\lambda(t)} = |\mu_{\gamma(t)}(X_{11}(t), \dots, X_{kn_k}(t))|,$$

$$(5.8) \quad e^{g_{\lambda'}(t)} = |\mu_{\gamma'(t)}(X'_{11}(t), \dots, X'_{kn_k}(t))|.$$

Recall that $\{X_{a1}\}_{a=1}^k$ (resp. $\{X'_{a1}\}_{a=1}^k$) is an orthonormal basis for $\mathcal{D}_{\gamma(t)}$ (resp. $\mathcal{D}'_{\gamma'(t)}$). Since the linear map $\phi_*|_{\gamma(t)} : \mathcal{D}_{\gamma(t)} \rightarrow \mathcal{D}'_{\gamma'(t)}$ is an isometry

for small $t \geq 0$, there exists a family of orthogonal $k \times k$ matrices $O(t)$ such that

$$X'_{a1}(t) = \sum_{b=1}^k O_{ab}(t)\phi_*(X_{b1}(t)), \quad \text{for } a = 1, \dots, k.$$

Moreover using Lemma 5.6 we have for $i > 1$

$$\begin{aligned} X'_{ai}(t) &= (-1)^{i-1} \mathcal{L}_{\Gamma'}^{(i-1)}(X'_{a1}(t)) \pmod{\mathcal{F}_{\gamma'(t)}^{i-1}} \\ (5.9) \quad &= (-1)^{i-1} \mathcal{L}_{\Gamma'}^{(i-1)} \left(\sum_{b=1}^k O(t)_{ab} \phi_*(X_{b1}(t)) \right) \pmod{\mathcal{F}_{\gamma'(t)}^{i-1}} \\ &= (-1)^{i-1} \sum_{b=1}^k O(t)_{ab} \mathcal{L}_{\Gamma'}^{(i-1)}(\phi_*(X_{b1}(t))) \pmod{\mathcal{F}_{\gamma'(t)}^{i-1}}, \end{aligned}$$

where the last identity follows by the chain rule. Indeed, when one differentiates the matrix $O(t)$, one obtains elements of $\mathcal{F}_{\gamma'(t)}^{i-1}$. Then

$$\begin{aligned} X'_{ai}(t) &= (-1)^{i-1} \sum_{b=1}^k O(t)_{ab} \phi_* \mathcal{L}_{\Gamma}^{(i-1)}(X_{b1}(t)) \pmod{\mathcal{F}_{\gamma'(t)}^{i-1}} \\ (5.10) \quad &= \sum_{b=1}^{d_i} O(t)_{ab} \phi_* X_{bi}(t) \pmod{\mathcal{F}_{\gamma'(t)}^{i-1}}, \end{aligned}$$

where the sum is restricted to those indices b such that $bi \in D$. This proves that there exists an orthogonal transformation that sends $\phi_* X_{ai}$ in X'_{ai} . Therefore

$$\begin{aligned} e^{g_{\lambda'}(t)} &= \left| \mu_{\gamma'(t)}(X'_{1,1}(t), \dots, X'_{kn_k}(t)) \right| \\ (5.11) \quad &= \left| \mu_{\gamma'(t)}(\phi_* X_{1,1}(t), \dots, \phi_* X_{kn_k}(t)) \right| \\ &= \left| (\phi^* \mu)_{\gamma(t)}(X_{1,1}(t), \dots, X_{kn_k}(t)) \right| = e^{g_{\lambda}(t)}, \end{aligned}$$

and the proof is complete. □

Actually, from the previous proof it follows that the invariant ρ depends only on the 1-jet of the one-parameter family of symbols (and the volume form μ) along the geodesics.

Remark 5.8 (On the volume form ω , II). — The volume form ω depends only on the symbol of the structure along the geodesic, that represents the microlocal nilpotent approximation of the structure at x along $\gamma(t)$. Symbols at different points along any geodesic in the Riemannian manifold are isomorphic, while in the general case this symbol could depend on the point on the curve. This is analogous to what happens for the nilpotent

approximation for a distribution (see for instance the discussion contained in [1]).

LEMMA 5.9. — *Let $\gamma(t) = \pi(\lambda(t))$ be an ample and equiregular geodesic. Assume that \mathbb{T} is an admissible extension of its tangent vector such that $e^{t\mathbb{T}}$ is an isometry of the distribution along $\gamma(t)$. Then $\operatorname{div}_\omega \mathbb{T}|_{\gamma(t)} = 0$ and $\rho(\lambda(t)) = \operatorname{div}_\mu \mathbb{T}$.*

Proof. — If we show that $\operatorname{div}_\omega \mathbb{T}|_{\gamma(t)} = 0$, then from (5.5) it immediately follows that $\rho(\lambda(t)) = \operatorname{div}_\mu \mathbb{T}$ and ρ depends only on the variation of the volume μ along the curve.

Let $\{X_{ai}(t)\}_{ai \in D}$ be the basis of $T_{\gamma(t)}M$ induced by the canonical frame along the curve $\lambda(t)$. The divergence is computed as

$$\begin{aligned}
 (5.12) \quad (\operatorname{div}_\omega \mathbb{T})|_{\gamma(t)} \omega_{\gamma(t)}(X_{11}(t), \dots, X_{kn_k}(t)) &= \mathcal{L}_{\mathbb{T}} \omega(X_{11}, \dots, X_{kn_k})|_{\gamma(t)} \\
 &= \frac{d}{d\epsilon} \Big|_{\epsilon=0} \omega_{\gamma(t+\epsilon)}(e_*^{\epsilon\mathbb{T}} X_{11}, \dots, e_*^{\epsilon\mathbb{T}} X_{kn_k}).
 \end{aligned}$$

Since the flow of \mathbb{T} is an isometry of the graded structure that defines the symbol, the last quantity is equal to 0, which proves that $\operatorname{div}_\omega \mathbb{T} = 0$ along the curve. □

LEMMA 5.10. — *The function $\rho : \mathcal{A} \rightarrow \mathbb{R}$ is a rational function.*

Proof. — Since H is a quadratic function on fibers, then the vector field \vec{H} is fiber-wise polynomial. Therefore for any vector field $V(t) \in T_{\lambda(t)}(T^*M)$, the quantity $\dot{V} = [\vec{H}, V]$ is a rational function of the initial covector λ . It follows that both E and F are rational as functions of λ , and so are also the projections $X(t) = \pi_* F(t)$. We conclude that

$$e^{g\lambda(t)} = |\mu_{\gamma(t)}(X_{a1}(t), \dots, X_{kn_k}(t))|,$$

and the coefficients of its Taylor expansion, are rational expressions in λ . □

Remark 5.11. — If the symbol is constant along the trajectory (i.e., symbols at different points are isomorphic) through a diffeomorphism ϕ and μ is preserved by ϕ , then $\rho(\lambda(t)) = 0$.

Indeed it is sufficient to apply Proposition 5.7 to $\gamma_t(s) := \gamma(t + s)$ for every s and one gets for $s, t \geq 0$ small $g(t) = g(t + s)$, that means that g is constant and $\rho = 0$.

DEFINITION 5.12. — *We say that an Hamiltonian of our class (2.3) is unimodular if there exists a volume form μ such that $\rho = 0$ on \mathcal{A} .*

It is easy to see that if an Hamiltonian is unimodular with respect to some volume form μ , then μ is unique.

6. A formula for ρ

In this section we provide a formula to compute ρ in terms only of the volume μ and the linear maps \mathcal{L}_\top^i . This will give another proof of the fact that the quantity $\rho(\lambda(t))$ depends only on the symbol and on μ along $\gamma(t) = \pi(\lambda(t))$.

Fix a smooth volume μ on M and let Y_1, \dots, Y_k be an orthonormal basis of \mathcal{D} in a neighborhood of x_0 . Choose vector fields Y_{k+1}, \dots, Y_n such that Y_1, \dots, Y_n is a local basis satisfying $\mu(Y_1, \dots, Y_n) = 1$ and define an auxiliary inner product on the tangent space declaring that this basis is orthonormal.

Let $\gamma(t) = \pi(e^{t\bar{H}}(\lambda))$ be an ample and equiregular curve, with initial covector $\lambda \in T_{x_0}^*M$. Recall that, according to the definition of $g_\lambda(t)$, it holds

$$(6.1) \quad g_\lambda(t) = \log |\mu(P_t)|,$$

where P_t is the parallelotope whose edges are the projections $\{X_{ai}(t)\}_{ai \in D}$ of the horizontal part of the canonical frame $X_{ai}(t) = \pi_* \circ e^{t\bar{H}} F_{ai}(t) \in T_{\gamma(t)}M$, namely

$$(6.2) \quad P_t = \bigwedge_{ai \in D} X_{ai}(t).$$

By Lemma 5.6 we can write the adapted frame $\{X_{ai}\}_{ai \in D}$ in terms of the smooth linear maps \mathcal{L}_\top , and we obtain the following identity

$$(6.3) \quad P_t = \bigwedge_{i=1}^m \bigwedge_{a_i=1}^{d_i} X_{a_i i}(t) = \bigwedge_{i=1}^m \bigwedge_{a_i=1}^{d_i} \mathcal{L}_\top^{i-1}(X_{a_i 1}(t)).$$

Consider the flag $\{\mathcal{F}_{\gamma(t)}^i\}_{i=1}^m$ and, using the auxiliary inner product induced by the choice of the basis, define the following sequence of subspaces of $T_{\gamma(t)}M$: for every $i \geq 1$ set (with the understanding that $\mathcal{F}^0 = \{0\}$)

$$(6.4) \quad V_i := \mathcal{F}_{\gamma(t)}^i \cap (\mathcal{F}_{\gamma(t)}^{i-1})^\perp.$$

The subspace V_i has dimension $\dim V_i = \dim \mathcal{F}_{\gamma(t)}^i - \dim \mathcal{F}_{\gamma(t)}^{i-1}$. Therefore there exists an isomorphism between $\mathcal{F}_{\gamma(t)}^i / \mathcal{F}_{\gamma(t)}^{i-1}$ and V_i , such that every $Y \in \mathcal{F}_{\gamma(t)}^i / \mathcal{F}_{\gamma(t)}^{i-1}$ is associated with the element of its equivalent class that lies in V_i . In conclusion, for the computation of $g_\lambda(t)$ in (6.1), one can

replace the vector $\mathcal{L}_T^{i-1}(X_{a_{i1}}(t))$ of the parallelotope in (6.3) with the corresponding equivalent element in V_i .

Now consider the surjective map $\mathcal{L}_T^{i-1} : \mathcal{D}_{\gamma(t)} \rightarrow \mathcal{F}_{\gamma(t)}^i / \mathcal{F}_{\gamma(t)}^{i-1}$. For every $i = 1, \dots, m$ this map descends to an isomorphism $\mathcal{L}_T^{i-1} : \mathcal{D}_{\gamma(t)} / \ker \mathcal{L}_T^{i-1} \rightarrow \mathcal{F}_{\gamma(t)}^i / \mathcal{F}_{\gamma(t)}^{i-1} \simeq V_i$. Then, thanks to the inner product structure on V_i , we can consider the map

$$(\mathcal{L}_T^{i-1})^* \circ \mathcal{L}_T^{i-1} : \mathcal{D}_{\gamma(t)} / \ker \mathcal{L}_T^{i-1} \rightarrow \mathcal{D}_{\gamma(t)} / \ker \mathcal{L}_T^{i-1}$$

obtained by composing \mathcal{L}_T^{i-1} with its adjoint $(\mathcal{L}_T^{i-1})^*$. This composition is a symmetric invertible operator and we define the smooth family of symmetric operators

$$(6.5) \quad M_i(t) := (\mathcal{L}_T^{i-1})^* \circ \mathcal{L}_T^{i-1} : \mathcal{D}_{\gamma(t)} / \ker \mathcal{L}_T^{i-1} \rightarrow \mathcal{D}_{\gamma(t)} / \ker \mathcal{L}_T^{i-1},$$

$$i = 1, \dots, m.$$

Recall in particular that for every $v_1, v_2 \in \mathcal{D}_{\gamma(t)} / \ker \mathcal{L}_T^{i-1}$ it holds the identity

$$\langle (\mathcal{L}_T^{i-1})^* \circ \mathcal{L}_T^{i-1} v_1, v_2 \rangle_{\mathcal{D}_{\gamma(t)}} = \langle \mathcal{L}_T^{i-1} v_1, \mathcal{L}_T^{i-1} v_2 \rangle_{V_i}.$$

By the expression of the parallelotope P_t with elements of the subspaces V_i and the definition of μ as the dual of an orthonormal basis of $T_{\gamma(t)}M$, we have

$$(6.6) \quad |\mu(P_t)| = \left| \mu \left(\bigwedge_{i=1}^m \bigwedge_{a_i=1}^{d_i} \mathcal{L}_T^{i-1}(X_{a_{i1}}(t)) \right) \right| = \sqrt{\prod_{i=1}^m \det M_i(t)}.$$

This formula does not depend on the chosen extension Y_{k+1}, \dots, Y_n of the orthonormal basis of \mathcal{D} , since in the computations we only used that the volume μ evaluated at this basis is equal to 1. For $\rho(\lambda) = \frac{d}{dt} \Big|_{t=0} \log |\mu(P_t)|$, a simple computation shows that

$$(6.7) \quad \rho(\lambda) = \frac{1}{2} \sum_{i=1}^m \text{tr} (M_i(0)^{-1} \dot{M}_i(0)).$$

7. Sub-Riemannian manifolds

In this section we specialize our construction to sub-Riemannian manifolds and we investigate in more details the properties of the invariant ρ for these structures.

Recall that a sub-Riemannian structure on a smooth manifold M is given by a completely non-integrable vector distribution \mathcal{D} endowed with an inner

product on it. In particular \mathcal{D} has constant rank but does not need to be equiregular. An *admissible curve* is a curve that is almost everywhere tangent to \mathcal{D} and for such a curve γ we can compute its length by the classical formula

$$\ell(\gamma) = \int_0^T \|\dot{\gamma}(s)\| \, ds.$$

Once we fix a local orthonormal frame X_1, \dots, X_k for g on \mathcal{D} , the problem of finding the geodesics in a sub-Riemannian manifold, namely the problem of minimizing the length of a curve between two fixed points, is equivalent to the minimization of the energy (with $T > 0$ fixed) and can be rewritten as the control problem

$$(7.1) \quad \begin{cases} \dot{x} = \sum_{i=1}^k u_i X_i(x) \\ J_T(u) = \frac{1}{2} \int_0^T \|u(s)\|^2 \, ds \rightsquigarrow \min, \end{cases}$$

This is an affine control problem, with zero drift field and quadratic cost. The complete non-integrability assumption on the distribution $\mathcal{D} = \text{span}\{X_1, \dots, X_k\}$ implies that the assumptions (H0)–(H1) are satisfied. The Hamiltonian function is fiber-wise quadratic and convex on fibers. In coordinates it is written as

$$H(p, x) = \frac{1}{2} \sum_{i=1}^k \langle p, X_i(x) \rangle^2, \quad (p, x) \in T^*M.$$

Denote by \mathcal{A} the set of $\lambda \in T^*M$ such that the corresponding trajectory is ample and equiregular. For a fixed $x \in M$ we set $\mathcal{A}_x = \mathcal{A} \cap T_x^*M$.

PROPOSITION 7.1. — *The set \mathcal{A} is a non-empty open dense subset of T^*M .*

Proof. — Denote by $\mathcal{F}_\lambda^i := \mathcal{F}_{\gamma(0)}^i$ where γ is the trajectory associated with initial covector λ and set $k_i(\lambda) = \dim \mathcal{F}_\lambda^i$. By semicontinuity of the rank the integer valued and bounded function $\lambda \mapsto k_i(\lambda)$ is locally constant on an open dense set Ω_i of T^*M . Since the intersection of a finite number of open dense sets is open and dense, it follows that the set $\Omega = \bigcap_i \Omega_i$ where the growth vector is locally constant is open and dense in T^*M . To prove that it is non-empty fix an arbitrary point $x \in M$ and consider a $\lambda \in T_x^*M$ such that the corresponding trajectory is ample for all t (the existence of such a trajectory is proved in [4, Section 5.2]). Since the functions $t \mapsto \dim \mathcal{F}_{\lambda(t)}^i$ are lower semi-continuous and bounded with respect to t , repeating the previous argument we have that they are locally constant on an open dense set of $[0, T]$, then the curve is equiregular at these points. □

We stress once more that for a fixed $x \in M$ one can have $\mathcal{A}_x = \emptyset$, as for instance in the non equiregular case. On the other hand, for every fixed x the set λ such that the corresponding trajectory is ample is open and dense and on each of these trajectories we can find equiregular points arbitrarily close to x .

7.1. Homogeneity properties

For all $c > 0$, let $H_c := H^{-1}(c/2)$ be the level set of the Hamiltonian function. In particular H_1 is the unit cotangent bundle: the set of initial covectors associated with unit-speed geodesics. Since the Hamiltonian function is fiber-wise quadratic, we have the following property for any $c > 0$

$$(7.2) \quad e^{t\vec{H}}(c\lambda) = ce^{ct\vec{H}}(\lambda).$$

Let $\delta_c : T^*M \rightarrow T^*M$ be the dilation along the fibers $\delta_c(\lambda) = c\lambda$ (if we write $\lambda = (p, x)$ this means $\delta_c(p, x) = (cp, x)$). Indeed $\alpha \mapsto \delta_{e^\alpha}$ is a one-parameter group of diffeomorphisms. Its generator is the Euler vector field $\epsilon \in \Gamma(\mathcal{V})$, and is characterized by $\delta_c = e^{(\ln c)\epsilon}$. We can rewrite (7.2) as the following commutation rule for the flows of \vec{H} and ϵ :

$$(7.3) \quad e^{t\vec{H}} \circ \delta_c = \delta_c \circ e^{ct\vec{H}}.$$

Observe that δ_c maps H_1 diffeomorphically on H_c . Let $\lambda \in H_1$ be associated with an ample, equiregular geodesic with Young diagram D . Clearly also the geodesic associated with $\lambda^c := c\lambda \in H_c$ is ample and equiregular, with the same Young diagram. This corresponds to a reparametrization of the same curve: in fact $\lambda^c(t) = e^{t\vec{H}}(c\lambda) = c(\lambda(ct))$, hence $\gamma^c(t) = \pi(\lambda^c(t)) = \gamma(ct)$. The canonical frame associated with $\lambda^c(t)$ can be recovered by the one associated with $\lambda(t)$ as shown in the following proposition. Its proof can be found in [11].

PROPOSITION 7.2. — *Let $\lambda \in H_1$ and $\{E_{ai}, F_{ai}\}_{ai \in D}$ be the associated canonical frame along the extremal $\lambda(t)$. Let $c > 0$ and define, for $ai \in D$*

$$(7.4) \quad E_{ai}^c(t) := \frac{1}{c^i} (d_{\lambda(ct)} P_c) E_{ai}(ct), \quad F_{ai}^c(t) := c^{i-1} (d_{\lambda(ct)} \delta_c) F_{ai}(ct).$$

*The moving frame $\{E_{ai}^c(t), F_{ai}^c(t)\}_{ai \in D} \in T_{\lambda^c(t)}(T^*M)$ is a canonical frame associated with the initial covector $\lambda^c = c\lambda \in H_c$, with matrix*

$$(7.5) \quad R_{ai,bj}^{\lambda^c}(t) = c^{i+j} R_{ai,bj}^\lambda(ct).$$

By Proposition 7.2, it follows the following homogeneity property of g_λ , and as a consequence of the function ρ .

LEMMA 7.3. — For every $\lambda \in \mathcal{A}_x$ and $c > 0$ one has $c\lambda \in \mathcal{A}_x$. Moreover

$$e^{g_{c\lambda}(t)} = c^{\mathcal{Q}-n} e^{g_\lambda(ct)},$$

where $n = \dim M$ and $\mathcal{Q} := \sum_{i=1}^m id_i$.

Proof. — Let $X_{ai}^c(t)$ and $X_{ai}(ct)$ be the basis of $T_{\gamma^c(t)}M = T_{\gamma(ct)}M$ induced by the canonical frame. Then by Proposition 7.2 it holds the identity $X_{ai}^c(t) = c^{i-1}X_{ai}(ct)$. Therefore by the definition of g_λ and $g_{c\lambda}$ we have

$$\begin{aligned} e^{g_{c\lambda}(t)} &= |\mu_{\gamma(ct)}(X_{11}^c(t), \dots, X_{kn_k}^c(t))| \\ &= \prod_{i=1}^m \prod_{j=1}^{d_i} c^{i-1} |\mu_{\gamma(ct)}(X_{11}(ct), \dots, X_{kn_k}(ct))| \\ (7.6) \qquad &= \prod_{i=1}^m c^{(i-1)d_i} e^{g_\lambda(ct)} \\ &= c^{\mathcal{Q}-n} e^{g_\lambda(ct)}. \qquad \square \end{aligned}$$

Lemma 7.3 gives $g_{c\lambda}(t) = g_\lambda(ct) + (\mathcal{Q} - n) \log(c)$ and differentiating at $t = 0$ we obtain

COROLLARY 7.4. — For every $\lambda \in \mathcal{A}_x$ and $c > 0$ one has

$$(7.7) \qquad \rho(c\lambda) = c\rho(\lambda).$$

Remark 7.5. — The function ρ is homogeneous of degree one but, in general, it might not be smooth. Indeed using formula (6.7) and denoting by $M_i^c(t)$ the matrices associated with the reparametrized curve γ^c , one can show from the homogeneity properties of Proposition 7.2 that

$$(7.8) \qquad M_i^c(t) = c^{2i-2}M_i(ct), \qquad \dot{M}_i^c(t) = c^{2i-1}\dot{M}_i(ct)$$

from which it follows that ρ is a rational function in λ with the degree of the denominator which is at most λ^{2m-2} . Notice that using (7.8) at $t = 0$ one can obtain another proof of (7.7) by

$$\rho(c\lambda) = \frac{1}{2} \sum_{i=1}^m \text{tr} (M_i^c(0)^{-1} \dot{M}_i^c(0)) = \frac{1}{2} \sum_{i=1}^m \text{tr} (cM_i(0)^{-1} \dot{M}_i(0)) = c\rho(\lambda).$$

7.2. Contact manifolds

In this section we focus on the special case of a contact sub-Riemannian manifold. Recall that a sub-Riemannian manifold (M, \mathcal{D}, g) of odd dimension $2n + 1$ is *contact* if there exists a non degenerate 1-form $\omega \in \Lambda^1(M)$, such that $\mathcal{D}_x = \ker \omega_x$ for every $x \in M$ and $d\omega|_{\mathcal{D}}$ is non degenerate. In this case \mathcal{D} is called *contact distribution*.

Remark 7.6. — Given a sub-Riemannian contact manifold, the contact form ω is not unique. Indeed if ω is a contact form then also $\alpha\omega$ is a contact form for any non-vanishing smooth function α . Once a contact form ω is fixed we can associate the *Reeb vector field*, X_0 , which is the unique vector field such that $\omega(X_0) = 1$ and $d\omega(X_0, \cdot) = 0$. Since the Reeb vector field X_0 is transversal to \mathcal{D} , we normalize ω so that $\|X_0\|_{\mathcal{D}^2/\mathcal{D}} = 1$.

The contact form ω induces a fiber-wise linear map $J : \mathcal{D} \rightarrow \mathcal{D}$, defined by

$$\langle JX, Y \rangle = d\omega(X, Y) \quad \forall X, Y \in \mathcal{D}.$$

Observe that the restriction $J_x := J|_{\mathcal{D}_x}$ is a linear skew-symmetric operator on (\mathcal{D}_x, g_x) .

Let X_1, \dots, X_{2n} be a local orthonormal frame of \mathcal{D} , then X_1, \dots, X_{2n}, X_0 is a local frame adapted to the flag of the distribution. Let $\nu^1, \dots, \nu^{2n}, \nu^0$ be the associated dual frame. The *Popp volume* μ on M (see [9] for more details) is the volume

$$(7.9) \quad \mu = \nu^1 \wedge \dots \wedge \nu^{2n} \wedge \nu^0.$$

On contact sub-Riemannian manifolds, every non constant geodesic $\gamma(t) = \pi(e^{t\vec{H}}(\lambda))$ is ample and equiregular with the same growth vector $(2n, 2n + 1)$. Moreover, it is possible to compute explicitly the value of the associated smooth function $g_\lambda(t)$ and the constant C_λ of Theorem 1.2 (cf. Remark 8.2).

We compute now the value of the function $g_\lambda(t)$ with respect to the Popp’s volume and a given geodesic.

PROPOSITION 7.7. — *Let $\gamma(t) = \pi(e^{t\vec{H}}(\lambda))$ be a geodesic on a contact manifold. Then*

$$(7.10) \quad \rho(\lambda) = \left. \frac{d}{dt} \right|_{t=0} \log \|J\dot{\gamma}(t)\|,$$

In particular if $J^2 = -1$ then $\rho = 0$.

Proof. — Recall that, by definition of g_λ (cf. (5.3)), one has

$$(7.11) \quad g_\lambda(t) = \log |\mu(P_t)|$$

where P_t is the parallelotope whose edges are given by the projections $X_{ai}(t)$ of the fields $F_{ai}(t)$ of a canonical basis along $\lambda(t)$ on $T_{\gamma(t)}M$.

Let \mathbb{T} be an horizontal extension of the tangent vector field $\dot{\gamma}(t)$ and consider the map $\mathcal{L}_\mathbb{T} : \mathcal{D}_{\gamma(t)} \rightarrow \mathcal{D}_{\gamma(t)}^2/\mathcal{D}_{\gamma(t)}$. Since the manifold is contact, this map is surjective. and its kernel is a subspace of $\mathcal{D}_{\gamma(t)}$ of dimension $2n - 1$. Let X_1, \dots, X_{2n} be an orthonormal basis of $\mathcal{D}_{\gamma(t)}$ such that $X_1 \in (\ker \mathcal{L}_\mathbb{T})^\perp$

and $X_2, \dots, X_{2n} \in \ker \mathcal{L}_T$. Then there exists an orthogonal map that transforms the first $2n$ vectors projections of the canonical basis, in this basis.

Notice that the definition (7.11) of $g_\lambda(t)$ does not change if we replace the first $2n$ edges of the parallelotope by X_1, \dots, X_{2n} . Moreover, by Lemma 5.6, the last projected vector $X_{ai} = X_{1,2}$ can be written as

$$X_{1,2}(t) = -\mathcal{L}_T X_1(t) \pmod{\mathcal{D}}.$$

Notice that X_1 is not in the kernel of \mathcal{L}_T , thus this basis is also adapted to the Young diagram of γ . Thanks to (7.9), the Popp volume of the parallelotope is equal to the length of the component of $\mathcal{L}_T X_1(t)$ with respect to X_0 , namely

$$|\mu(P_t)| = |\langle [T, X_1], X_0 \rangle_{\gamma(t)}|.$$

This quantity can be written equivalently in terms of the map J . Indeed

$$(7.12) \quad \begin{aligned} |\mu(P_t)| &= |\langle [T, X_1], X_0 \rangle_{\gamma(t)}| = |\omega_{\gamma(t)}([T, X_1])| = |d\omega_{\gamma(t)}(T, X_1)| \\ &= |\langle J_{\gamma(t)} T, X_1 \rangle_{\gamma(t)}|. \end{aligned}$$

Since $\langle JT, Y \rangle = -\omega(\mathcal{L}_T Y)$ for every horizontal field Y , then $\ker \mathcal{L}_T = JT^\perp$. This implies that JT is a multiple of X_1 , i.e., $JT = \|JT\|X_1$. Then we simplify the formula for $|\mu(P_t)|$ as

$$|\mu(P_t)| = |\langle J_{\gamma(t)} T, X_1 \rangle_{\gamma(t)}| = \|JT_{\gamma(t)}\|.$$

Notice that if $J^2 = -1$, then J is an isometry, hence $\|JT_{\gamma(t)}\| = \|T_{\gamma(t)}\| = 1$. □

Remark 7.8. — If $\dim M = 3$, then $\rho(\lambda(t)) = 0$ for every t . Indeed $\ker \mathcal{L}_T$ has dimension 1 and $T = \|T\|X_2$. Moreover, if we denote by c_{ij}^k the structure constants such that $[X_i, X_j] = \sum_{k=0}^2 c_{ij}^k X_k$, then the normalization of ω implies $c_{12}^0 = -1$ and

$$(7.13) \quad \begin{aligned} |\mu(P_t)| &= |\langle [T, X_1], X_0 \rangle| = \|T_{\gamma(t)}\| |\langle [X_2, X_1], X_0 \rangle| \\ &= \|T_{\gamma(t)}\| = 1. \end{aligned}$$

Remark 7.9. — Notice that even in the contact case, not every structure is unimodular (in the sense of Definition 5.12). When $J^2 = -1$ then the structure is unimodular, choosing μ as the Popp volume. See also [3] for the computation of the curvature in the contact case.

8. Proof of the main result

In this section we prove the following proposition, that is Theorem 1.2 written along the canonical frame.

PROPOSITION 8.1. — Let $\gamma(t) = \pi(e^{t\bar{H}}(\lambda))$ be an ample equiregular geodesic and let $\omega_{\gamma(t)}$ be the n -form defined in (5.2). Given a volume μ on M , define implicitly the smooth function $g_\lambda : [0, T] \rightarrow \mathbb{R}$ by $\mu_{\gamma(t)} = e^{g_\lambda(t)}\omega_{\gamma(t)}$. Then we have the following Taylor expansion

$$(8.1) \quad \left\langle \left(\pi \circ e^{t\bar{H}}\right)^* \mu, E(0) \right\rangle \Big|_\lambda = C_\lambda t^{\mathcal{N}(\lambda)} e^{g_\lambda(t)} \left(1 - t^2 \frac{\text{tr}\mathcal{R}_\lambda}{6} + o(t^2) \right)$$

where E is the n -dimensional row vector introduced in Remark 4.2 and C_λ depends only on the structure of the Young diagram. In particular we have the identity

$$\rho(\lambda)\mu_\lambda^* = \frac{d}{dt} \Big|_{t=0} \log \left(t^{-\mathcal{N}(\lambda)} \left(\pi \circ e^{t\bar{H}} \right)^* \mu \Big|_{\mathcal{V}_\lambda} \right) = \dot{g}_\lambda(0)\mu_\lambda^*.$$

Remark 8.2. — As it follows from the proof, the constant C_λ is explicitly computed by

$$(8.2) \quad C_\lambda = \prod_{a=1}^k \frac{\prod_{j=0}^{n_a-1} j!}{\prod_{j=n_a}^{2n_a-1} j!} > 0.$$

In the contact case, since the Young diagram is equal for all λ , with $2n$ rows of length 1, and one row of length 2, the leading constant is $C_\lambda = \frac{1}{12}$.

Proof. — The left hand side of the equation (8.1) can be rewritten as

$$\left\langle \left(\pi \circ e^{t\bar{H}}\right)^* \mu, E(0) \right\rangle \Big|_\lambda = \left\langle e^{g(t)}\omega, \left(\pi \circ e^{t\bar{H}}\right)_* E(0) \right\rangle \Big|_{\gamma(t)}.$$

For every $ai \in D$, the field $e_*^{t\bar{H}} E_{ai}(0)$ is a Jacobi field, so in coordinates with respect to the canonical frame we can write

$$e_*^{t\bar{H}} E(0) = E(t)M(t) + F(t)N(t)$$

for $n \times n$ matrices M and N , that satisfy the Jacobi equations (4.11). More explicitly we have the system

$$(8.3) \quad \begin{cases} \dot{N}_{ai,bj} = N_{ai-1,bj} & \text{if } i \neq 1 \\ \dot{N}_{a1,bj} = M_{a1,bj} \\ \dot{M}_{ai,bj} = -R(t)_{ai,ch} N_{ch,bj} - M_{ai+1,bj} & \text{if } i \neq n_a \\ \dot{M}_{an_a,bj} = -R(t)_{an_a,ch} N_{ch,bj}. \end{cases}$$

Moreover $M(0) = \text{Id}$ and $N(0) = 0$. It follows that

$$\left\langle \left(\pi \circ e^{t\bar{H}}\right)^* \mu, E(0) \right\rangle \Big|_\lambda = e^{g(t)} \det N(t).$$

In what follows we compute the Taylor expansion of the matrix $N(t)$ at 0.

Let us first discuss the proof in the case of a Young diagram made of a single row. In this case, for simplicity, we will omit the index a in the notation for N and M . Fix integers $1 \leq i, j \leq n$. The coefficients N_{ij} can be computed by integrating M_{1j} . So let us find the asymptotic expansion of M_{1j} . Notice that

$$\begin{aligned}
 M_{1j}(0) &= \delta_{1j} \\
 \dot{M}_{1j} &= -R_{1h}N_{hj} - M_{2j}(1 - \delta_{1n}) \\
 (8.4) \quad \ddot{M}_{1j} &= -\dot{R}_{1h}N_{hj} - \sum_{h \neq 1} R_{1h}N_{h-1j} - R_{11}M_{1j} \\
 &\quad + (1 - \delta_{1n})(R_{2h}N_{hj} + M_{3j}(1 - \delta_{2n}))
 \end{aligned}$$

In these equations the only non-vanishing component at $t = 0$ is $M_{jj}(0) = 1$, that can be obtained only by differentiating terms M_{ij} with $i < j$. Thus, in the expansion of $M_{1j}(t)$, the element M_{jj} appears first at $(j - 1)$ -th derivative. Next, it appears, multiplied by $R_{11}(0)$, at $(j + 1)$ -th derivative. We can conclude that the asymptotics of M_{1j} at $t = 0$ is

$$M_{1j}(t) = (-1)^{j-1} \frac{t^{j-1}}{(j-1)!} - (-1)^{j-1} R_{11}(0) \frac{t^{j+1}}{(j+1)!} + o(t^{j+1}).$$

Since M_{1j} is the i -th derivative of N_{ij} and $N(0) = 0$, we have also the expansion for N :

$$N_{ij}(t) = (-1)^{j-1} \frac{t^{i+j-1}}{(i+j-1)!} - (-1)^{j-1} R_{11}(0) \frac{t^{i+j+1}}{(i+j+1)!} + o(t^{i+j+1}).$$

Let us now consider a general distribution of dimension $k > 1$. Now we have to study the whole system in (8.3). Fix indices $ai, bj \in D$. Again, to find $N_{ai,bj}$ it's enough to determine the expansion of $M_{a1,bj}$, by integration. To compute the latter, notice that

$$\begin{aligned}
 M_{a1,bj}(0) &= \delta_{ab}\delta_{1j} \\
 \dot{M}_{a1,bj} &= -R_{a1,ch}N_{ch,bj} - M_{a2,bj}(1 - \delta_{1n_a}) \\
 M_{a1,bj}^{(2)} &= -\dot{R}_{a1,ch}N_{ch,bj} - \sum_{h \neq 1} R_{a1,ch}N_{ch-1,bj} - R_{a1,c1}M_{c1,bj} \\
 &\quad + (1 - \delta_{1n_a})(R_{a2,ch}N_{ch,bj} + M_{a3,bj}(1 - \delta_{2n_a}))
 \end{aligned}$$

When $a = b$, the argument is similar to the one discussed above when $k = 1$ (in this case every derivative generates also terms like $M_{ch,aj}$, but

these terms, when $c \neq a$, need higher order derivatives to generate non-vanishing terms). One obtains:

$$M_{a1,aj}(t) = (-1)^{j-1} \frac{t^{j-1}}{(j-1)!} - (-1)^{j-1} R_{a1,a1} \frac{t^{j+1}}{(j+1)!} + o(t^{j+1}),$$

$$N_{ai,aj}(t) = (-1)^{j-1} \frac{t^{i+j-1}}{(i+j-1)!} - (-1)^{j-1} R_{a1,a1} \frac{t^{i+j+1}}{(i+j+1)!} + o(t^{i+j+1}).$$

On the other hand, if $a \neq b$, then the first term different from zero of $M_{a1,bj}$ appears at $j + 1$ -th derivative, multiplied by $R_{a1,b1}$, Therefore the Taylor expansions of $M_{ai,bj}$ and of a generic element of the matrix N can be derived as

$$M_{a1,bj}(t) = \delta_{ab} (-1)^{j-1} \frac{t^{j-1}}{(j-1)!} - (-1)^{j-1} R_{a1,b1}(0) \frac{t^{j+1}}{(j+1)!} + o(t^{j+1}),$$

$$N_{ai,bj}(t) = \tilde{N}_{ai,bj} t^{i+j-1} - G_{ai,bj} t^{i+j+1} + o(t^{i+j+1}).$$

where the constant matrices \tilde{N} and G are defined by

$$\tilde{N}_{ai,bj} := (-1)^{j-1} \frac{\delta_{ab}}{(i+j-1)!}, \quad G_{ai,bj} := (-1)^{j-1} \frac{R_{a1,b1}(0)}{(i+j+1)!}.$$

To find the asymptotics of the left hand side of (8.1), we need only to determine the asymptotic of $\det N(t)$. Let $I_{\sqrt{t}}$ be a n -dimensional diagonal matrix, whose j -th element is equal to \sqrt{t}^{2i-1} , for $k_{i-1} < j \leq k_i$. Then the Taylor expansion of N can be written as $N(t) = I_{\sqrt{t}}(\tilde{N} - t^2 G + O(t^3))I_{\sqrt{t}}$ and its determinant is

$$\det N(t) = \det \tilde{N} t^{\mathcal{N}} \left(1 - \text{tr} \left(\tilde{N}^{-1} G \right) t^2 + o(t^2) \right),$$

where $\mathcal{N} = \mathcal{N}(\lambda)$ is the geodesic dimension given in Definition 3.7. The main coefficient is computed in the following lemma, whose proof is contained in Appendix A.

LEMMA 8.3. — *The determinant of \tilde{N} is given by*

$$(8.5) \quad C_\lambda = \det \tilde{N} = \prod_{a=1}^k \frac{\prod_{j=0}^{n_a-1} j!}{\prod_{j=n_a}^{2n_a-1} j!}.$$

Since the matrix \tilde{N} is block-wise diagonal, to find the trace of $\tilde{N}^{-1} G$ we just need the elements of G with $a = b$. Thanks to (4.14), that relates the curvature operator \mathcal{R}_λ with the elements of the matrix $R_{a1,b1}$, we have

$$\mathcal{R}_{aa} = 3 \frac{n_a}{4n_a^2 - 1} R_{a1,a1}(0).$$

Moreover we have the following identity

$$(8.6) \quad \text{tr} \left(\tilde{N}^{-1}G \right) = \sum_{a=1}^k \left(\sum_{i,j=1}^{n_a} [\tilde{N}^{-1}]_{ai,aj} \frac{(-1)^{j-1}}{(i+j+1)!} \right) R_{a1,a1}(0).$$

where we denoted by $[\tilde{N}^{-1}]_{ai,bj}$ the (ai, bj) entry of the matrix \tilde{N}^{-1} . The proof of the statement is then reduced to the following lemma, whose proof is postponed in Appendix B.

LEMMA 8.4. — *Let \hat{N} and \hat{G} be $n \times n$ matrices, whose elements are $\hat{N}_{ij} = \frac{(-1)^{j-1}}{(i+j-1)!}$ and $\hat{G}_{ij} = \frac{(-1)^{j-1}}{(i+j+1)!}$. Then*

$$(8.7) \quad \text{tr} \left(\hat{N}^{-1}\hat{G} \right) = \frac{1}{2} \frac{n}{4n^2 - 1}. \quad \square$$

Appendix A. Proof of Lemma 8.3

We compute the value of the leading constant $C_\lambda := \det \tilde{N}$. Recall that \tilde{N} is a block matrix, whose only non-vanishing blocks are the diagonal ones. Moreover, every aa -block of the diagonal is the matrix \hat{N} of dimension n_a . Thus, to find the determinant of \tilde{N} , it is sufficient to evaluate the determinant of the generic matrix \hat{N} of dimension n defined by

$$\hat{N}_{ij} = \frac{(-1)^{j-1}}{(i+j-1)!}$$

The matrix \hat{N} has already been studied in [4], Section 7.3 and Appendix G, and its inverse can be expressed as a product of two matrices $\hat{N}_{ij}^{-1} = \left(\hat{S}^{-1}\hat{A}^{-1} \right)_{ij}$, where

$$(A.1) \quad \hat{A}_{ij}^{-1} := \frac{(-1)^{i-j}}{(i-j)!} \quad i \geq j,$$

$$(A.2) \quad \hat{S}_{ij}^{-1} := \frac{1}{i+j-1} \binom{n+i-1}{i-1} \binom{n+j-1}{j-1} \frac{(n!)^2}{(n-i)!(n-j)!}.$$

Therefore the inverse of \hat{N} is

$$(A.3) \quad \hat{N}_{ij}^{-1} = \sum_{k=j}^n \frac{(-1)^{k-j}}{(i+k-1)(k-j)!} \binom{n+i-1}{i-1} \binom{n+k-1}{k-1} \frac{(n!)^2}{(n-i)!(n-k)!}.$$

By Cramer’s rule one obtains

$$(A.4) \quad \hat{N}_{ij}^{-1} = (-1)^{i+j} \frac{\det \hat{N}_{ji}^0}{\det \hat{N}},$$

where \widehat{N}_{ji}^0 is the matrix of dimension $n - 1$ obtained from \widehat{N} by removing the j -th row and the i -th column. Applying (A.4) for $i = j = n$ we reduce the computation of the determinant of the matrix \widehat{N} of dimension n as the product of the (n, n) -entry of the matrix \widehat{N}^{-1} and the determinant of the matrix \widehat{N} of dimension $n - 1$, namely we get the recursive formula:

$$\det \widehat{N}_{(n)} = \frac{\det \widehat{N}_{(n-1)}}{\widehat{N}_{nn}^{-1}} = \det \widehat{N}_{(n-1)} \frac{(n - 1)!^2}{(2n - 2)!(2n - 1)!},$$

where the last equality follows from equation (A.3). Using that, for $n = 1$, the determinant is equal to 1, we obtain the general formula

$$(A.5) \quad \det \widehat{N}_{(n)} = \frac{\prod_{j=0}^{n-1} j!}{\prod_{j=n}^{2n-1} j!}.$$

The value of the constant C_λ is then obtained as the product

$$(A.6) \quad C_\lambda = \prod_{a=1}^k \det \widehat{N}_{(n_a)} = \prod_{a=1}^k \frac{\prod_{j=0}^{n_a-1} j!}{\prod_{j=n_a}^{2n_a-1} j!}.$$

This concludes the proof of Lemma 8.3.

Appendix B. Proof of Lemma 8.4

The goal of this appendix is to prove Lemma 8.4, i.e., the identity

$$(B.1) \quad \text{tr} \left(\widehat{N}^{-1} \widehat{G} \right) = \sum_{i,j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji} = \frac{1}{2} \frac{n}{4n^2 - 1}.$$

where we recall that the matrices \widehat{N} and \widehat{G} are defined by

$$(B.2) \quad \widehat{N}_{ij} = \frac{(-1)^{j-1}}{(i + j - 1)!}, \quad \widehat{G}_{ij} = \frac{(-1)^{j-1}}{(i + j + 1)!}.$$

Using the formula (A.3) for the expression of \widehat{N}_{ij}^{-1} , we are reduced to prove the following combinatorial identity

$$(B.3) \quad \sum_{i=1}^n \sum_{j=1}^n \sum_{k=j}^n \frac{(-1)^{k-j}}{(i + k - 1)(k - j)!} \binom{n + i - 1}{i - 1} \binom{n + k - 1}{k - 1} \\ \times \frac{(n!)^2}{(n - i)!(n - k)!} \frac{(-1)^{i-1}}{(i + j + 1)!} = \frac{1}{2} \frac{n}{4n^2 - 1}.$$

It is immediate to check that for $n = 1$ the identity (B.3) is true. Then in what follows we assume that $n \geq 2$.

Notice that for $i = 1, \dots, n - 2$, we have $\widehat{G}_{ji} = \widehat{N}_{j(i+2)}$ therefore this sum reduces to the sum of the components with $i = n - 1$ and $i = n$:

$$\begin{aligned} \sum_{i,j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji} &= \sum_{i=1}^{n-2} \sum_{j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji} + \sum_{i=n-1}^n \sum_{j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji} \\ &= \sum_{i=1}^{n-2} \sum_{j=1}^n \widehat{N}_{ij}^{-1} \widehat{N}_{j(i+2)} + \sum_{i=n-1}^n \sum_{j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji} \\ &= \sum_{i=1}^{n-2} \delta_{i(i+2)} + \sum_{i=n-1}^n \sum_{j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji} \\ &= \sum_{i=n-1}^n \sum_{j=1}^n \widehat{N}_{ij}^{-1} \widehat{G}_{ji}, \end{aligned}$$

where δ_{ij} is the Kronecker symbol. In particular, our initial claim (8.7) follows by summing the next two combinatorial identities, that are valid for $n \geq 2$:

$$\begin{aligned} \text{(B.4)} \quad \sum_{j=1}^n \sum_{k=j}^n \frac{(-1)^{k-j+n}}{(n+k-2)(k-j)!} \binom{2n-2}{n-2} \binom{n+k-1}{k-1} \\ \times \frac{(n!)^2}{(n-k)!(n+j)!} = -\frac{n-1}{4(2n-1)}, \end{aligned}$$

$$\begin{aligned} \text{(B.5)} \quad \sum_{j=1}^n \sum_{k=j}^n \frac{(-1)^{k-j+n+1}}{(n+k-1)(k-j)!} \binom{2n-1}{n-1} \binom{n+k-1}{k-1} \\ \times \frac{(n!)^2}{(n-k)!(n+j+1)!} = \frac{n+1}{4(2n+1)}. \end{aligned}$$

Before proving (B.4) and (B.5), let us first simplify them. Using the following binomial identity, for $n \geq 2$ and $1 \leq j \leq k \leq n$

$$\frac{1}{(k-j)!} \binom{2n-2}{n-2} \frac{(n!)^2}{(n-k)!(n+j)!} \frac{2(2n-1)}{n-1} = \binom{2n}{n+k} \binom{n+k}{n+j},$$

one obtains that (B.4) is equivalent to

$$\text{(B.6)} \quad \sum_{j=1}^n \sum_{k=j}^n \frac{(-1)^{n+k+j}}{n+k-2} \binom{2n}{n+k} \binom{n+k}{n+j} \binom{n+k-1}{k-1} = -\frac{1}{2}.$$

LEMMA B.1. — For $k, n \geq 1$, one has the following combinatorial identity

$$(B.7) \quad \sum_{j=1}^k (-1)^j \binom{n+k}{n+j} = -\binom{n+k-1}{k-1}.$$

Proof of Lemma B.1. — We have the following chain of identities

$$(B.8) \quad \begin{aligned} \sum_{j=1}^k (-1)^j \binom{n+k}{n+j} &= (-1)^n \sum_{j=n+1}^{n+k} (-1)^j \binom{n+k}{j} \\ &= -(-1)^n \sum_{j=0}^n (-1)^j \binom{n+k}{j} \\ &= -\binom{n+k-1}{n} = -\binom{n+k-1}{k-1}, \end{aligned}$$

where in the first one we used a change of variable $j \rightarrow j+n$ in the sum, while in the second one we used the general identity $0 = (-1+1)^N = \sum_{j=0}^N (-1)^j \binom{N}{j}$. The last equality follows from the identity

$$\sum_{j=0}^n (-1)^j \binom{n+k}{j} = (-1)^n \binom{n+k-1}{n},$$

that can be easily proved for every fixed k , by induction on $n \geq 1$. □

Applying Lemma B.1 to the left hand side of (B.6), we have

$$\begin{aligned} &\sum_{j=1}^n \sum_{k=j}^n \frac{(-1)^{n+k+j}}{n+k-2} \binom{2n}{n+k} \binom{n+k}{n+j} \binom{n+k-1}{k-1} \\ &= \sum_{k=1}^n \left[\sum_{j=1}^k (-1)^j \binom{n+k}{n+j} \right] \frac{(-1)^{n+k}}{n+k-2} \binom{2n}{n+k} \binom{n+k-1}{k-1} \\ &= -\sum_{k=1}^n \frac{(-1)^{n+k}}{n+k-2} \binom{2n}{n-k} \binom{n+k-1}{k-1}^2. \end{aligned}$$

Thus we have proved that (B.6), which is equivalent to (B.4), is also equivalent to

$$(B.9) \quad \sum_{k=1}^n \frac{(-1)^{n+k}}{n+k-2} \binom{2n}{n-k} \binom{n+k-1}{k-1}^2 = \frac{1}{2}.$$

Performing analogous transformations, one proves that (B.5) is equivalent to

$$(B.10) \quad \sum_{k=1}^n \frac{(-1)^{n+k}}{(n+k)(n+k-1)} \binom{2n+1}{n-k} \binom{n+k}{k-1}^2 = \frac{1}{2}.$$

The proof is then completed thanks to the next lemma.

LEMMA B.2. — *For $n \geq 2$, the following combinatorial identities hold*

$$(B.11) \quad \sum_{k=1}^n \frac{(-1)^{n+k}}{n+k-2} \binom{2n}{n-k} \binom{n+k-1}{k-1}^2 = \frac{1}{2},$$

$$(B.12) \quad \sum_{k=1}^n \frac{(-1)^{n+k}}{(n+k)(n+k-1)} \binom{2n+1}{n-k} \binom{n+k}{k-1}^2 = \frac{1}{2}.$$

Proof of Lemma B.2. — Let us first prove (B.11). Denote by

$$\beta_{k,n} := \frac{(-1)^{n+k}}{n+k-2} \binom{2n}{n-k} \binom{n+k-1}{k-1}^2,$$

the coefficient appearing in the sum (B.11), we want to prove $\sum_{k=1}^n \beta_{k,n} = \frac{1}{2}$. To this aim, we apply Lemma B.3 to two different matrices.

Let us first consider the Hilbert matrix $H_1 := [\frac{1}{i+j-1}]_{ij}$. It is a matrix of type (B.17), with $a_i := i$ and $b_j := -j + 1$. We compute the coefficients of the $(n-1)$ -th row of H_1^{-1} :

$$(B.13) \quad \begin{aligned} (H_1^{-1})_{n-1,j} &= \frac{1}{-n+2-j} \frac{\prod_k (-n+2-k)(j+k-1)}{\prod_{k \neq j} (j-k) \prod_{l \neq n-1} (-n+2+l-1)} \\ &= \frac{n(n-1)^2}{2(2n-1)} (n+j) \frac{(-1)^{n+j+1}}{n+j-2} \binom{2n}{n-j} \binom{n+j-1}{j-1}^2 \\ &= -\frac{n(n-1)^2}{2(2n-1)} (n+j) \beta_{j,n} \end{aligned}$$

Then consider $H_2 := [\frac{1}{a_i - b_j}]$, with $a_i = i$ for $i < n$ and $a_n = -n$, while $b_j = -j + 1$. We compute the coefficients of the $(n-1)$ -th row of H_2^{-1} . For $j < n$ we have

$$(B.14) \quad \begin{aligned} (H_2^{-1})_{n-1,j} &= \frac{1}{-n+2-j} \frac{\prod_{k \neq n} (-n+2-k)(-n+2+n) \prod_k (j+k-1)}{\prod_{k \neq j, k \neq n} (j-k) (n+j) \prod_{l \neq n-1} (-n+2+l-1)} \\ &= \frac{n(n-1)}{2(2n-1)} (n-j) \frac{(-1)^{n+j+1}}{n+j-2} \binom{2n}{n-j} \binom{n+j-1}{j-1}^2 \\ &= -\frac{n(n-1)}{2(2n-1)} (n-j) \beta_{j,n}, \end{aligned}$$

while for $j = n$ we get

$$\begin{aligned}
 & (H_2^{-1})_{n-1,n} \\
 \text{(B.15)} \quad &= \frac{1}{-n+2+n} \frac{\prod_{k \neq n} (-n+2-k)(-n+2+n) \prod_k (-n+k-1)}{\prod_{k \neq n} (-n-k)(n+j) \prod_{l \neq n-1} (-n+2+l-1)} \\
 &= \frac{n^2(n-1)}{2(2n-1)}.
 \end{aligned}$$

Setting

$$\alpha_1 := \sum_{j=1}^n (H_1^{-1})_{n-1,j}, \quad \alpha_2 := \sum_{j=1}^n (H_2^{-1})_{n-1,j}$$

and summing over j the identities (B.13) and (B.14) one gets

$$\sum_{j=1}^n \beta_{j,n} = -\frac{1}{2n} \left[\frac{2(2n-1)}{n(n-1)^2} \alpha_1 + \frac{2(2n-1)}{n(n-1)} \left(\alpha_2 - \frac{n^2(n-1)}{2(2n-1)} \right) \right].$$

Now the proof of equation (B.11) is completed once we use formula (B.18) to find the values

$$\alpha_1 = -\frac{(2n-2)!}{(n-2)!^2}, \quad \alpha_2 = \frac{2(2n-3)!}{(n-2)!^2}.$$

Equation (B.12) can be proved along the same lines. More precisely, define

$$\gamma_{k,n} := \frac{(-1)^{n+k}}{(n+k)(n+k-1)} \binom{2n+1}{n-k} \binom{n+k}{k-1}^2$$

as the coefficients appearing in the sum (B.12), and consider the Hilbert matrix H_1 , and the matrix H_3 obtained by $a_j = j$ if $j < n$ and $a_n = -n-1$, and $b_j = -j+1$. Then by Lemma B.3

$$(H_1^{-1})_{n,j} = (n+j+1) \frac{n(n+1)^2}{2(2n+1)} \gamma_{j,n}.$$

Moreover, for $j < n$

$$(H_3^{-1})_{n,j} = (n-j) \frac{n(n+1)^2}{(2n+1)(2n-1)} \gamma_{j,n},$$

while for $j = n$

$$(H_3^{-1})_{n,n} = -\frac{n(n+1)^2}{2(2n-1)}.$$

One can also compute

$$\eta_1 := \sum_{j=1}^n (H_1^{-1})_{n,j} = \frac{(2n-1)!}{(n-1)!^2}, \quad \eta_3 := \sum_{j=1}^n (H_3^{-1})_{n,j} = -\frac{2(2n-2)!}{(n-1)!^2}.$$

Then the sum in (B.12) is given by

$$(B.16) \quad \sum_{j=1}^n \gamma_{j,n} = \frac{1}{2n+1} \left[\frac{2(2n+1)}{n(n+1)^2} \eta_1 + \frac{(2n+1)(2n-1)}{n(n+1)^2} \left(\eta_3 + \frac{n(n+1)^2}{2(2n-1)} \right) \right] = \frac{1}{2}.$$

that completes the proof of the lemma. □

The following lemma concerns the inverse of the generalized Hilbert matrix.

LEMMA B.3 (see [20]). — *Let $a_1, \dots, a_n, b_1, \dots, b_n$ be $2n$ distinct reals and define the $n \times n$ matrix*

$$(B.17) \quad H_{ij} = \frac{1}{a_i - b_j}.$$

Then we have

$$(B.18) \quad (H^{-1})_{ij} = \frac{1}{b_i - a_j} \frac{\prod_{k=1}^n (b_i - a_k)(a_j - b_k)}{\prod_{k \neq j} (a_j - a_k) \prod_{l \neq i} (b_i - b_l)},$$

$$\sum_{j=1}^n (H^{-1})_{ij} = - \frac{\prod_{k=1}^n (b_i - a_k)}{\prod_{k \neq i} (b_i - b_k)}.$$

BIBLIOGRAPHY

- [1] A. AGRACHEV, D. BARILARI & U. BOSCAIN, “On the Hausdorff volume in sub-Riemannian geometry”, *Calc. Var. Partial Differ. Equ.* **43** (2012), no. 3-4, p. 355-388.
- [2] ———, *A Comprehensive Introduction to Sub-Riemannian Geometry*, Monograph, Cambridge University Press, 2019.
- [3] A. AGRACHEV, D. BARILARI & L. RIZZI, “Sub-Riemannian curvature in contact geometry”, *J. Geom. Anal.* **27** (2017), no. 1, p. 366-408.
- [4] ———, “Curvature: a variational approach”, *Mem. Am. Math. Soc.* **1225** (2018), no. iii-vi, p. 146.
- [5] A. AGRACHEV & Y. L. SACHKOV, *Control Theory from the Geometric Viewpoint*, Encyclopaedia of Mathematical Sciences, vol. 87, Springer, 2004, xiv+412 pages.
- [6] D. BARILARI, “Trace heat kernel asymptotics in 3D contact sub-Riemannian geometry”, *J. Math. Sci., New York* **195** (2013), no. 3, p. 391-411.
- [7] D. BARILARI & F. BOAROTTO, “Kolmogorov-Fokker-Planck operators in dimension two: heat kernel and curvature”, *J. Evol. Equ.* **18** (2018), no. 3, p. 1115-1146.
- [8] D. BARILARI & E. PAOLI, “Curvature terms in small time heat kernel expansion for a model class of hypoelliptic Hörmander operators”, *Nonlinear Anal., Theory Methods Appl.* **164** (2017), p. 118-134.
- [9] D. BARILARI & L. RIZZI, “A formula for Popp’s volume in sub-Riemannian geometry”, *Anal. Geom. Metr. Spaces* **1** (2013), p. 42-57.
- [10] ———, “Comparison theorems for conjugate points in sub-Riemannian geometry”, *ESAIM, Control Optim. Calc. Var.* **22** (2016), no. 2, p. 439-472.

- [11] ———, “On Jacobi fields and a canonical connection in sub-Riemannian geometry”, *Arch. Math., Brno* **53** (2017), no. 2, p. 77-92.
- [12] J.-M. BISMUT, *Large deviations and the Malliavin calculus*, Progress in Mathematics, vol. 45, Birkhäuser, 1984, viii+216 pages.
- [13] S. GALLOT, D. HULIN & J. LAFONTAINE, *Riemannian geometry*, third ed., Universitext, Springer, 2004, xvi+322 pages.
- [14] B. JAKUBCZYK, “Introduction to geometric nonlinear control; controllability and Lie bracket”, in *Mathematical control theory (Trieste, 2001)*, ICTP Lecture Notes, vol. 8, The Abdus Salam International Centre for Theoretical Physics, 2002, p. 107-168.
- [15] V. JURDJEVIC, *Geometric control theory*, Cambridge Studies in Advanced Mathematics, vol. 52, Cambridge University Press, 1997, xviii+492 pages.
- [16] R. MONTGOMERY, “Abnormal Minimizers”, *SIAM J. Control Optimization* **32** (1994), no. 6, p. 1605-1620.
- [17] ———, *A tour of subriemannian geometries, their geodesics and applications*, Mathematical Surveys and Monographs, vol. 91, American Mathematical Society, 2002, xx+259 pages.
- [18] E. PAOLI, “Small time asymptotics on the diagonal for Hörmander’s type hypoelliptic operators”, *J. Dyn. Control Syst.* **23** (2017), no. 1, p. 111-143.
- [19] L. S. PONTRYAGIN, V. G. BOLTYANSKIĬ, R. V. GAMKRELIDZE & E. F. MISHCHENKO, *Selected works. Vol. 4*, Classics of Soviet Mathematics, Gordon and Breach Science Publishers, 1986, xxiv+360 pages.
- [20] S. SCHECHTER, “On the Inversion of Certain Matrices”, *Math. Tables Aids Comput.* **13** (1959), no. 66, p. 73-77.
- [21] C. VILLANI, *Optimal transport. Old and new*, Grundlehren der Mathematischen Wissenschaften, vol. 338, Springer, 2009, xxii+973 pages.
- [22] I. ZELENKO & C. LI, “Differential geometry of curves in Lagrange Grassmannians with given Young diagram”, *Differ. Geom. Appl.* **27** (2009), no. 6, p. 723-742.

Manuscrit reçu le 18 octobre 2016,
révisé le 15 janvier 2018,
accepté le 13 mars 2018.

Andrei A. AGRACHEV
SISSA
Via Bonomea 265
Trieste (Italy)
and
Steklov Math. Inst.
Moscow (Russia)
agrachev@sissa.it

Davide BARILARI
IMJ-PRG, UMR CNRS 7586
Université Paris-Diderot
Batiment Sophie Germain, Case 7012
75205 Paris Cedex 13 (France)
davide.barilari@imj-prg.fr

Elisa PAOLI
SISSA
Via Bonomea 265
Trieste (Italy)
epaoli@sissa.it