

PUBLISHED VERSION

L. Graves, S. P. Garnett, C. T. Cowell, L. A. Baur, A. Ness, N. Sattar and D. A. Lawlor
Waist-to-height ratio and cardiometabolic risk factors in adolescence: findings from a prospective birth cohort

Pediatric Obesity, 2014; 9(5):327-338


© 2013 The Authors. Pediatric Obesity published by John Wiley & Sons Ltd on behalf of International Association for the Study of Obesity. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Originally published at:

<http://doi.org/10.1111/j.2047-6310.2013.00192.x>

PERMISSIONS

<http://creativecommons.org/licenses/by/3.0/>



The image shows a Creative Commons Attribution 3.0 Unported (CC BY 3.0) license banner. It features the CC logo and a person icon in a circle. The text reads "Attribution 3.0 Unported (CC BY 3.0)". Below this, it states "This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#)." The banner is divided into two main sections: "You are free to:" and "Under the following terms:". Under "You are free to:", it lists "Share" (copy and redistribute) and "Adapt" (remix, transform, and build upon). A circular seal on the right says "Free Cultural Works APPROVED FOR". Under "Under the following terms:", it lists "Attribution" (give credit, link to license, indicate changes) and "No additional restrictions" (no legal terms or technological measures).

Attribution 3.0 Unported (CC BY 3.0)

This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#).

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.

1 November 2017

<http://hdl.handle.net/2440/107060>

Waist-to-height ratio and cardiometabolic risk factors in adolescence: findings from a prospective birth cohort

L. Graves^{1,2}, S. P. Garnett^{2,3,4}, C. T. Cowell^{2,3,4}, L. A. Baur², A. Ness⁵, N. Sattar⁶ and D. A. Lawlor⁷

¹Sydney Medical School, University of Sydney, Sydney, Australia; ²The Children's Hospital at Westmead Clinical School, University of Sydney, Sydney, Australia; ³Kids Research Institute, The Children's Hospital at Westmead, Westmead, Australia; ⁴Institute of Endocrinology and Diabetes, The Children's Hospital at Westmead, Westmead, Australia; ⁵School of Oral and Dental Sciences, University of Bristol, Bristol, UK; ⁶Metabolic Medicine, University of Glasgow, Glasgow, UK; ⁷MRC Centre for Causal Analyses in Translational Epidemiology, School of Social and Community Medicine, University of Bristol, Bristol, UK

Received 12 December 2012; revised 12 June 2013; accepted 20 June 2013

What is already known about this subject

- In adults, associations between body mass index (BMI), waist-to-height ratio (WHtR) and cardiometabolic outcomes are similar.
- In children and adolescents, results from cross-sectional studies examining the associations between BMI z scores, WHtR and cardiometabolic outcomes are conflicting and there is a paucity of prospective data.

What this study adds

- This is the first study to demonstrate the *prospective* association between WHtR in childhood and cardiometabolic outcomes in adolescent boys.
- WHtR is a simple calculation that can be used to identify children and adolescents for cardiometabolic risk without the need for reference growth charts.
- The WHtR cut-point of ≥ 0.5 was highly specific in identifying cardiometabolic risk co-occurrence but has poor sensitivity.

Summary

Objective: To examine the associations between body mass index (BMI) and waist-to-height ratio (WHtR) measured in childhood and adolescence and cardiometabolic risk factors in adolescence.

Methods: Secondary data analysis of the Avon Longitudinal Study of Parents and Children, a population based cohort. Data from 2858 adolescents aged 15.5 (standard deviation 0.4) years and 2710 of these participants as children aged 7–9 years were used in this analysis. Outcome measures were cardiometabolic risk factors, including triglycerides, low density lipoprotein cholesterol, high density lipoprotein cholesterol, insulin, glucose and blood pressure at 15 years of age.

Results: Both BMI and WHtR measured at ages 7–9 years and at age 15 years were associated with cardiometabolic risk factors in adolescents. A WHtR ≥ 0.5 at 7–9 years increased the odds by 4.6 [95% confidence interval 2.6 to 8.1] for males and 1.6 [0.7 to 3.9] for females of having three or more cardiometabolic risk factors in adolescence. Cross-sectional analysis indicated that adolescents who had a WHtR ≥ 0.5 , the odds ratio of having three or more cardiometabolic risk factors was 6.8 [4.4 to 10.6] for males and 3.8 [2.3 to 6.3] for females. The WHtR cut-point was highly specific in identifying cardiometabolic risk co-occurrence in male children and adolescents as well as female children (90 to 95%), but had poor sensitivity (17 to 53%). Similar associations were observed when BMI was used to define excess adiposity.

Address for correspondence: A/Professor SP Garnett, Institute of Endocrinology and Diabetes, The Children's Hospital at Westmead, Locked Bag 4001, Westmead, NSW 2145, Australia. E-mail: sarah.garnett@health.nsw.gov.au

© 2013 The Authors. Pediatric Obesity published by John Wiley & Sons Ltd on behalf of International Association for the Study of Obesity. **Pediatric Obesity** 9, 327–338

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Conclusions: WHtR is a simple alternative to age and sex adjusted BMI for assessing cardiometabolic risk in adolescents.

Keywords: Waist-to-height ratio, cardiometabolic risk, ALSPAC longitudinal study.

Introduction

Obesity in childhood is associated with adverse levels of cardiometabolic risk factors, including higher blood pressure (BP), triglycerides, total and low density lipoprotein cholesterol (LDLc) and insulin, and lower high density lipoprotein cholesterol (HDLc) (1–3). Furthermore, childhood obesity is positively and linearly associated with obesity and related cardiovascular disease in adulthood (4). It is important to identify children who are at increased risk of developing comorbidities associated with obesity, to potentially intervene and prevent the development of chronic disease including type 2 diabetes.

We have previously reported, using data from the Avon Longitudinal Study of Parents and Children (ALSPAC), that childhood body mass index (BMI), waist circumference and total fat mass are positively associated with cardiovascular risk factors in adolescence and the magnitudes of these associations are similar for all measures of adiposity (2). There has been recent interest in the use of the waist-to-height ratio (WHtR) for identifying excessive central adiposity in children and adolescents (5–8). It has been suggested that a WHtR ≥ 0.5 , irrespective of age, sex or ethnicity (5,7,9), is a valid predictor of higher cardiometabolic risk (10–12). WHtR may be a more straightforward anthropometric index to apply in the clinical setting where BMI centile charts may not be readily available.

There have been two systematic reviews in adults examining the associations between BMI and WHtR and cardiometabolic outcomes which have indicated broadly similar associations between the different anthropometric measures and cardiometabolic risk factors (7,13). However, the utility of WHtR in identifying young people at cardiometabolic risk is unclear. Results from cross-sectional studies are conflicting some indicate that WHtR is more strongly associated with cardiometabolic risk factors than BMI (10,12,14–16), while others have found either a weaker (17) or a similar (18,19) association. To our knowledge, there is only one prospective study looking at this issue in children and adolescents. That study reported similar correlation between BMI and WHtR assessed in children when they were aged 7 to 15 years and the metabolic syndrome in young adulthood (20).

Therefore the aims of this study were to:

1. Examine the cross-sectional association between WHtR and cardiometabolic risk factors in adolescents.
2. Examine the prospective association between WHtR assessed in childhood and cardiometabolic risk factors assessed in adolescence.
3. Examine these associations in relevance to the respective associations with BMI.

Methods

Participants

This study is a secondary analysis of data from the ALSPAC (<http://www.bristol.ac.uk/alspac>). The design and methods have been published (21). In brief, ALSPAC is a longitudinal population-based cohort that recruited 14 541 pregnant women who were expected to deliver between 1 April 1991 and 31 December 1992 (21). All members of the ALSPAC cohort were invited to annual follow-up clinics from the age of 7 to 13 years and then approximately every 2 years thereafter. In addition, when the oldest children were approximately 7 years of age, the sample size was increased by inviting eligible cases who had failed to join the study originally resulting in an additional 713 children being enrolled; 8297, 7725 and 5509 participants attended the 7-, 9- and 15-year clinics, respectively. In the present study, baseline 'childhood' anthropometric data were taken from a combination of the 7- and 9-year clinics. If available, data from the 7-year clinic were used preferentially ($n = 2540$) and data from an additional 168 children were used from the 9-year clinic. Anthropometric and cardiometabolic risk factors were taken from the 15-year clinic ($n = 2858$): the 'adolescent' group. The eligibility criteria for these analyses were adolescents who had their waist circumference, height and serum lipid levels measured at the 15 year clinic. There were no childhood anthropometric measurements for 150 adolescents who had both anthropometric and cardiometabolic risk factors measured at the 15-year clinic; consequently, data are missing for 5% (148 of 2858) of the children. The final sample used in the analyses here were 2858 adolescents (cross-sectional analysis) and 2710 children (prospective analysis). The majority

(98%) of the mothers of the participants self-identified their ethnicity as white. Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

Anthropometry

Standard protocols for assessing anthropometry were used, with the participants in light clothing and no shoes. Age was recorded in months. Weight was measured to the nearest 0.1 kg using Tanita THF 300GS (Tanita UK Ltd, Yewsey, Middlesex, UK). Height was measured using a Harpenden stadiometer (Holtain Ltd, Crymych, Pembrokeshire, UK) to the nearest 1 mm. Waist circumference was measured at the mid-point between the lower rib and the iliac crest to the nearest 1 mm with a flexible tape measure. Height, weight and BMI z scores were determined using age and sex specific national reference values for the UK (22). WHtR was determined by dividing waist circumference by height.

Cardiometabolic risk factors

The participants were requested to fast before attending the 15-year clinic. For those attending morning clinics they were asked to fast overnight. For those attending afternoon clinics, they were asked to fast for a minimum of 6 h prior to attendance. Blood samples were taken from the cubital fossa and immediately spun and plasma was frozen at -80°C . Approximately 3 to 9 months later, the samples were assayed. Total cholesterol, triglyceride and HDLc concentrations were measured using a modified Lipid Research Clinics Protocol with enzymatic reagents for lipid determination (2). LDLc was determined with the Friedewald equation (23). An automated assay was used to measure blood glucose concentration. Insulin was measured using an enzyme-linked immunosorbent assay (Mercodia, Uppsala, Sweden) which does not cross react with proinsulin. BP was measured using a Dinamap 9301 Vital Signs Monitor (Morton Medical, London, UK) with appropriate cuff size. Each participant was at rest and his/her arm was supported. For analysis, the mean of the two measurements that were taken was used.

Statistical analysis

Data were analysed using IBM SPSS Statistics 19.0 (IBM, Chicago, IL, USA) and MedCalc version 12.5.0 (Ostend, Belgium). The data were examined cross-sectionally and prospectively and explored as continuous variables and as binary categorical variables.

Relationships between continuous variables were examined by Spearman correlation coefficients. χ^2 test was used as a measure of association between categorical variables and odds ratios were used to examine the strength of associations. The cut-points used in this analysis to indicate cardiometabolic risk were $\geq 1.7 \text{ mmol L}^{-1}$ for triglycerides, $< 1.03 \text{ mmol L}^{-1}$ for HDLc, $\geq 5.6 \text{ mmol L}^{-1}$ for plasma glucose, $\geq 130 \text{ mmHg}$ for systolic BP, and $\geq 85 \text{ mmHg}$ for diastolic BP, as recommended by the International Diabetes Federation for children and adolescents (10 to 16 years) (24). The cut-points for LDLc and insulin were $\geq 2.79 \text{ mmol L}^{-1}$ and $\geq 16.95 \text{ IU L}^{-1}$, respectively, which is ≥ 90 th centile for the cohort (2). Cardiometabolic risk factor co-occurrence was defined as having three or more cardiometabolic risk factors using the binary outcome thresholds listed above. Participants were classified as overweight or obese based on sex- and age- specific International Obesity Task Force (IOTF) BMI criteria (25) and ≥ 0.5 for WHtR (7). Receiver operator characteristic (ROC) curves were used to identify the optimal WHtR cut-points, sensitivity and specificity.

Results

The anthropometric characteristics of participants are shown in Table 1. The prevalence of overweight or obese was lower in childhood (13.8%; $n = 375$) than in adolescence (17.2%; $n = 490$). The correlation between BMI z scores in childhood and adolescence was $r = 0.72$, $P < 0.001$, and 63.6% of overweight and obese children remained overweight or obese at 15 years. The proportion of participants who had a high WHtR (≥ 0.5) was also lower in childhood (6.8%; $n = 185$) compared with adolescence (17.2%; $n = 492$). The correlation between WHtR in childhood and adolescence was $r = 0.57$, $P < 0.001$ and 69.2% of children with a high WHtR had a high WHtR as adolescents.

BMI z scores and WHtR were highly correlated in both childhood and adolescence: $r = 0.80$ and $r = 0.78$ (both $P < 0.001$), respectively. The majority (91.9%) of children with a high WHtR were also overweight or obese. Conversely, only 45.1% of overweight or obese children had a high WHtR. In adolescence 72.0% of adolescents with a high WHtR were overweight or obese, and 71.8% of overweight or obese adolescents had a high WHtR.

The prevalence of elevated cardiometabolic risk factors in the adolescents varied from 2.2% ($n = 62$) for high diastolic BP, to 29.0% ($n = 806$) with high systolic BP, Table 2. In general, elevated cardiometabolic risk factors were more common in

Table 1 Anthropometric characteristics in childhood and adolescence. Results are expressed as median and interquartile range unless otherwise indicated

Childhood	Male		Female		Total	
	<i>n</i>		<i>n</i>		<i>n</i>	
Age (months)	1317	89 [89, 90]	1393	89 [89, 90]	2710	89 [89, 90]
Height (cm)	1315	127.0 [123.1, 130.9]	1393	125.7 [122.0, 130.1]	2708	126.5 [122.5, 130.6]
Height z score	1315	0.38 [-0.36, 1.01]	1393	0.23 [-0.49, 0.90]	2708	0.28 [-0.42, 0.95]
Weight (kg)	1317	25.4 [23.2, 28.6]	1393	25.2 [22.8, 28.8]	2710	25.4 [23.0, 28.6]
Weight z score	1317	0.26 [-0.35, 0.94]	1393	0.15 [-0.45, 0.86]	2710	0.20 [-0.41, 0.90]
BMI	1315	15.77 [14.92, 16.88]	1393	15.94 [14.90, 17.40]	2708	15.87 [14.91, 17.14]
BMI z score	1315	0.08 [-0.54, 0.71]	1393	0.04 [-0.58, 0.76]	2708	0.06 [-0.57, 0.74]
Overweight and obese, <i>n</i> (%)	1315	149 (11.3)	1393	226 (16.2)	2708	375 (13.8)
WHtR*	1314	0.44 [0.43, 0.46]	1392	0.44 [0.42, 0.46]	2706	0.44 [0.42, 0.46]
WHtR* ≥ 0.5 , <i>n</i> (%)	1314	82 (6.2)	1392	103 (7.4)	2706	185 (6.8)
Adolescence	Male		Female		Total	
	<i>n</i>		<i>n</i>		<i>n</i>	
Age (months)	1376	184 [183, 186]	1482	184 [183, 187]	2858	184 [183, 187]
Height (cm)	1376	174.9 [170.0, 180.0]	1482	164.6 [161, 168.7]	2858	169.0 [163.5, 175.5]
Height z score	1376	0.47 [-0.16, 1.10]	1482	0.29 [-0.32, 0.96]	2858	0.37 [-0.22, 1.03]
Weight (kg)	1375	62.6 [56.8, 69.7]	1477	57.4 [52.2, 63.8]	2852	60.1 [53.9, 66.8]
Weight z score	1375	0.46 [-0.10, 1.03]	1477	0.36 [-0.29, 1.04]	2852	0.42 [-0.19, 1.03]
BMI	1375	20.38 [8.90, 22.24]	1477	21.10 [19.41, 23.23]	2852	20.71 [19.12, 22.80]
BMI z score	1375	0.32 [-0.30, 0.98]	1477	0.34 [-0.32, 1.01]	2852	0.33 [-0.31, 0.99]
Overweight or obese, <i>n</i> (%)		223 (16.2)		267 (18.1)		490 (17.2)
WHtR*	1376	0.43 [0.41, 0.46]	1482	0.46 [0.43, 0.49]	2858	0.44 [0.41, 0.48]
WHtR* ≥ 0.5 , <i>n</i> (%)		152 (11.0)		340 (22.9)		492 (17.2)

*Waist-to-height ratio.

BMI, body mass index; WHtR, waist-to-height ratio.

males than females, with approximately twice as many males having low HDLc, high glucose levels or high systolic BP compared to females. Co-occurrence of three or more cardiometabolic risk factors was more common in males than females. Correlations were stronger between cardiometabolic risk factors and anthropometric outcomes measured in adolescence compared to those measured in childhood, Table 3.

The proportions of overweight and obese adolescents, or adolescents with a WHtR ≥ 0.5 , that had elevated LDLc, reduced HDLc, elevated glucose, elevated insulin and/or high BP are shown in Table 4. The odds ratios between measures of adiposity and risk factors are shown in Fig. 1. Adolescent males with WHtR ≥ 0.5 had an increased odds of elevated triglycerides, LDLc, glucose, insulin and systolic BP and low HDLc compared to males with a WHtR < 0.5 . Furthermore, males with WHtR ≥ 0.5 had an increased odds ratio of co-occurrence of cardiometabolic risk factors compared to males with a WHtR < 0.5 (odds ratio = 6.8 [95% confidence

interval {CI} 4.4 to 10.6]). Overweight and obese males had correspondingly increased odds of a similar magnitude, compared with males who were not overweight or obese. Females with a high WHtR had increased odds ratio for elevated triglycerides, LDLc, insulin and systolic BP and low HDLc, and were approximately four times more likely (3.8 [2.3 to 6.3]) to have co-occurrence of cardiometabolic risk factors than females with a WHtR < 0.5 . Overweight and obese females had correspondingly increased odds of the same cardiometabolic risk factors and co-occurrence of these, (Fig. 1).

The specificity and sensitivity of a WHtR cut-point of 0.5 in male adolescents for identifying cardiometabolic risk factor co-occurrence was 90.0 (95% CI 88.2 to 91.6) and 41.3 (31.8 to 51.4), respectively. For female adolescents the specificity and sensitivity of a WHtR cut-point of 0.5 was 76.2 (73.7 to 78.2) and 53.1 (40.2 to 65.7). Using ROC analysis the optimal WHtR cut point for identifying co-occurrence of risk factors was 0.47 (specificity 83.5 [81.3 to 85.5]; sensitivity 51.9 [41.9 to 61.8]) for

Table 2 Biochemistry and blood pressure in adolescence. Data are expressed as median [interquartile range] and the number (%) above cut-points, or below cut-point for HDLc

	Male		Female		Total		P†
	n		n		n		
Triglycerides (mmol L ⁻¹)	1376	0.73 [0.57, 0.96]	1482	0.77 [0.62, 0.99]	2858	0.75 [0.59, 0.98]	
≥1.7 mmol L ⁻¹		43 (3.1)		43 (2.9)		86 (3.0)	0.727
LDLc (mmol L ⁻¹)	1376	1.94 [1.62, 2.30]	1482	2.12 [1.79, 2.52]	2858	2.03 [1.70, 2.41]	
≥2.79 mmol L ⁻¹		104 (7.6)		221 (14.9)		325 (11.4)	<0.001
HDLc (mmol L ⁻¹)	1376	1.20 [1.01, 1.38]	1482	1.33 [1.16, 1.54]	2858	1.27 [1.09, 1.47]	
<1.03 mmol L ⁻¹		347 (25.2)		178 (12.0)		525 (18.4)	<0.001
Glucose (mmol L ⁻¹)	1376	5.3 [5.0, 5.5]	1482	5.1 [4.9, 5.3]	2858	5.2 [5.0, 5.4]	
≥5.6 mmol L ⁻¹		313 (22.7)		159 (10.7)		472 (16.5)	0.004
Insulin (IU L ⁻¹)	1374	8.17 [6.00, 11.16]	1480	9.65 [7.25, 12.76]	2854	8.97 [6.63, 12.05]	
≥16.95 IU L ⁻¹		102 (7.4)		155 (10.5)		257 (9.0)	<0.001
Systolic BP (mmHg)	1335	126.5 [119.0, 133.5]	1444	120.5 [113.0, 127.5]	2779	123.5 [115.5, 131.0]	
≥130 mmHg		517 (38.7)		289 (20.0)		806 (29.0)	<0.001
Diastolic BP (mmHg)	1335	66.5 [60.5, 73.5]	1444	66.0 [60.5, 71.0]	2779	66.0 [60.5, 72.0]	
≥85 mmHg		40 (3.0)		22 (1.5)		62 (2.2)	0.009
Risk co-occurrence*		104 (7.6)		64 (4.3)		168 (5.9)	<0.001

*Risk co-occurrence is the presence of ≥3 of the following: triglycerides ≥1.7 mmol L⁻¹, LDLc ≥2.79 mmol L⁻¹, HDLc <1.03 mmol L⁻¹, plasma glucose ≥5.6 mmol L⁻¹, insulin ≥16.95 IU L⁻¹, systolic blood pressure ≥130 mmHg, or diastolic blood pressure ≥85 mmHg.

†Pearson χ^2 .

BP, blood pressure; HDLc, high density lipoprotein cholesterol; LDLc, low density lipoprotein cholesterol.

male adolescents and 0.48 (specificity 65.8 [63.3 to 68.3]; sensitivity 67.2 [54.3 to 78.4] for female adolescents.

The proportions of overweight and obese children or those with a WHtR ≥0.5 who had elevated LDLc, reduced HDLc, elevated glucose, elevated insulin and/or high BP during adolescence are shown in Table 5. Like the cross-sectional analysis, male children who had a WHtR ≥ 0.05 or who were overweight or obese had increased odds of cardiometabolic risk factors in adolescence, Fig. 2. However, the CIs around the odds ratio tended to be wider. Male children with WHtR ≥0.5 had approximately five times higher probability (4.6 [2.6 to 8.1]) of co-occurrence of risk factors during adolescence than male children with a WHtR <0.5. Overweight and obese male children had approximately four times higher probability (3.6 [2.2 to 5.8]) of co-occurrence of risk factors during adolescence. Similar associations between both anthropometric measures and cardiometabolic risk factors were also observed in female children, although the odds ratios were generally small, and the lower limits of the CIs below one, Fig. 2.

The specificity and sensitivity of a WHtR cut-point of 0.5 in male children for identifying cardiometabolic risk factor co-occurrence in adolescents was 94.7 (93.3 to 95.9) and 21.1 (13.4 to 30.6), respectively. For

female children the specificity and sensitivity of a WHtR cut-point of 0.5 was 91.4 (89.8 to 92.9) and 17.0 (8.1 to 29.8). Using ROC analysis the optimal WHtR cut point for identifying co-occurrence of risk factors in male children was the same as it was in male adolescents 0.47 (specificity 83.3 [81.0 to 85.3]; sensitivity 37.9 [28.1 to 48.4]), but lower in female children compared to female adolescents 0.44 (specificity 54.4 [51.7 to 57.1]; sensitivity 71.7 [57.7 to 83.2]).

Discussion

To our knowledge this is the first study to demonstrate a prospective association between having a high WHtR in childhood and cardiometabolic risk co-occurrence in adolescent boys. The results demonstrate that the of risk for having a WHtR ≥0.5 was comparable to being overweight or obese as defined by sex- and age- specific IOTF BMI criteria. For example, children with a WHtR ≥0.5, had two to five times higher odds of cardiometabolic risk co-occurrence compared to children who had a WHtR <0.5 and children who were overweight or obese had two to four times higher odds compared to normal weight children. However, there was a sex difference and the association between anthropometric measures in childhood and

Table 3 Spearman correlations (*P* value) between BMI z score and waist-to-height ratio measured in childhood and adolescence and cardiometabolic risk factors measured in adolescence

Cardiometabolic risk factors	Childhood				Adolescence			
	BMI z score		WHtR		BMI z score		WHtR	
	Male	Female	Male	Female	Male	Female	Male	Female
Triglycerides (mmol L ⁻¹)	0.083 (0.003)	0.052 (0.055)	0.077 (0.005)	0.079 (0.003)	0.270 (<0.001)	0.145 (<0.001)	0.321 (<0.001)	0.185 (<0.001)
LDLc (mmol L ⁻¹)	0.073 (0.008)	0.102 (<0.001)	0.096 (<0.001)	0.120 (<0.001)	0.119 (<0.001)	0.158 (<0.001)	0.148 (<0.001)	0.161 (<0.001)
HDLc (mmol L ⁻¹)	-0.137 (<0.001)	-0.101 (<0.001)	-0.143 (<0.001)	-0.096 (<0.001)	-0.271 (<0.001)	-0.242 (<0.001)	-0.211 (<0.001)	-0.260 (<0.001)
Glucose (mmol L ⁻¹)	0.001 (0.964)	-0.015 (0.570)	0.012 (0.661)	-0.014 (0.597)	0.077 (0.004)	-0.004 (0.888)	0.092 (0.001)	-0.116 (0.542)
Insulin (μU L ⁻¹)	0.119 (<0.001)	0.111 (<0.001)	0.099 (<0.001)	0.105 (<0.001)	0.284 (<0.001)	0.192 (<0.001)	0.352 (<0.001)	0.189 (<0.001)
Systolic BP (mmHg)	0.129 (<0.001)	0.139 (<0.001)	0.060 (0.033)	0.103 (<0.001)	0.228 (<0.001)	0.228 (<0.001)	0.148 (<0.001)	0.183 (<0.001)
Diastolic BP (mmHg)	0.025 (0.368)	0.003 (0.912)	-0.044 (0.119)	-0.063 (0.021)	0.031 (0.257)	0.021 (0.417)	-0.022 (0.429)	0.015 (0.576)

BMI, body mass index; BP, blood pressure; HDLc, high density lipoprotein cholesterol; LDLc, low density lipoprotein cholesterol; WHtR, waist-to-height ratio.

cardiometabolic risk co-occurrence in adolescence was significantly stronger in boys than girls. In the clinical environment, using a WHtR cut-point of 0.5 has several advantages over defining a child as overweight or obese by BMI; it is a simpler index to calculate, easily understood by adolescents and families and does not require sex- and age- specific centiles.

The findings of this study are broadly consistent with another prospective study that examined correlations between BMI and WHtR in Australian children and adult cardiometabolic risk factors (20). While direct comparisons between studies is difficult, it is interesting to note that the correlations between WHtR and BMI measured at 7 to 11 years with triglycerides ($\rho = 0.07$ and 0.05 for WHtR and BMI, respectively) and insulin ($\rho = 0.11$ for both WHtR and BMI) measured in adulthood were of a similar magnitude to what we report between WHtR and BMI measured at 7 to 9 years and triglycerides and insulin measured in adolescence.

The association between WHtR and cardiometabolic risk, both measured in adolescence, support a number of other cross-sectional studies (26,27), which have been recently reviewed (7). In general, all studies showed good agreement in the magnitude and outcome of their analysis between WHtR and BMI and outcomes of cardiometabolic risk. Consistent with our study, both WHtR and BMI had a stronger association with triglycerides and HDLc, than LDLc. The association between anthropometric measures and BP tended to be stronger for systolic BP compared to diastolic BP and both WHtR and BMI were associated with fasting insulin and a 'cardiometabolic risk score', albeit defined differently among the different studies.

There is no consensus on the cross-sectional association of WHtR and BMI and fasting glucose (3,15,17). In part, this may be explained by the differing age of the cohorts (range from 4 to 17 years) and methods of analysing the data. Findings from our study suggest that adolescent males with a high WHtR or who are overweight or obese have an increased risk of an elevated fasting glucose. However, it is of note that in our study a high proportion of adolescents (23% of males; 11% of females) had fasting blood glucose levels ≥ 5.6 mmol L⁻¹. In the absence of evidence to indicate that this population has disturbances in glucose metabolism, these findings could reflect the diurnal variation in blood glucose (28), since some adolescents had blood taken in the afternoon after a 6 h fast and/or inadequate fasting. The results pertaining

Table 4 The proportion of adolescents with cardiometabolic risk factors, stratified by sex and anthropometric status (cross sectional analysis)

Outcome and exposure	Male		Female		P [†]	
	% with increased adiposity* and risk factor/s	% without increased adiposity* and with risk factor/s	P ^b	% with increased adiposity* and risk factor/s		% without increased adiposity* and with risk factor/s
Triglycerides (≥1.7 mmol L ⁻¹)						
BMI	9.9	1.8	<0.001	5.6	2.2	0.003
WtHR	12.5	2.0	<0.001	2.9	2.0	0.001
LDLc (≥2.79 mmol L ⁻¹)						
BMI	12.1	6.6	0.004	20.2	13.7	0.007
WtHR	12.5	6.9	0.013	20.3	13.3	0.002
HDLc (<1.03 mmol L ⁻¹)						
BMI	43.5	21.7	<0.001	23.2	9.4	<0.001
WtHR	44.1	22.9	<0.001	21.2	9.3	<0.001
Glucose (≥5.6 mmol L ⁻¹)						
BMI	30.0	21.4	0.005	11.2	10.5	0.72
WtHR	34.2	21.3	<0.001	10.9	10.7	0.97
Insulin (≥16.95 IU L ⁻¹)						
BMI	22.9	4.4	<0.001	8.2	7.4	<0.001
WtHR	28.3	4.8	<0.001	18.9	8.0	<0.001
Systolic BP (≥130 mmHg)						
BMI	50.2	36.6	<0.001	26.5	18.6	0.003
WtHR	52.7	37.0	<0.001	25.6	18.3	0.004
Diastolic BP (≥85 mmHg)						
BMI	3.3	2.9	0.81	2.3	1.4	0.26
WtHR	2.7	3.0	0.85	2.1	1.3	0.31
Risk co-occurrence [‡]						
BMI	20.6	5.0	<0.001	9.7	3.1	<0.001
WtHR	27.0	5.1	<0.001	9.7	2.7	<0.001


*BMI was categorized as overweight/obese or not overweight/obese and WtHR was categorized as <0.5 or ≥0.5.

[†]Pearson χ^2 .

[‡]Risk co-occurrence is the presence of ≥3 of the following: triglycerides ≥1.7 mmol L⁻¹, LDLc ≥2.79 mmol L⁻¹, HDLc <1.03 mmol L⁻¹, plasma glucose ≥5.6 mmol L⁻¹, insulin ≥16.95 IU L⁻¹, systolic blood pressure ≥130 mmHg, or diastolic blood pressure ≥85 mmHg.

BMI, body mass index; BP, blood pressure; HDLc, high density lipoprotein cholesterol; LDLc, low density lipoprotein cholesterol; WtHR, waist-to-height ratio.

Figure 1 Odds ratio (95%CI) for risk factors associated with overweight and obesity defined by BMI (symbol: black triangle) and WHtR of ≥ 0.5 (symbol: white triangle) in adolescents (cross-sectional analysis). Risk factors were defined as follows: triglycerides ≥ 1.7 mmol/L, LDLc ≥ 2.79 mmol/L, HDLc < 1.03 mmol/L, plasma glucose ≥ 5.6 mmol/L, insulin ≥ 16.95 IU/L, systolic blood pressure ≥ 130 mmHg, or diastolic blood pressure ≥ 85 mmHg. ^bRisk co-occurrence was defined as ≥ 3 of the above.



to fasting glucose in this study need to be interpreted with caution.

More children were classified as overweight or obese using BMI criteria compared to those classified as having a high WHtR. Children with a WHtR ≥ 0.5 were more severely obese compared to children were defined as overweight or obese using BMI criteria, mean \pm standard deviation BMI z-score was 2.05 ± 0.9 and 1.82 ± 0.9 , respectively. The most appropriate cut-point for WHtR to identify children and adolescents at cardiometabolic risk is not clear. In the present study we used ≥ 0.5 which has demonstrated utility in adults (7) and children (10) but other cut-points have been suggested (8). Our results indicated that a cut-point ≥ 0.5 was highly specific in identifying cardiometabolic risk co-occurrence in male children and adolescents as well as female children (90 to 95%), but had poor sensitivity (17 to 53%); the risk of being wrongly

labelled as having co-occurrence of cardiometabolic risk factors was small, but many of those who were at risk were not identified. Statistically optimum cut-points were identified as 0.47 for male children and male adolescents and 0.44 and 0.48 for female children and female adolescents, respectively. However, by increasing the sensitivity, the specificity decreased particularly for female children (from 91 to 54%); using lower cut-points more children and adolescents would be incorrectly labelled as being as having co-occurrence of cardiometabolic risk factors. Given the stigma of being incorrectly labelled and the limited resources for managing children and adolescents at cardiometabolic risk it could be prudent to use a higher but more specific cut-point.

There are a number of limitations to the study including the number of participants which had complete data. Initially over 14 000 pregnant women

Table 5 The proportion of adolescents with cardiometabolic risk factors, stratified by sex and anthropometric status in childhood (prospective analysis)

Outcome and exposure	Male		Female	
	% with increased adiposity* and risk factor/s	P†	% with increased adiposity* and risk factor/s	P†
Triglycerides (≥1.7 mmol L ⁻¹)				
BMI	6.7	0.002	1.3	0.17
WtHR	11.0	<0.001	1.0	0.27
LDLc (≥2.79 mmol L ⁻¹)				
BMI	10.0	0.22	17.0	0.58
WtHR	11.0	0.22	16.5	0.58
HDLc (<1.03 mmol L ⁻¹)				
BMI	40.7	<0.001	16.6	0.02
WtHR	42.7	<0.001	16.5	0.12
Glucose (≥5.6 mmol L ⁻¹)				
BMI	24.7	0.50	10.4	0.92
WtHR	28.0	0.23	11.7	0.63
Insulin (≥16.95 IU L ⁻¹)				
BMI	16.1	<0.001	14.5	0.001
WtHR	19.8	<0.001	13.7	0.22
Systolic BP (≥130 mmHg)				
BMI	53.1	<0.001	27.5	0.006
WtHR	57.0	0.001	25.3	0.13
Diastolic BP (≥85 mmHg)				
BMI	3.4	0.71	1.8	0.65
WtHR	3.8	0.66	1.0	0.69
Risk co-occurrence‡				
BMI	18.1	<0.001	6.3	0.04
WtHR	24.4	<0.001	6.1	0.27


*BMI was categorized as overweight/obese or not overweight/obese and WtHR was categorized as <0.5 or ≥0.5.

†χ² (Pearson).

‡Risk co-occurrence is the presence of ≥3 of the following: triglycerides ≥1.7 mmol L⁻¹, LDLc ≥2.79 mmol L⁻¹, HDLc <1.03 mmol L⁻¹, plasma glucose ≥5.6 mmol L⁻¹, insulin ≥16.95 IU L⁻¹, systolic blood pressure ≥130 mmHg, or diastolic blood pressure ≥85 mmHg.

BMI, body mass index; BP, blood pressure; HDLc, high density lipoprotein cholesterol; LDLc, low density lipoprotein cholesterol; WtHR, waist-to-height ratio.

Figure 2 Odds ratio (95% confidence interval) for risk factors associated with overweight and obesity defined by body mass index (▲) and waist-to-height ratio of ≥ 0.5 (▽) in childhood (prospective analysis). Risk factors were defined as follows: triglycerides ≥ 1.7 mmol/L, low density lipoprotein cholesterol (LDLc) ≥ 2.79 mmol/L, high density lipoprotein cholesterol (HDLc) < 1.03 mmol/L, plasma glucose ≥ 5.6 mmol/L, insulin ≥ 16.95 IU/L, systolic blood pressure (BP). ^bRisk co-occurrence was defined as ≥ 3 of the above.



were recruited, with over 5500 participants attending the 15 years clinic, yet only 2710 met our inclusion criteria for our prospective analysis and 2858 for our cross-sectional analysis. The implications of attrition are not clear and may affect the generalizability of the findings. Nevertheless, this is the largest study in children and adolescents addressing the association between WHtR and cardiometabolic risk factors, as well as the first in the UK.

Another potential limitation relates to measuring waist circumference and applies to both the location of the measurement and sensitivity around measuring waist circumference in heavier children. There is no universally accepted method of measuring waist circumference. In this study the waist circumference was measured at the mid-point between the lowest rib and the iliac crest, which has been demonstrated to be reproducible and highly correlated with total

body and trunk adiposity (29). It is recognized that removal of clothing and placement of the measuring tape may be awkward or embarrassing for the overweight or obese children and needs to be handled sensitively.

In conclusion, we found that BMI z scores and WHtR were highly correlated and that they have similar associations with cross-sectional and prospective cardiometabolic risk factors in adolescence. WHtR is a simple calculation that could be used to identify male children and adolescents for cardiometabolic risk.

Conflicts of interest statement

The authors declare no conflict of interest related to this manuscript.

Acknowledgements and funding

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The United Kingdom Medical Research Council (Grant ref: 74882) the Wellcome Trust (Grant ref: 076467) and the University of Bristol provide core support for ALSPAC. SPG is funded by a Cancer Institute NSW Early Career Development Fellowship Grant (10/ECF/2-11).

The concept of the study was developed by SPG and CTC. NS managed laboratory assays. Data analysis and drafting the manuscript were primarily done by LG and SPG. CTC, LAB, AN, NS and DAL participated in data interpretation and preparation of the manuscript. We would like to thank Professor Jennifer Peat for statistical support and review of the manuscript.

References

1. Falaschetti E, Hingorani AD, Jones A, *et al.* Adiposity and cardiovascular risk factors in a large contemporary population of pre-pubertal children. *Eur Heart J* 2010; 31: 3063–3072.
2. Lawlor DA, Benfield L, Logue J, *et al.* Association between general and central adiposity in childhood, and change in these, with cardiovascular risk factors in adolescence: prospective cohort study. *BMJ* 2010; 341: c6224–c6224.
3. Botton J, Heude B, Kettaneh A, *et al.* Cardiovascular risk factor levels and their relationships with overweight and fat distribution in children: the Fleurbaix Laventie Ville Santé II study. *Metabolism* 2007; 56: 614–622.
4. Venn AJ, Thomson RJ, Schmidt MD, *et al.* Overweight and obesity from childhood to adulthood: a follow-up of participants in the 1985 Australian Schools Health and Fitness Survey. *Med J Aust* 2007; 186: 458–460.
5. Goulding A, Taylor RW, Grant AM, Parnell WR, Wilson NC, Williams SM. Waist-to-height ratios in relation to BMI z-scores in three ethnic groups from a representative sample of New Zealand children aged 5–14 years. *Int J Obes (Lond)* 2010; 34: 1188–1190.
6. Xiong F, Garnett SP, Cowell CT, *et al.* Waist circumference and waist-to-height ratio in Han Chinese children living in Chongqing, south-west China. *Public Health Nutr* 2011; 14: 20–26.
7. Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev* 2010; 23: 247–269.
8. Nambiar S, Hughes I, Davies PS. Developing waist-to-height ratio cut-offs to define overweight and obesity in children and adolescents. *Public Health Nutr* 2010; 13: 1566–1574.
9. Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 2005; 56: 303–307.
10. Garnett SP, Baur LA, Cowell CT. Waist-to-height ratio: a simple option for determining excess central adiposity in young people. *Int J Obes* 2008; 32: 1028–1030.
11. Taylor RW, Williams SM, Grant AM, Taylor BJ, Goulding A. Predictive ability of waist-to-height in relation to adiposity in children is not improved with age and sex-specific values. *Obesity* 2011; 19: 1062–1068.
12. McCarthy HD, Ashwell M. A study of central fatness using waist-to-height ratios in UK children and adolescents over two decades supports the simple message – ‘keep your waist circumference to less than half your height. *Int J Obes (Lond)* 2006; 30: 988–992.
13. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obes Rev* 2012; 13: 275–286.
14. Savva SC, Tornaritis M, Savva ME, *et al.* Waist circumference and waist-to-height ratio are better predictors of cardiovascular disease risk factors in children than body mass index. *Int J Obes Relat Metab Disord* 2000; 24: 1453–1458.
15. Kahn H, Imperatore G, Cheng Y. A population-based comparison of BMI percentiles and waist-to-height ratio for identifying cardiovascular risk in youth. *J Pediatrics* 2005; 146: 482–488.
16. Hara M, Saitou E, Iwata F, Okada T, Harada K. Waist-to-height ratio is the best predictor of cardiovascular disease risk factors in Japanese schoolchildren. *J Atheroscler Thromb* 2002; 9: 127–132.
17. Sung RYT, Yu CCW, Choi KC, *et al.* Waist circumference and body mass index in Chinese children: cutoff values for predicting cardiovascular risk factors. *Int J Obes (Lond)* 2006; 31: 550–558.
18. Freedman DS, Kahn HS, Mei Z, *et al.* Relation of body mass index and waist-to-height ratio to cardiovascular disease risk factors in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 2007; 86: 33–40.
19. Manios Y, Kourlaba G, Kafatos A, Cook TL, Spyridaki A, Fragiadakis GA. Associations of several anthropometric indices with insulin resistance in children: the Children Study. *Acta Paediatr* 2008; 97: 494–499.
20. Schmidt MD, Dwyer T, Magnussen CG, Venn AJ. Predictive associations between alternative measures of childhood adiposity and adult cardio-metabolic health. *Int J Obes (Lond)* 2011; 35: 38–45.
21. Boyd A, Golding J, Macleod J, *et al.* Cohort profile: the ‘Children of the 90s’ – the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 2013; 42: 111–127.

22. Pan H, Cole TJ. LMS Growth Program. 2010. [WWW document]. URL <http://www.healthforallchildren.com/?product=lmsgrowth> (accessed January 2011).
23. Friedewald W, Levy R, Fredrickson D. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972; 18: 499–502.
24. Zimmet P, Alberti G, Kaufman F, et al. The metabolic syndrome in children and adolescents. *Lancet* 2007; 369: 2059–2061.
25. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000; 320: 1240–1243.
26. Kondaki K, Grammatikaki E, Pavón DJ, et al. Comparison of several anthropometric indices with insulin resistance proxy measures among European adolescents: the Helena Study. *Eur J Pediatr* 2011; 170: 731–739.
27. Androutsos O, Grammatikaki E, Moschonis G, et al. Neck circumference: a useful screening tool of cardiovascular risk in children. *Pediatr Obes*. 2012; 7: 187–195.
28. Troisi RJ, Cowie CC, Harris MI. Diurnal variation in fasting plasma glucose: implications for diagnosis of diabetes in patients examined in the afternoon. *JAMA* 2000; 284: 3157–3159.
29. Wang J. Waist circumference: a simple, inexpensive, and reliable tool that should be included as part of physical examinations in the doctor's office. *Am J Clin Nutr* 2003; 78: 902–903.