

Warming of the Indian Ocean threatens eastern and southern African food security but could be mitigated by agricultural development

Chris Funk^{*†}, Michael D. Dettinger[‡], Joel C. Michaelsen^{*}, James P. Verdin[§], Molly E. Brown[¶], Mathew Barlow^{||}, and Andrew Hoell^{||}

^{*}United States Geological Survey, Geography Department, University of California, Santa Barbara, CA 93106-1120; [†]United States Geological Survey, Scripps Institution of Oceanography, La Jolla, CA 92093-0224; [‡]United States Geological Survey, Center for Earth Resources Observation and Science, SD 57198-0001; [§]Science Systems and Applications, Code 614.4, National Aeronautics and Space Administration–Goddard Space Flight Center, Greenbelt, MD 20771; and ^{||}Environmental, Earth, and Atmospheric Sciences, University of Massachusetts, Lowell, MA 01854

Edited by Stephen H. Schneider, Stanford University, Stanford, CA, and approved May 27, 2008 (received for review August 29, 2007)

Since 1980, the number of undernourished people in eastern and southern Africa has more than doubled. Rural development stalled and rural poverty expanded during the 1990s. Population growth remains very high, and declining per-capita agricultural capacity retards progress toward Millennium Development goals. Analyses of *in situ* station data and satellite observations of precipitation have identified another problematic trend: main growing-season rainfall receipts have diminished by $\approx 15\%$ in food-insecure countries clustered along the western rim of the Indian Ocean. Occurring during the main growing seasons in poor countries dependent on rain-fed agriculture, these declines are societally dangerous. Will they persist or intensify? Tracing moisture deficits upstream to an anthropogenically warming Indian Ocean leads us to conclude that further rainfall declines are likely. We present analyses suggesting that warming in the central Indian Ocean disrupts onshore moisture transports, reducing continental rainfall. Thus, late 20th-century anthropogenic Indian Ocean warming has probably already produced societally dangerous climate change by creating drought and social disruption in some of the world's most fragile food economies. We quantify the potential impacts of the observed precipitation and agricultural capacity trends by modeling "millions of undernourished people" as a function of rainfall, population, cultivated area, seed, and fertilizer use. Persistence of current tendencies may result in a 50% increase in undernourished people by 2030. On the other hand, modest increases in per-capita agricultural productivity could more than offset the observed precipitation declines. Investing in agricultural development can help mitigate climate change while decreasing rural poverty and vulnerability.

climate change | drought | famine | precipitation

Sustainability science often examines trends and the interaction of trends (1, 2) through the lens of "use-inspired basic research" (1). We present here an analysis of rainfall and agricultural capacity trends inspired by our research supporting earlier early warning for the U.S. Agency for International Development's Famine Early Warning System [supporting information (SI) Table S1]. Most of this work has focused on monitoring and predicting emerging drought conditions in eastern and southern Africa (3–5), one way of improving adaptive capacity (6) recommended by the Intergovernmental Panel on Climate Change (IPCC). In 2003, however, a routine attempt to use early-season (March to April) rainfall as a predictor of late-season (September to October) crop production produced a disturbing discovery: substantial rainfall declines had occurred within critical crop-growing areas in Ethiopia. The location and timing of these trends made them dangerous. Further analysis suggested links to drought across much of eastern Africa during the long-rains season and potential anthropogenic links to the central Indian Ocean (3, 5). In addition to climate analysis, this work also included evaluations of population, agriculture, and

water security. We find that in eastern and southern Africa, the combination of declining per-capita agricultural capacity (7) and increasing aridity is exacerbating vulnerability (8) and rural poverty (9). Declining investments in rural development, rapidly increasing rural populations, the depletion of soil nutrients through erosion, and the cultivation of most cultivatable areas limit agricultural productivity growth (10). Low per-capita agricultural production and rural poverty go hand in hand, slowing progress toward Millennium Development goals (MDGs). A recent overview of these goals (11) indicated differential progress on different fronts. Education and economic conditions have improved. Population growth remains high (2.3% per year). Progress on hunger and health has been halting. Our concern is that post-1980 rainfall declines during the main eastern African long rains (March–April–May, MAM) and the main southern African summer rains (December–January–February, DJF) may be contributing to African food insecurity. We believe that these declines may be associated with anthropogenic warming in the Indian Ocean. If so, these precipitation declines, acting in concert with declining agricultural capacity, likely constitute a prime example of societally dangerous climate change.

In this report we quantify the likely impact of observed precipitation and agricultural capacity trends by using empirical food-balance indicator models. These models estimate the millions of undernourished people as a function of population, cropped area, seed use, fertilizer applications, and main growing-season precipitation. These models show that continued declines in rainfall and per-capita agricultural capacity will produce increasing food insecurity. We next use observations and climate model simulations to argue that recent declines in eastern and southern African growing-season rainfall are linked to anthropogenic warming in the Indian Ocean. This link to global warming implies that these precipitation declines are likely to continue or intensify. For eastern Africa, this result is at odds with the most recent IPCC assessment (12, 13), which anticipates precipitation increases. For southern Africa, this result is consistent with previous analyses (13, 14), which anticipate rainfall declines. Here we argue that warming in the

Author contributions: C.F. designed research; C.F., M.B., and A.H. performed research; C.F., M.D.D., and M.B. contributed new reagents/analytic tools; C.F., M.E.B., and M.B. analyzed data; and C.F., M.D.D., J.C.M., J.P.V., and M.E.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

[†]To whom correspondence should be addressed at: University of California, Geography Department, 1629 Ellison Hall, Climate Hazard Group, Santa Barbara, CA 93106-1120. E-mail: chris@geog.ucsb.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0708196105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. Food-security analysis. (A) Rainfall, population, cropped area, seed use, and fertilizer use for eastern and southern Africa. Rainfall is expressed as a percentage of the 1951–1980 average. The other variables are expressed as percentages of 1979–1981 averages. Global radiative forcing is shown with a stippled line on an inverted axis. (B) The black vectors denote recent observed rainfall and agricultural capacity tendencies. Agricultural data were obtained from the Food and Agriculture Organization. (C) Empirical food-balance model results expressed as percentages of the millions undernourished in 2000.

south-central Indian Ocean (0–15°S, 60–90°E) has a similar effect on near-coastal eastern and southern Africa during boreal winter and spring. In each season, oceanic warming seems to reduce onshore moisture transports (5, 15–17) while increasing continental atmospheric stability. This may explain recent drought tendencies in gauge data, satellite-observed precipitation,** lake levels (5, 16), and vegetation indices in densely populated water-insecure regions of eastern Africa. If a common anthropogenic Walker-cell-like disruption has caused declines in both southern and eastern African rainfall, then this disruption is likely to persist. Furthermore, recent climate-change impact assessments (18), based on optimistic precipitation simulations over eastern Africa, may underestimate yield reductions (4). Our food-balance modeling of millions of undernourished people, however, suggests that these impacts could be mitigated by agricultural development.

Modeling the Impacts of Agricultural Capacity and Rainfall

Over the past 25 years the population of food-insecure eastern and southern Africa has doubled while per-capita cropped area has declined by 33% and the millions of undernourished people increased by 80% (19). Today, one in three small children is dangerously underweight, and 40% of 308 million people are undernourished. Average national per-capita cropped-area values are often below 1,000 m² per person.†† Food aid has become chronic, with 2005 World Food Program aid valued at U.S. \$200 million. Poor farmers in these countries, often trapped in cycles of displacement, division, and degradation (20), depend on rain-fed agriculture (21). Per-capita crop production is an important metric of food availability and security. Limited technological inputs create a strong dependence between national average cropped area and national average production ($r = 0.95$).

Recent agricultural capacity trends tend to be dominated by cropped-area and population-growth increases.** In general, increased food production has matched growing population through increased labor inputs and through extensification, with area under cultivation increasing by 50% over the past 25 years while population has almost doubled (Fig. 1A). Seed and fertilizer use has also lagged population growth. Unfortunately, since 1990, overseas agricultural assistance has declined from 12% to 4% of total foreign aid (7), and only 4% of African public spending goes to agriculture (7). This declining investment in agriculture has contributed to stalling increases in agricultural

inputs. As the gap between population growth and structural agricultural components continues to grow, vulnerability and rural poverty will increase, amplifying the impact of agricultural droughts. These droughts have seemed to be more severe in recent years (Fig. 1A), and main growing-season rainfall** has declined by $\approx 15\%$ across eastern and southern Africa, in step with recent increases in radiative forcing (22). The interaction of agricultural and rainfall trends may be represented by empirical models estimating the millions of undernourished people.

The five components of national per-capita production (rainfall, population, cropped area, and seed and fertilizer use) can be combined into two variables: rainfall and per-capita agricultural capacity. Agricultural capacity represents the slowly varying nonweather component of a national food balance and is a function of cropped area and seeds and fertilizer divided by population with units of per-capita crop production (kg per person per year). Fig. 1B shows main growing-season rainfall and per-capita agricultural capacity trends for 10 eastern and southern African nations. Somalia has been excluded as a result of limited data. The downward-pointing arrows indicate countries that experienced key growing-season rainfall declines between 1979 and 2005 (Ethiopia, Kenya, Burundi, Tanzania, Malawi, Zambia, and Zimbabwe). Note the geographic clustering of observed rainfall declines along Africa's eastern seaboard. Countries with left-pointing vectors (Ethiopia, Uganda, Zambia, and Zimbabwe) have experienced 1994–2003 declines in per capita agricultural capacity. In these countries population has grown faster than agricultural infrastructure, leading to increased rural poverty and vulnerability to climatic shocks.

In these semiarid countries a strong dependence on rain-fed smallholder farming practices results in quasilinear relationships between grain yields, seasonal rainfall receipts, and food deficits. Hence, agricultural capacity multiplied by percent normal rainfall is strongly related to per-capita production. The inverse of this measure (food imbalance) is related to food insecurity.** For each country (except Ethiopia, because of data limitations), the food-imbalance measure was regressed against “millions of undernourished” statistics obtained from the United Nations Food and Agriculture Organization. These regressions give us a pragmatic means of translating changes in rainfall, cropped area, and seed and fertilizer use into an index of food insecurity that is grounded empirically by historic observations. The resulting model performed well at a national/interannual scale for eastern and southern Africa ($r = 0.86$, $P = 1.6 \times 10^{-5}$). To stress the semiquantitative nature of these projections, we present our results as percent changes from a 2000 baseline, which also allows comparison with the MDG of halving the number of undernourished people by 2015.

**Please see our online *SI Materials and Methods* for a specific description of our methods.

††Agriculture, population, food-security, and food-aid statistics were obtained from the United Nations Food and Agriculture Organization.

Table 1. Correlations between climate, rainfall, and food-security indices

	DJF, <i>R</i> (<i>P</i>)	MAM, <i>R</i> (<i>P</i>)	JJA, <i>R</i> (<i>P</i>)
CC _P and CC _U			
Interannual (1950–2005)	0.45 (5.6 × 10 ⁻⁴)	0.90 (2.3 × 10 ⁻³)	0.81 (6.5 × 10 ⁻¹⁴)
7-yr time scale (1950–2005)	0.97 (6.6 × 10 ⁻⁵)	0.64 (1.4 × 10 ⁻⁸)	0.90 (2.3 × 10 ⁻³)
CC _U and growing season rainfall			
Interannual (1950–2005)	-0.39 (3.2 × 10 ⁻³) (southern Africa)	-0.82 (4.6 × 10 ⁻²) (eastern Africa)	-0.50 (1.0 × 10 ⁻⁴) (Sahel countries)
7-yr time scale (1950–2005)	-0.92 (9.3 × 10 ⁻³) (southern Africa)	-0.57 (2.7 × 10 ⁻⁵) (eastern Africa)	-0.92 (9.3 × 10 ⁻³) (Sahel countries)
CC _P and CMIP3 oceanic rainfall: 7-yr time scale (1966–2005)	0.97 (1.2 × 10 ⁻³)	0.91 (1.6 × 10 ⁻²)	0.86 (2.8 × 10 ⁻²)
CC _P and global temperature: 7-yr time scale (1966–2005)	0.88 (2.0 × 10 ⁻²)	0.85 (3.2 × 10 ⁻²)	0.84 (3.6 × 10 ⁻²)
CC _P and CC _U	0.45 (5.6 × 10 ⁻⁴)	0.90 (2.3 × 10 ⁻³)	0.81 (6.5 × 10 ⁻¹⁴)

can be explored empirically by canonical correlation analysis (CCA). CCA is a statistical technique that finds dominant patterns of covariability in two sets of multivariate data. The two sets of data examined were seasonal 1950–2005 zonal reanalysis moisture transports over Africa's eastern seaboard (20°N–20°S, 25°E–45°E) and seasonal tropical Indian Ocean precipitation (15°N–15°S, 55°E–90°E). Tropical oceanic precipitation is linearly related to diabatic atmospheric heating, which can produce east–west Walker-cell-like circulation anomalies (31). Our hypothesis is that these anomalous circulations control decadal rainfall fluctuations that endanger food-insecure countries along the eastern seaboard of Africa (Fig. 1*B*).

Fig. 2*C* shows our CCA results for MAM (other seasons are presented in Fig. S1). In each season, we find that the first canonical correlate identifies a first CCA pattern describing shared Indian Ocean precipitation and moisture-transport variations. This approach produces a time series describing covariant Indian Ocean precipitation (CC_P) and moisture transports (CC_M). Increasing tropical Indian Ocean precipitation (blue circles) is associated with atmospheric ridging and anticyclonic moisture circulation over eastern Africa (arrows), which disrupt the main onshore moisture flows (shown with blue shading). University of Delaware precipitation deficits for the strongest six oceanic precipitation events are shown with red boxes. Although only boreal spring results are shown here, the patterns for other seasons are similar; a Walker-cell-like dipole emerges as the dominant (first canonical correlate) relationship. The transport correlations indicate reduced onshore moisture fluxes between 0 and 15°S. This region supplies 66% of Indian Ocean-sourced zonal moisture transport into tropical Africa, and 83% of the annual mean zonal moisture transport entering tropical Africa originates in the Indian Ocean. The northern flanks of the CCA patterns are associated with transport anomalies flowing from the drier Sahelian regions. Thus, changes in central Indian Ocean precipitation are correlated with transitions between dry continental and moist oceanic air masses over tropical Africa.

At low (decadal) frequencies, correlations between the CC_M and CC_P approach unity, and the zonal transport indices are negatively correlated ($r \approx -0.9$) with regional rainfall (Table 1). In total the CCA-identified transports can explain 85% of the low frequency variations in eastern MAM, southern DJF, and Sahelian June–July–August (JJA) rainfall. All three seasons/regions experienced a 1960–1970 decline in observed rainfall and CCA transport forcing, in synchrony with anthropogenic 1960–1970 increases in Indian Ocean heat content (24–26). A second heat-content increase in the early 1980s was associated with MAM and DJF transport disruptions and drying tendencies in southern and eastern Africa. Slower increases in JJA transport disruptions, combined with north/south shifts in Atlantic SSTs,

have likely led to a modest recovery in Sahelian rainfall (14, 32), albeit with continued declines across the eastern Sahel and southern Ethiopia (32).

To examine the influence of diabatic heating on moisture transports in a simulation framework, we have also conducted experiments with the National Center for Atmospheric Research Community Atmosphere Model,** a fully nonlinear atmospheric model with explicitly resolved moist processes. An additional diabatic heating term is added over the central Indian Ocean, which increases the local rainfall, as in the work by Barlow *et al.* (33). The added heating is sinusoidal in latitude and longitude for a half period over the domain -20°S to 5°N and 50 to 100°E with a magnitude of 0.02 W·kg⁻¹. This southern central Indian Ocean domain was chosen to match our previous observational analyses (5, 16) and the early assertion that enhanced cyclonic activity helped create the tragic 1984 Ethiopian famine (30). A clear influence on moisture transport over the interior of Africa is observed (Fig. 2*D*), in good agreement with the expectations of simple Gill-type dynamics, and is associated with decreases in local rainfall. These MAM circulation anomalies are similar to previous modeling results for JFM (17), and the observed low-level circulation anomalies associated with warm Indian Ocean warming seasons for boreal spring (Fig. 2*C*). Thus, although the relevance of the simple Gill model might be questioned, especially because it neglects the mean wind (34), the general link between increased central Indian Ocean precipitation and decreased onshore moisture transport seems robust even in a full atmospheric model.

In summary, the strong, probably anthropogenic (24–26), warming of the south-central Indian Ocean (Fig. 2*A*) increases maritime precipitation, which is statistically and dynamically related to continental rainfall declines and moisture-transport reductions (Fig. 2*B* and *C*). Nonlinear climate simulations driven by diabatic heating over the Indian Ocean produce similar transport anomalies (Fig. 2*D*) (17). In this context, what do projected 21st-century climate changes suggest over the Indian Ocean and tropical Africa?

Climate-Change Simulations Suggest Continued Indian Ocean Warming

In this section we present a multimodel analysis of CMIP3 precipitation simulations. If the consensus of these models suggests continued increases in oceanic precipitation, then we should be concerned that continued continental rainfall declines, similar to those shown in Fig. 2*B*, will lead to increased undernourishment unless mitigated (Fig. 1*C*). We selected 11 models having both 20th-century and 21st-century climate simulations. Because near-term (2000–2030) projections of radiative forcing (22) are quite similar across the various emission projections

Vulnerability arises through a complex interplay of exposures, sensitivities, and resiliencies that can either dampen or amplify the impact of climatic shocks (8). The recent IPCC assessment identifies many means of enhancing adaptive capacity (6) in Africa. Improved forecasts and earlier early warning can help (4), as can improving existing rain-fed agriculture through farm management (42) (cf. table 2 in ref. 6), as can enhanced biotechnological (43) inputs. Recent research has suggested that investments in African agricultural development will lead to substantial propoor economic growth (43–45). Our analysis suggests that the continuation of recent tendencies will be likely to result in continued increases in undernourishment; but this is not a *fait accompli*. The range of possible human outcomes is large and depends primarily on the choices that we make. By 2025 rates of child malnutrition, for example, could range from 9.4 to 41.9 million (46). Our agricultural capacity models suggest that a 15% increase in yields per decade (equivalent to a 2 kg per person per year increase in agricultural capacity) could come close to achieving the MDG of halving the number of undernourished people, albeit by 2030 rather than 2015.

Although notable achievements in economic growth and education have been made since 1999, recent progress on hunger in Africa has been very slow (11) and has fallen short of keeping pace with the relatively rapid population growth. The past few years have seen an increased governmental appreciation of the

role of agricultural development (7), but public investments in agriculture and international donor assistance have lagged behind MDG commitments (11). More and higher quality assistance and agricultural policies, guided by objective policy design and better governance, can help eastern and southern Africa achieve sustainability transition. Investments in agricultural capacity seem warranted, given the large financial and humanitarian costs of business as usual in a changing climate system. These investments may also be just, given the strong evidence for anthropogenic drought and the disproportionate emissions by middle- and high-income countries (41).

ACKNOWLEDGMENTS. This research was supported by the U.S. Agency for International Development Famine Early Warning System Network under U.S. Geological Survey Cooperative Agreement 04HQAG0001, the National Aeronautics and Space Administration under Precipitation Science Grant NNX07AG26G, National Aeronautics and Space Administration decision support grant NNG07HW21G, and by the National Science Foundation under Grants ATM 0603555 and ATM 0621237. We thank these agencies for their support. We also thank and acknowledge the CMIP3 modeling groups for providing data for analysis; the Program for Climate Model Diagnosis and Intercomparison for collecting and archiving the model output; and the Johnson Space Center Climate Variability and Predictability Working Group on Coupled Modeling for organizing the model data-analysis activity. The multimodel data archive is supported by the Office of Science, U.S. Department of Energy. We would like to sincerely thank Stephen Schneider and the paper reviewers for their efforts.

- Clark WC (2007) Sustainability science: A room of its own. *Proc Natl Acad Sci USA* 104:1737–1738.
- Kates RW, Parris TM (2003) Long-term trends and a sustainability transition. *Proc Natl Acad Sci USA* 100:8062–8067.
- Funk C, et al. (2005) Recent drought tendencies in Ethiopia and equatorial-subtropical eastern Africa *Famine Early Warning System Network Special Report* (US Agency for International Development, Washington, DC).
- Brown ME, Funk CC, Galu G, Choularton R (2007) Earlier famine warning possible using remote sensing and models. *Trans Am Geophys Union* 88:381–382.
- Verdin J, Funk C, Senay G, Choularton R (2005) Climate science and famine early warning. Cautionary tales: Adaptation and the global poor. *Philos Trans R Soc London Ser B* 360:2155–2168.
- Boko M, et al. (2007) Climate change: Impacts, vulnerabilities, and adaptation in developing countries. *Climate Change 2007: Impacts, Adaptation and Vulnerability. The Working Group II Contribution to the IPCC Fourth Assessment Report*, eds Parry ML, et al. (Cambridge Univ Press, Cambridge UK), pp 433–467.
- WorldBank (2008) *World Development Report 2008: Agriculture for Development* (International Bank for Reconstruction and Development/World Bank, Washington, DC).
- Turner BL, II, et al. (2003) A framework for vulnerability analysis in sustainability science. *Proc Natl Acad Sci USA* 100:8074–8079.
- Oxfam (2006) *Causing hunger. Oxfam Briefing Paper* (OXFAM, Washington, DC).
- Kates RW, Parris TM (2003) Long-term trends and a sustainability transition. *Proc Natl Acad Sci USA* 100:8062–8067.
- United Nations (2007) *The Millennium Development Goals Report, 2007* (United Nations Department of Public Information, Washington, DC).
- Christensen JH, et al. (2007) Regional climate projections, Ch. 11 *Climate Change 2007: The physical science basis*, eds Solomon S, et al. (Cambridge Univ Press, Cambridge, UK; New York, NY).
- Solomon S, et al. (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge Univ Press, Cambridge, UK).
- Hoerling M, Hurrell J, Eischeid J, Phillips A (2006) Detection and attribution of twentieth-century northern and southern African rainfall change. *J Clim* 19:3989–4008.
- Cook KH (2000) The South Indian convergence zone and interannual rainfall variability over southern Africa. *J Clim* 13:3789–3804.
- Funk C, et al. (2006) El Niño and Indian Ocean dipole conditions likely into early 2007, with drought and flooding implications for Southern and Eastern Africa (US Agency for International Development Famine Early Warning System Network, Washington, DC).
- Washington R, Preston A (2006) Extreme wet years over southern Africa: Role of Indian Ocean sea surface temperatures. *J Geophys Res* 111:D15104.
- Lobell DB, et al. (2008) Prioritizing climate change adaptation needs for food security in 2030. *Science* 319:607–610.
- Food and Agriculture Organization (2006) *State of food insecurity in the world (Food and Agriculture Organization of the United Nations, Rome, Italy)*.
- Kates RW (2000) Cautionary tales: Adaptation and the global poor. *Clim Change* 45:5–17.
- Rockstrom J (2000) Water resources management in smallholder farms in eastern and southern Africa: An overview. *Phys Chem Earth B Hydrol Oceans Atmos* 25:275–283.
- Forster P, et al. (2007) Changes in Atmospheric Constituents and in Radiative Forcing, Ch. 2 *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Solomon S, et al. (Cambridge Univ Press, Cambridge, UK).
- Hansen J, et al. (2006) Global temperature change. *Proc Natl Acad Sci USA* 103:14288–14293.
- Barnett TP, Pierce DW, Schnur R (2001) Detection of anthropogenic climate change in the world's oceans. *Science* 292:270–274.
- Cai W, et al. (2007) Atlantic interocean exchange in a regional model. *Geophys Res Lett* 34:L14611.
- Alory G, Wijffels S, Meyers G (2007) Observed temperature trends in the Indian Ocean over 1960–1999 and associated mechanisms. *Geophys Res Lett* 34:L02606.
- Meehl GA, et al. (2007) The WCRP CMIP3 multimodel dataset: A new era in climate change research. *Bull Am Meteorol Soc* 88:1383–1394.
- Adler RF, et al. (2003) Thermodynamic and hydrological impacts of increasing greenness in northern high latitudes. *J Hydrometeorol* 4:1147–1167.
- Kalnay E, et al. (1996) The NCEP/NCAR 40-year reanalysis project. *Bull Am Meteorol Soc* 77:437–471.
- Shanko D, Camberlin D (1998) The effects of the Southwest Indian Ocean tropical cyclones on Ethiopian drought. *Int J Climatol* 18:1373–1388.
- Gill A (1982) *Atmosphere-Ocean Dynamics* (Academic, London, UK).
- Held IM, et al. (2005) Simulation of Sahel drought in the 20th and 21st centuries. *Proc Natl Acad Sci USA* 102:17891–17896, and erratum (2006) 103:1152–1153.
- Barlow M, Hoell A, Colby F (2007) Examining the wintertime response to tropical convection over the Indian Ocean by modifying convective heating in a full atmospheric model. *Geophys Res Lett* 34:L19702.
- Ting M, Sardeshmukh P (1993) Factors determining the extratropical response to equatorial diabatic heating anomalies. *J Atmos Sci* 34:280–296.
- Dai A, et al. (2001) Climates of the 20th and 21st centuries simulated by the NCAR Climate System Model. *J Clim* 14:485–519.
- Dettinger M (2006) A component-resampling approach for estimating probability: Distributions from small forecast ensembles. *Clim Change* 76:149–168.
- Easterling W, et al. (2007) Food, Fibre and Forest Products, Ch. 5. *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Parry ML, et al. (Cambridge Univ Press, Cambridge, UK), pp 273–313.
- Funk C, Michaelsen J (2003) A simplified diagnostic model of orographic rainfall for enhancing satellite-based rainfall estimates in data-poor regions. *J Appl Met* 43:1366–1378.
- Cook KH, Viny EK (2006) Coupled model simulations of the west African monsoon system: Twentieth- and twenty-first-century simulations. *J Clim* 19:3681–3703.
- Brown ME, Funk CC (2008) Climate: Food security under climate change. *Science* 319:580–581.
- Srinivasana UT, et al. (2008) The debt of nations and the distribution of ecological impacts from human activities. *Proc Natl Acad Sci USA* 105:1768–1773.
- Rockström J (2003) Resilience building and water demand management for drought mitigation. *Phys Chem Earth* 28:869–877.
- Delmer DP (2005) Agriculture in the developing world: Connecting innovations in plant research to downstream applications. *Proc Natl Acad Sci USA* 102:15739.
- Thirtle C, Lin L, Piesse J (2003) The impact of research-led agricultural productivity growth on poverty reduction in Africa, Asia and Latin America. *World Dev* 31:1959–1975.
- Tiffin R, Xavier I (2006) Is agriculture the engine of growth? *Agric Econ* 35:79–89.
- Rosegrant MW, et al. (2005) *Looking ahead: Long term prospects for Africa's agricultural development and food security* (International Food Policy Research Institute, Washington, DC).