

## Water-Use Efficiency and Carbon Isotope Discrimination in Peanut under Water Deficit Conditions

G C Wright, R C Nageswara Rao,\* and G D Farquhar

### ABSTRACT

Because of its relationship with water-use efficiency (W), carbon isotope discrimination in leaves ( $\Delta$ ) was proposed to be useful for identifying genotypes with greater water-use efficiency. In this study we examined the relationship between W and  $\Delta$  in four peanut (*Arachis hypogaea* L.) genotypes. The genotypes were grown in and around mini-lysimeters embedded in soil and were subjected to two drought regimes, intermittent and prolonged water deficit conditions, by varying the irrigation timing and amount. Automated rain-out shelters prevented any rain from reaching the experimental plots during the treatment period. The mini-lysimeters allowed accurate measurement of water use and total dry matter (including roots) in a canopy environment. Water-use efficiency, which ranged from 1.81 to 3.15 g kg<sup>-1</sup>, was negatively correlated with  $\Delta$ , which ranged from 19.1 to 21.8‰. Difton-8 had the highest W (3.15 g kg<sup>-1</sup>) and Chico the lowest (1.81 g kg<sup>-1</sup>), representing a variation in W of 74% among genotypes. Variation in W arose mainly from genotypic differences in total dry matter production rather than from differences in water use. It is concluded that  $\Delta$  is a useful trait for selecting genotypes of peanut with improved W under drought conditions in the field. A strong negative relationship existed between W and specific leaf area (SLA, cm<sup>2</sup> g<sup>-1</sup>) and between  $\Delta$  and SLA, indicating that genotypes with thicker leaves had greater W. SLA could therefore be used as a rapid and inexpensive selection index for high W in peanut where mass spectrometry facilities are not available.

IDENTIFICATION OF PHYSIOLOGICAL TRAITS contributing to superior performance of crop plants under drought conditions has been a long-term goal of plant scientists. Water-use efficiency is one such trait which can contribute to productivity when water resources are scarce. Reviews of the literature have suggested that intraspecific variations in W are small and are likely to be increased only by crop management (Fischer, 1979) or modifying the environment (Tanner and Sinclair, 1983). However, variation in W was shown to exist between and within species (Briggs and Shantz, 1914). More recently, variation in W was observed among genotypes of grass species (Farquhar and Richards, 1984; Frank et al., 1985), cotton (Hubick and Farquhar, 1987), and peanut (Wright et al., 1988; ICRISAT, 1990). While potentially useful, W cannot be easily exploited because of practical difficulties involved in measurement of transpiration and root biomass in the field. Spot measurements of transpiration ratio, i.e., CO<sub>2</sub> assimilation rate divided by transpiration rate, only give instantaneous estimates of transpiration efficiency, which may not necessarily correlate with long-term differences.

Farquhar et al. (1982) predicted that at constant vapor pressure difference, discrimination against <sup>13</sup>C during CO<sub>2</sub> assimilation ( $\Delta$ ) will give an estimate of the ratio of the

internal CO<sub>2</sub> concentration in the leaf (C<sub>i</sub>) to ambient CO<sub>2</sub> concentration (C<sub>a</sub>) in plants with the C<sub>3</sub> photosynthetic pathway. Lower  $\Delta$  was associated with lower C<sub>i</sub>/C<sub>a</sub> and greater W (Farquhar et al., 1989). In agreement with the theory,  $\Delta$  in leaf tissue was negatively correlated with W in wheat (Farquhar and Richards, 1984), peanut (Hubick et al., 1986; Wright et al., 1988), cotton (Hubick and Farquhar, 1987), barley (Hubick and Farquhar, 1989), and C<sub>3</sub> grasses (Johnson and Basset, 1991), suggesting that measurement of  $\Delta$  could potentially be used to screen genotypes for greater W.

Much of the research into genotypic variation in W has been conducted in pots under glasshouse conditions where accurate monitoring of water use and above and below ground biomass production could be made. Measurement of W in the field conditions is considerably more difficult. Problems in accurate measurement of water use and its apportionment into transpiration (T) and soil evaporation (E<sub>s</sub>) components, and difficulty in recovering roots for root mass determination can lead to erroneous estimates of W. In addition, the microclimate in field canopies can be considerably different from that of isolated plants in pots, leading to potential differences in stomatal control of transpiration and hence, a breakdown in the relationship between W and  $\Delta$  (Cowan, 1988; Farquhar et al., 1988).

Wright et al. (1988) overcame some of the above mentioned difficulties by using a mini-lysimeter facility which enabled accurate measurement of transpiration and shoot and root mass in peanut genotypes grown in closed canopies. Under adequately watered conditions, substantial variation in W (range of 2.46–3.76 g kg<sup>-1</sup>) among peanut genotypes, and a strong negative correlation between W and  $\Delta$  was demonstrated ( $r = -0.82$ ,  $P < 0.01$ ). The situation under water-limited conditions is less clear. Wright et al. (1993) showed that severe plant water deficits can result in a break down in the relationship between W and  $\Delta$  for peanut genotypes grown in small pots. Greater respiratory losses of carbon was suggested as a possible reason for this discrepancy (Masle et al., 1990).

The objective of the present study was to examine the relationship between W and  $\Delta$  in four peanut genotypes grown under intermittent and long-term water deficit conditions in the field. Such information is essential before large-scale use of this trait as a selection criterion in breeding programs can be recommended.

### MATERIALS AND METHODS

The field experiment was conducted on a deep red clay loam or oxisol (Soil Survey Staff, 1975) at the Bjelke Petersen Research Station, Kingaroy (151 °E, 26 °S), Queensland, during

G C Wright, Queensland Dep Primary Ind., PO Box 23, Kingaroy, Q 4610, R C Nageswara Rao, Crop Physiology, Legumes Program, ICRISAT, Patancheru 502 324, A P., India, G D Farquhar, Res. School of Biological Sci., Canberra, ACT 2601. Submitted as ICRISAT Journal Article no. 1389. Received on 31 Aug 1992. \*Corresponding author.

Published in *Crop Sci.* 34:92–97 (1994)

**Abbreviations:** W, water use efficiency,  $\Delta$ , carbon isotope discrimination, T, transpiration, ET, evapotranspiration, E<sub>s</sub>, soil evaporation, RUE, radiation use efficiency, SLA, specific leaf area, C<sub>i</sub>, internal CO<sub>2</sub> concentration in leaf, C<sub>a</sub>, ambient CO<sub>2</sub> concentration, H<sub>1</sub> and H<sub>2</sub>, initial and final harvests, respectively, I, and C, intermittent and continuous water deficit, respectively, PAR, photosynthetically active radiation, VPD, air saturation vapor pressure deficit.

Table 1 Schedule of Irrigation followed in the two treatments

Date	Irrigation applied	
	I <sub>1</sub>	I <sub>2</sub>
	mm	
7 Jan 1991	70	70.0
11 Jan 1991	—	7.5
18 Jan 1991	—	12.0
25 Jan 1991	—	10.4
31 Jan 1991	100	10.0
9 Feb 1991	—	15.0
15 Feb 1991	—	15.0
22 Feb 1991	—	15.0

the 1990–1991 summer season (November–April). Four peanut genotypes, i.e., Chico, McCubbin, Shulamit, and Tifton 8, were selected based on previously measured differences in  $W$  and  $\Delta$  (Hubick et al., 1986; Wright et al., 1988). Each genotype was grown under two irrigation regimes (I<sub>1</sub> and I<sub>2</sub>) in a split-plot design with genotypes as sub- and irrigation regimes as main treatments, replicated twice under each of two rain-out shelters (four replicates). Each rain-out shelter, covering an area of 6 by 19 m (Hatfield et al., 1989), prevented rain from reaching the experimental plots throughout the growing period. Staggered plantings were made to match the timing of imposition of treatments with the phenology of genotypes. Planting dates were selected based on the thermal time requirement for flowering (Bell et al., 1991). Tifton-8 and Shulamit (long-duration types) were planted on 19 Nov 1990, and McCubbin (medium duration type) and Chico (short-duration type) were planted on 22 and 28 Nov 1990 respectively. A basal fertilizer of P and K, each at a rate of 30 kg ha<sup>-1</sup> was incorporated into the top soil. Seeds, treated with captan<sup>1</sup> [N-(trichloromethylthio)-4-cyclohexene 1,2-dicarboximide] at 3 g kg<sup>-1</sup> (to prevent seedling diseases) and ethephon [2 (chloroethyl)phosphonic acid] (to ensure seed dormancy was broken), were hand-sown in excess and later thinned to the required population. Each plot consisted of four rows each of 6 m length. A density of 22 plants m<sup>-2</sup> was achieved with an interrow spacing of 30 cm and an intra plant spacing of 15 cm. Main treatments were separated by planting two guard rows. Plots were maintained weed free by hand weeding.

About 3 wk before sowing, two intact soil cores 0.8 m deep and 0.3-m diam were excavated in each plot at a distance of 2.5 m from each other with a Proline coring machine (Evans Deakin Proline, S.A. Australia). Mini-lysimeters were prepared with PVC storm water pipe (0.3-m internal diam and 0.8 m deep) and the intact soil cores, as described by Wright et al. (1988). Each plot consisted of two mini lysimeters, hereafter referred to as *pots*, to monitor transpiration and biomass production. A basal fertilizer of P and K, each at a rate of 30 kg ha<sup>-1</sup> was mixed in the top soil of each pot before planting. Five seeds of the appropriate genotype were planted and later thinned to two plants per pot.

In addition to the two cores in each plot, three cores were similarly prepared in each shelter in the guard rows between the main treatments. These pots were left unplanted (hereafter referred to as *bare pots*) and were used to monitor potential soil evaporation in the absence of plants.

After planting, irrigation of the bulk crop area was accomplished with trickle-irrigation (T-tape with outlets every 20 cm), and irrigation input into each main treatment was measured by water meters (having an accuracy of  $\pm 2\%$ ).

Plants in pots were irrigated at a depth of 20 cm to minimize soil evaporation. Before sowing in pots, a perforated plastic pipe (25-mm diam) was made into a circular shape by connecting each end to a T-junction. Another plastic feeder pipe

of 50 cm length (25 mm diam) was connected to the other end of the T-junction. The ring was buried at a depth of 20 cm in the pot allowing the feeder pipe to reach about 30 cm above the soil surface through which desired amounts of water were added by hand. The bulk crop, as well as plants in the pots, were kept well watered until the imposition of treatments.

Two treatments, an intermittent (I<sub>1</sub>) and a continuous water deficit (I<sub>2</sub>), of 50-d duration (7 January–25 February) were imposed after flowering by varying the timing and amount of irrigation (Table 1). In I<sub>1</sub>, two drought episodes of 25-d duration each, were imposed, with a single release from drought on 31 January. For I<sub>2</sub>, plants were subjected to continuous water deficit by irrigating weekly, with an amount equal to 25% of weekly cumulative pan evaporation during the treatment period. One of the three bare pots in each shelter received irrigation similar to I<sub>1</sub>, while the other two received irrigation similar to I<sub>2</sub>. The experiment was terminated on 25 February to avoid any confounding effects from maturity differences among genotypes.

An initial harvest (H<sub>1</sub>) was performed on 6 Jan 1991 in which ten uniform plants from each plot and one plant from each pot were harvested to provide dry matter estimates at the beginning of the treatments. Plants from the bulk crop were harvested at ground level while one of the two plants from each pot was harvested, with care in order to recover as much of the root system as possible. The final harvest (H<sub>2</sub>) was made on 25 February in which plants in the two middle rows of the bulk crop (2 m length) were harvested, and the remaining plant in each pot was harvested separately.

Plants from each pot and a sub-sample of three plants from the bulk crop were partitioned into leaf, stem, pods, and roots prior to drying (at 80 °C for 48 h) and weighing. Specific leaf area (SLA) was determined as the ratio of the leaf area of a leaf sub sample (about 40 leaflets) to the oven-dry weight of the leaf sub sample. At H<sub>2</sub>, total root biomass was measured, following recovery of roots from the soil cores as described by Wright et al. (1988).

Water loss from pots was estimated by weighing them at weekly intervals. An electronic load cell (accuracy of  $\pm 0.1$  kg) mounted on a tractor driven gantry was used to weigh pots by lifting them approximately 10 cm above the ground. Transpiration (T), during the treatment period was then determined as

$$T = I + (Wt_1 - Wt_2) - E_s,$$

where  $I$  was irrigation (in kilograms),  $Wt_1$  and  $Wt_2$  were the weights of each pot at the beginning and end of the week, respectively, and  $E_s$  was soil evaporation during the period of measurement. The value of  $E_s$  was estimated as  $e(1-f)$ , where  $e$  was the potential soil evaporation determined from the changes in water content in bare pots and  $f$  was the fractional radiation interception by the foliage (Cooper et al., 1983). The value of  $W$  was then estimated as the ratio of total dry matter production (leaf + stem + pod + roots) and transpiration between H<sub>1</sub> and H<sub>2</sub>.

Fractional photosynthetically active radiation (PAR) intercepted by the crop ( $f$ ) was measured at 1400 h AST at H<sub>1</sub>, and at both 0900 and 1400 h at H<sub>2</sub> with a line quantum sensor (LI 1915, LI COR, Lincoln, NE). Radiation use efficiency (RUE) of the crop was estimated as the ratio of dry matter produced and the cumulative PAR intercepted by the crop between H<sub>1</sub> and H<sub>2</sub>. Leaf temperature was measured at weekly intervals at 1400 h on fully expanded leaves (third or fourth leaf on main axis) with a copper-constantan thermocouple fixed to a leaf clip. In each plot, measurements were made on two plants in the bulk crop and on one plant in each pot.

Dried leaf samples used for SLA measurements at H<sub>1</sub> and H<sub>2</sub> were ground to pass through a 100  $\mu$ m sieve. The  $\Delta$  of these samples were measured by ratio mass spectrometry using

<sup>1</sup> Mention of commercial products or companies does not imply endorsement or recommendation by ICRISAT over others of similar nature.




Fig. 1. Pan evaporation (E, 1A), and transpiration (T) of Chico (—), McCubbin (---), Shulamit (---) and Tifton-8 (---) grown in pots under (1B) and intermittent (I<sub>1</sub>) and (1C) continuous drought (I<sub>2</sub>) treatments. Vertical bars indicate LSD ( $P < 0.05$ ) to compare means within a given sampling time.

techniques as described by Hubick et al. (1986). The  $\Delta$ , which occurred between  $H_1$  and  $H_2$ , was derived as follows. If at  $H_1$ , plant dry weight was  $M_1$ , and the associated discrimination in fixing the carbon to that point was  $\Delta_1$ , and there followed a period where dry weight increased to  $M_2$ , during which the carbon was fixed with discrimination  $\Delta$ , then the average discrimination for the total mass,  $\Delta_2$ , would be,

$$\Delta_2 = [M_1 \Delta_1 + (M_2 - M_1)\Delta] / M_2$$

Solving the above equation for  $\Delta$  would be

$$\Delta = (\Delta_2 M_2 - \Delta_1 M_1) / (M_2 - M_1)$$

where  $\Delta_1$  and  $\Delta_2$  are the isotope discrimination, and  $M_1$  and  $M_2$  are total plant dry weights, at  $H_1$  and  $H_2$ , respectively.

Maximum and minimum temperatures, relative humidity, incident solar radiation and Class A pan evaporation were collected daily from an automatic weather station located adjacent to the experimental site. Air saturation vapor pressure deficit (VPD) was calculated with air temperature and RH.

## RESULTS

Mean maximum and minimum air temperatures during the growing season were  $30 \pm 4^\circ\text{C}$  and  $15 \pm 3^\circ\text{C}$  respectively while incident solar radiation ranged from 20 to 35  $\text{MJ m}^{-2} \text{d}^{-1}$ , and VPD ranged from 1.7 to 2.2 kPa during the treatment period.

There was an initial increase in transpiration, T, in response to higher evaporative demand in both treatments at 2 wk following the start of irrigation treatments (Fig. 1). This was followed by a steady decline in T as drought progressed. In general, seasonal T of Chico was lower compared to other genotypes in both treatments. However, genotype differences in T were only significant ( $P < 0.05$ ) at 2 and 6 wk after imposition of I<sub>1</sub>, with Chico having significantly lower T compared to the other genotypes. These differences were also associated with higher soil evaporation (E) losses in Chico due to relatively smaller ground cover compared to other genotypes. In I<sub>2</sub>, T was not significantly different among genotypes, and declined gradually as drought progressed until it reached a stable rate of  $0.5 \text{ kg wk}^{-1}$ , on a pot basis (Fig. 1). Evapotranspiration (ET) was generally lower in I<sub>2</sub> compared to I<sub>1</sub>.

Total dry matter (TDM, including roots) which accumulated during the treatment period varied significantly ( $P < 0.05$ ) among genotypes and ranged from 20 to 37 and 18 to 31  $\text{g plant}^{-1}$  in I<sub>1</sub> and I<sub>2</sub>, respectively (Table 2). In both treatments, Chico produced the least, and Tifton-8 the greatest TDM during the treatment period. The difference in TDM between McCubbin, Shulamit, and Tifton-8 were not significant.

Significant ( $P < 0.01$ ) genotypic variation in root weight of the order of 250% was evident in both irrigation treatments. These differences in root weight and root shoot ratios were associated with relatively small variation in T between genotypes (Table 2).

It was interesting to note that while differences in TDM between genotypes were of the order of 72 to 85%, corresponding differences in T were only 12 to 19%. Thus, variation in TDM seems to dominate, although

Table 2. Total dry weight (TDM, including roots), root weight, transpiration (T), water-use efficiency (W), root shoot ratio (RS), radiation-use efficiency (RUE), and carbon isotope discrimination in leaf ( $\Delta \times 10^3$ ) for four peanut genotypes grown under two drought treatments

Trt	Genotype	TDM	Root	l	W	RS	RUE	$\Delta$ pot	$\Delta$ crop
		g plant <sup>-1</sup>							
I <sub>1</sub>	Chico	20.5	3.3	11.3	1.81	0.15	0.66	21.1	21.1
	McCubbin	36.3	5.7	13.4	2.72	0.15	0.80	20.2	20.7
	Shulamit	35.7	7.2	12.8	2.78	0.21	0.79	19.1	19.9
	Tifton-8	37.5	6.8	12.2	3.05	0.19	0.86	19.3	19.4
I <sub>2</sub>	Chico	17.8	2.3	8.9	2.02	0.11	0.62	21.8	20.9
	McCubbin	26.8	4.6	10.0	2.68	0.16	0.64	20.2	19.4
	Shulamit	29.0	5.9	9.9	2.93	0.20	0.66	19.5	18.8
	Tifton-8	31.3	6.4	10.0	3.15	0.21	0.75	18.6	18.6
	LSD 5%	5.5	1.2	1.1	0.38	0.033	0.09	0.41	0.93




Fig. 2. Relationship between  $W$  and  $\Delta$  (A) and  $W$  and SLA (B) in leaves of Chico ( $\circ$ ), McCubbin ( $\square$ ), Shulamit ( $\Delta$ ) and Tifton-8 ( $\nabla$ ) under intermittent (closed symbols) and continuous (open symbols) drought treatments.  $W = 11.31 - 0.43 \Delta$ , ( $r^2 = 0.89$ ,  $P < 0.01$ );  $W = 5.4 - 0.2 \text{ SLA}$ , ( $r^2 = 0.84$ ,  $P < 0.01$ ).

considerable variation exists for  $T$  among peanut genotypes.

Genotypic differences in  $W$  ranged from 1.81 to 3.05  $\text{g kg}^{-1}$  in  $I_1$  and from 2.02 to 3.15  $\text{g kg}^{-1}$  in  $I_2$ , representing variation in  $W$  of the order of 56 to 69% (Table 2). Differences in TDM among genotypes accounted for up to 80 and 90% of the variation in  $W$  measured in  $I_1$  and  $I_2$ , respectively. Genotype  $\times$  treatment interaction for  $W$  was not significant, and the ranking of genotypes for  $W$  was similar in the two drought treatments. Radiation-use efficiency (RUE) ranged from 0.66 to 0.86  $\text{g MJ}^{-1}$  in  $I_1$  and was lower in  $I_2$  (0.62–0.75  $\text{g MJ}^{-1}$ ), with Tifton-8 and Chico having the maximum and minimum RUE, respectively, in both treatments (Table 2). There was a strong positive relationship between  $W$  and RUE in both  $I_1$  ( $r^2 = 0.99$ ,  $P < 0.01$ ) and  $I_2$  ( $r^2 = 0.66$ ,  $P < 0.05$ ) treatments (not shown). Although RUE of all genotypes declined in  $I_2$ , ranking of genotypes for  $W$  and RUE was consistent in both treatments.

The  $\Delta$  varied significantly among genotypes (Table 2) with Tifton-8 and Chico having the least ( $19.0 \pm 0.17\%$ ) and the greatest ( $21.5 \pm 0.17\%$ ) values, re-

spectively (when averaged over both treatments). There was a strong negative relationship between  $\Delta$  and  $W$  ( $r^2 = 0.89$ ,  $P < 0.01$ , Fig. 2A). Interestingly,  $W$  was also negatively related to SLA ( $r^2 = 0.84$ ,  $P < 0.01$ , Fig. 2B).

The leaf temperature measurements done at weekly intervals indicated that the difference between plants in pots and adjacent crop was less than 1 °C during the most of the treatment period. In  $I_1$ , although leaf temperature in pots increased relative to the crop at 5 wk after imposition of treatments, the difference was less than 2 °C at 7 wk when plants were harvested. In  $I_2$ , leaf temperature in pots and the adjacent crop was similar throughout the treatment period, although leaf temperature of Shulamit and Tifton-8 was marginally higher (by 2–3 °C) in pots compared to the bulk crop at 7 wk.

The  $\Delta$  in pots was well correlated with that of the bulk crop [ $\Delta(\text{crop}) = 4.34 + 0.77\Delta(\text{pot})$ ,  $r^2 = 0.61$ ,  $P < 0.05$ ,  $n = 64$ ]. Similarly, the SLA in pots was well correlated with that of the bulk crop [ $\text{SLA}(\text{crop}) = -46.4 + 1.61 \text{ SLA}(\text{pot})$ ,  $r^2 = 0.73$ ,  $P < 0.01$ ,  $n = 32$ ]. Significant correlation of  $\Delta$  and SLA between plants grown in pots and those in adjacent crop showed that the relative ranking of genotypes for  $\Delta$  and SLA was consistent in both situations.

The relationship between SLA and  $\Delta$  was positive and significant ( $r^2 = 0.81$ ,  $P < 0.01$ , Fig. 3) confirming the earlier observation (Fig. 2B) that SLA could be used to identify genotypes with lower  $\Delta$  and greater  $W$ .

## DISCUSSION

Genotypes used in this study varied in their phenology, with Chico being a short duration, and McCubbin a medium duration spanish type (var. *vulgaris* of ssp. *fastigiata*), and Shulamit and Tifton-8 being long duration virginia bunch and virginia runner types (i.e., var. *hypogaea* of ssp. *hypogaea*), respectively. Staggered planting of genotypes based on differing thermal time requirements for flowering (Bell et al., 1991) enabled us to match crop phenology with the timing of drought. More than 50% of the plants in all plots had flowered by the beginning of the drought treatments. Genotypes received similar amounts of water in a given drought




Fig. 3. Relationship between the mean SLA (of  $H_1$  and  $H_2$ ) and  $\Delta$  in leaves of four peanut cultivars under the two drought treatments.  $\Delta = 14.2 + 0.04 \text{ SLA}$ , ( $r^2 = 0.81$ ,  $P < 0.01$ ).

treatment (Table 1), yet produced differing amounts of total dry matter, TDM (Table 2). Variation in transpiration,  $T$  among genotypes was relatively small compared to variation in TDM with the latter accounting for 80 and 90% of the variation in water use efficiency,  $W$  in  $I_1$  and  $I_2$ , respectively. The  $W$  was measured on single plants grown in mini-lysimeter pots of 58-L capacity which were placed within canopies in the field. Canopies of plants in pots and adjacent crops were well intermixed. Earlier studies have indicated that evapotranspiration was not significantly different among large ( $3.0 \text{ m}^2$ ), medium ( $0.75 \text{ m}^2$ ) and small ( $0.18 \text{ m}^2$ ) lysimeters after adjusting evapotranspiration for leaf area differences (Dugas and Bland, 1989). The  $\Delta$  and SLA of plants in pots were well correlated with that of the bulk crop suggesting consistency in relative ranking of genotypes for  $\Delta$  and SLA in the pot and canopy situations. However, the fact that the slopes and  $Y$  Intercepts of these relationships were substantially different from unity and zero, respectively, raises the possibility that some differences in microclimate between pots and adjacent crop may have been present.

There was significant genotypic variation in production of root biomass during the treatment period, with genotypes with high  $W$  having greater root mass (Table 2). Similar observations have been reported for peanut grown in pots (Hubick et al., 1986), and in the field under well-watered conditions (Wright et al., 1988). It is also of interest that while Tifton-8 produced greater root biomass than Chico, there was no significant difference in transpiration during the treatment period. Whether this difference is related to variation in root length density (Cooper et al., 1987), or superior water uptake per unit root length is unknown. This observation does raise the question as to whether some genotypes may produce roots in excess of what is required. Further information on this aspect may have important implications for peanut yield performance under drought conditions.

Changes in  $C_i/C_a$  and  $\Delta$  can arise from changes in the balance between leaf stomatal conductance and photosynthetic capacity. Where  $C_i/C_a$  changes because of stomatal movements, significant boundary layer resistances to fluxes of water vapor and heat may occur and cause the relationship between  $W$  and  $\Delta$  to break down in plants grown in canopies in the field (Cowan, 1988; Farquhar et al., 1988). Where  $C_i/C_a$  changes in response to variation in photosynthetic capacity the problem of poor coupling between the crop canopy and atmosphere may be less important (Farquhar et al., 1988). A strong correlation between  $W$  and  $\Delta$  demonstrated in this study under water-limited conditions in the field therefore suggests that variation in  $C_i/C_a$  and  $\Delta$  arose mainly from variation in photosynthetic capacity. The negative association between  $\Delta$  and TDM for the genotypes tested here (Table 2) indeed suggests that variation in photosynthetic capacity, rather than variation in stomatal conductance (and water use), predominated. This finding is in agreement with earlier studies in peanut (Hubick et al., 1986, 1988; Wright et al., 1988) and sunflower (Virgona et al., 1990).

The RUE was significantly less in  $I_2$  compared to  $I_1$ , with little change in  $W$  (Table 2), although  $W$  and RUE were linearly related in both treatments. This result again

reinforces the hypothesis that genotypic variation in  $W$  occurs due to effects of photosynthetic capacity on dry matter production, rather than to effects of stomatal conductance on water use in peanut. Thus, RUE in all genotypes was significantly less in  $I_2$  than in  $I_1$ , presumably because of lowered leaf (and canopy) conductance arising from greater levels of soil and crop water deficits reducing photosynthetic activity. This observation suggests the reduction in conductance did not influence the genotypic variation in photosynthetic capacity, as genotypic ranking in RUE was maintained in  $I_1$  and  $I_2$ .

Significant differences in RUE among peanut genotypes grown under water-limited conditions have been shown previously (Matthews et al., 1988). Our results confirm such variability and suggest that selection for high  $W$  (via  $\Delta$  or SLA) may concurrently improve RUE, which is an important trait for high biomass production under non-limiting conditions.

The strong negative relationship between  $W$  and SLA (Fig. 2B) suggests that the genotypes with lower SLA (greater leaf thickness) had greater  $W$ . The significant relationship between  $\Delta$  and SLA (Fig. 3) suggests that genotypes with thicker leaves (low SLA) had greater photosynthetic capacity and thereby assimilated more carbon per unit leaf area, as observed in other crops (Dornhoff and Shibles, 1976; Wolf and Blaser, 1972; Bowes et al., 1972). Substantial variation in photosynthetic rates per unit leaf area have been demonstrated in peanut (Pallas and Samish, 1974; Bhagsari and Brown, 1976; Pallas, 1982). The strong correlation between SLA and  $W$  (and  $\Delta$ ) suggests that SLA could be used as a rapid and inexpensive selection index to identify genotypes with high  $W$  (low  $\Delta$ ), where mass spectrometry facilities are not available. However, SLA is influenced by environmental factors such as temperature and water deficit which affect cell elongation and multiplication (Vivekanandan and Gunasena, 1976) and starch accumulation in chloroplasts (Araus et al., 1989). Similarly, Wright et al. (1988) and Hubick and Farquhar (1989) have shown that  $\Delta$  was strongly influenced by various environmental factors. The  $G \times E$  interaction for the relationship between SLA and  $\Delta$  in peanut is further described by Nageswara Rao and Wright (1993).

#### ACKNOWLEDGMENTS

This research was done under ACIAR Project no. 8834. Support by ICRISAT and ACIAR for the R. C. Nageswara Rao's sabbatical at ODPI, Kingaroy is gratefully acknowledged. Authors thank Bill Tapsall and his staff for maintaining the experimental site and Blair Winders for technical assistance.

#### REFERENCES

- Araus, J. L., L. Sauque, J. Matas, and M. D. Serret. 1989. Seasonal changes in the photosynthetic capacity and leaf structure of *Festuca japonica* leaves grown in the shade. *J. Hort. Sci.* 64: 189-197.
- Bell, M. J., R. Shorter, and R. Mayer. 1991. Cultivar and environmental effects of growth and development of peanuts (*Arachis hypogaea* L.) I. Emergence and flowering. *Field Crops Res.* 27: 17-33.
- Bhagsari, A. S., and R. H. Brown. 1976. Photosynthesis of peanut (*Arachis*) genotypes. *Peanut Sci.* 3: 1-9.
- Bowes, G., W. L. Orgen, and R. H. Hageman. 1972. Light saturation, photosynthesis rate, RUDP carboxylase activity and specific leaf weight in soybeans grown under different light intensities. *Crop Sci.* 12: 77-79.

- Briggs I.J., and H.L. Shantz. 1914. Relative water requirements of plants. *J. Agric. Res.* 1:64.
- Cooper, P.J.M., P.J. Gregory, J.D.H. Keatings, and S.C. Brown. 1987. Effects of fertilizer, variety and location on barley production under rainfed conditions in Northern Syria. 2. Soil water dynamics and crop water use. *Field Crops Res.* 16:67-84.
- Cooper, P.J.M., J.D.H. Keatings, and G. Hughes. 1983. Crop evapotranspiration—A technique for calculation of its components by field measurements. *Field Crops Res.* 7:299-312.
- Cowan, I.R. 1988. Stomatal physiology and gas exchange in the field. In O.T. Denmead (ed.) *Flow and transport in the natural environment: advances and application*. Springer-Verlag, New York.
- Dornhoff, G.M., and R.M. Shibles. 1976. Leaf morphology and anatomy in relation to CO<sub>2</sub> exchange rate of soybean leaves. *Crop Sci.* 16:377-381.
- Dugas, W.A., and W.L. Bland. 1989. The accuracy of evaporation measurements from small lysimeters. *Agric. For. Meteorol.* 1-2:119-120.
- Farquhar, G.D., J.R. Ehleringer, and K.T. Hubick. 1989. Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40:503-537.
- Farquhar, G.D., K.T. Hubick, A.G. Condon, and R.A. Richards. 1988. Carbon isotope fractionation and plant water-use efficiency. p. 21-40. In P.W. Rundel et al. (ed.) *Applications of stable isotope ratios to ecological research*. Springer-Verlag, New York.
- Farquhar, G.D., M.H. O'Leary, and J.A. Berry. 1982. On the relationship between carbon isotope discrimination and intercellular carbon dioxide concentration in leaves. *Aust. J. Plant Physiol.* 9:121-137.
- Farquhar, G.D., and R.A. Richards. 1984. Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. *Aust. J. Plant Physiol.* 11:539-552.
- Fischer, R.A. 1979. Growth and water limitation to dryland wheat yield in Australia. A physiological framework. *J. Aust. Inst. Agric. Sci.* 45:83-94.
- Frank, A.B., J.D. Berdahl, and R.E. Barker. 1985. Morphological development of and water use in clonal lines for forage grasses. *Crop Sci.* 25:339-344.
- Hatfield, P.M., G.C. Wright, and W.R. Tapsall. 1989. A large, retractable, low cost and re-locatable rainout shelter design. *Expt. Agric.* 26:57-62.
- Hubick, K.T., and G.D. Farquhar. 1987. Carbon isotope discrimination—selecting for water-use efficiency. *Aust. Cotton Grower.* 8:66-68.
- Hubick, K.T., and G.D. Farquhar. 1989. Carbon isotope discrimination and the ratio of carbon gained to water lost in barley cultivars. *Plant Cell Environ.* 12:795-804.
- Hubick, K.T., G.D. Farquhar, and R. Shorter. 1986. Correlation between water-use efficiency and carbon discrimination in diverse peanut (*Arachis*) germplasm. *Aust. J. Plant Physiol.* 13:803-816.
- Hubick, K.T., R. Shorter, and G.D. Farquhar. 1988. Heritability and genotype  $\times$  environment interactions of carbon isotope discrimination and transpiration efficiency in peanut (*Arachis hypogaea* L.). *Aust. J. Plant Physiol.* 15:799-813.
- ICRISAT (International Crops Research Institute for the Semi-Arid Tropics) 1990. Annual Report 1989. Patancheru, A.P. 502-324, India, ICRISAT, P. 143.
- Johnson, R.C., and I.M. Bassett. 1991. Carbon isotope discrimination and water-use efficiency in four cool season grasses. *Crop Sci.* 31:157-162.
- Masle, J., G.D. Farquhar, and R.M. Gifford. 1990. Growth and carbon economy of wheat seedlings as affected by soil resistance to penetration and ambient partial pressure of CO<sub>2</sub>. *Aust. J. Plant Physiol.* 17:465-487.
- Mathews, R.B., D. Harris, J.H. Williams, and R.C. Nageswara Rao. 1988. The physiological basis for yield differences between four genotypes of groundnut (*Arachis hypogaea* L.) in response to drought. II. Solar radiation interception and leaf movement. *Exp. Agric.* 24:203-213.
- Nageswara Rao, R.C., and G.C. Wright. 1993. Stability of the relationship between specific leaf area and carbon isotope discrimination across environments in peanut. *Crop Sci.* 34:98-103 (this issue).
- Pallas, J.E., Jr. 1982. Photosynthetic traits of selected peanut genotypes. *Peanut Sci.* 9:14-17.
- Pallas, J.E., Jr., and Y.B. Samish. 1974. Photosynthetic response of peanut. *Crop Sci.* 14:478-482.
- Soil Survey Staff. 1975. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. USDA Agric. Handb. 436. U.S. Gov. Print. Office, Washington, DC.
- Fanner, C.B., and T.R. Sinclair. 1983. Efficient water use in crop production: research or re-search. p. 1-28. In H. Taylor (ed.) *Limitations to efficient water use in crop production*. ASA, CSSA, and SSSA, Madison, WI.
- Virgona, J.M., K.T. Hubick, H.M. Rawson, and G.D. Farquhar. 1990. Genotypic variation in transpiration efficiency, carbon isotope discrimination and dry matter partitioning during early growth in sunflower. *Aust. J. Plant Physiol.* 17:207-214.
- Vivekanandan, A.S., and H.P.M. Gunesena. 1976. Lysimetric studies on the effect of soil moisture tension on the growth and yield of maize (*Zea mays* L.) and groundnut (*Arachis hypogaea*). *Beitr. Trop. Landwirtschaft. Veterinarmedizin* 14:369-378.
- Wolf, D.D., and R.E. Blaser. 1972. Growth rate and physiology of Alfalfa as influenced by canopy and light. *Crop Sci.* 12:23-26.
- Wright, G.C., K.T. Hubick, and G.D. Farquhar. 1988. Discrimination in carbon isotopes of leaves correlates with water-use efficiency of field-grown peanut cultivars. *Aust. J. Plant Physiol.* 15:815-825.
- Wright, G.C., K.T. Hubick, G.D. Farquhar, and R.C. Nageswara Rao. 1993. Genetic and environmental variation in transpiration efficiency and its correlation with carbon isotope discrimination and specific leaf area in peanut. p. 247-267. In J. Ehleringer et al. (ed.) *Stable isotopes and plant carbon-water relations*. Academic Press, San Diego, CA.