

WEAK CONVERGENCE OF MULTIDIMENSIONAL EMPIRICAL PROCESSES FOR STATIONARY ϕ -MIXING PROCESSES¹

BY PRANAB KUMAR SEN

University of North Carolina, Chapel Hill

For a stationary ϕ -mixing sequence of stochastic $p(\geq 1)$ -vectors, weak convergence of the empirical process (in the J_1 -topology on $D^p[0, 1]$) to an appropriate Gaussian process is established under a simple condition on the mixing constants $\{\phi_n\}$. Weak convergence for random number of stochastic vectors is also studied. Tail probability inequalities for Kolmogorov-Smirnov statistics are provided.

1. Introduction. Let $\{X_i = (X_{i1}, \dots, X_{ip})', -\infty < i < \infty\}$ be a stationary ϕ -mixing sequence of stochastic vectors defined on a probability space (Ω, \mathcal{A}, P) with each X_i having (marginally) a continuous distribution function (df) $F(\mathbf{x})$, $\mathbf{x} \in R^p$, the $p(\geq 1)$ dimensional Euclidean space. Thus, if $\mathcal{M}_{-\infty}^k$ and $\mathcal{M}_{k+n}^{\infty}$ be respectively the σ -fields generated by $\{X_i, i \leq k\}$ and $\{X_i, i \geq k+n\}$, and if, $A \in \mathcal{M}_{-\infty}^k$ and $B \in \mathcal{M}_{k+n}^{\infty}$, then for all $k: -\infty < k < \infty$,

$$(1.1) \quad |P(A \cap B) - P(A)P(B)| \leq \phi_n P(A), \quad \phi_n \geq 0, \text{ for all nonnegative } n,$$

where ϕ_n is \downarrow in n and $\lim_{n \rightarrow \infty} \phi_n = 0$. We denote the marginal df of X_{ij} by $F_{[j]}$, let $Y_{ij} = F_{[j]}(X_{ij}), j = 1, \dots, p; Y_i = (Y_{i1}, \dots, Y_{ip})', -\infty < i < \infty$, and denote the df of Y_i by

$$(1.2) \quad G(\mathbf{t}) = P\{Y_i \leq \mathbf{t}\}, \quad \mathbf{t} \in E^p \text{ (so that } G_{[j]}(t) = P\{Y_{ij} \leq t\} = t; \\ 0 \leq t \leq 1; j = 1, \dots, p),$$

where $E^p = \{\mathbf{t}: \mathbf{0} \leq \mathbf{t} \leq \mathbf{1}\}$ is the p -dimensional unit cube, $\mathbf{0} = (0, \dots, 0)$, $\mathbf{1} = (1, \dots, 1)$ and $\mathbf{a} \leq \mathbf{b}$ means that $a_j \leq b_j, 1 \leq j \leq p$. Note that $G(\mathbf{t}) = 0$ if at least one coordinate of \mathbf{t} is 0. For a sample X_1, \dots, X_n of size n , the empirical df for Y_1, \dots, Y_n is defined by

$$(1.3) \quad G_n(\mathbf{t}) = n^{-1} \sum_{i=1}^n c(\mathbf{t} - Y_i), \quad \mathbf{t} \in E^p, n \geq 1,$$

where $c(\mathbf{u}) = 1$ iff $\mathbf{u} \geq \mathbf{0}$, and 0, otherwise. Also, $G_n(\mathbf{t}) = 0$, when at least one coordinate of \mathbf{t} is 0. The empirical process $W_n = \{W_n(\mathbf{t}), \mathbf{t} \in E^p\}$ is then defined by

$$(1.4) \quad W_n(\mathbf{t}) = n^{1/2}[G_n(\mathbf{t}) - G(\mathbf{t})], \quad \mathbf{t} \in E^p, n \geq 1.$$

For every $n \geq 1$, the process W_n belongs to the space $D^p[0, 1]$ of all real valued

Received October 1972; revised February 2, 1973.

¹ Research sponsored by the Aerospace Research Laboratories, Air Force Systems Command, U.S. Air Force, Contract F33615-71-C-1927. Reproduction in whole or in part permitted for any purpose of the U.S. Government.

AMS 1970 subject classifications. 60F05, 60B10.

Key words and phrases. $D^p[0, 1]$ space, empirical processes, Gaussian process, random sample size, Skorokhod J_1 -topology, tightness, weak convergence.

functions on E^p with no discontinuities of the second kind, and with $D^p[0, 1]$, we associate the (extended) Skorokhod J_1 -topology. For excellent expositions of weak convergence of processes on $D^p[0, 1]$, we may refer to [1], [5], [12]. Also, for $p = 1$, a detailed account is given in Billingsley (1968).

When the X_i are independent and identically distributed (i.i.d.), i.e., $\phi_n = 0$ for $n \geq 1$, W_n converges in distribution (in the J_1 -topology on $D^p[0, 1]$) to an appropriate Gaussian process ([1], [5]) (for $p = 1$, Brownian bridge). For ϕ -mixing processes and $p = 1$, weak convergence of W_n to an appropriate Gaussian function has been studied by Billingsley ((1968) page 197) and Sen (1971). Our first objective is to show that for general $p \geq 1$, the weak convergence of W_n to an appropriate Gaussian function holds under identical conditions.

For i.i.d. random variables, Pyke (1968) has studied the weak convergence of empirical processes for random sample sizes. Related results for multidimensional empirical processes are treated in [1], [11], [13]. Our second objective is to extend these results for ϕ -mixing processes.

For i.i.d. random vectors, Kiefer (1961) has obtained an exponential bound for the tail probability of the Kolmogorov–Smirnov statistics. Bounds for the tail probability of the Kolmogorov–Smirnov statistics for multivariate ϕ -mixing processes are studied in the last section of the paper.

2. Weak convergence of empirical processes. Let us write

$$(2.1) \quad A_k(\phi) = \sum_{n=0}^{\infty} (n + 1)^k \phi_n^{\frac{1}{2}} \quad \text{and} \quad A_k(\phi^2) = \sum_{n=0}^{\infty} (n + 1)^k \phi_n, \quad k \geq 0.$$

Then, $[A_k(\phi) < \infty] \Rightarrow [A_q(\phi) < \infty], \forall q \leq k$, and $[A_q(\phi^2) < \infty], \forall q \leq 2k$. Consider now a p -dimensional Gaussian process $W = \{W(t), t \in E^p\}$, where $EW(t) = 0, t \in E^p$, and for every $s, t \in E^p$,

$$(2.2) \quad \begin{aligned} \gamma(s, t) &= E[W(s)W(t)] = E\{[c(s - Y_1)c(t - Y_1)] - G(s)G(t)\} \\ &+ \sum_{k=2}^{\infty} E\{c(s - Y_1)c(t - Y_k) + c(s - Y_k)c(t - Y_1) \\ &- 2G(s)G(t)\}. \end{aligned}$$

Note that by Theorem 20.1 of Billingsley ((1968) page 174), the series on the right hand side (rhs) of (2.2) converges when $A_0(\phi) < \infty$. The following theorem extends Theorem 22.1 of Billingsley ((1968) page 197) to the case of $p \geq 1$ under the conditions of Sen (1971).

THEOREM 2.1. *Under (1.1) and $A_1(\phi) < \infty$, W_n converges in law (in the Skorokhod J_1 -topology on $D^p[0, 1]$) to a Gaussian process W for which (2.2) holds.*

PROOF. We need to establish

- (a) the convergence of the finite dimensional distributions of $\{W_n\}$ to those of W , and
- (b) the tightness of $\{W_n\}$.

Under $A_0(\phi) < \infty$ and (1.1), the proof of (a) follows along the same line as in

Billingsley ((1968) page 197), so that in the remaining of the proof, we only establish the tightness of $\{W_n\}$. Also, it is understood that $p \geq 2$. Since, the usual techniques for proving the tightness of multiparameter stochastic processes [viz., Neuhaus (1971) and Bickel and Wichura (1971)] are not directly applicable for ϕ -mixing processes, we adapt a modified approach.

For general ϕ -mixing processes, it has been observed in Sen (1972a) that empirical processes behave quite smoothly for large n . This, along with the treatment of the univariate case in Billingsley ((1968) pages 198–199), suggests the following approach. First, using the basic inequality between the moduli of continuity for $C^p[0, 1]$ and $D^p[0, 1]$ spaces [cf. Billingsley (1968) page 110 and Neuhaus (1971) page 1288] and the fact that $W_n(\mathbf{t}) = 0$ when at least one coordinate of \mathbf{t} is 0, it suffices to show that for every $\varepsilon > 0$ and $\eta > 0$, there exist a $\delta > 0$ and an integer n_0 , such that

$$(2.3) \quad P\{\omega_\delta(W_n) > \varepsilon\} < \eta, \quad n \geq n_0,$$

where for every $0 < \delta < 1$ and $n \geq 1$,

$$(2.4) \quad \omega_\delta(W_n) = \sup \{|W_n(\mathbf{t}) - W_n(\mathbf{s})| : \mathbf{s}, \mathbf{t} \in E^p \text{ and } |\mathbf{t} - \mathbf{s}| < \delta\}.$$

Second, by a direct multiparameter extension of Theorem 8.3 of Billingsley ((1968) page 56), it suffices to show that for every $\mathbf{0} \leq \mathbf{b}_0 \leq \mathbf{1}$, $\varepsilon > 0$ and $\eta > 0$, there exist a $\delta: 0 < \delta < 1$ and an integer n_0 , such that for $B = \{\mathbf{t}: \mathbf{b}_0 \leq \mathbf{t} \leq \mathbf{b}_0 + \delta \mathbf{1}\}$ and $n \geq n_0$,

$$(2.5) \quad P\{\sup_{\mathbf{t} \in B} |W_n(\mathbf{t}) - W_n(\mathbf{b}_0)| > \varepsilon\} < \eta[\mu(B) + \delta^p]/2,$$

where $\mu(B) = P\{\mathbf{Y}_1 \in B\}$. [Note that $[\mu(A) + \|A\|]/2$ (where $\|A\|$ is the Lebesgue measure of A) is ≤ 1 for every $A \in E^p$, and as $\mu(A)$ is bounded from above by any side of A , we have $\delta^p \leq \mu(B) + \delta^p \leq \delta + \delta^p \rightarrow 0$ as $\delta \rightarrow 0$.]

For a given $\varepsilon > 0$ and $\delta > 0$ (to be chosen later on), select n_0 so large that $\delta > n_0^{-1/2} \varepsilon/2p$. Let then (for $n \geq n_0$),

$$(2.6) \quad \mathbf{b}_n(\mathbf{i}) = \mathbf{b}_0 + (n^{-1/2} \varepsilon/2p) \mathbf{i}, \quad \mathbf{0} \leq \mathbf{i} \leq \mathbf{m}_n;$$

$$(2.7) \quad \mathbf{m}_n = m_n \cdot \mathbf{1} \quad \text{and} \quad m_n = [2p\delta n^{1/2}/\varepsilon] + 1 (\geq 1).$$

Also, let

$$(2.8) \quad B(\mathbf{i}, n) = \{\mathbf{t}: \mathbf{b}_n(\mathbf{i}) \leq \mathbf{t} \leq \mathbf{b}_n(\mathbf{i} + \mathbf{1})\}, \quad \mathbf{i} \geq \mathbf{0}.$$

Note that by (1.2) and (2.6), for every $\mathbf{i} \geq \mathbf{0}$,

$$(2.9) \quad n^{1/2} [G(\mathbf{b}_n(\mathbf{i} + \mathbf{1})) - G(\mathbf{b}_n(\mathbf{i}))] \leq \sum_{j=1}^p n^{1/2} [G_{[j]}(i_j + 1) - G_{[j]}(i_j)] \\ = \sum_{j=1}^p (\varepsilon/2p) = \varepsilon/2.$$

Hence, on using the fact that for $\mathbf{t} \in B(\mathbf{i}, n)$, $G_n(\mathbf{b}_n(\mathbf{i})) \leq G_n(\mathbf{t}) \leq G_n(\mathbf{b}_n(\mathbf{i} + \mathbf{1}))$ and $G(\mathbf{b}_n(\mathbf{i})) \leq G(\mathbf{t}) \leq G(\mathbf{b}_n(\mathbf{i} + \mathbf{1}))$, we obtain by (1.4), (2.9) and a few routine steps [as in Billingsley (1968) page 199] that

$$(2.10) \quad \sup_{\mathbf{t} \in B} |W_n(\mathbf{t}) - W_n(\mathbf{b}_0)| \leq \max_{\mathbf{0} \leq \mathbf{i} \leq \mathbf{m}_n} |W_n(\mathbf{b}_n(\mathbf{i})) - W_n(\mathbf{b}_0)| + \varepsilon/2.$$

Thus, it suffices to show that for every $\varepsilon > 0$ and $\eta > 0$, there exist a $\delta: 0 < \delta < 1$ and an integer n_0 , such that for $n \geq n_0$ and every $\mathbf{b}_0 \in E^p$.

$$(2.11) \quad P\{\max_{0 \leq i \leq m_n} |W_n(\mathbf{b}_n(\mathbf{i})) - W_n(\mathbf{b}_0)| > \frac{1}{2}\varepsilon\} < \frac{1}{2}\eta[\mu(B) + \delta^p].$$

Now, by (1.4) and (2.6), for every $\mathbf{i} \geq \mathbf{0}$,

$$(2.12) \quad W_n(\mathbf{b}_n(\mathbf{i} + \mathbf{1})) - W_n(\mathbf{b}_0) = \sum_{0 \leq j \leq i} V_n(B(\mathbf{j}, n));$$

$$(2.13) \quad V_n(B) = n^{\frac{1}{2}}[\mu_n(B) - \mu(B)]; \quad \mu_n(B) = [\# \text{ of } \mathbf{Y}_i \in B, i = 1, \dots, n]/n,$$

for every $B \in E^p$, where by Lemma 2.1 of Sen (1971), under $A_1(\phi) < \infty$,

$$(2.14) \quad E[V_n^4(B(\mathbf{i}, n))] \leq K_\phi[\mu^2(B(\mathbf{i}, n)) + n^{-1}\mu(B(\mathbf{i}, n))].$$

Unfortunately $\mu(B(\mathbf{i}, n))$, though bounded from above by $e/2n^{\frac{1}{2}}$, can be arbitrarily close to 0. For example, if $G(\mathbf{t})$ is degenerate on a lower dimensional space, then $\mu(B)$ may be equal to 0 for some $B \in E^p$. To overcome this difficulty, we define

$$(2.15) \quad \lambda_n(\mathbf{i}) = \max\{\mu(B(\mathbf{i}, n)), (\varepsilon/2pn^{\frac{1}{2}})^p\}, \quad \forall \mathbf{i} \geq \mathbf{0}.$$

Now, (2.15) implies that $\lambda_n(\mathbf{i}) \geq (\varepsilon/2pn^{\frac{1}{2}})^p$ i.e., $(2p/\varepsilon)^2 \lambda_n^{2/p}(\mathbf{i}) \geq n^{-\frac{1}{2}}$. Hence, from (2.14) we have under $A_1(\phi) < \infty$,

$$(2.16) \quad E[V_n^4(B(\mathbf{i}, n))] \leq K_\phi\{\lambda_n^2(\mathbf{i}) + (2p/\varepsilon)^2[\lambda_n(\mathbf{i})]^{1+2/p}\} \leq K_{\phi,\varepsilon}[\lambda_n(\mathbf{i})]^\beta,$$

where

$$(2.17) \quad \beta = 1 + 2/p > 1 \quad \text{and} \quad K_{\phi,\varepsilon} \leq K_\phi\{1 + (2p/\varepsilon)^2\} < \infty, \quad \forall \varepsilon > 0.$$

By virtue of (2.17) and Theorem 1 of Bickel and Wichura (1971), we obtain that for every $\varepsilon > 0$.

$$(2.18) \quad P\{\max_{0 \leq i \leq m_n-1} |S_n(\mathbf{i})| > \frac{1}{2}\varepsilon\} \leq (16K_{\phi,\varepsilon}^*/\varepsilon^4)(\sum_{0 \leq i \leq m_n-1} \lambda_n(\mathbf{i}))^\beta,$$

where $K_{\phi,\varepsilon}^* (< \infty)$ depends on ε through $K_{\phi,\varepsilon}$ in (2.17). Now, $\lambda_n(\mathbf{i}) \leq \mu(B(\mathbf{i}, n)) + (\varepsilon/2pn^{\frac{1}{2}})^p$, $\forall \mathbf{i} \geq \mathbf{0}$, so that the rhs of (2.18) is bounded by

$$(2.19) \quad (16K_{\phi,\varepsilon}^*/\varepsilon^4)[\mu(B_n) + \delta_n^p]^\beta, \quad \beta = 1 + 2/p > 1,$$

where $B_n = \{\mathbf{t}: \mathbf{b}_0 \leq \mathbf{t} \leq \mathbf{b}_0 + (\varepsilon/2pn^{\frac{1}{2}})\mathbf{m}_n\}$ and $\delta_n = m_n(\varepsilon/2pn^{\frac{1}{2}})$. By (2.6), $0 \leq \mu(B_n) - \mu(B) \leq \varepsilon/2n^{\frac{1}{2}}$ and $\delta \leq \delta_n \leq \delta + \varepsilon/2pn^{\frac{1}{2}}$. Thus, using the fact that $\delta_n^p \leq \mu(B_n) + \delta_n^p \leq \delta_n + \delta_n^p$, we obtain that for every $\eta > 0$, there exist a $\delta > 0$ and an n_0 , such that

$$(2.20) \quad \mu(B_n) + \delta_n^p \leq \frac{3}{2}(\mu(B) + \delta^p), \quad n \geq n_0,$$

$$(2.21) \quad (16K_{\phi,\varepsilon}^*/\varepsilon^4)[\mu(B_n) + \delta_n^p]^{2/p} < \eta/3, \quad n \geq n_0,$$

which completes the proof of (2.11). Hence the proof of Theorem 2.1 is complete.

Consider now a sequence of stochastic processes $\{W_n^* = [W_n^*(\mathbf{t}, u), \mathbf{t} \in E^p, 0 \leq u \leq 1]; n \geq 1\}$, defined on the $D^{p+1}[0, 1]$ space, where

$$(2.22) \quad W_n^*(\mathbf{t}, u) = [nu]^{\frac{1}{2}}W_{[nu]}(\mathbf{t})/n^{\frac{1}{2}}, \quad 0 \leq u \leq 1 \text{ and } \mathbf{t} \in E^p,$$

[s] being the largest integer contained in s . Also, let $W^* = [W^*(\mathbf{t}, u), \mathbf{t} \in E^p, 0 \leq u \leq 1]$ be a Gaussian function with $EW^*(\mathbf{t}, u) = 0$ and

$$(2.23) \quad E[W^*(\mathbf{s}, v)W^*(\mathbf{t}, u)] = \min(u, v)\gamma(\mathbf{s}, \mathbf{t}), \quad \text{for every } \mathbf{s}, \mathbf{t} \in E^p, \\ 0 \leq u, v \leq 1,$$

where $\gamma(\mathbf{s}, \mathbf{t})$ is defined by (2.2). Then, we have the following theorem.

THEOREM 2.2. *Under (1.1) and $A_1(\phi) < \infty$, W_n^* converges in law (in the extended Skorokhod J_1 -topology on $D^{p+1}[0, 1]$ space) to W^* for which (2.23) holds.*

PROOF. Here also, the convergence of the finite dimensional laws of W_n^* to those of W^* poses no problem, so we shall only prove the tightness of $\{W_n^*\}$. As in (2.3)–(2.5), it suffices to show that for every $\mathbf{b}_0 \in E^p, 0 \leq u_0 \leq 1, \varepsilon > 0$ and $\eta > 0$, there exist a $\delta: 0 < \delta < 1$ and an integer n_0 , such that for $B^* = \{(\mathbf{t}, u): \mathbf{b}_0 \leq \mathbf{t} \leq (\mathbf{b}_0 + \delta \mathbf{1}) \wedge \mathbf{1}, u_0 \leq u \leq (u_0 + \delta) \wedge 1\}$ and $n \geq n_0$,

$$(2.24) \quad P[\sup_{(\mathbf{t}, u) \in B^*} |W_n^*(\mathbf{t}, u) - W_n^*(\mathbf{b}_0, u_0)| > \varepsilon] < \delta\eta[\mu(B) + \delta^p]/2,$$

where $B = \{\mathbf{t}: \mathbf{b}_0 \leq \mathbf{t} \leq (\mathbf{b}_0 + \delta \mathbf{1}) \wedge \mathbf{1}\}, \mu(B) = P\{Y_1 \in B\}$ and $\mathbf{a} \wedge \mathbf{b} = (\min(a_1, b_1), \dots, \min(a_p, b_p))$. Let now $k_n = [(\varepsilon/4)n^{\frac{1}{2}}] + 1$, and

$$(2.25) \quad b_n^*(\mathbf{i}, j) = (\mathbf{b}_0, u_0) + ((\varepsilon/4pn^{\frac{1}{2}})\mathbf{i}, n^{-\frac{1}{2}}jk_n), \quad \mathbf{0} \leq \mathbf{i} \leq \mathbf{m}_n^*, 0 \leq j \leq m_n^{**},$$

where $\mathbf{m}_n^* = m_n^* \mathbf{1}, m_n^* = [4p\delta n^{\frac{1}{2}}/\varepsilon] + 1$ and $m_n^{**} = \max\{j: jn^{-1}k_n \leq \delta\} + 1$. Then, by the same technique as in (2.10), we have by a few standard steps that

$$(2.26) \quad \sup_{(\mathbf{t}, u) \in B^*} |W_n^*(\mathbf{t}, u) - W_n^*(\mathbf{b}_0, u_0)| \\ \leq \max_{1 \leq j \leq m_n^{**}} \max_{\mathbf{0} \leq \mathbf{i} \leq \mathbf{m}_n^*} |W_n^*(b_n^*(\mathbf{i}, j)) - W_n^*(\mathbf{b}_0, u_0)| + \varepsilon/2,$$

where by Lemma 2.1 of Sen (1971), under $A_1(\phi) < \infty$,

$$(2.27) \quad E[W_n^*(b_n^*(\mathbf{i}, j)) - W_n^*(\mathbf{b}_0, u_0)]^4 \\ \leq K_\phi\{(n^{-1}jk_n)^2\mu^2(B(\mathbf{i}, n)) + n^{-2}jk_n\mu(B(\mathbf{i}, n))\}, \quad K_\phi < \infty,$$

and $B(\mathbf{i}, n)$ is defined by (2.8). Repeating then the steps (2.14) through (2.21) and replacing $\lambda_n(\mathbf{i})$ by

$$(2.28) \quad \lambda_n(\mathbf{i}, j) = (n^{-1}jk_n)\lambda_n(\mathbf{i}) \quad \text{for } 0 \leq j \leq m_n^{**}, \quad \mathbf{0} \leq \mathbf{i} \leq \mathbf{m}_n^*,$$

the proof of (2.24) follows along the same line as of the proof of (2.11). Hence, the details are omitted. \square

Let now $\{N_\nu, \nu \geq 1\}$ be a sequence of positive integer valued random variables such that as $\nu \rightarrow \infty, \nu^{-1}N_\nu \rightarrow \xi$, in probability, where ξ is a positive random variable defined on the same probability space (Ω, \mathcal{A}, P) . Then, by virtue of Theorem 2.2, we conclude that $\{W_{N_\nu}\}$ converges weakly to W as $\nu \rightarrow \infty$.

3. Tail probability inequalities for Kolmogorov–Smirnov statistics. For i.i.d. stochastic $p(\geq 1)$ vectors, Kiefer (1961) has shown that for every $\varepsilon > 0$, there exists a positive $c_p(\varepsilon)$, such that for every $n > 1$ and $\lambda > 0$.

$$(3.1) \quad P[\sup_{\mathbf{t} \in E^p} |W_n(\mathbf{t})| > \lambda] \leq c_p(\varepsilon)\{\exp[-(2 - \varepsilon)\lambda^2]\}.$$

For ϕ -mixing processes, such a strong result is not known. We provide here with certain alternative bounds depending on $A_k(\phi) < \infty$ for some positive k . For this, we consider first the following lemma which extends Lemma 2.1 of Sen (1971) to general $k \geq 1$. Let $\{T_i = T(\mathbf{X}_i), -\infty < i < \infty\}$ be stationary mixing such that (1.1) holds and

$$(3.2) \quad \begin{aligned} ET = 0, \quad ET_i^2 = \tau; \quad 0 \leq \tau \leq 1, \\ P\{|T_i| > 1\} = 0 \quad \text{and} \quad E|T_i| \leq c\tau, \quad 0 < c < \infty. \end{aligned}$$

Let then, $S_n = T_1 + \dots + T_n$ for $n \geq 1$.

LEMMA 3.1. *Under (1.1) and (3.2) if $A_k(\phi) < \infty$ for some $k(\geq 1)$, then for every $n \geq 1$,*

$$(3.3) \quad E(S_n^{2(k+1)}) \leq K_\phi \{n\tau + \dots + (n\tau)^{k+1}\}, \quad K_\phi < \infty,$$

where K_ϕ depends only on $\{\phi_n\}$ and c .

PROOF. Note that for $k \geq 0$,

$$(3.4) \quad E(S_n^{2(k+1)}) \leq [(2k+2)!]n \sum_{n,2k+1} |E(T_1 T_{i_1} \dots T_{i_{2k+1}})|$$

where the summation $\sum_{n,2k+1}$ extends over all $1 \leq i_1 \leq \dots \leq i_{2k+1} \leq n$. Also, note that if ξ and η be $\mathcal{M}_{-\infty}^k$ and \mathcal{M}_{k+n}^∞ measurable, $E|\xi| < \infty$ and $P\{|\eta| > 1\} = 0$, then [cf. Billingsley (1968) page 171]

$$(3.5) \quad |E(\xi\eta) - E(\xi)E(\eta)| \leq 2\phi_n E|\xi|, \quad \forall n.$$

Proceeding as in the proof of Lemma 2.1 of Sen (1971) and using (3.5), we obtain that if $A_0(\phi) < \infty$, under (1.1) and (3.2),

$$(3.6) \quad n \sum_{n,1} |E(T_1 T_{i_1})| \leq [2cA_0(\phi^2)](n\tau),$$

and if $A_1(\phi) < \infty$ [$\Rightarrow A_2(\phi^2) < \infty$], then

$$(3.7) \quad n \sum_{n,2} |E(T_1 T_{i_1} T_{i_2})| \leq 6nc\tau \sum_{i=0}^{n-1} (i+1)^2 \phi_i < [6cA_2(\phi^2)](n\tau),$$

$$(3.8) \quad n \sum_{n,3} |E(T_1 T_{i_1} \dots T_{i_3})| \leq K_\phi [n\tau + (n\tau)^2], \quad K_\phi < \infty,$$

Let us now assume that for $1 \leq a \leq 2k-1$, $k \geq 1$, $n \geq 1$,

$$(3.9) \quad n \sum_{n,a} |E(T_1 T_{i_1} \dots T_{i_a})| \leq K_{\phi,a} \{n\tau + \dots + (n\tau)^{a*}\}, \quad K_{\phi,a} < \infty,$$

where $a^* = t$ for $a = 2t$ or $2t-1$, $t \geq 1$. Then, we shall show that $A_k(\phi) < \infty$ implies that (3.9) also holds for $a = 2k$ and $2k+1$. We consider only the case of $a = 2k+1$ (as the other case follows similarly). For this, we let $i_0 = 1$, $i_j = i_{j-1} + r_j$, $r_j \geq 0$, $1 \leq j \leq 2k+1$, and let $\sum_{n,2k+1}^{(j)}$ be the summation over all $1 \leq i_1 \leq \dots \leq i_{2k+1} \leq n$ for which $r_j = \max\{r_1, \dots, r_{2k+1}\}$, for $j = 1, \dots, 2k+1$. Then

$$(3.10) \quad n \sum_{n,2k+1} |E(T_1 T_{i_1} \dots T_{i_{2k+1}})| \leq \sum_{j=1}^{2k+1} \{n \sum_{n,2k+1}^{(j)} |E(T_1 \dots T_{i_{2k+1}})|\},$$

where by (3.2) and (3.5), for each j : $1 \leq j \leq 2k+1$,

$$(3.11) \quad \begin{aligned} n \sum_{n,2k+1}^{(j)} |E(T_1 \dots T_{i_{2k+1}})| \leq n \sum_{n,2k+1}^{(j)} |E(T_1 \dots T_{i_{j-1}})E(T_{i_j} \dots T_{i_{2k+1}})| \\ + 2n \sum_{n,2k+1}^{(j)} \phi_{r_j} E|T_1 \dots T_{i_{j-1}}|, \end{aligned}$$

and the second term on the rhs of (3.11) is bounded by

$$(3.12) \quad 2nE|T_1| \sum_{n,2k+1}^{(j)} \phi_{r_j} \leq 2nc\tau \sum_{r_j=0}^{n-1} (r_j + 1)^{2k} \phi_{r_j} < 2nc\tau(\sum_{i=0}^{\infty} (i + 1)^{2k} \phi_i) = [2cA_{2k}(\phi^2)](n\tau),$$

and $A_{2k}(\phi^2) < \infty$. For $j = 1$ or $2k + 1$, the first term on the rhs of (3.11) vanishes [by (3.2)], while for $2 \leq j \leq 2k$, we have

$$(3.13) \quad n \sum_{n,2k+1}^{(j)} |E(T_1 - T_{i_{j-1}})E(T_{i_j} \cdots T_{i_{2k+1}})| \leq n \sum_{i_j=1}^n \{ \sum_{i_j, j-1} |E(T_1 \cdots T_{i_{j-1}})| \} \times \{ \sum_{n-i_j+1, 2k+1-j} |E(T_{i'_0} \cdots T_{i'_{2k+1-j}})| \},$$

where $i'_l = i_{j+l} - i_j + 1, l = 0, \dots, 2k + 1 - j$. Since for $2 \leq j \leq 2k, 1 \leq j - 1, 2k + 1 - j \leq 2k - 1$, and by assumption, (3.9) holds for $a \leq 2k - 1$, we obtain from (3.9), (3.13) and the inequality that for $a \geq 0, b \geq 0, \sum_{i=1}^n i^a(n - i + 1)^b \leq c(n + 1)^{a+b+1} \leq c^*n^{a+b+1}, c^* < \infty$, that the rhs of (3.13) is bounded by

$$(3.14) \quad K_{\phi, j}[n\tau + \cdots + (n\tau)^{k^*}], \quad K_{\phi, j} < \infty; \quad k^* = k, \quad j = \text{odd}, \\ = k + 1, \quad j = \text{even}.$$

Thus, from (3.10) through (3.14), we conclude that (3.9) holds for $a = 2k + 1$. Using then (3.6)—(3.8), the proof for general $k \geq 1$ follows by the method of induction.

THEOREM 3.2. Under (1.1) and $A_k(\phi) < \infty$ for some $k \geq 1$,

$$(3.15) \quad \sup_n P\{\sup_{t \in E} |W_n(t)| > \lambda\} \leq C_{\phi} \lambda^{-2(k+1)}, \quad \forall \lambda \geq 1,$$

where $C_{\phi} (< \infty)$ depends on $\{\phi_n\}$.

PROOF. Virtually, we repeat the steps in (2.5) through (2.19) with the following changes: (i) in (2.6), (2.7) and (2.15), we let $\varepsilon = \delta = 1$, (ii) in (2.14), with the aid of Lemma 3.1, we use the moment of order $2(k + 1)$, and (iii) in (2.18), we take $\varepsilon = \lambda, \lambda \geq 1$. Then, the corresponding bound in (2.19) appears to be

$$(3.16) \quad [2^{2(k+1)} K_{\phi, 1}^*][1 + (1 + 1/pn^{\beta})^{\beta} \lambda^{-2(k+1)}], K_{\phi, 1}^* \leq K_{\phi}[1 + 4p^2] < \infty,$$

where $1 < \beta \leq \max(k + 1, 1 + 2k/p)$, so that (3.15) follows. \square

Note that (3.15) implies that under (1.1) and $A_k(\phi) < \infty$ for some $k \geq 1$,

$$(3.17) \quad \sup_n E\{\sup_{t \in E^p} |W_n(t)|^{2k+\delta}\} < \infty, \quad \forall 0 \leq \delta < 1.$$

Acknowledgment. The author is grateful to the referee for his useful comments on the manuscript.

REFERENCES

[1] BICKEL, P. J. and WICHURA, M. J. (1971). Convergence criteria for multiparameter stochastic processes and some applications. *Ann. Math. Statist.* **42** 1656-1670.
 [2] BILLINGSLEY, P. (1968). *Convergence of Probability Measures*. Wiley, New York.

- [3] BLUM, J. R., HANSON, D. L., and ROSENBLATT, J. I. (1963). On the central limit theorem for the sum of a random number of independent random variables. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **1** 389–393.
- [4] KIEFER, J. (1961). On large deviations of the empirics D.F. of vector chance variables and a law of iterated logarithm. *Pacific J. Math.* **11** 649–660.
- [5] NEUHAUS, G. (1971). On weak convergence of stochastic processes with multi-dimensional time parameter. *Ann. Math. Statist.* **42** 1285–1295.
- [6] PYKE, R. (1968). The weak convergence of empirical process with random sample sizes. *Proc. Cambridge Philos. Soc.* **64** 155–160.
- [7] RÉNYI, A. (1958). On mixing sequences of sets. *Acta Math. Acad. Sci. Hungar.* **9** 215–228.
- [8] SEN, P. K. (1971). A note on weak convergence of empirical processes for sequences of ϕ -mixing random variables. *Ann. Math. Statist.* **42** 2131–2133.
- [9] SEN, P. K. (1972a). On the Bahadur representation of sample quantiles for sequences of ϕ -mixing random variables. *J. Mult. Anal.* **2** 77–95.
- [10] SEN, P. K. (1972 b). Limiting behaviour of regular functionals of empirical distributions for stationary \ast -mixing processes. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **25** 71–82.
- [11] SEN, P. K. (1973). On weak convergence of empirical processes for random number of independent stochastic vectors. *Proc. Cambridge Philos. Soc.* **73** 139–145.
- [12] STRAF, M. L. (1972). Weak convergence of stochastic processes with several parameters. *Proc. Sixth Berkeley Symp. Math. Statist. Prob.* **2** 187–222.
- [13] WICHURA, M. J. (1968). On the weak convergence of non-Borel probabilities on a metric space. Ph. D. dissertation, Columbia Univ.

DEPARTMENT OF BIostatISTICS
UNIVERSITY OF NORTH CAROLINA
CHAPEL HILL, NORTH CAROLINA 27514