

## NATIVE FRESHWATER FISH

# Whitebait can't jump

Cindy Baker

Richard Allibone

*Fish-pass design needs to take into account the abilities of the fish species likely to be using these structures to get past obstacles in rivers.*

Most New Zealanders are aware of the annual spring migration of swarms of tiny fish commonly known as whitebait. It is these migrations that prompt many a hopeful angler to sit with nets set on the margins of river mouths for hours at a time. The fish that do not end up on our dinner tables face the task of looking for freshwater habitat to grow and produce the offspring that may, next year, be part of our whitebait fritters.

During this upstream journey the whitebait must surmount any dams, culverts, weirs and other in-stream man-made structures that might block their passage. If the obstacles are unsurpassable, whitebait distributions could become restricted and this could ultimately lead to a decline in adult stocks.

### Climbers and non-climbers

Many people are unaware that the whitebait catch contains five different species of fish. Some are good climbers and can scale the wetted margins of waterfalls, rapids and spillways. On the other hand, inanga – the main whitebait species – cannot climb and must swim past such obstacles. To do this, they swim in bursts to get past high-velocity areas and then rest in low-velocity areas. During floods or high flows, many obstacles become drowned allowing the passage of non-climbing fish. However, at low or moderate stream flows, swimming fish may be unable to get past. It is not uncommon to see fish such as inanga congregating in large numbers below obstacles. Here they are vulnerable to high mortality from predation, competition and disease.

### Are some weirs easier to pass than others?


In a recent study, we looked at the effects of weirs of different shapes and heights on the upstream passage of inanga.

Three different weir shapes were evaluated: the vee-notch, square notch and circular notch. Each weir shape was tested with a vertical drop of 5, 10 and 15 cm and a constant flow rate of 3.5 l/s. For each trial 50 fish were given 90 minutes to pass the weir. Four trials were undertaken for each vertical drop and weir shape.

The shape of the weir did not affect the ability of inanga to pass (top graph, below). Instead, the vertical drop was the determining factor: as the vertical drop was increased, fewer inanga passed the weir.

The size of the fish did not significantly influence their ability to pass the weir (lower graph, below). As no size effect was found relative to vertical drop, this suggests that, given more time, more fish would eventually pass the weir at the greater heights.


Another factor affecting passage appeared to be motivation. We found that when fish were



Proportion of inanga passing the weirs of different shapes with different vertical drops.

**Teachers:** this article can be used for Bio L8 A.O. 8.1. See other curriculum connections at [www.niwa.co.nz/pubs/wa/resources](http://www.niwa.co.nz/pubs/wa/resources)

*Cindy Baker is based at NIWA in Hamilton; Richard Allibone was formerly with NIWA in Hamilton and is now with the Department of Conservation in Wellington.*




Proportion of inanga of different sizes found at the base of the weir and passing the weir for **A** 15-cm vertical drop and **B** 5-cm vertical drop. Data from all notch shapes are combined; for each vertical drop there were 50 fish per 90 min trial over 12 trials.

freshly caught and placed below the weir, only a few fish would swim past. After we held fish in live bins for 24 hours, their motivation increased. Thus, there was a marked increase in the number of fish passing the weir during the experimental period. One explanation could be reduced stress. In the later experiments, the fish had had a day to recover from the stress of catching, handling and transportation.

Inanga that successfully passed the weir did so by burst swimming – as shown right. Fish that jumped out of the water in an attempt to pass the weir were unsuccessful (right, bottom photograph).

We also checked the effect of changing flows on the ability of inanga to pass the weir. Using the square notch weir only (since notch shape was not a crucial factor) and a vertical drop of 10 cm, we found that doubling the flow resulted in significantly fewer fish passing the weir.

Our experiments showed that inanga are capable of passing a weir with a 15-cm vertical drop. However, these fish are hindered by increasing flow and increasing drop height. Thus, whitebait can't jump but they sure can swim! This finding has implications for the construction of flow-monitoring weirs to ensure migratory fish passage. The results of experiments such as these can be used to determine the design of weirs in particular situations. An example is given on pages 25–26 in this issue. ■



Fish attempting to pass the vee-notch weir with a 10-cm fall.

**A** Sequence of frames showing an inanga swimming over the weir; each frame 1/25 s.

**B** An unsuccessful attempt by an inanga to pass the weir by jumping from below.

## Recent publications by NIWA staff

The following list includes papers in refereed journals, books and book chapters reported between November 2001 and January 2002. Please note that NIWA staff papers appear in a range of journals and are not published by NIWA. Your local library will be able to obtain copies through the interloan system if required.

### Climatology / Atmospheric sciences

**Bodeker, G.E.; Scott, J.C.; Kreher, K.; McKenzie, R.L.** (2001). Global ozone trends in potential vorticity coordinates using TOMS and GOME intercompared against the Dobson network, 1978–1998. *Journal of Geophysical Research* 106(D19): 23,029–23,042.

**Clarkson, T.C.; Wood, S.W.; Fisher, G.W.** (2001). The state of the Ross Sea region atmosphere. In: Waterhouse, E.J. (ed.). *Ross Sea region 2001: a state of the environment report for the Ross Sea region of Antarctica*, pp. 6.1–6.26. Antarctica New Zealand, Christchurch.

**Fisher, G.W.** (2001). Air quality. In: Waterhouse, E. (ed.). *Ross Sea region State of the Environment Report*, pp. 6.14–16.26. Antarctica New Zealand, Christchurch.

**Harris, C.; Given, D.; Bassett, J.; Patrick, M.; Wood, S.** (2001). Key pressures on the Ross Sea region environment. In: Waterhouse, E.J. (ed.). *Ross Sea region 2001: a state of the environment report for the Ross Sea region of Antarctica*, pp. 3.1–3.63. Antarctica New Zealand, Christchurch.

**Irie, H.; Koike, M.; Kondo, Y.; Bodeker, G.E.; Danilin, M.Y.; Sasano, Y.** (2001). Redistribution of nitric acid in the Arctic

lower stratosphere during the winter of 1996–1997. *Journal of Geophysical Research* 106(D19): 23,139–23,150.

**McKenzie, R.L.; Seckmeyer, G.; Bais, A.; Madronich, S.** (2001). Satellite retrievals of erythemal UV dose compared with ground-based measurements at Northern and Southern mid-latitudes. *Journal of Geophysical Research* 106(D20): 24,051–24,062.

**Reid, S.; Turner, R.** (2001). Correlations of real and model wind speeds in different terrains. *Weather and Forecasting* 16: 620–627.

**Revell, M.J.; Kidson, J.W.; Kiladis, G.N.** (2001). Interpreting low-frequency modes of Southern Hemisphere atmospheric variability as the rotational response to divergent forcing. *Monthly Weather Review* 129: 2416–2425.

**Salinger, M.J.** (2001). Climate variation in New Zealand and the southwest Pacific. In: Sturman, A.P.; Spronken-Smith, R.A. (eds). *The physical environment, a New Zealand perspective*, pp. 130–149. Oxford University Press, Melbourne.

**Salinger, M.J.; Griffiths, G.M.** (2001). Trends in New Zealand daily temperature and rainfall extremes. *International Journal of Climatology* 21: 1437–1452.

**Salinger, M.J.; Renwick, J.A.; Mullan, A.B.** (2001). Interdecadal Pacific Oscillation

and South Pacific climate. *International Journal of Climatology* 21: 1705–1722.

### Fresh water

**Broad, T.L.; Townsend, C.R.; Closs, G.; Jellyman, D.J.** (2001). Microhabitat use by longfin eels in New Zealand streams with contrasting riparian vegetation. *Journal of Fish Biology* 59: 1385–1400.

**Fenwick, G.D.** (2001). *Paracrangonyx* Stebbing, 1899, a genus of New Zealand subterranean amphipods (Crustacea: Amphipoda: Gammaridea). *Journal of the Royal Society of New Zealand* 31(3): 457–479.

**Glova, G.J.; Jellyman, D.J.; Bonnett, M.L.** (2001). Spatiotemporal variation in the distribution of eel (*Anguilla* spp.) populations in three New Zealand lowland streams. *Ecology of Freshwater Fish* 10: 147–153.

**Jowett, I.G.** (2001). Effect of floods and droughts on fish in a New Zealand gravel-bed river. In: Mosley, M.P. (ed.). *Gravel-bed rivers V*, pp. 451–464. New Zealand Hydrological Society, Wellington.

**Jowett, I.G.; Boustead, N.C.** (2001). Effects of substrate and sedimentation on the abundance of upland bullies (*Gobiomorphus breviceps*). *New Zealand Journal of Marine and Freshwater Research* 35: 605–613.

**Kinnison, M.T.; Unwin, M.J.; Hendry, A.P.; Quinn, T.P.** (2001). Migratory costs and the evolution of egg size and number allocation in new and indigenous salmon populations. *Evolution* 55: 1656–1667.

**McDowall, R.M.** (2001). Diadromy, divergence and diversity: implications for speciation processes in fishes. *Fish and Fisheries* 2: 278–286.

**McDowall, R.M.** (2001). Getting the measure of freshwater fish habitat in New Zealand. *Aquatic Ecosystem Health: Research and Management* 4: 343–355.

**McDowall, R.M.** (2001). The principal caudal fin ray count: a fundamental character in the galaxioid fishes. *New Zealand Journal of Zoology* 28: 395–405.

**Nagels, J.W.; Davies-Colley, R.J.; Donnison, A.M.; Muirhead, R.W.** (2001). Faecal contamination over flood events in a pastoral catchment in New Zealand. In: Tsuno, H. (ed.). *Proceedings of the Fifth International Symposium on Waste Management Problems in Agro-Industries: IWA AGRO-2001*, Shiga, Japan, pp. 33–40. Research Centre for Environmental Quality Control, Kyoto University.

**Purdy, K.J.; Hawes, I.; Bryant, C.L.; Fallick, A.E.; Nedwell, D.B.** (2001). Estimates of sulphate reduction rates in Lake Vanda, Antarctica, support the proposed recent history of the lake. *Antarctic Science* 13(4): 393–399.

continued on page 28