

Article

Whole-Genome Characterisation of ESBL-Producing *E. coli* Isolated from Drinking Water and Dog Faeces from Rural Andean Households in Peru

Maria Luisa Medina-Pizzali ¹, Apoorva Venkatesh ^{2,3}, Maribel Riveros ⁴, Diego Cuicapuza ⁵, Gabriela Salmon-Mulanovich ^{1,6}, Daniel Mäusezahl ^{1,2,3} and Stella M. Hartinger ^{1,2,3,*}

- ¹ Research Unit in Integral Development, Environment and Health, School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Lima 15102, Peru; maria.medina.p@upch.pe (M.L.M.-P.); gsalmonm@pucp.edu.pe (G.S.-M.); daniel.maeusezahl@unibas.ch (D.M.)
- ² Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, 4123 Allschwil, Switzerland; apoorva.venkatesh@swisstph.ch
- ³ University of Basel, 4001 Basel, Switzerland
- ⁴ School of Medicine, Universidad Peruana Cayetano Heredia, Lima 15102, Peru; maribel.riveros@upch.pe
- ⁵ Laboratory of Microbial Genomics, Department of Cellular and Molecular Sciences, School of Science and Philosophy, Universidad Peruana Cayetano Heredia, Lima 15102, Peru; diego.cuicapuza@upch.pe
- ⁶ Institute for Earth, Nature and Energy, Pontificia Universidad Católica del Perú, Lima 15088, Peru
- * Correspondence: stella.hartinger.p@upch.pe



Citation: Medina-Pizzali, M.L.; Venkatesh, A.; Riveros, M.; Cuicapuza, D.; Salmon-Mulanovich, G.; Mäusezahl, D.; Hartinger, S.M. Whole-Genome Characterisation of ESBL-Producing *E. coli* Isolated from Drinking Water and Dog Faeces from Rural Andean Households in Peru. *Antibiotics* **2022**, *11*, 692. <https://doi.org/10.3390/antibiotics11050692>

Academic Editor: Jie Fu

Received: 2 April 2022

Accepted: 11 May 2022

Published: 20 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: *E. coli* that produce extended-spectrum β -lactamases (ESBLs) are major multidrug-resistant bacteria. In Peru, only a few reports have characterised the whole genome of ESBL enterobacteria. We aimed to confirm the identity and antimicrobial resistance (AMR) profile of two ESBL isolates from dog faeces and drinking water of rural Andean households and determine serotype, phylogroup, sequence type (ST)/clonal complex (CC), pathogenicity, virulence genes, ESBL genes, and their plasmids. To confirm the identity and AMR profiles, we used the VITEK[®]2 system. Whole-genome sequencing (WGS) and bioinformatics analysis were performed subsequently. Both isolates were identified as *E. coli*, with serotypes -:H46 and O9:H10, phylogroups E and A, and ST/CC 5259/- and 227/10, respectively. The isolates were ESBL-producing, carbapenem-resistant, and not harbouring carbapenemase-encoding genes. Isolate 1143 ST5259 harboured the *astA* gene, encoding the EAST₁ heat-stable toxin. Both genomes carried ESBL genes (*bla*_{EC-15}, *bla*_{CTX-M-8}, and *bla*_{CTX-M-55}). Nine plasmids were detected, namely IncR, IncFIC(FII), IncI, IncFIB(AP001918), Col(pHAD28), IncFII, IncFII(pHN7A8), IncI1, and IncFIB(AP001918). Finding these potentially pathogenic bacteria is worrisome given their sources and highlights the importance of One-Health research efforts in remote Andean communities.

Keywords: phylogenomic analysis; one health; ESBL-producing *E. coli*; carbapenem resistance; whole-genome sequencing

1. Introduction

Antimicrobial resistance (AMR)—particularly in the *Enterobacteriaceae* family—has become a problem of great relevance worldwide due to its increasing prevalence, and the emergence of multiple-drug-resistant strains AMR can affect everyone, irrespective of age. It is estimated that, in 2019, 4.95 million people died from illnesses associated with bacterial AMR. Of those, 1.27 million deaths, mostly in low- and middle-income countries (LMICs), were directly attributable to bacterial AMR [1]. Enterobacteria are the most important etiological agents of serious hospital-acquired and community-onset bacterial infections in humans [2–5]. In South America, resistance to β -lactam antibiotics and fluoroquinolones is a major problem when treating enterobacterial infections [6]. In addition, reports of the emergence of colistin resistance in this region have been recently published [7,8]. South

American countries show some of the highest rates of AMR in *Enterobacteriaceae* worldwide. Huge socioeconomic differences, broad access to antimicrobials, and ineffective health systems and sanitation problems favour the emergence and spread of resistant bacterial strains [6]. In this region, antibiotic use is widespread in human medicine and animal production; thus antibiotic residues and resistant enterobacteria and AMR genes have been detected in soil, water from different environs, agricultural products (produce), and livestock. In rural Latin America, anthropogenic activities such as animal husbandry, fish farming, and agriculture were identified as drivers for antimicrobial resistance dissemination [9]. The most frequent animal contributions were the carriage and/or transfer of antimicrobial resistance determinants, the inadequate or unregulated use of veterinary antimicrobials, and the spread of resistant bacteria throughout the food supply chain via foods of animal origin [10]. Water was the most frequently identified among the environmental contributors to AMR spread in Andean Peru [9,11,12].

β -Lactamases are the leading cause of resistance to β -lactam antibiotics, and β -lactamase production is the primary mechanism of antibiotic resistance in *Escherichia coli* [13]. Among β -lactamases, extended-spectrum β -lactamases (ESBLs) that mediate resistance to all penicillins, third-generation cephalosporins, and aztreonam are of utmost importance, given that they substantially reduce the available antibiotic treatment options [14]. ESBLs include the extended-spectrum TEM-, SHV-, OXA-, and cefotaxime (CTX)-M type enzymes [15], while *E. coli* that produce CTX-Ms are most commonly associated with ESBLs [16]. AmpC β -lactamases—which function as cephalosporinases—are also clinically important. They hydrolyse cephamycins and other extended-spectrum cephalosporins and are poorly inhibited by clavulanic acid; in some cases, they confer resistance to carbapenems [17].

In Peru, previous studies have investigated ESBLs in enterobacteria isolated from clinical, community, and peri-urban settings. However, few reports have characterised ESBL-producing enterobacteria at the molecular level, and if so, mostly in clinical settings [18–22]. Using molecular methods, we previously characterised ESBL-producing enterobacteria from rural Andean communities [12,23]. However, literature on genomic characterisation of ESBL bacteria in Peru is very scarce [24–26]. The present study is the first report of the whole-genome characterisation of ESBL-producing *E. coli* isolates from water and farm/companion animals in a rural Andean setting in Peru. This study aims to confirm the identity and AMR profile and determine the serotype, phylogroup, sequence type (ST) and clonal complex (CC), pathogenicity, virulence genes, ESBL genes, and type of plasmids for ESBL-producing *E. coli* isolates.

2. Materials and Methods

2.1. Study Setting and Design

Hartinger et al. studied the dissemination pathways of AMR in humans, animals, and the environment in the Cajamarca Region in the northern highlands of Peru. The cross-sectional study design and sampling scheme are described elsewhere [23]. In brief, they sampled 40 households. Their study collected two rectal or cloacal swabs of fresh stool samples from a domestic animal (dog, cat) and/or a farm animal (cow, pig, fowl) from each household ($n = 80$). Additionally, two water samples per household were collected, one from the household's drinking water (DW) source and/or one from the household's primary water source ($n = 80$). The two ESBL-positive isolates analysed in the present study originated from two out of 40 households sampled. They were isolated from one water sample and one dog faeces sample, as shown in Supplementary Materials Figure S1.

2.2. Households' Characteristics

Hartinger et al. found only one ESBL isolate from a dog and an ESBL isolate from the DW source from two homes [23]. In these households, animals were allowed to roam freely in the kitchen and courtyard, and their faeces were found in both areas. Both homes used

piped water from the public water network system but had no sewage system, as they had installed latrines instead.

2.3. Microbiological Identification and Antimicrobial Susceptibility Testing

The microbiological identification and the antimicrobial susceptibility testing were performed using a VITEK[®] 2 Compact automated system (bioMérieux, Marcy l'Etoile, France). A colorimetric reagent card (GN) to identify the most significant fermenting and non-fermenting Gram-negative bacilli was used according to the manufacturer's instructions. AST-N249 cards were prepared according to the manufacturer's instructions. The minimum inhibitory concentrations for the following antimicrobials were recorded (ampicillin/sulbactam (SAM), piperacillin/tazobactam (TZP), cefazolin (CFZ), cefuroxime (CXM), cefotaxime (CTX), ceftazidime (CAZ), cefepime (FEP), ertapenem (ETP), imipenem (IPM), meropenem (MEM), amikacin (AMK), gentamicin (GEN), ciprofloxacin (CIP), colistin (CST), and tigecycline (TGC). Previously [23], antimicrobial susceptibility testing for other antimicrobials has been performed using commercial discs [27]; *Escherichia coli* ATCC 25922 was used as a reference strain for quality assurance of the susceptibility testing of *E. coli*.

2.4. Phenotypical and Molecular Identification of ESBL-Producing Ability

The phenotypical ESBL-producing ability of the isolates was determined using the Jarlier method [28] for the following antibiotics: aztreonam (5 µg disk), ceftazidime (30 µg disk), cefotaxime (30 µg disk), ceftriaxone (30 µg disk), amoxicillin/clavulanic acid (30 µg disk) and cefepime (30 µg disk), and confirmed by the combined disc method [27]. Further molecular confirmation was performed using conventional polymerase chain reaction PCR. These procedures have been described elsewhere [23]. Briefly, genes encoding TEM-, SHV-, OXA-, and CTX-M-type β-lactamases (*bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA}, *bla*_{CTX-M-U}, *bla*_{CTX-M-2}, *bla*_{CTX-M-3}, *bla*_{CTX-M-8}, *bla*_{CTX-M-9}, *bla*_{CTX-M-10}) were PCR-amplified using the primers shown previously [23].

2.5. Genome Sequencing and Bioinformatic Analysis

The genomic DNA was extracted from overnight cultures using the GeneJET genomic DNA purification kit (Thermo Fisher Scientific, Waltham, MA, USA), according to the manufacturer's instructions. DNA concentration was evaluated by Qubit[®] 4.0 fluorometer (Life Technologies, Carlsbad, CA, USA). The genomic library was constructed using a Nextera XT DNA Library Preparation Kit (Illumina, Inc., San Diego, CA, USA) and subsequently sequenced using a 2 × 250 paired-end library on a MiSeq platform (Illumina). The quality control of each sequence was evaluated using Fastqc v0.11.5 [29], and Trimmomatic v0.38 [30] was used to remove adapters and filter low-quality reads. The reads were assembled de novo using SPAdes v.3.15.2 [31]. The bioinformatic analysis was carried out using the tools available at the Center of Genomic Epidemiology (www.genomicepidemiology.org) (accessed on 1 October 2021), setting identity at ≥90% and coverage at ≥90%. We performed *E. coli* serotyping (SerotypeFinder v.2.0), in silico phylogenetic typing (ClermonTyping (<https://github.com/A-BN/ClermonTyping>)) (accessed on 1 January 2022), multilocus sequence typing (MLST v.2.0), pathogenicity (Pathogen Finder v.1.1), virulence genes detection (VirulenceFinder v.2.0), plasmid replicon typing (PlasmidFinder v.2.1 and MobileElementFinder v.1.0), and ESBL genes detection using ABRicate v.1.0.1 (<https://github.com/tseemann/abricate>) (accessed on 1 January 2022) NCBI and ResFinder databases. Raw Illumina reads were uploaded to GenBank under BioProject PRJNA816508. The complete outputs for all the results of the bioinformatic tools are shown in Supplementary Materials Table S2.

3. Results

3.1. Microbiological Identification and Antimicrobial Susceptibility Testing of the *E. coli* Isolates

For this analysis, only ESBL-phenotypically-positive isolates obtained in the mother study were included, one from dog faeces (Isolate 1143) and one from the household's DW

source (Isolate 1144). Isolates 1143 and 1144 were identified as *E. coli* strains with a 93% and 96% probability using VITEK[®]2 (bioMérieux, Marcy l’Etoile, France) automated reading and further confirmed by WGS.

Based on the VITEK[®]2 system, both isolates were resistant to SAM (combined effect); cephalosporins CFZ, CXM, and CTX; TZP (β -lactam/ β -lactamase inhibitor combination); and MEM (a carbapenem). However, only Isolate 1143 was resistant to important cephalosporins CAZ and FEP and the combination antibiotic SXT. On the other hand, only Isolate 1144 was resistant to GEN and CIP, while 1143 was not. We compared the VITEK[®]2 automated readings for antimicrobial susceptibility to the disk diffusion assay results, as shown in Supplementary Materials Table S1. It is noteworthy that Isolate 1144 showed resistance to FEP by the disk diffusion method, contrary to the results obtained using the automated VITEK[®]2 platform. The VITEK[®]2 system identified Isolate 1143 as an ESBL producer. Isolate 1144, however, was characterised as ESBL negative based on the MIC results, which were interpreted automatically by the system based on the VITEK[®]2 breakpoints (see Supplementary Materials Table S1).

3.2. Detection of ESBL-Encoding Genes by PCR and Whole-Genome Sequencing Analysis (WGS)

Detection of the most common β -lactamase-encoding genes (*bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA}, and *bla*_{CTX-M}) was carried out using PCR. None of the isolates showed amplification for *bla*_{SHV}, and *bla*_{OXA} genes, but both harboured *bla*_{TEM} and *bla*_{CTX-M} genes. WGS analysis using Resfinder and NCBI identified the presence of genes encoding resistance to cephalosporins (*bla*_{CTX-M-8}, *bla*_{CTX-M-55}, *bla*_{EC-15}), broad-spectrum β -lactamases (*bla*_{TEM-1}), tetracyclines (*tet*(B), *tet*(M)), aminoglycosides (*aac*(3)-*Iid*, *aph*(3’)-*Iia*, *aph*(3’)-*Ib*, *aph*(6)-*Id*, *aadA1* and *aadA2*), sulphonamides (*sul3*), phenicols (*floR*, *cmlA1*), fosfomycin (*fosA3*), quinolone (*qnrB19*), and trimethoprim (*dfrA12*), as shown in Table 1. The genotypic resistance of both isolates correlated with their phenotypical resistance, except for MEM; the resistance to this carbapenem was only phenotypically observed.

Table 1. Characterisation of the ESBL-producing *E. coli* isolates according to origin and source, sequence type and clonal complex, serotype, virulence genes, resistance phenotype and genotype, and plasmid type.

ID <i>E. coli</i> Strains	Source	ST/CC	Serotype	Phylogroup	Virulence Genes	Resistance Phenotype	Resistance Genotype	Plasmid Type
1143	dog faeces	5259/-	-:H46	E	<i>astA</i> , <i>chuA</i> , <i>cia</i> , <i>gad</i> , <i>ompT</i> , <i>terC</i>	SAM, TZP, KZ, CXM, CTX, CAZ, FEP, MEM, SXT	<i>aac</i> (3)- <i>IId</i> , <i>aph</i> (3’)- <i>Ib</i> , <i>aph</i> (6)- <i>Id</i> , <i>aadA1</i> , <i>aadA2</i> , <i>floR</i> , <i>bla</i> _{TEM-1} , <i>bla</i> _{EC-15} , <i>bla</i> _{CTX-M-8} , <i>tet</i> (M), <i>tet</i> (B) <i>dfrA12</i>	IncI1-IAAlpha, IncR, Inc- FIB(AP001918), IncFIC(FII)
1144	drinking water	227/10	O9:H10	A	<i>capU</i> , <i>cea</i> , <i>cia</i> , <i>fyuA</i> , <i>iutA</i> , <i>sitA</i> , <i>irp2</i> , <i>iroN</i> , <i>terC</i>	SAM, TZP, KZ, CXM, CTX, MEM, CN, CIP	<i>aph</i> (3’)- <i>Iia</i> , <i>cmlA1</i> , <i>bla</i> _{CTX-M-55} , <i>bla</i> _{EC-15} , <i>fosA3</i> , <i>qnrB19</i> , <i>sul3</i> , <i>tet</i> (B)	IncFIB, Col440I, IncFII, IncI1-IAAlpha, IncFII(pHN7A8)

Abbreviations: CC, clonal complex; ST, sequence type. Resistance phenotype: SAM, ampicillin/sulbactam; TZP, piperacillin/tazobactam; KZ, ceftazidime; CXM, cefuroxime; CTX, cefotaxime; CAZ, ceftazidime; FEP, cefepime; MEM, meropenem; SXT, trimethoprim/sulfamethoxazole; CN, gentamicin; CIP, ciprofloxacin.

Genomic analysis using MLST 2.0 indicated that the *E. coli* Isolates 1143 and 1144 belonged to ST5259 and ST227, respectively, and 1144 belonged to CC 10. According to the genomes’ analysis using the ClermontTyping tool, the 1143 and 1144 isolates corresponded to the phylogenetic groups E and A, respectively. Using SerotypeFinder v.2.0, Isolate 1143 was classified as serotype -:H46, whereas Isolate 1144 belonged to serotype O9:H10, as presented in Table 1.

3.3. Pathogenicity, Virulence Genes, and Type of Plasmids

Using Pathogenfinder 1.1 and VirulenceFinder 2.0, both isolates were identified as likely human pathogens, but Shiga-toxin genes were not detected in their genomes. A

total of 28 different virulence genes were detected from the *E. coli* genomes. The colicin *ia* (*cia*) virulence gene was detected in both *E. coli* isolates. Other prevalent virulent genes included: *astA*, *iroN*, *chuA*, *gad*, *ompT*, *capU*, *cea*, *fyuA*, *sitA*, *irp2*, *iutA*, and *terC*. The *astA* gene (gene product: EAST₁ heat-stable toxin) was found in Isolate 1143 only.

Using the PlasmidFinder 2.1 and the MobileElementFinder 1.0 tools, four plasmids were detected for Isolate 1143, IncR (harbouring the *aph(3'')*-Ib and *aph(6)*-Id aminoglycoside resistance genes), IncFIC(FII) (harbouring the florfenicol/chloramphenicol resistance *floR* gene), IncI1 (carrying the *cia* virulence gene), and IncFIB(AP001918). On the other hand, for Isolate 1144, five plasmids were identified, Col(pHAD28) (harbouring the *qnrB19* ciprofloxacin resistance gene), IncFII, IncFII(pHN7A8), IncI1, and IncFIB(AP001918).

4. Discussion

We confirmed the identity of the two enterobacteria isolates (1143 and 1144) as *E. coli*; Isolate 1143—obtained from dog faeces—was an atypical non-lactose fermenter. Lactose-negative uropathogenic *E. coli* strains have been recently isolated from dogs in Brazil [32], and they cause urinary tract infections in humans [33]. Further, lactose-negative *E. coli* are usually enteroinvasive (EIEC), are closely related to *Shigella* spp., and produce a dysenteric form of diarrhoea in humans [34]. Both isolates were likely human pathogens, with serotypes -:H46 and O9:H10, respectively. The *E. coli* O9:H10 serotype has been listed among the typical enteroaggregative (EAEC) *E. coli* serotypes in Brazil [35], and the same serotype was identified among samples of Shiga-toxin-producing *E. coli* (STEC) and Enteroaggregative *E. coli* (EAEC) from Mexico [36].

Both isolates were confirmed as ESBL producers given their genomes encoded ESBL genes (*bla*_{CTX-M-8}, *bla*_{CTX-M-55}, *bla*_{EC-15}), contrary to the VITEK[®]2 automatised system's classification of Isolate 1144 as a non-ESBL producer. The VITEK[®]2 ESBL test has a varying sensitivity for detecting ESBL-producing enterobacteria correctly; it ranges between 86% and 98.1% [37,38]. The VITEK[®]2 AST N249 card used in our study identified Isolate 1143 as an ESBL producer, but not Isolate 1144, since it was not resistant to FEP and CAZ. However, Isolate 1144 showed a phenotypical resistance to other oxyimino cephalosporins CXM and CTX.

Isolate 1143 and Isolate 1144 belonged to phylogroups E and A, respectively. Phylogroup E includes the O157:H7 EHEC lineage, which is only a small subset of the whole genetic diversity found within the group. Phylogroup-E strains are rarely isolated compared to other phylogroup strains. They have been isolated from animals, the environment, and humans, and virulent/resistant strains are linked to humans [39]. Thus, finding an ESBL-producing, potentially pathogenic, phylogroup-E isolate in dog faeces points to a probable transmission route from humans. On the other hand, most phylogroup-A strains are human commensals and can be found in wastewater [40] and animals, especially poultry [41]. Notably, in our setting, backyard chicken farming is widespread (83% of households), and farm animals, in general, are administered antibiotics by the owners without prescription nor technical supervision (unpublished data from our research group). Other studies carried out in rural locations in Peru show similar findings [42,43]. Chickens roam freely, and household members are in contact with their animals and their scattered faeces. On the other hand, the piped water supply and the soil near the houses were contaminated with faecal coliforms originating from the environment, humans, or animals [23]. The fact that both isolates in this study displayed resistance to carbapenems is of great concern, given that this “third line” antimicrobial medication is exclusively for human use [44]. Thus, the ESBL/AmpC carbapenem-resistant *E. coli* isolates found in DW and dog faeces could originate from other farm animals or humans and be transmitted by different pathways, as shown by Hartinger et al. in our study area [23].

Isolates 1143 and 1144 belonged to ST/CC 5259/- and 227/10, respectively. We found that *E. coli* ST227 are considerably common and have been frequently reported around the world. In the public database Enterobase (<https://enterobase.warwick.ac.uk>) (accessed on 1 February 2022), we found 84 publicly available *E. coli* ST227 genome assemblies from

human, animal, and environmental sources. In Tunisia, an ST227 carbapenem-resistant *E. coli* isolate of human origin was reported in 2016, carrying the *bla*_{OXA-48}, *bla*_{CTX-M-15}, *bla*_{TEM}, and *bla*_{OXA-1} genes [45]. A carbapenem-resistant *E. coli* isolate of the ST227 lineage was isolated from clinical samples in Lebanon, carrying the *bla*_{OXA-48} and the *bla*_{TEM-1} genes [46]. *E. coli* ST5259 are scarce; only two were found in EnteroBase, one of them originated from poultry in Ecuador and the other from humans in the United States of America. In China, an *E. coli* ST5259 genome of human origin was mentioned as part of BioProject PRJNA400107, being reported along with other *E. coli* genomes as carriers of the *mcr-1* gene, which is a plasmid-mediated colistin resistance gene [47].

Many of the identified virulence genes were associated with iron acquisition systems, and a small part was associated with toxin production, metal resistance (tellurite), and other virulence determinants, as shown in Supplementary Materials Table S2. The *cia* gene—involved in killing other bacteria—was found in both isolates. The iron uptake genes detected were: *iroN*, *chuA*, *fyuA*, *iutA*, and *sitA*. The *chuA* gene encodes an outer membrane hemin receptor, and the *fyuA* gene encodes the yersiniabactin receptor, and both contribute most during infection of the urinary tract by uropathogenic *E. coli* [48]. The *chuA* gene was detected in Isolate 1143, whereas the *fyuA* gene was found in Isolate 1144 along with the other iron acquisition genes. This flags Isolate 1144 as a possible uropathogenic strain, given it harboured many genes that enable iron uptake, best suited for iron-deprived environments such as human urine [49]. On the other hand, Isolate 1143 harboured the *astA* gene, encoding the EAST₁ heat-stable toxin, which is associated with enteroaggregative *E. coli* (EAEC) in humans and enterotoxigenic *E. coli* (ETEC) in porcines [50]. None of the isolates' genomes harboured Shiga-toxin genes.

The *bla*_{CTX-M-55} is one of the most abundant ESBL genes in the *Enterobacteriaceae* family, with a rising prevalence in *E. coli* from livestock and humans, especially in China [51,52]. It is usually carried by plasmids, but it has also been found chromosomally [52]. In Canada, enterobacteria isolated from turkeys carried the *bla*_{CTX-M-55} gene mediated by *IncF* plasmids [53]. In Japan, the *bla*_{CTX-M-8} gene was found in *E. coli* from humans and retail chicken meat, and its transmission was associated with *IncI1* plasmids [54].

Unlike *bla*_{CTX-M-55} and *bla*_{CTX-M-8}, which code for Class A ESBL, the *bla*_{EC-15} gene product belongs to the Class C β-lactamases [55], also known as AmpC-type β-lactamases. In comparison to ESBL (including Class A β-lactamases), AmpC-type β-lactamases hydrolyse broad- and extended-spectrum cephalosporins (cephamycins as well as oxymino-β-lactams) but are not inhibited by clavulanic acid or other β-lactamase inhibitors. EC β-lactamases—encoded in *bla*_{EC} genes—are a specific type of AmpC β-lactamases found in *E. coli* [56]. In a recent One-Health study in Canada, the *bla*_{EC-15} gene was detected in *E. coli* isolates from different human, animal, and environmental sources. It did not occur in all isolates, having a prevalence of 16% [57]. The fact that both isolates in our study carried genes for Class A ESBL and AmpC β-lactamases has great clinical importance, given that AmpC/ESBL *E. coli* cannot be treated with clavulanic acid, which complicates the treatment [58]. However, we did not include a cephalosporin/clavulanic acid combination in the disc diffusion assay, so resistance to clavulanic acid was not phenotypically confirmed. Further, both isolates expressed resistance to other antibiotics such as fluoroquinolones and a carbapenem, narrowing treatment options even further. According to Guzmán-Blanco et al. [59], in Latin America, the rates of nosocomial infection caused by ESBL-producing enterobacteria—especially CTX-M enzyme producers—are higher than in other regions. A One-Health-based study in Brazil—the biggest country in the region—showed that ESBL-producing *E. coli* carrying a diversity of *bla*_{CTX-M} gene variants and *mcr-1* genes are endemic across their territory at the interface between humans and animals [60]. ESBL-producing enterobacteria can also be resistant to cefepime and exhibit resistance to fluoroquinolones and ampicillin/sulbactam, as well as to aminoglycosides and piperacillin/tazobactam. Fortunately, more than 90% of ESBL-producing enterobacteria are still susceptible to carbapenems [59], although reports of carbapenem-resistant enterobacteria alarmingly increase in Latin American countries [61].

Despite the phenotypic resistance to meropenem of both isolates in our study, we did not find evidence of well-known carbapenemase-encoding genes—*bla*_{KPC}, *bla*_{NDM}, *bla*_{NDM}, *bla*_{IMP}, *bla*_{VIM} and *bla*_{OXA-48}—or their variants [62]. Thus, non-carbapenemase mechanisms could be suspected for the isolates in our study. For instance, point mutations in porin genes reduce membrane permeability to carbapenems, in combination with hyperproduction of the β -lactamase AmpC or ESBL. Therefore, the isolates could be characterised as non-carbapenemase, carbapenem-resistant *E. coli* [63,64]. In Singapore, the emergence of this type of bacteria in clinical settings was suggested to result from selective antibiotic pressure along with *de novo* mutations or genetic reassortment in carbapenem-sensitive enterobacteria [65].

Plasmids play a major role in the dissemination of AMR genes in Gram-negative bacteria. Usually, they harbour multiple physically connected genetic determinants, conferring resistance to different classes of antibiotics simultaneously, such as extended-spectrum β -lactams, carbapenems aminoglycosides, sulfonamides, and quinolones [66,67]. Plasmid families, including IncF, IncI1, IncI2, IncX, IncA/C, and IncHI2, play an important part in ESBL gene spread [67]. It is noteworthy that both *E. coli* genomes carried IncI1 plasmid. Our findings show that for Isolate 1143, some AMR genes were harboured in plasmids or other mobile genetic elements, e.g., aminoglycoside resistance genes in the IncR plasmid and the *bla*_{TEM-1} gene (resistance to ampicillins) were found to be carried in the Tn2 transposon (<https://transposon.lstmed.ac.uk/tn-registry>) (accessed on 1 January 2022). Conversely, for Isolate 1144, the ciprofloxacin resistance gene (*qnrB19*) was carried by Col440I; this is a plasmid recently associated with enterobacteria-carrying ESBL and carbapenemase genes isolated from environment and humans in South Africa [68]. In our analysis, we found that many AMR genes appeared to be inserted in the chromosome—such as *bla*_{CTX-M-55} and *bla*_{CTX-M-8}—pointing to the mobility of these genes from mobile genetic elements to the chromosome and vice versa and between different plasmids [69].

Our study has some limitations. The presence of AMR genes harboured in plasmids points to horizontal transfer of AMR determinants. However, conjugation assays should be performed for these isolates to confirm the transfer of genes via plasmids to other bacteria. In addition, isolates were not tested to phenotypically confirm AmpC, pathogenicity, and/or EAST₁ toxin production. Due to laboratory limitations and restrictions, it was impossible to perform these confirmatory assays. Another limitation is the low number of isolates available for characterisation, which may have limited the scope of detection of important resistance/virulence genes and plasmids.

5. Conclusions

Isolates 1143 and 1144 were identified as ESBL-producing *E. coli*, with serotypes -:H46 and O9:H10; phylogroups E and A, and ST/CC 5259/- and 227/10, respectively. They were characterised as potential human pathogens, and they carried multiple virulence genes; Isolate 1143 ST5259 harboured the *astA* gene, encoding the EAST₁ heat-stable toxin. Both genomes were carriers of a *bla*_{EC-15} gene (AmpC), displaying carbapenem resistance, but not harbouring carbapenemase genes. Other ESBL genes (*bla*_{CTX-M-8} and *bla*_{CTX-M-55}) among various AMR genes were detected, mainly located in plasmids or the chromosome. *E. coli* of the ST227 lineage were reported in other countries, while *E. coli* ST5259 reports were rare. Both isolates were found in a remote area in the highlands of Peru, which is of public health concern considering the likely anthropogenic origin derived from incorrect and often unrestricted use of antibiotics in both humans and livestock. Our results can help identify and track *E. coli* strains that pose a risk to human, animal, and environmental health in rural Andean communities. The *E. coli* isolates originated from an animal and DW, highlighting the importance of comprehensive research for preventive action along the One-Health continuum in isolated Andean communities. The sharing of living spaces between humans and animals and the use of contaminated DW could facilitate the transfer of these pathogens to all environs in the community. Thus, new research paths to limit

the spread of AMR should focus on the epidemiology of ESBL/AmpC *E. coli*, particularly carbapenem-resistant strains.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/antibiotics11050692/s1>, Figure S1: Obtention of the *E. coli* isolates 1143 and 1144; Table S1: Antibiotic susceptibility^a profile for *E. coli* Isolates 1143 and 1144 determined by the VITEK[®]2 system and compared to the disk diffusion assay results^b; Table S2: Outputs of bioinformatic tools.

Author Contributions: Conceptualisation, S.M.H., M.R. and D.C.; methodology, M.R. and D.C.; bioinformatic analysis, D.C. and M.L.M.-P.; visualisation, D.C.; writing—original draft preparation, M.L.M.-P., A.V. and D.C.; writing—review and editing, M.L.M.-P., A.V., D.C., M.R., S.M.H., G.S.-M. and D.M.; supervision, D.M., S.M.H. and G.S.-M.; funding acquisition, S.M.H. and D.M. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Novartis Foundation (Grant Number 18A059), and D.C. was supported by a scholarship from Emerge, the Emerging Diseases Epidemiology Research Training D43 TW007393 training grant awarded by the Fogarty International Center of the US National Institutes of Health.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Institutional Review Board for humans and animal subjects of the Universidad Peruana Cayetano Heredia, Protocol Codes 418-16-18 and 010-03-20.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The raw reads and assembled data have been submitted to National Center for Biotechnology Information (NCBI) under BioProject Number PRJNA816508.

Acknowledgments: We thank Pablo Tsukayama for his comments on the manuscript.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Murray, C.J.; Ikuta, K.S.; Sharara, F.; Swetschinski, L.; Robles Aguilar, G.; Gray, A.; Han, C.; Bisignano, C.; Rao, P.; Wool, E.; et al. Global Burden of Bacterial Antimicrobial Resistance in 2019: A Systematic Analysis. *Lancet* **2022**, *399*, 629–655. [[CrossRef](#)]
2. D’Agostino, M.; Cook, N. Foodborne Pathogens. In *Encyclopedia of Food and Health*; Caballero, B., Finglas, P., Toldrá, F., Eds.; Academic Press: Cambridge, MA, USA, 2015; pp. 83–86.
3. Guentzel, M.N. *Escherichia*, *Klebsiella*, *Enterobacter*, *Serratia*, *Citrobacter*, and *Proteus*. In *Medical Microbiology*; Baron, S., Ed.; University of Texas Medical Branch at Galveston: Galveston, TX, USA, 1996.
4. Larsson, D.G.J.; Flach, C.-F. Antibiotic Resistance in the Environment. *Nat. Rev. Microbiol.* **2021**, *20*, 257–269. [[CrossRef](#)] [[PubMed](#)]
5. Laxminarayan, R.; Duse, A.; Wattal, C.; Zaidi, A.K.M.; Wertheim, H.F.L.; Sumpradit, N.; Vlieghe, E.; Hara, G.L.; Gould, I.M.; Goossens, H.; et al. Antibiotic Resistance—the Need for Global Solutions. *Lancet Infect. Dis.* **2013**, *13*, 1057–1098. [[CrossRef](#)]
6. Bonelli, R.R.; Moreira, B.M.; Picão, R.C. Antimicrobial Resistance among Enterobacteriaceae in South America: History, Current Dissemination Status and Associated Socioeconomic Factors. *Drug Resist. Updat.* **2014**, *17*, 24–36. [[CrossRef](#)]
7. Mendes Oliveira, V.R.; Paiva, M.C.; Lima, W.G. Plasmid-Mediated Colistin Resistance in Latin America and Caribbean: A Systematic Review. *Travel Med. Infect. Dis.* **2019**, *31*, 101459. [[CrossRef](#)]
8. Rapoport, M.; Faccone, D.; Pasteran, F.; Ceriana, P.; Albornoz, E.; Petroni, A.; Corso, A.; Maldonado, M.L.; Lucero, C.; Amalfa, F.; et al. First Description of Mcr-1-Mediated Colistin Resistance in Human Infections Caused by *Escherichia coli* in Latin America. *Antimicrob. Agents Chemother.* **2016**, *60*, 4412–4413. [[CrossRef](#)]
9. Medina-Pizzali, M.L.; Hartinger, S.M.; Salmon-Mulanovich, G.; Larson, A.; Riveros, M.; Mäusezahl, D. Antimicrobial Resistance in Rural Settings in Latin America: A Scoping Review with a One Health Lens. *Int. J. Environ. Res. Public Health* **2021**, *18*, 9837. [[CrossRef](#)]
10. Van Boeckel, T.P.; Pires, J.; Silvester, R.; Zhao, C.; Song, J.; Criscuolo, N.G.; Gilbert, M.; Bonhoeffer, S.; Laxminarayan, R. Global Trends in Antimicrobial Resistance in Animals in Low- And Middle-Income Countries. *Science* **2019**, *365*, eaaw1944. [[CrossRef](#)]
11. Ahmad, I.; Malak, H.A.; Abulreesh, H.H. Environmental Antimicrobial Resistance and Its Drivers: A Potential Threat to Public Health. *J. Glob. Antimicrob. Resist.* **2021**, *27*, 101–111. [[CrossRef](#)]

12. Larson, A.; Hartinger, S.M.; Riveros, M.; Salmon-Mulanovich, G.; Hattendorf, J.; Verastegui, H.; Huaylinos, M.L.; Mäusezahl, D. Antibiotic-Resistant *Escherichia coli* in Drinking Water Samples from Rural Andean Households in Cajamarca, Peru. *Am. J. Trop. Med. Hyg.* **2019**, *100*, 1363–1368. [[CrossRef](#)]
13. Pitout, J.D.D. Extraintestinal Pathogenic *Escherichia coli*: An Update on Antimicrobial Resistance, Laboratory Diagnosis and Treatment. *Expert Rev. Anti Infect. Ther.* **2012**, *10*, 1165–1176. [[CrossRef](#)] [[PubMed](#)]
14. World Health Organization. *Antimicrobial Resistance: Global Report on Surveillance*; World Health Organization: Geneva, Switzerland, 2014.
15. Sawa, T.; Kooguchi, K.; Moriyama, K. Molecular Diversity of Extended-Spectrum β -Lactamases and Carbapenemases, and Antimicrobial Resistance. *J. Intensive Care* **2020**, *8*, 13. [[CrossRef](#)] [[PubMed](#)]
16. Peirano, G.; Pitout, J.D.D. Extended-Spectrum β -Lactamase-Producing Enterobacteriaceae: Update on Molecular Epidemiology and Treatment Options. *Drugs* **2019**, *79*, 1529–1541. [[CrossRef](#)] [[PubMed](#)]
17. Mohamudha, P.R.; Harish, B.N.; Parija, S.C. AmpC Beta Lactamases among Gram Negative Clinical Isolates from a Tertiary Hospital, South India. *Braz. J. Microbiol.* **2010**, *41*, 596–602. [[CrossRef](#)] [[PubMed](#)]
18. Colquechagua Aliaga, F.; Andrade Sevilla, C.; Escalante Gonzales, E. Extended-Spectrum Beta-Lactamase (ESBL)-producing Enterobacteriaceae in Fecal Samples at the National Institute of Child Health, Peru. *Rev. Peru Med. Exp. Salud Publica* **2015**, *32*, 26–32. [[CrossRef](#)] [[PubMed](#)]
19. Loyola, S.; Gutierrez, L.R.; Horna, G.; Petersen, K.; Agapito, J.; Osada, J.; Rios, P.; Lescano, A.G.; Tamariz, J. Extended-Spectrum β -Lactamase-Producing Enterobacteriaceae in Cell Phones of Health Care Workers from Peruvian Pediatric and Neonatal Intensive Care Units. *Am. J. Infect. Control* **2016**, *44*, 910–916. [[CrossRef](#)] [[PubMed](#)]
20. Garcia, C.; Hinostrroza, N.; Astocondor, L.; Ochoa, T.; Jacobs, J. Characterization of ESBL-Producing Salmonella Enterica Serovar Infantis Infection in Humans, Lima, Peru. *Am. J. Trop. Med. Hyg.* **2019**, *101*, 746–748. [[CrossRef](#)]
21. León-Luna, D.; Fajardo-Loyola, A.; Yareta-Yareta, J.; Burgos-Espejo, A.; Peralta-Siesquen, C.; Galarza-Pérez, M.; Marcos-Carbajal, P. Molecular Characterization of Multi-Resistant Enterobacteria in Two Departments of the Peruvian Jungle. *Biomedica* **2021**, *41*, 180–187. [[CrossRef](#)]
22. Benavides, J.A.; Shiva, C.; Virhuez, M.; Tello, C.; Appelgren, A.; Vendrell, J.; Solassol, J.; Godreuil, S.; Streicker, D.G. Extended-Spectrum Beta-Lactamase-Producing *Escherichia coli* in Common Vampire Bats *Desmodus Rotundus* and Livestock in Peru. *Zoonoses Public Health* **2018**, *65*, 454–458. [[CrossRef](#)]
23. Hartinger, S.M.; Medina-Pizzali, M.L.; Salmon-Mulanovich, G.; Larson, A.J.; Pinedo-Bardales, M.; Verastegui, H.; Riberos, M.; Mäusezahl, D. Antimicrobial Resistance in Humans, Animals, Water and Household Environs in Rural Andean Peru: Exploring Dissemination Pathways through the One Health Lens. *Int. J. Environ. Res. Public Health* **2021**, *18*, 4604. [[CrossRef](#)]
24. Quino, W.; Hurtado, C.V.; Escalante-Maldonado, O.; Flores-León, D.; Mestanza, O.; Vences-Rosales, F.; Zamudio, M.L.; Gavilán, R.G. Multidrug Resistance of Salmonella infantis in Peru: A Study through next Generation Sequencing. *Rev. Peru Med. Exp. Salud Publica* **2019**, *36*, 37–45. [[CrossRef](#)] [[PubMed](#)]
25. Benavides, J.A.; Godreuil, S.; Opazo-Capurro, A.; Mahamat, O.O.; Falcon, N.; Oravcova, K.; Streicker, D.G.; Shiva, C. Long-Term Maintenance of Multidrug-Resistant *Escherichia coli* Carried by Vampire Bats and Shared with Livestock in Peru. *Sci. Total Environ.* **2022**, *810*, 152045. [[CrossRef](#)] [[PubMed](#)]
26. Murray, M.; Salvatierra, G.; Dávila-Barclay, A.; Ayzanoa, B.; Castillo-Vilcahuaman, C.; Huang, M.; Pajuelo, M.J.; Lescano, A.G.; Cabrera, L.; Calderón, M.; et al. Market Chickens as a Source of Antibiotic-Resistant *Escherichia coli* in a Peri-Urban Community in Lima, Peru. *Front. Microbiol.* **2021**, *12*, 635871. [[CrossRef](#)] [[PubMed](#)]
27. Clinical Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing*, 30th ed.; Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2020.
28. Lezameta, L.; Gonzáles-Escalante, E.; Tamariz, J.H. Comparison of four phenotypic methods to detect Extended-Spectrum Betalactamases. *Rev. Peru Med. Exp. Salud Publica* **2010**, *27*, 345–351. [[CrossRef](#)] [[PubMed](#)]
29. Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data. Available online: <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/> (accessed on 28 February 2022).
30. Bolger, A.M.; Lohse, M.; Usadel, B. Trimmomatic: A Flexible Trimmer for Illumina Sequence Data. *Bioinformatics* **2014**, *30*, 2114–2120. [[CrossRef](#)] [[PubMed](#)]
31. Bankevich, A.; Nurk, S.; Antipov, D.; Gurevich, A.A.; Dvorkin, M.; Kulikov, A.S.; Lesin, V.M.; Nikolenko, S.I.; Pham, S.; Pribelski, A.D.; et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J. Comput. Biol.* **2012**, *19*, 455–477. [[CrossRef](#)] [[PubMed](#)]
32. Siqueira, F.M.; de Carli, S.; Lopes, C.E.; Machado, L.; Vieira, T.R.; Pöppl, G.; Cardoso, M.R.I.; Zaha, A. Non-Lactose-Fermenting Uropathogenic *Escherichia coli* from Dogs: Virulence Profile Characterization. *Lett. Appl. Microbiol.* **2021**, *72*, 596–603. [[CrossRef](#)]
33. Gajdács, M.; Ábrók, M.; Lázár, A.; Burián, K. Differential Epidemiology and Antibiotic Resistance of Lactose-Fermenting and Non-Fermenting *Escherichia coli*: Is It Just a Matter of Taste? *Biol. Futur.* **2020**, *71*, 175–182. [[CrossRef](#)]
34. Food and Drug Administration. Chapter 4A: Diarrheagenic *Escherichia coli*. Available online: <https://www.fda.gov/food/laboratory-methods-food/bam-chapter-4a-diarrheagenic-escherichia-coli> (accessed on 1 January 2022).
35. Ori, E.L.; Takagi, E.H.; Andrade, T.S.; Miguel, B.T.; Cergole-Novella, M.C.; Guth, B.E.C.; Hernandez, R.T.; Dias, R.C.B.; Pinheiro, S.R.S.; Camargo, C.H.; et al. Diarrhoeagenic *Escherichia coli* and *Escherichia Albertii* in Brazil: Pathotypes and Serotypes over a 6-Year Period of Surveillance. *Epidemiol. Infect.* **2019**, *147*, e10. [[CrossRef](#)]

36. Navarro, A.; van der Ploeg, C.; Rogé, A.; Licono-Moreno, D.; Delgado, G.; Morales-Espinosa, R.; Cravioto, A.; Eslava, C. Diversity of Potentially Pathogenic *Escherichia coli* O104 and O9 Serogroups Isolated before 2011 from Fecal Samples from Children from Different Geographic Regions. *Microorganisms* **2021**, *9*, 2227. [[CrossRef](#)]
37. Spanu, T.; Sanguinetti, M.; Tumbarello, M.; D'Inzeo, T.; Fiori, B.; Posteraro, B.; Santangelo, R.; Cauda, R.; Fadda, G. Evaluation of the New VITEK 2 Extended-Spectrum Beta-Lactamase (ESBL) Test for Rapid Detection of ESBL Production in Enterobacteriaceae Isolates. *J. Clin. Microbiol.* **2006**, *44*, 3257–3262. [[CrossRef](#)] [[PubMed](#)]
38. Wiegand, I.; Geiss, H.K.; Mack, D.; Stürenburg, E.; Seifert, H. Detection of Extended-Spectrum Beta-Lactamases among Enterobacteriaceae by Use of Semiautomated Microbiology Systems and Manual Detection Procedures. *J. Clin. Microbiol.* **2007**, *45*, 1167–1174. [[CrossRef](#)] [[PubMed](#)]
39. Clermont, O.; Condamine, B.; Dion, S.; Gordon, D.M.; Denamur, E. The E Phylogroup of *Escherichia coli* Is Highly Diverse and Mimics the Whole E. Coli Species Population Structure. *Environ. Microbiol.* **2021**, *23*, 7139–7151. [[CrossRef](#)] [[PubMed](#)]
40. De Castro Stoppe, N.; Silva, J.S.; Carlos, C.; Sato, M.I.Z.; Saraiva, A.M.; Ottoboni, L.M.M.; Torres, T.T. Worldwide Phylogenetic Group Patterns of *Escherichia coli* from Commensal Human and Wastewater Treatment Plant Isolates. *Front. Microbiol.* **2017**, *8*, 2512. [[CrossRef](#)]
41. Coura, F.M.; Diniz, S.D.A.; Silva, M.X.; Mussi, J.M.S.; Barbosa, S.M.; Lage, A.P.; Heinemann, M.B. Phylogenetic Group Determination of *Escherichia coli* Isolated from Animals Samples. *Sci. World J.* **2015**, *2015*, 258424. [[CrossRef](#)]
42. Alberca, V.; León, D.; Falcón, N. Keeping Backyard Animals and Evaluation of Knowledge and Practices Associated with Exposure to Zoonotic Agents in La Coipa, Cajamarca, Peru. *Rev. Inv. Vet. Perú* **2020**, *31*, e18733. [[CrossRef](#)]
43. Benavides, J.A.; Streicker, D.G.; Gonzales, M.S.; Rojas-Paniagua, E.; Shiva, C. Knowledge and Use of Antibiotics among Low-Income Small-Scale Farmers of Peru. *Prev. Vet. Med.* **2021**, *189*, 105287. [[CrossRef](#)]
44. World Organisation for Animal Health. *OIE List Of Antimicrobial Agents Of Veterinary Importance*; World Organisation for Animal Health: Paris, France, 2007.
45. Tanfous, F.B.; Alonso, C.A.; Achour, W.; Ruiz-Ripa, L.; Torres, C.; Hassen, A.B. First Description of KPC-2-Producing *Escherichia coli* and ST15 OXA-48-Positive *Klebsiella pneumoniae* in Tunisia. *Microb. Drug Resist.* **2017**, *23*, 365–375. [[CrossRef](#)]
46. Beyrouthy, R.; Robin, F.; Dabboussi, F.; Mallat, H.; Hamzé, M.; Bonnet, R. Carbapenemase and Virulence Factors of Enterobacteriaceae in North Lebanon between 2008 and 2012: Evolution via Endemic Spread of OXA-48. *J. Antimicrob. Chemother.* **2014**, *69*, 2699–2705. [[CrossRef](#)]
47. Peng, Z.; Hu, Z.; Li, Z.; Zhang, X.; Jia, C.; Li, T.; Dai, M.; Tan, C.; Xu, Z.; Wu, B.; et al. Antimicrobial Resistance and Population Genomics of Multidrug-Resistant *Escherichia coli* in Pig Farms in Mainland China. *Nat. Commun.* **2022**, *13*, 1116. [[CrossRef](#)]
48. Spurbeck, R.R.; Dinh, P.C.; Walk, S.T.; Stapleton, A.E.; Hooton, T.M.; Nolan, L.K.; Kim, K.S.; Johnson, J.R.; Mobley, H.L.T. *Escherichia coli* Isolates That Carry Vat, Fyua, Chua, and Yfcv Efficiently Colonize the Urinary Tract. *Infect. Immun.* **2012**, *80*, 4115–4122. [[CrossRef](#)] [[PubMed](#)]
49. Gonzales-Rodriguez, A.O.; Varillas, S.F.I.; Pastor, H.J.B.; Mitma, Y.L.; Canales, D.H.; Chero, P.A.W.; Gutierrez, C.; Cunza, S.S. Immunological and Biochemical Response from Older Adults with Urinary Tract Infection to Uropathogenic *Escherichia coli* Virulence Factors. *Rev. Peru Med. Exp. Salud Publica* **2020**, *37*, 527–531. [[CrossRef](#)] [[PubMed](#)]
50. Zhang, W.; Berberov, E.M.; Freeling, J.; He, D.; Moxley, R.A.; Francis, D.H. Significance of Heat-Stable and Heat-Labile Enterotoxins in Porcine Colibacillosis in an Additive Model for Pathogenicity Studies. *Infect. Immun.* **2006**, *74*, 3107–3114. [[CrossRef](#)] [[PubMed](#)]
51. Zeng, S.; Luo, J.; Li, X.; Zhuo, C.; Wu, A.; Chen, X.; Huang, L.S. Molecular Epidemiology and Characteristics of CTX-M-55 Extended-Spectrum β -Lactamase-Producing *Escherichia coli* From Guangzhou, China. *Front. Microbiol.* **2021**, *12*, 730012. [[CrossRef](#)] [[PubMed](#)]
52. Zhang, C.Z.; Ding, X.M.; Lin, X.L.; Sun, R.Y.; Lu, Y.W.; Cai, R.M.; Webber, M.A.; Ding, H.Z.; Jiang, H.X. The Emergence of Chromosomally Located BlaTX-M-55 in *Salmonella* from Foodborne Animals in China. *Front. Microbiol.* **2019**, *10*, 1268. [[CrossRef](#)] [[PubMed](#)]
53. Moffat, J.; Chalmers, G.; Reid-Smith, R.; Mulvey, M.R.; Agunos, A.; Calvert, J.; Cormier, A.; Ricker, N.; Weese, J.S.; Boerlin, P. Resistance to Extended-Spectrum Cephalosporins in *Escherichia coli* and Other Enterobacteriales from Canadian Turkeys. *PLoS ONE* **2020**, *15*, e0236442. [[CrossRef](#)]
54. Norizuki, C.; Wachino, J.I.; Suzuki, M.; Kawamura, K.; Nagano, N.; Kimura, K.; Arakawa, Y. Specific BlaCTX-M-8/IncI1 Plasmid Transfer among Genetically Diverse *Escherichia coli* Isolates between Humans and Chickens. *Antimicrob. Agents Chemother.* **2017**, *61*, e00663-17. [[CrossRef](#)]
55. National Library of Medicine (US). *Reference Gene Catalog*; National Library of Medicine: Bethesda, MD, USA, 2022.
56. Alcock, B.P.; Raphenya, A.R.; Lau, T.T.Y.; Tsang, K.K.; Bouchard, M.; Edalatmand, A.; Huynh, W.; Nguyen, A.L.V.; Cheng, A.A.; Liu, S.; et al. CARD 2020: Antibiotic Resistome Surveillance with the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Res.* **2020**, *48*, D517. [[CrossRef](#)]
57. Adator, E.H.; Walker, M.; Narvaez-Bravo, C.; Zaheer, R.; Goji, N.; Cook, S.R.; Tymensen, L.; Hannon, S.J.; Church, D.; Booker, C.W.; et al. Whole Genome Sequencing Differentiates Presumptive Extended Spectrum Beta-Lactamase Producing *Escherichia coli* along Segments of the One Health Continuum. *Microorganisms* **2020**, *8*, 448. [[CrossRef](#)]
58. Tekele, S.G.; Teklu, D.S.; Tullu, K.D.; Birru, S.K.; Legese, M.H. Extended-Spectrum Beta-Lactamase and AmpC Beta-Lactamases Producing Gram Negative Bacilli Isolated from Clinical Specimens at International Clinical Laboratories, Addis Ababa, Ethiopia. *PLoS ONE* **2020**, *15*, e0241984. [[CrossRef](#)]

59. Guzmán-Blanco, M.; Labarca, J.A.; Villegas, M.V.; Gotuzzo, E. Extended Spectrum β -Lactamase Producers among Nosocomial Enterobacteriaceae in Latin America. *Braz. J. Infect. Dis.* **2014**, *18*, 421–433. [[CrossRef](#)] [[PubMed](#)]
60. Fuga, B.; Sellera, F.P.; Cerdeira, L.; Esposito, F.; Cardoso, B.; Fontana, H.; Moura, Q.; Cardenas-Arias, A.; Sano, E.; Ribas, R.M.; et al. WHO Critical Priority *Escherichia coli* as One Health Challenge for a Post-Pandemic Scenario: Genomic Surveillance and Analysis of Current Trends in Brazil. *Microbiol. Spectr.* **2022**, *10*, 1–18. [[CrossRef](#)] [[PubMed](#)]
61. Escandón-Vargas, K.; Reyes, S.; Gutiérrez, S.; Villegas, M.V. The Epidemiology of Carbapenemases in Latin America and the Caribbean. *Expert Rev. Anti Infect. Ther.* **2017**, *15*, 277–297. [[CrossRef](#)] [[PubMed](#)]
62. Cui, X.; Zhang, H.; Du, H. Carbapenemases in Enterobacteriaceae: Detection and Antimicrobial Therapy. *Front. Microbiol.* **2019**, *10*, 1823. [[CrossRef](#)] [[PubMed](#)]
63. Centers for Disease Control and Prevention (CDC). CRE Technical Information. Available online: <https://www.cdc.gov/hai/organisms/cre/technical-info.html> (accessed on 19 February 2022).
64. Temkin, E.; Adler, A.; Lerner, A.; Carmeli, Y. Carbapenem-Resistant Enterobacteriaceae: Biology, Epidemiology, and Management. *Ann. NY Acad. Sci.* **2014**, *1323*, 22–42. [[CrossRef](#)]
65. Marimuthu, K.; Ng, O.T.; Cherng, B.P.Z.; Fong, R.K.C.; Pada, S.K.; De, P.P.; Ooi, S.T.; Smitasin, N.; Thoon, K.C.; Krishnan, P.U.; et al. Antecedent Carbapenem Exposure as a Risk Factor for Non-Carbapenemase-Producing Carbapenem-Resistant Enterobacteriaceae and Carbapenemase-Producing Enterobacteriaceae. *Antimicrob. Agents Chemother.* **2019**, *63*, e00845-19. [[CrossRef](#)]
66. Carattoli, A. Plasmids and the Spread of Resistance. *Int. J. Med. Microbiol.* **2013**, *303*, 298–304. [[CrossRef](#)]
67. Carattoli, A. Resistance Plasmid Families in Enterobacteriaceae. *Antimicrob. Agents Chemother.* **2009**, *53*, 2227–2238. [[CrossRef](#)]
68. Ramsamy, Y.; Mlisana, K.P.; Amoako, D.G.; Abia, A.L.K.; Ismail, A.; Allam, M.; Mbanga, J.; Singh, R.; Essack, S.Y. Mobile Genetic Elements-Mediated Enterobacterales-Associated Carbapenemase Antibiotic Resistance Genes Propagation between the Environment and Humans: A One Health South African Study. *Sci. Total Environ.* **2022**, *806*, 150641. [[CrossRef](#)]
69. Partridge, S.R.; Kwong, S.M.; Firth, N.; Jensen, S.O. Mobile Genetic Elements Associated with Antimicrobial Resistance. *Clin. Microbiol. Rev.* **2018**, *31*, e00088-17. [[CrossRef](#)]