

RESEARCH ARTICLE

Open Access



Whole genome sequencing and function prediction of 133 gut anaerobes isolated from chicken caecum in pure cultures

Matej Medvecký¹, Darina Cejková¹, Ondřej Polansky¹, Daniela Karasová¹, Tereza Kubasová¹, Alois Cizek^{2,3} and Ivan Rychlík^{1*}

Abstract

Background: In order to start to understand the function of individual members of gut microbiota, we cultured, sequenced and analysed bacterial anaerobes from chicken caecum.

Results: Altogether 204 isolates from chicken caecum were obtained in pure cultures using Wilkins-Chalgren anaerobe agar and anaerobic growth conditions. Genomes of all the isolates were determined using the NextSeq platform and subjected to bioinformatic analysis. Among 204 sequenced isolates we identified 133 different strains belonging to seven different phyla - *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria*, *Verrucomicrobia*, *Elusimicrobia* and *Synergistetes*. Genome sizes ranged from 1.51 Mb in *Elusimicrobium minutum* to 6.70 Mb in *Bacteroides ovatus*. Clustering based on the presence of protein coding genes showed that isolates from phyla *Proteobacteria*, *Verrucomicrobia*, *Elusimicrobia* and *Synergistetes* did not cluster with the remaining isolates. *Firmicutes* split into families *Lactobacillaceae*, *Enterococcaceae*, *Veillonellaceae* and order *Clostridiales* from which the *Clostridium perfringens* isolates formed a distinct sub-cluster. All *Bacteroidetes* isolates formed a separate cluster showing similar genetic composition in all isolates but distinct from the rest of the gut anaerobes. The majority of *Actinobacteria* clustered closely together except for the representatives of genus *Gordonibacter* showing that the genome of this genus differs from the rest of *Actinobacteria* sequenced in this study. Representatives of *Bacteroidetes* commonly encoded proteins (collagenase, hemagglutinin, hemolysin, hyaluronidase, heparinases, chondroitinase, mucin-desulfating sulfatase or glutamate decarboxylase) that may enable them to interact with their host. Aerotolerance was recorded in *Akkermansia* and *Cloacibacillus* and was also common among representatives of *Bacteroidetes*. On the other hand, *Elusimicrobium* and the majority of *Clostridiales* were highly sensitive to air exposure despite their potential for spore formation.

Conclusions: Major gut microbiota members utilise different strategies for gut colonisation. High oxygen sensitivity of *Firmicutes* may explain their commonly reported decrease after oxidative burst during gut inflammation.

Keywords: Microbiota, Microbiome, Chicken, Whole genome sequencing, Metabolic pathway, Host-microbiota interactions, Anaerobe, Butyrate, Propionate

* Correspondence: rychlik@vri.cz

¹Veterinary Research Institute, Hudcova 70, 621 00 Brno, Czech Republic
Full list of author information is available at the end of the article



Background

Characterisation of gut microbiota is nowadays relatively simple due to recent advances in next generation sequencing. However, though DNA sequencing is useful for monitoring dynamic changes in microbiota composition, it does not enable the understanding of biological functions of individual microbiota members. Shotgun sequencing of total DNA or RNA/cDNA sequencing can partly indicate the metabolic potential of microbial communities but is limited in addressing biological functions of individual microbiota members [1–3]. Even in the cases of analysis of microbial communities with low complexity when long DNA contigs can be assembled and associated with a particular bacterium, such approaches do not allow for subsequent experimental verification of observed data due to the unavailability of pure bacterial cultures. Isolation of gut anaerobes in pure cultures is therefore the best way to examine the characteristics of individual microbiota members experimentally [4].

Isolation of bacterial gut microbiota members in pure cultures is usually an issue since the majority of bacterial isolates colonising the intestinal tract are strict anaerobes. Although it is commonly reported that between 10 and 50% of bacterial species colonising the intestinal tract can be grown *in vitro*, recent studies proposed that up to 90% of major gut colonisers may be grown *in vitro* if multiple culture conditions are tested [5, 6]. Despite this, the isolation of a particular gut anaerobe may remain an issue since even the most abundant microbiota members at species level only rarely form more than 1% of the total population [7, 8]. This means that the desired bacterial species may be represented by a single colony growing on an agar plate together with hundreds or thousands of others and the likelihood of picking up the particular species is indeed rather low [6].

Chickens represent a specific case for studies focused on host - microbiota interactions. Chickens in commercial production are hatched in a clean environment of hatcheries without any contact with adult hens and their colonisation is dependent on environmental sources. Moreover, newly hatched chickens can be colonised by microbiota of wide a range of compositions [7] and colonisation of the intestinal tract of commercially hatched chickens may therefore differ from the colonisation of intestinal tract of chickens which would hatch in a nest. Perhaps not surprisingly, commercially hatched chickens are highly sensitive to colonisation with different pathogens, e.g. *Salmonella*, nevertheless, their resistance can be increased by providing them with a complex microbiota of adult hens [9, 10].

In our recent studies, we characterised the development of gut microbiota in commercially hatched chickens throughout their whole life [8], identified proteins

expressed by the main microbiota members [7] and verified that gut microbiota may protect chickens against *Salmonella* Enteritidis infection [9]. However, which bacterial species are the protective ones is not known and more detailed knowledge of individual microbiota members is needed. One way forward is to obtain well-characterised pure cultures of gut anaerobes [11]. In this study we therefore cultured hundreds of isolates of anaerobes from chicken caecum and sequenced more than 200 of them. Analysis of their genomic sequences showed that we collected isolates from 7 different phyla. The aims of the subsequent comparisons focused mainly on the representatives from phyla *Bacteroidetes* and *Firmicutes* was to reveal differences in (poly)saccharide utilisation, propionate and butyrate fermentation and interactions with the host including motility or the ability to degrade host derived proteins.

Results

Altogether 204 isolates were obtained in pure culture and sequenced. Since in several cases we sequenced isolates exhibiting higher than 99% sequence similarity between their genomes, the final number of different isolates decreased to 133 (Table 1, Fig. 1). The lowest sequencing coverage was 43 fold for *Drancourtella massiliensis* An12, the highest 1526 fold for *Lactobacillus gasseri* An197, and median coverage over all sequenced isolates was 312 fold (see Additional file 1 for all details). The isolates belonged to 7 different phyla – *Firmicutes* (84 isolates), *Bacteroidetes* (29 isolates), *Actinobacteria* (15 isolates), *Proteobacteria* (1 isolate each of *Escherichia* and *Desulfovibrio*), *Verrucomicrobia* (1 isolate of *Akkermansia*), *Elusimicrobia* (1 isolate of *Elusimicrobium*) and *Synergistetes* (1 isolate of *Cloacibacillus*). The similarity of whole sequences of 16S rRNA to the to the most homologous GenBank entries was lower than 94% for 15 isolates. Considering taxonomic recommendations [12], these isolates represent candidates for new genera and two *Muribaculum*-like isolates might belong to a novel bacterial family or even an order. Alternative analysis based on alignment of RpoB amino acid sequences [13] yielded similar clustering of individual isolates as that achieved by 16S rRNA comparison (Additional file 2).

When we compared the sequence of 16S rRNA genes of all 133 isolates with the operational taxonomic units (OTUs) combined from our previous studies [7, 8], rather unexpectedly, 7 isolates were excluded from the analysis by QIIME at the chimera removal step. Three isolates formed OTUs which we did not detect among the microbiota of the two studies. The rest of the isolates were assigned into particular OTUs. Nineteen isolates were assigned to 11 OTUs from the 100 most frequent OTUs, and out of the 1000 most common OTUs, we obtained 76 isolates belonging to 42 OTUs. Exact

Table 1 List of strains isolated, sequenced and analysed in this study

Phylum	Family	Species	ID	rRNA % sim. ^a	genome (bp)	GC (%)
Actinobacteria	Coriobacteriaceae	<i>Collinsella intestinalis</i>	An268	94	2,375,164	65.28
Actinobacteria	Coriobacteriaceae	<i>Collinsella intestinalis</i>	An7	95	2,368,937	65.33
Actinobacteria	Coriobacteriaceae	<i>Collinsella intestinalis</i>	An307	96	2,218,890	62.95
Actinobacteria	Coriobacteriaceae	<i>Collinsella tanakaei</i>	An271	96	2,790,023	64.34
Actinobacteria	Coriobacteriaceae	<i>Collinsella tanakaei</i>	An2	94	2,542,639	62.39
Actinobacteria	Coriobacteriaceae	<i>Enorma massiliensis</i>	An70	99	2,374,054	62.09
Actinobacteria	Coriobacteriaceae	<i>Enorma timonensis</i>	An5	96	2,299,978	65.99
Actinobacteria	Coriobacteriaceae	<i>Gordonibacter urolithinifaciens</i>	An234A	99	3,542,488	65.99
Actinobacteria	Coriobacteriaceae	<i>Gordonibacter urolithinifaciens</i>	An232A	94	3,457,314	65.16
Actinobacteria	Coriobacteriaceae	<i>Gordonibacter urolithinifaciens</i>	An230	94	3,885,330	64.16
Actinobacteria	Coriobacteriaceae	<i>Olsenella uli</i>	An188	96	2,138,506	68.68
Actinobacteria	Coriobacteriaceae	<i>Olsenella uli</i>	An290	96	2,158,343	68.09
Actinobacteria	Coriobacteriaceae	<i>Olsenella uli</i>	An285	96	2,413,556	66.98
Actinobacteria	Coriobacteriaceae	<i>Olsenella uli</i>	An293	96	2,318,914	67.87
Actinobacteria	Coriobacteriaceae	<i>Olsenella uli</i>	An270	96	2,014,154	67.55
Bacteroidetes	Bacteroidaceae	<i>Bacteroides clarus</i>	An43	99	4,226,284	45.22
Bacteroidetes	Bacteroidaceae	<i>Bacteroides clarus</i>	An189	99	4,166,078	45.36
Bacteroidetes	Bacteroidaceae	<i>Bacteroides dorei</i>	An16	99	5,376,103	41.79
Bacteroidetes	Bacteroidaceae	<i>Bacteroides dorei</i>	An41	99	5,463,463	42.00
Bacteroidetes	Bacteroidaceae	<i>Bacteroides ovatus</i>	An161	99	6,700,861	42.08
Bacteroidetes	Bacteroidaceae	<i>Bacteroides salanitronis</i>	An322	92	3,449,415	44.84
Bacteroidetes	Bacteroidaceae	<i>Bacteroides uniformis</i>	An67	99	4,590,834	46.36
Bacteroidetes	Bacteroidaceae	<i>Bacteroides xylanisolvens</i>	An109	99	5,713,108	41.57
Bacteroidetes	Bacteroidaceae	<i>Bacteroides xylanisolvens</i>	An107	99	5,745,201	41.89
Bacteroidetes	Bacteroidaceae	<i>Mediterranea massiliensis</i>	An20	91	3,968,548	49.41
Bacteroidetes	Porphyromonadaceae	<i>Barnesiella viscericola</i>	An22	98	3,264,150	53.04
Bacteroidetes	Porphyromonadaceae	<i>Barnesiella viscericola</i>	An55	98	3,040,531	51.37
Bacteroidetes	Porphyromonadaceae	<i>Butyricimonas paravirosa</i>	An62	98	5,176,855	42.63
Bacteroidetes	Porphyromonadaceae	<i>Muribaculum intestinale</i>	An289	83	2,336,263	49.09
Bacteroidetes	Porphyromonadaceae	<i>Muribaculum intestinale</i>	An287	83	2,349,712	49.04
Bacteroidetes	Porphyromonadaceae	<i>Odoribacter splanchnicus</i>	An45	99	4,610,398	43.75
Bacteroidetes	Porphyromonadaceae	<i>Odoribacter splanchnicus</i>	An39	99	4,722,515	43.48
Bacteroidetes	Porphyromonadaceae	<i>Parabacteroides distasonis</i>	An199	99	5,145,110	45.11
Bacteroidetes	Porphyromonadaceae	<i>Parabacteroides johnsonii</i>	An42	99	4,430,164	45.03
Bacteroidetes	Porphyromonadaceae	<i>Parabacteroides merdae</i>	An277	92	3,709,857	46.69
Bacteroidetes	Rikenellaceae	<i>Alistipes onderdonkii</i>	An90	100	3,488,443	58.52
Bacteroidetes	Rikenellaceae	<i>Alistipes senegalensis</i>	An31A	97	2,626,508	61.85
Bacteroidetes	Rikenellaceae	<i>Alistipes senegalensis</i>	An116	97	3,264,205	58.57
Bacteroidetes	Rikenellaceae	<i>Alistipes senegalensis</i>	An66	97	3,064,564	59.37
Bacteroidetes	Rikenellaceae	<i>Alistipes shahii</i>	An54	96	3,272,633	58.38
Bacteroidetes	unclassified_"Bacteroidales"	<i>Bacteroides salanitronis</i>	An269	92	4,466,522	45.87
Bacteroidetes	unclassified_"Bacteroidales"	<i>Bacteroides salanitronis</i>	An279	92	3,976,735	45.82
Bacteroidetes	unclassified_"Bacteroidales"	<i>Bacteroides salanitronis</i>	An19	92	4,779,606	45.88

Table 1 List of strains isolated, sequenced and analysed in this study (Continued)

Phylum	Family	Species	ID	rRNA % sim. ^a	genome (bp)	GC (%)
Bacteroidetes	unclassified_“Bacteroidales”	Bacteroides salanitronis	An51A	92	4,415,476	45.79
Firmicutes	Enterococcaceae	Enterococcus cecorum	An69	100	2,010,800	36.66
Firmicutes	Enterococcaceae	Enterococcus cecorum	An144	100	2,521,030	36.10
Firmicutes	Lactobacillaceae	Lactobacillus gallinarum	An119	99	2,068,702	36.50
Firmicutes	Lactobacillaceae	Lactobacillus gallinarum	An153	99	2,042,196	36.47
Firmicutes	Lactobacillaceae	Lactobacillus gallinarum	An115	99	2,039,377	36.52
Firmicutes	Lactobacillaceae	Lactobacillus gallinarum	An101	99	2,051,080	36.54
Firmicutes	Lactobacillaceae	Lactobacillus gasserii	An197	99	1,786,561	34.60
Firmicutes	Lactobacillaceae	Lactobacillus reuteri	An71	99	2,330,171	38.47
Firmicutes	Lactobacillaceae	Lactobacillus reuteri	An166	99	2,293,009	38.50
Firmicutes	Lactobacillaceae	Lactobacillus salivarius	An63	99	1,826,390	32.76
Firmicutes	Lactobacillaceae	Lactobacillus salivarius	An84	99	2,115,320	32.74
Firmicutes	Lactobacillaceae	Lactobacillus salivarius	An128	99	1,900,005	32.75
Firmicutes	Erysipelotrichaceae	[Clostridium] spiroforme	An158	99	2,749,214	28.90
Firmicutes	Erysipelotrichaceae	[Clostridium] spiroforme	An15	94	3,043,058	30.24
Firmicutes	Erysipelotrichaceae	[Clostridium] spiroforme	An26	99	2,686,825	28.58
Firmicutes	Erysipelotrichaceae	[Clostridium] spiroforme	An149	99	2,786,011	28.86
Firmicutes	Erysipelotrichaceae	[Clostridium] spiroforme	An173	94	2,963,161	30.12
Firmicutes	Erysipelotrichaceae	[Eubacterium] cylindroides	An64	99	1,822,768	34.80
Firmicutes	Erysipelotrichaceae	[Eubacterium] cylindroides	An178	99	1,979,688	34.75
Firmicutes	Erysipelotrichaceae	Massiliomicrobiota timonensis	An13	100	2,808,053	31.46
Firmicutes	Erysipelotrichaceae	Massiliomicrobiota timonensis	An142	98	2,805,108	31.51
Firmicutes	Erysipelotrichaceae	Massiliomicrobiota timonensis	An134	98	2,583,105	31.58
Firmicutes	Erysipelotrichaceae	Massiliomicrobiota timonensis	An105	98	2,657,304	31.69
Firmicutes	Erysipelotrichaceae	Massiliomicrobiota timonensis	An80	98	2,632,118	31.88
Firmicutes	Clostridiaceae 1	<i>Clostridium perfringens</i>	An68	99	3,279,733	28.07
Firmicutes	Clostridiaceae 1	<i>Clostridium perfringens</i>	An185	99	3,267,175	28.06
Firmicutes	Lachnospiraceae	[Clostridium] glycyrrhizinilyticum	An169	96	4,874,615	49.44
Firmicutes	Lachnospiraceae	[Clostridium] glycyrrhizinilyticum	An298	96	3,190,622	46.76
Firmicutes	Lachnospiraceae	[Clostridium] glycyrrhizinilyticum	An76	96	3,392,322	50.34
Firmicutes	Lachnospiraceae	[Clostridium] lactatifermentans	An114	95	2,765,027	31.21
Firmicutes	Lachnospiraceae	[Clostridium] lactatifermentans	An75	99	3,241,294	44.78
Firmicutes	Lachnospiraceae	[Clostridium] oroticum	An181	95	2,778,317	43.19
Firmicutes	Lachnospiraceae	[Clostridium] saccharolyticum	An14	95	4,319,160	54.42
Firmicutes	Lachnospiraceae	[Clostridium] saccharolyticum	An168	99	3,501,825	50.31
Firmicutes	Lachnospiraceae	[Clostridium] saccharolyticum	An196	93	3,261,385	50.05
Firmicutes	Lachnospiraceae	[Eubacterium] contortum	An118	95	3,480,620	52.25
Firmicutes	Lachnospiraceae	[Eubacterium] fissicatena	An131	94	3,529,099	50.87
Firmicutes	Lachnospiraceae	[Eubacterium] fissicatena	An138	94	3,702,317	50.44
Firmicutes	Lachnospiraceae	[Eubacterium] hallii	An3	95	4,260,545	46.43
Firmicutes	Lachnospiraceae	[Eubacterium] hallii	An11	95	3,929,296	46.78
Firmicutes	Lachnospiraceae	<i>Blautia coccoides</i>	An46	94	3,844,348	44.62
Firmicutes	Lachnospiraceae	<i>Blautia producta</i>	An81	95	4,340,741	44.56

Table 1 List of strains isolated, sequenced and analysed in this study (*Continued*)

Phylum	Family	Species	ID	rRNA % sim. ^a	genome (bp)	GC (%)
Firmicutes	Lachnospiraceae	<i>Blautia schinkii</i>	An249	93	3,969,115	45.08
Firmicutes	Lachnospiraceae	<i>Drancourtella massiliensis</i>	An210	95	3,049,918	45.07
Firmicutes	Lachnospiraceae	<i>Drancourtella massiliensis</i>	An177	95	2,986,142	44.76
Firmicutes	Lachnospiraceae	<i>Drancourtella massiliensis</i>	An57	95	3,307,621	45.72
Firmicutes	Lachnospiraceae	<i>Drancourtella massiliensis</i>	An12	96	3,637,323	46.18
Firmicutes	Ruminococcaceae	<i>Anaerofilum agile</i>	An201	91	3,232,216	60.34
Firmicutes	Ruminococcaceae	<i>Anaeromassilibacillus senegalensis</i>	An200	94	3,738,663	54.30
Firmicutes	Ruminococcaceae	<i>Anaeromassilibacillus senegalensis</i>	An250	97	3,582,839	53.28
Firmicutes	Ruminococcaceae	<i>Anaeromassilibacillus senegalensis</i>	An172	92	2,820,032	41.40
Firmicutes	Ruminococcaceae	<i>Anaerotruncus colihominis</i>	An174	99	4,104,028	53.46
Firmicutes	Ruminococcaceae	<i>Anaerotruncus colihominis</i>	An175	99	4,098,164	53.45
Firmicutes	Ruminococcaceae	<i>Anaerotruncus colihominis</i>	An251	99	3,446,606	54.60
Firmicutes	Ruminococcaceae	<i>Butyricoccus pullicaecorum</i>	An179	99	3,474,625	53.61
Firmicutes	Ruminococcaceae	<i>Butyricoccus pullicaecorum</i>	An180	99	3,016,034	54.43
Firmicutes	Ruminococcaceae	<i>Faecalibacterium prausnitzii</i>	An121	95	2,728,163	61.06
Firmicutes	Ruminococcaceae	<i>Faecalibacterium prausnitzii</i>	An77	96	3,092,450	59.48
Firmicutes	Ruminococcaceae	<i>Faecalibacterium prausnitzii</i>	An192	97	3,520,066	58.80
Firmicutes	Ruminococcaceae	<i>Faecalibacterium prausnitzii</i>	An122	96	3,272,200	59.47
Firmicutes	Ruminococcaceae	<i>Faecalibacterium prausnitzii</i>	An58	96	2,956,350	60.63
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An91	97	3,629,120	57.60
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An52	97	2,845,789	59.13
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An92	98	3,498,453	62.13
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An112	97	2,967,986	59.27
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An135	98	3,907,586	61.45
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An82	97	3,683,590	58.85
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An248	100	3,776,725	60.99
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An4	97	3,239,670	58.22
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An9	97	3,357,728	58.20
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An100	95	3,054,748	57.15
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An306	98	3,967,264	59.00
Firmicutes	Ruminococcaceae	<i>Flavonifractor plautii</i>	An10	96	3,867,419	61.74
Firmicutes	Ruminococcaceae	<i>Gemmiger formicilis</i>	An50	94	3,597,344	58.67
Firmicutes	Ruminococcaceae	<i>Gemmiger formicilis</i>	An194	95	3,080,663	59.30
Firmicutes	Ruminococcaceae	<i>Gemmiger formicilis</i>	An87	94	3,355,244	58.58
Firmicutes	Ruminococcaceae	<i>Gemmiger formicilis</i>	An120	95	3,406,814	60.15
Firmicutes	Ruminococcaceae	<i>Pseudoflavonifractor capillosus</i>	An176	96	2,574,287	58.19
Firmicutes	Ruminococcaceae	<i>Pseudoflavonifractor capillosus</i>	An187	96	2,623,188	57.97
Firmicutes	Ruminococcaceae	<i>Pseudoflavonifractor capillosus</i>	An44	96	2,764,660	57.41
Firmicutes	Ruminococcaceae	<i>Pseudoflavonifractor capillosus</i>	An85	96	3,018,026	56.20
Firmicutes	Ruminococcaceae	<i>Pseudoflavonifractor capillosus</i>	An184	96	3,617,258	59.87
Firmicutes	Veillonellaceae	<i>Megamonas hypermegale</i>	An288	99	2,165,576	33.61
Firmicutes	Veillonellaceae	<i>Megasphaera elsdenii</i>	An286	95	2,396,433	53.29
Proteobacteria	Desulfovibrionaceae	<i>Desulfovibrio desulfuricans</i>	An276	92	3,230,576	55.37

Table 1 List of strains isolated, sequenced and analysed in this study (Continued)

Phylum	Family	Species	ID	rRNA % sim. ^a	genome (bp)	GC (%)
Proteobacteria	Enterobacteriaceae	<i>Escherichia fergusonii</i>	An190	99	5,352,565	50.48
Synergistetes	Synergistaceae	<i>Cloacibacillus porcorum</i>	An23	93	2,902,045	57.89
Verrucomicrobia	Verrucomicrobiaceae	<i>Akkermansia muciniphila</i>	An78	100	2,734,062	55.68
Elusimicrobia	Elusimicrobiaceae	<i>Elusimicrobium minutum</i>	An273	94	1,505,722	51.73

^aPercent similarity along the whole 16S rRNA sequence to the GenBank entry with the highest similarity

ranking of each individual isolate among the most frequent OTUs present in the chicken caecum can be found in Additional file 1.

Genome sizes ranged from 1.51 Mb in *Elusimicrobium* to 6.70 Mb in *Bacteroides ovatus*. Larger genomes were usually recorded in isolates from phylum *Bacteroidetes* (Additional file 1). *Actinobacteria* possessed small genomes ranging from 2.1 to 2.5 Mb. Genomes of *Firmicutes* ranged mostly from 3 to 4 Mb although genomes of *Enterococcus*, [*Eubacterium*] *cylindroides* and *Lactobacillus* were among the smallest ones with genome sizes around 2 Mb. Genomic GC content ranged from 28.0 to 62.1% in *Firmicutes*, from 41.6 to 61.9% in *Bacteroidetes* and from 62.1 to 68.7% in *Actinobacteria*. GC content of individual isolates belonging to the remaining 4 phyla ranged from 50.5 to 57.9% (Additional file 1).

Whole genome comparison

Network analysis based on the correlation of individual gene counts in individual isolates confirmed similarities observed by 16S rRNA gene alignment. Individual isolates from phyla *Proteobacteria*, *Verrucomicrobia*, *Elusimicrobia* and *Synergistetes* formed disconnected vertices of the network (Fig. 2). *Firmicutes* split into families *Lactobacillaceae*, *Enterococcaceae*, *Veillonellaceae* (genera *Megasphaera* and *Megamonas*) and order *Clostridiales* (families *Erysipelotrichaceae*, *Ruminococcaceae* and *Lachnospiraceae*) except for *Clostridium perfringens*. Members of the family *Erysipelotrichaceae* formed a slightly eccentric cluster at the periphery of other isolates from the order *Clostridiales* indicating their slightly different coding capacity when compared to isolates belonging to families *Lachnospiraceae* and *Ruminococcaceae*. All *Bacteroidetes* formed a disconnected network cluster showing similar genetic composition in all isolates. The majority of *Actinobacteria* formed a single network cluster except for the representatives of genus *Gordonibacter* showing that the genome of this genus differed from the rest of *Actinobacteria* sequenced in this study.

Basic biological processes

Since gut microbiota is formed mainly by representatives of phyla *Bacteroidetes* and *Firmicutes*, we specifically focused on the comparison of genomes of isolates

belonging to these two phyla. Representatives belonging to phylum *Bacteroidetes* (Gram negative bacteria) coded for proteins required for the biosynthesis of a Gram negative cell wall while representatives of phylum *Firmicutes* (Gram positive bacteria) coded for proteins required for the biosynthesis of a Gram positive cell wall. However, *Megamonas* and *Megasphaera* (family *Veillonellaceae*, phylum *Firmicutes*, i.e. Gram positive bacteria) harboured genes required for the synthesis of Gram negative cell wall type (Fig. 3). *Bacteroidetes* and *Firmicutes* differed in their mode of transport across the cell wall and cytoplasmic membrane. In *Bacteroidetes*, genes belonging to Ton and Tol transport systems were the most frequent whilst *Firmicutes* encoded ABC transporters, ECF class transporters and TRAP transporters (Fig. 3). Genes enabling sporulation were specific to Gram positive *Firmicutes* except for *Lactobacillaceae*, *Enterococcaceae*, *Veillonellaceae* and two [*Eubacterium*] *cylindroides* isolates (Fig. 3). However, there were differences in the composition and distribution of genes necessary for spore formation among the isolates with sporulation potential. Members of the family *Erysipelotrichaceae* (*Massiliomicrobiota* sp. and [*Clostridium*] *spiroforme*) did not code for stage III sporulation proteins and *Faecalibacterium*, *Anaerofilum* and *Gemmiger* (family *Ruminococcaceae*) did not code for spore proteins though the rest of the genes required for sporulation were present in their genomes (Fig. 3). None of the *Bacteroidetes* isolates encoded proteins required for sporulation and their oxygen tolerance could be dependent on *batABCDE* operon (*Bacteroides aerotolerance*). *batABE* genes were present in all representatives of *Bacteroidetes* and *batCD* were present in all *Bacteroidetes* except for two *Muribaculum* isolates. When we examined the survival rate of the anaerobes after exposure to aerobic conditions experimentally, the most sensitive isolates were single isolates of *Muribaculum intestinale* (phylum *Bacteroidetes*), [*Clostridium*] *glycyrrhizinilyticum*, [*Clostridium*] *saccharolyticum*, *Anaeromassilibacillus senegalensis* and *Flavonifractor plautii* (all phylum *Firmicutes*) which did not survive an hour long exposure to the air. Bacteria which did not survive 8 or 24-h-long exposure to the air belonged mainly to the order *Clostridiales*. Within *Clostridiales*, 45% of the tested isolates did not survive 8-h




Fig. 1 Phylogenetic tree with selected functional properties of 133 sequenced isolates obtained from chicken caecum based on the Bayesian analysis of the full-length sequences of 16S rRNA genes. Families within the phylum *Firmicutes* are shown in light blue, green and yellow. Families within the phylum *Bacteroidetes* are shown in shades of purple. *Actinobacteria* (family *Coriobacteriaceae* – Cor) are highlighted in red. In the cases when only one or two isolates were sequenced per phylum, these isolates are described by phylum name – Prot – *Proteobacteria*, Elu – *Elusimicrobia*, Ver – *Verrucomicrobia*, Syn – *Synergistetes*. In the remaining cases, branches with different families are highlighted with different colors. Rik – *Rikenellaceae*, Porp - *Porphyromonadaceae*, Bact – *Bacteroidaceae*, Lach – *Lachnospiraceae*, Clos - *Clostridiaceae*, Ery – *Erysipelotrichaceae*, Entcoc – *Enterococcaceae*, Lact – *Lactobacillaceae*, Rum – *Ruminococcaceae*, Veil – *Veillonellaceae*. BUK – butyrate kinase, PBT – phosphate butyryltransferase, BTR – butyryl-CoA transferase, AcCo – acetyl CoA pathway, LYS – lysine fermentation pathway, SUC – succinate fermentation pathway, M-MLN presence of methylmalonyl mutase, epimerase and decarboxylase required for conversion of succinate to methyl-malonyl CoA and propionate. AcGen – presence of genes required for reductive acetogenesis

long exposure to the air, and an additional 25% died between 8 to 24-h exposure. Only 25% of tested isolates from the order *Clostridiales* survived 24-h air exposure. On the other hand, representatives of *Bacteroidetes* and *Actinobacteria* were usually tolerant to sudden air exposure since 62 and 73% of tested isolates survived 24-h air

exposure, respectively (Fig. 4 and Additional file 3). *Bacteroides* sp. encoded a high number of proteins involved in polysaccharide and monosaccharide metabolism while the number of genes required for metabolism of di- and oligosaccharides was similar in both *Bacteroidetes* and *Firmicutes* (Fig. 3).




Fig. 2 Network analysis of isolates from chicken caecum based on their gene composition. *Firmicutes* split into families *Lactobacillaceae*, *Enterococcaceae*, *Veillonellaceae* (genera *Megasphaera* and *Megamonas*) and the order *Clostridiales* except for *Clostridium perfringens*. Separation of *Erysipelotrichaceae* from *Lachnospiraceae* and *Ruminococcaceae* (all belonging to order *Clostridiales*) was observed. All *Bacteroidetes* formed a separated cluster showing similar genetic composition in all isolates. The majority of *Actinobacteria* formed a single cluster except for the representatives of the genus *Gordonibacter* indicating that genetic composition of these isolates differed from the rest of *Actinobacteria*




Fig. 3 Distribution of genes in selected categories among representatives of major gut colonisers belonging to phyla *Bacteroidetes* and *Firmicutes*. X axes indicate the numbers of genes in a given category per genome. For a scalable figure see Additional file 5 online




Fig. 4 Sensitivity of gut anaerobes to air exposure. Most of isolates from the order *Clostridiales*, phylum *Firmicutes* ($n = 69$) died during 1–8 long hour exposure to the air. On the other hand, majority of the tested representatives of *Bacteroidetes* ($n = 29$) and *Actinobacteria* ($n = 15$) survived 24-h-long exposure to air. For full information on survival of individual isolates see Additional file 3 online

Production of butyrate, propionate and acetogenesis


Short chain fatty acids and butyrate in particular, are acknowledged as important metabolites of bacterial origin [14, 15]. All genes required for butyrate production from pyruvate and acetyl-CoA were present in the genomes of *Ruminococcaceae* (genera *Butyricoccus*, *Pseudoflavonifractor*, *Flavonifractor*, *Anaeromassilibacillus*, *Anaerotruncus* and *Faecalibacterium*) (Fig. 1). In addition, this pathway was also present in *Megasphaera elsdenii*, [*Eubacterium*] *cylindroides*, [*Clostridium*] *lactatifermentans*, [*Clostridium*] *saccharolyticum* and [*Eubacterium*] *hallii*, the latter three belonging to the family *Lachnospiraceae* (Fig. 1). *Butyricimonas paravirosa* was the only isolate from phylum *Bacteroidetes* coding for all genes required for butyrate production from pyruvate and acetyl-CoA. Genes coding for enzymes of complete lysine fermentation pathway leading to butyrate production were present in three genera of the phylum *Bacteroidetes* (*Muribaculum*, *Butyricimonas* and *Odoribacter*) and the majority of *Flavonifractor* isolates belonging to the phylum *Firmicutes*. *Butyricimonas* and *Odoribacter* also encoded the whole pathway required for the conversion of succinate and 4-hydroxybutyrate into butyrate (Fig. 1). Terminal steps in butyrate production were dependent on transferases transferring CoA moiety from butyryl-CoA to acetate, acetoacetate or 4-hydroxybutyrate, or phosphate butyryltransferase (PBT) and butyrate kinase (BUK). Surprisingly, we did not find butyrate kinase in all isolates using the PBT - BUK pathway for butyrate release from butyryl-CoA. In such a case, butyrate-phosphate may serve for substrate phosphorylation in enzymatic reactions similar to acetate-phosphate.

Propionate production via a succinate-methylmalonate pathway was characteristic of *Bacteroidetes* (Fig. 1) as genes for methylmalonyl-CoA mutase, epimerase and decarboxylase were detected in genomes of all isolates from this phylum. This pathway was quite rare in *Firmicutes* since methylmalonyl-CoA mutase, epimerase and decarboxylase were encoded only by *Megamonas* and two *Flavonifractor* isolates (Fig. 1).

Fermentation of carbohydrates results in the production of H_2 which has to be removed from the community since its increased concentration suppresses glycolysis [16–18]. H_2 can be removed by methanogens, acetogens or sulphate reducing bacteria. We did not isolate a single methanogen. *Desulfovibrio* (phylum *Proteobacteria*) encoded key genes for sulphate reduction to H_2S (adenylylsulphate reductase, sulphate adenylyltransferase, dissimilatory sulphite reductase and sulphite reduction-associated complex DsrMKJOP). Potential for H_2 removal by acetogenesis was recorded in [*Clostridium*] *saccharolyticum*, [*Eubacterium*] *fissicatena* and all *Blautia* isolates (*B. coccoides*, *B. producta*, *B. schinkii*) since all these bacteria encoded corrinoid iron-sulfur acetyl-CoA synthase and 5-methyltetrahydrofolate methyltransferase (Fig. 1).

Genes encoding proteins mediating interactions with the host

Since gut microbiota may interact with its host, finally we searched for the presence of genes which may facilitate such interactions (Additional file 4). Genes encoding collagenase precursor, hemagglutinin or hemolysin A were present in all 29 *Bacteroidetes* isolates but not in *Firmicutes* (Fig. 5). Hyaluronidase was present in 16 isolates from the phylum *Bacteroidetes* and five *Firmicutes*. Different heparinases were detected in 14 isolates from the phylum *Bacteroidetes* and seven representatives of *Firmicutes* – out of these all five *Faecalibacterium* isolates encoded heparinase II/III-like and outside this genus, heparinases were present only in two isolates from the phylum *Firmicutes* (i.e. *Gemmiger formicilis* and [*Clostridium*] *glycyrrhizinilyticum*). Chondroitinase was present in nine isolates from phylum *Bacteroidetes* (genera *Alistipes* and *Bacteroides*) but in no isolate from *Firmicutes*. Mucin-desulfating sulfatase was present in the genomes of 20 *Bacteroidetes* isolates (mainly in genera *Alistipes*, *Parabacteroides* and *Bacteroides*) but was absent from the genomes of *Firmicutes*. A gene for endothelin-converting enzyme 1 precursor was present in the genomes of all *Bacteroidetes* but not in *Firmicutes*.



Except for two *Odoribacter* isolates and *Parabacteroides distatonis*, all the remaining representatives of the phylum *Bacteroidetes* encoded 3-oxo-5- α -steroid 4-dehydrogenase capable of modification of bile or steroid hormones. This gene was not detected in *Firmicutes*. Glutamate decarboxylase catalysing production of γ -aminobutyrate (GABA) and glutamate/GABA antiporter was present in 20 different *Bacteroidetes* isolates but only in two *Firmicutes* and these were both pathogenic *Clostridium perfringens* isolates. Histidine decarboxylase catalysing production of histamine was quite rare and was present only in two *Bacteroides dorei* isolates (Fig. 5). On the other hand, except for all *Lactobacilli*, the gene for fibronectin/fibrinogen-binding protein was present in all isolates from *Firmicutes* but in none representative from the phylum *Bacteroidetes*. Genes for flagellar motility were quite rare and were absent in all *Bacteroidetes* isolates. Complete flagellar operon was present only in two *Flavonifractor* isolates, one strain of *Anaeromassilibacillus senegalensis* and all three isolates of *Anaerotruncus colihominis* (Fig. 5).

Discussion

In this study we sequenced, annotated and analysed genomes of 133 different anaerobes cultured from the chicken caecum. Several isolates represented species which are available in only a few pure cultures worldwide – a single manuscript reported a pure culture of *Elusimicrobium* [19] and only two papers reported culture of *Cloacibacillus porcorum* [20, 21]. Although such isolates are of clear potential for future experiments, in this report we mainly focused on the comparison of genetic potential of isolates belonging to two main phyla inhabiting the intestinal tract of chickens, *Bacteroidetes* and *Firmicutes* [8, 18].

Whole genome sequencing showed that representatives of *Bacteroidetes* and *Firmicutes* have a genetic potential to follow different strategies of gut colonisation which may explain their coexistence in the intestinal tract. *Bacteroidetes* encoded genes for the biosynthesis of a Gram negative cell wall, Gram positive anaerobes encoded genes for the biosynthesis of a Gram positive cell wall. Only *Megamonas* and *Megasphaera*, though belonging to Gram positive *Firmicutes*, encoded genes for the biosynthesis of Gram negative cell wall type, in agreement with previous report [22]. The Ton/ExbD transport system of *Bacteroidetes* has been identified earlier as highly expressed in vivo [7] and ABC, ECF and TRAP transporters were described as characteristic of *Firmicutes* [23–25]. *Bacteroidetes* were reported to increase in gut microbiota with high fiber content in their diet [26, 27] and to forage on host derived polysaccharides in the absence of fiber [28–30]. Consistent with this, *Bacteroidetes* and the genus *Bacteroides* in particular encoded a high number of genes required for polysaccharide metabolism [28, 30–33]. Since the polysaccharides of feed and host origin consist of various (amino)monosaccharides, *Bacteroidetes* encoded also a wider range of genes required for monosaccharide degradation than *Firmicutes* (Fig. 3).

Butyrate is produced mainly by *Firmicutes*. Carbohydrate fermentation to pyruvate and acetyl-CoA was the most frequent butyrate production pathway, as proposed earlier [34]. Butyrate production was mainly associated with *Ruminococcaceae* and less frequently with *Lachnospiraceae* or *Erysipelotrichaceae* [35]. *Bacteroidetes* are not the main butyrate producers as acetyl-CoA conversion to butyrate was only found in *Butyricimonas*. However, *Bacteroidetes* were capable of butyrate production by alternative pathways, e.g. from 4-hydroxybutyrate as recorded in *Butyricimonas* and *Odoribacter* or by lysine fermentation as recorded in *Muribaculum*, *Butyricimonas* and *Odoribacter*. This is in line with conclusions derived from human microbiota studies [35].

Gut microbiota interacts with the host. The potential of *Firmicutes* to interact with the chicken host seems to be less extensive in comparison to *Bacteroidetes*. Except for *Lactobacilli*, *Firmicutes* isolates encoded fibronectin/fibrinogen-binding protein [36, 37]. Interestingly, we detected chicken fibrinogen-domain containing proteins as tightly associated with gut microbiota [38]. Such proteins aggregate bacteria [39, 40] and enable association of different gut microbiota members, based on current results, preferentially those belonging to the phylum *Firmicutes*. Binding to chicken fibrinogen-domain containing proteins may result in the formation of random bacterial aggregates and those with the most optimal composition, e.g. butyrate producers releasing H_2 with *Blautia* consuming H_2 for acetate production thus

allowing glycolysis in butyrate producers to proceed [16], will rapidly multiply and define final microbiota composition. Except for fibrinogen binding, representatives from the phylum *Bacteroidetes* had a greater potential to affect host behaviour than representatives from the phylum *Firmicutes*. *Bacteroidetes* encoded collagenase, hemagglutinin, hemolysin, hyaluronidase, heparinases, chondroitinase or mucin-desulfating sulfatase required for degradation of host structures. In addition, representatives from the phylum *Bacteroidetes* encoded endothelin-converting enzyme 1 precursor, which plays a significant role in cardiovascular diseases and Alzheimer's disease in humans [41], 3-oxo-5- α -steroid 4-dehydrogenase capable of modification of steroid hormones or bile [42], or glutamate decarboxylase catalysing production of γ -aminobutyrate (GABA), a mediator within the enteric nervous system [43]. However, it will have to be elucidated whether these genes are expressed in vivo and whether their products may reach host structures.

Finally, the rather counterintuitive conclusion came from the prediction of survival following air exposure. Although *Bacteroidetes* should be more sensitive to air exposure than *Firmicutes* which are capable of spore formation, our data clearly showed that *Clostridiales*, including all important butyrate producers, were highly sensitive to air exposure. Due to the experimental design, we likely determined sensitivity of vegetative cells and not of spores. Despite this, the extreme sensitivity of vegetative cells of *Clostridiales* may explain their commonly reported disappearance in inflammatory bowel disease patients [44]. Inflammation leads to locally increased oxygen levels due to the oxidative burst of granulocytes and macrophages [45, 46]. Disappearance of *Clostridiales* including major butyrate producers can therefore be a consequence rather than a cause of inflammatory bowel diseases. Similarly, a reported increase in the abundance of *Bacteroidetes* or *Megasphaera* during inflammatory diseases [35, 47] may be a mere consequence of their higher resistance to oxygen and the disappearance of oxygen sensitive bacterial species from order *Clostridiales*. An overgrowth of facultative anaerobes like those from family *Enterobacteriaceae* in acute colitis also fits in the proposed scenario [47–49].

Conclusions

In this study we isolated and sequenced 133 different strains originating from chickens intestinal tracts belonging to seven different phyla. Analysis of their genomic sequences showed that butyrate production was mainly associated with *Ruminococcaceae*, and less frequently with *Lachnospiraceae* or *Erysipelotrichaceae*, all belonging to phylum *Firmicutes*. Representatives of phylum *Bacteroidetes* commonly encoded proteins

(collagenase, hemagglutinin, hemolysin, hyaluronidase, heparinases, chondroitinase, mucin-desulfating sulfatase or glutamate decarboxylase) that may enable them to interact with their host. Even such a brief list of genes shows that representatives of *Bacteroidetes* and *Firmicutes* follow different strategies of gut colonisation which contributes to their coexistence in the intestinal tract.

Methods

Ethics statement

The handling of animals in this study was performed in accordance with current Czech legislation (Animal Protection and Welfare Act No. 246/1992 Coll. of the Government of the Czech Republic). Experiments performed in this study were approved by the Ethics Committee of the Veterinary Research Institute (permit number 4/2016) followed by the Committee for Animal Welfare of the Ministry of Agriculture of the Czech Republic.

Isolation and identification of caecal bacteria

The chickens were sacrificed under chloroform anesthesia by cervical dislocation. Whole caeca with their contents originating from 18 random healthy chickens or hens 4 to 40 weeks of age were removed during necropsy, chilled on ice and transported to an anaerobic chamber for further processing within one hour. The caeca were opened in an anaerobic chamber (10% CO₂, 5% H₂ and 85% N₂ atmosphere; Concept 400, Baker Ruskinn, USA) and 0.5 g of content was squeezed into 4.5 ml pre-reduced PRAS dilution blank (0.1 g magnesium sulfate heptahydrate, 0.2 g monobasic potassium phosphate, 0.2 g potassium chloride, 1.15 g dibasic sodium phosphate, 3.0 g sodium chloride, 1.0 g sodium thioglycolate, 0.5 g L-cysteine, 1000 ml distilled water; final pH 7.5 \pm 0.2 at 25 °C) and mixed thoroughly. All samples were serially diluted in pre-reduced PRAS dilution blank and plated on Wilkins-Chalgren anaerobe agar (WCHA) (Oxoid) supplemented with 30% of rumen fluid. The rumen fluid was collected from cows by an oral probe, filtered through cheesecloth, centrifuged at 8000 g for 30 min and sterilised by filtration through a 0.22 μ m filter. Aliquots of rumen fluid were stored at –20 °C. WCHA was additionally supplemented with 5 mg/l hemin, 1 mg/l cellobiose, 0.5 g/l soluble starch, 1 mg/ml maltose, 0.2 ml vitamin K1 solution (0.1 ml of filter sterilized vitamin K1 in 20 ml 95% ethanol) and 0.5 mg/ml L-cysteine. Approx. 10 well-separated colonies of different morphology were selected from each agar plate after a five-day incubation at 37 °C and purified by subculture on WCHA. All isolates were stored at –80 °C in pre-reduced PRAS dilution blank containing glycerol at 20% concentration and equal volume of sterile sheep blood. Sensitivity of pure anaerobe

cultures to air oxygen exposure was tested exactly as described elsewhere [6]. Briefly, bacterial cultures were serially diluted in anaerobic chamber and plated on 4 copies of WCHA. One copy of WCHA was left in the anaerobic chamber to determine initial counts of each anaerobe. The remaining 3 copies of WCHA plates were placed into a standard aerobic 37 °C incubator and after 1, 8 and 24 h, a single copy of agar plate was returned back to the anaerobic chamber to check for growth restoration.

Whole genome sequencing

DNA was purified using DNeasy Blood & Tissue Kit (Qiagen). Sequencing library was prepared from 1 ng of RNA-free genomic DNA using the Nextera XT DNA Sample Preparation kit (Illumina) and whole genome sequencing was performed using the NextSeq 500/550 High Output Kit v2 and Illumina NextSeq 500 sequencing platform in the paired-end modus (2 × 150 bp). Raw sequencing reads were quality trimmed using Trimmomatic v0.32 [50] with the sliding window of 4 bp and average quality threshold value equal to 17. Minimal read length was set to 48 bp. Trimmed paired-end reads were assembled via de novo assembler IDBA-UD v1.1.1 [51] with k-mer sizes ranging from 20 to 110 with an increment of 15. Contigs with coverage lower than 10% of average coverage of L50 contigs were filtered out and the remaining contigs were scaffolded employing SSPACE scaffolder v3.0 [52, 53]. All scaffolds containing N's in their sequences were split into N-free sequences. Finally, scaffolds with a length shorter than 500 bp were discarded.

Species definition and genome annotation

Species definitions used in this study are based on the BLAST comparison of whole 16S rRNA sequences against entries deposited in NCBI 16S rRNA sequence database performed on January 4, 2017. For clarity of the paper, we used the designation of the most similar bacterium based on the lowest E-value for description of our isolates, thus apparently ignoring the fact that in some cases there was 100% identity whilst in the opposite extreme, the sequence of 16S rRNA of one of our isolates was only 83% similar to the closest relative deposited in the NCBI 16S rRNA database (Additional file 1). All 16S rRNA sequences were compared also against RDP SeqMatch database (on January 10, 2017) which allowed for alternative taxonomy including classification of individual isolates into higher taxonomic units. In addition, ribosomal protein multilocus sequence typing (rMLST) [54] and GTDB organism identification (<http://gtdb.ecogenomic.org/>) databases were used to verify taxonomic classification. Taxonomic classification by these alternative protocols is provided in

Additional file 1. Gene predictions and functional annotations were performed by RAST [55]. Assembled and annotated genomes as well as raw sequencing data were deposited in NCBI under accession number PRJNA377666 and genomes with comprehensive RAST annotation are available upon request.

Genome comparison

To exclude genomes of the same isolates picked up on independent occasions from the subsequent analyses, whole genome sequences of all isolates were mutually compared using the QCAST tool v3.1 [56]. Two genomes were considered as identical if they shared more than 99% of genome content and had less than 1 indel per 100 kb. A single isolate was selected as a representative of each group of identical isolates for all downstream analyses.

Whole gene content similarity clustering was compared using Gene Co-Expression Network analysis. This protocol detects genes with similar transcriptional regulatory program (potential members of the same pathway, protein complex, etc.). In our case, the gene expression vector for particular gene was replaced by gene copy-number vector for particular bacteria and the protocol therefore detected bacterial isolates with similar gene content. Interconnected bacteria shared more than 50% of genes based on gene name designation provided by RAST annotation. Whole gene content similarity of individual isolates was analysed in R. At first, matrix of Spearman's correlation coefficients was calculated for all pairs of isolates using vectors of respective gene counts. The correlation matrix was then transformed to the adjacency matrix using threshold of +0.5 as a cut-off for two vertices to be considered as connected. The undirected network was constructed from such a matrix using igraph package (<http://igraph.org/r/>), and edge betweenness based community structure detection algorithm was then employed to identify separate network modules. Communities with more than three members were considered nontrivial and were highlighted in a network plot.

Analysis of 16S rRNA genes

Trimmed reads were aligned against SILVA bacterial 16S rRNA database using SortMeRNA v2.1 [57] and extracted 16S rRNA reads were assembled via de Bruijn graph-based de novo assembler SPAdes v3.6.0 [58]. Finally, sequences coding for 16S rRNA genes were predicted employing barrnap tool v0.6 (<https://github.com/tseemann/barrnap>). For the purposes of phylogenetic analysis, 16S rRNA genes were aligned by ClustalW v2.1 [59] using default gap penalties, DNA weight matrix IUB and transition weight 0.2. The phylogenetic tree topologies were inferred employing Bayesian statistics via MrBayes v3.2.6 [60] using the parameters as follows: mixed model of nucleotide substitution, gamma-distributed rates

among sites, four Monte Carlo Markov chains for 7,000,000 cycles which were sampled every 1000th generation and the first 25% of the samples were discarded as burn-in. The final tree topology was generated employing 50% majority-rule consensus. Average standard deviation of split frequencies was 0.0083, maximum standard deviation of split frequencies was 0.1095, average potential scale reduction factor was 1.000 and maximum potential scale reduction factor was 1.015. The final tree topology was visualized by iTOL v3.4.3 [61] and edited using Inkscape v0.91 (www.inkscape.org).

Additional files

Additional file 1: List of 133 different isolates characterised in this study. The file contains taxonomical classification using NCBI and RDP databases based on whole sequence of 16S rRNA, similarity to the closest relative in the NCBI database, genome size, number of contigs into which the genome was assembled, genomic and 16S rRNA GC content, consensus 16S rRNA sequences with SNP positions indicated with lower case letters, 16S rRNA copy number estimated based on sequencing coverage and 16S rRNA copy number determined by RT-PCR (only for the isolates in which sequencing coverage predicted more 10 copies of 16S rRNA genes). (XLSX 338 kb)

Additional file 2: Phylogenetic tree of 133 sequenced isolates obtained from chicken caecum based on the Clustal alignment of the full-length sequence of RpoB proteins. Families within the phylum *Firmicutes* are shown in light blue, green and yellow. Families within the phylum *Bacteroidetes* are shown in shades of purple. The whole genome size and genomic GC content of each isolate is shown external to the dendrogram. (PDF 576 kb)

Additional file 3: Aerobic survival of chicken gut anaerobes. (XLS 43 kb)

Additional file 4: List of proteins encoded by individual anaerobes. (XLSX 7930 kb)

Additional file 5: Distribution of genes in selected categories among representatives of major gut colonisers belonging to phyla *Bacteroidetes* and *Firmicutes*. X axes indicate the numbers of genes in a given category per genome. (PDF 23 kb)

Abbreviations

ABC transporter: ATP-binding cassette transporter; *batABCDE* operon: *Bacteroides* aerotolerance operon; BUK: Butyrate kinase; ECF class transporter: Energy-coupling factor transporter; GABA: γ -aminobutyrate; GTDB: Genome taxonomy database; NCBI: National Center for Biotechnology Information; OTU: Operational taxonomic unit; PBT: Phosphate butyryltransferase; PRAS dilution blank: Pre-reduced anaerobically sterilized dilution blank; RAST: Rapid annotations using subsystems technology; RDP: Ribosomal database project; rMLST: Ribosomal protein multilocus sequence typing; TRAP transporter: Tripartite ATP-independent periplasmic transporter; WCHA: Wilkins-Chalgren anaerobe agar

Acknowledgements

Authors would like to thank Peter Eggenhuizen for English language corrections.

Funding

This work has been supported by project RO0517 from the Czech Ministry of Agriculture, AdmireVet project CZ.1.05/2.1.00/01.0006-ED0006/01/01 from the Czech Ministry of Education, Czech Science Foundation project 15-11688S and project QJ1310019 from the Czech Ministry of Agriculture. AC was supported by CEITEC 2020 (LQ1601) from the Ministry of Education, Youth and Sports of the Czech Republic under the National Sustainability Programme II. The funders had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

Assembled and annotated genomes as well as raw sequencing data were deposited in NCBI under accession number PRJNA377666. Genomes with comprehensive RAST annotation are available upon request.

Authors' contributions

MM participated in the sequence alignment, performed bioinformatic analysis and drafted the manuscript. DC performed the whole genome sequencing and sequence alignment. OP analysed sequencing data for protein coding genes and their functions. DK and TK subcultured bacterial isolates, and purified and checked the DNA for whole genome sequencing. AC was responsible for anaerobic culture, preliminary strain identification by mass spectrometry and maintenance of laboratory strain collection. IR participated in the design of the study, analysed data and wrote the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The handling of animals in this study was performed in accordance with current Czech legislation (Animal protection and welfare Act No. 246/1992 Coll. of the Government of the Czech Republic). Experiments performed in this study were approved by the Ethics Committee of the Veterinary Research Institute (permit number 4/2016) followed by the Committee for Animal Welfare of the Ministry of Agriculture of the Czech Republic. The animals used in this study originated from commercial sources. Their use for anaerobic gut microbiota culture was based on verbal consent of private owners due to continuous collaboration between authors' laboratories and individual owners.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Veterinary Research Institute, Hudcova 70, 621 00 Brno, Czech Republic. ²Central European Institute of Technology (CEITEC), University of Veterinary and Pharmaceutical Sciences Brno, Brno, Czech Republic. ³Department of Infectious Diseases and Microbiology, Faculty of Veterinary Medicine, University of Veterinary and Pharmaceutical Sciences Brno, Brno, Czech Republic.

Received: 27 December 2017 Accepted: 24 July 2018

Published online: 31 July 2018

References

- Gosalbes MJ, Durban A, Pignatelli M, Abellan JJ, Jimenez-Hernandez N, Perez-Cobas AE, Latorre A, Moya A. Metatranscriptomic approach to analyze the functional human gut microbiota. *PLoS One*. 2011;6:e17447.
- Gill SR, Pop M, Deboy RT, Eckburg PB, Turnbaugh PJ, Samuel BS, Gordon JL, Relman DA, Fraser-Liggett CM, Nelson KE. Metagenomic analysis of the human distal gut microbiome. *Science*. 2006;312:1355–9.
- Kurokawa K, Itoh T, Kuwahara T, Oshima K, Toh H, Toyoda A, Takami H, Morita H, Sharma VK, Srivastava TP, Taylor TD, Noguchi H, Mori H, Ogura Y, Ehrlich DS, Itoh K, Takagi T, Sakaki Y, Hayashi T, Hattori M. Comparative metagenomics revealed commonly enriched gene sets in human gut microbiomes. *DNA Res*. 2007;14:169–81.
- Marx V. Microbiology: the return of culture. *Nat Methods*. 2016;14:37–40.
- Lau JT, Whelan FJ, Herath I, Lee CH, Collins SM, Bercik P, Surette MG. Capturing the diversity of the human gut microbiota through culture-enriched molecular profiling. *Genome Med*. 2016;8:72.
- Browne HP, Forster SC, Anonye BO, Kumar N, Neville BA, Stares MD, Goulding D, Lawley TD. Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation. *Nature*. 2016;533:543–6.
- Polansky O, Sekelova Z, Faldynova M, Sebkova A, Sisak F, Rychlik I. Important metabolic pathways and biological processes expressed by chicken cecal microbiota. *Appl Environ Microbiol*. 2016;82:1569–76.

8. Vidsenska P, Sedlar K, Lukac M, Faldynova M, Gerzova L, Cejkova D, Sisak F, Rychlik I. Succession and replacement of bacterial populations in the caecum of egg laying hens over their whole life. *PLoS One*. 2014;9:e115142.
9. Varmuzova K, Kubasova T, Davidova-Gerzova L, Sisak F, Havlickova H, Sebkova A, Faldynova M, Rychlik I. Composition of gut microbiota influences resistance of newly hatched chickens to *Salmonella* Enteritidis infection. *Front Microbiol*. 2016;7:957.
10. Rantala M, Nurmi E. Prevention of the growth of *Salmonella* infantis in chicks by the flora of the alimentary tract of chickens. *Br Poult Sci*. 1973;14:627–30.
11. Duggett NA, Kay GL, Sergeant MJ, Bedford M, Constantinidou CI, Penn CW, Millard AD, Pallen MJ. Draft genome sequences of six novel bacterial isolates from chicken caeca. *Genome Announc*. 2016;4:e00448–16.
12. Yarza P, Yilmaz P, Pruesse E, Glockner FO, Ludwig W, Schleifer KH, Whitman WB, Euzéby J, Amann R, Rossello-Mora R. Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Microbiol*. 2014;12:635–45.
13. Adekambi T, Drancourt M, Raoult D. The *rhoB* gene as a tool for clinical microbiologists. *Trends Microbiol*. 2009;17:37–45.
14. Fleming SE, Fitch MD, DeVries S, Liu ML, Kight C. Nutrient utilization by cells isolated from rat jejunum, cecum and colon. *J Nutr*. 1991;121:869–78.
15. Kelly CJ, Zheng L, Campbell EL, Saeedi B, Scholz CC, Bayless AJ, Wilson KE, Glover LE, Kominsky DJ, Magnuson A, Weir TL, Ehrentraut SF, Pickel C, Kuhn KA, Lanis JM, Nguyen V, Taylor CT, Colgan SP. Crosstalk between microbiota-derived short-chain fatty acids and intestinal epithelial HIF augments tissue barrier function. *Cell Host Microbe*. 2015;17:662–71.
16. Carbonero F, Benefiel AC, Gaskins HR. Contributions of the microbial hydrogen economy to colonic homeostasis. *Nat Rev Gastroenterol Hepatol*. 2012;9:504–18.
17. Rey FE, Gonzalez MD, Cheng J, Wu M, Ahern PP, Gordon JI. Metabolic niche of a prominent sulfate-reducing human gut bacterium. *Proc Natl Acad Sci U S A*. 2013;110:13582–7.
18. Sergeant MJ, Constantinidou C, Cogan TA, Bedford MR, Penn CW, Pallen MJ. Extensive microbial and functional diversity within the chicken cecal microbiome. *PLoS One*. 2014;9:e91941.
19. Geissinger O, Herlemann DP, Morschel E, Maier UG, Brune A. The ultramicrobacterium "*Elusimicrobium minutum*" gen. Nov., sp. nov., the first cultivated representative of the termite group 1 phylum. *Appl Environ Microbiol*. 2009;75:2831–40.
20. Domingo MC, Yansouni C, Gaudreau C, Lamothe F, Levesque S, Tremblay C, Garceau R. *Cloacibacillus* sp., a potential human pathogen associated with bacteremia in Quebec and New Brunswick. *J Clin Microbiol*. 2015;53:3380–3.
21. Looft T, Levine UY, Stanton TY. *Cloacibacillus porcorum* sp. nov., a mucin-degrading bacterium from the swine intestinal tract and emended description of the genus *Cloacibacillus*. *Int J Syst Evol Microbiol*. 2013;63:1960–6.
22. Antunes LC, Poppleton D, Klingl A, Criscuolo A, Dupuy B, Brochier-Armanet C, Beloin C, Gribaldo S. Phylogenomic analysis supports the ancestral presence of LPS-outer membranes in the *Firmicutes*. *Elife*. 2016;5:e14589.
23. Fischer M, Zhang QY, Hubbard RE, Thomas GH. Caught in a TRAP: substrate-binding proteins in secondary transport. *Trends Microbiol*. 2010;18:471–8.
24. Rodionov DA, Hebbeln P, Eudes A, ter Beek J, Rodionova IA, Erkens GB, Slotboom DJ, Gelfand MS, Osterman AL, Hanson AD, Eitinger T. A novel class of modular transporters for vitamins in prokaryotes. *J Bacteriol*. 2009;191:42–51.
25. Slotboom DJ. Structural and mechanistic insights into prokaryotic energy-coupling factor transporters. *Nat Rev Microbiol*. 2014;12:79–87.
26. Kubasova T, Davidova-Gerzova L, Merlot E, Medvecký M, Polansky O, Gardan-Salmon D, Quesnel H, Rychlik I. Housing systems influence gut microbiota composition of sows but not of their piglets. *PLoS One*. 2017;12:e0170051.
27. Ivarsson E, Roos S, Liu HY, Lindberg JE. Fermentable non-starch polysaccharides increases the abundance of *Bacteroides-Prevotella-Porphyrmonas* in ileal microbial community of growing pigs. *Animal*. 2014;8:1777–87.
28. Wu M, McNulty NP, Rodionov DA, Khoroshkin MS, Griffin NW, Cheng J, Latreille P, Kerstetter RA, Terrapon N, Henrissat B, Osterman AL, Gordon JI. Genetic determinants of in vivo fitness and diet responsiveness in multiple human gut *Bacteroides*. *Science*. 2015;350:aac5992.
29. Benjdia A, Martens EC, Gordon JI, Berteau O. Sulfatases and a radical S-adenosyl-L-methionine (AdoMet) enzyme are key for mucosal foraging and fitness of the prominent human gut symbiont, *Bacteroides thetaiotaomicron*. *J Biol Chem*. 2011;286:25973–82.
30. Martens EC, Chiang HC, Gordon JI. Mucosal glycan foraging enhances fitness and transmission of a saccharolytic human gut bacterial symbiont. *Cell Host Microbe*. 2008;4:447–57.
31. Nihira T, Suzuki E, Kitaoka M, Nishimoto M, Ohtsubo K, Nakai H. Discovery of beta-1,4-D-mannosyl-N-acetyl-D-glucosamine phosphorylase involved in the metabolism of N-glycans. *J Biol Chem*. 2013;288:27366–74.
32. Zhang M, Chekan JR, Dodd D, Hong PY, Radlinski L, Revindran V, Nair SK, Mackie RI, Cann I. Xylan utilization in human gut commensal bacteria is orchestrated by unique modular organization of polysaccharide-degrading enzymes. *Proc Natl Acad Sci U S A*. 2014;111:E3708–17.
33. Magnusdottir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, Greenhalgh K, Jager C, Baginska J, Wilmes P, Fleming RM, Thiele I. Generation of genome-scale metabolic reconstructions for 773 members of the human gut microbiota. *Nat Biotechnol*. 2017;35:81–9.
34. Vital M, Howe AC, Tiedje JM. Revealing the bacterial butyrate synthesis pathways by analyzing (meta)genomic data. *MBio*. 2014;5:e00889.
35. Anand S, Kaur H, Mande SS. Comparative in silico analysis of butyrate production pathways in gut commensals and pathogens. *Front Microbiol*. 2016;7:1945.
36. Barketi-Klai A, Hoys S, Lambert-Bordes S, Collignon A, Kansau I. Role of fibronectin-binding protein a in *Clostridium difficile* intestinal colonization. *J Med Microbiol*. 2011;60:1155–61.
37. Hennequin C, Janoir C, Barc MC, Collignon A, Karjalainen T. Identification and characterization of a fibronectin-binding protein from *Clostridium difficile*. *Microbiology*. 2003;149:2779–87.
38. Volf J, Polansky O, Varmuzova K, Gerzova L, Sekelova Z, Faldynova M, Babak V, Medvecký M, Smith AL, Kaspers B, Velge P, Rychlik I. Transient and prolonged response of chicken cecum mucosa to colonization with different gut microbiota. *PLoS One*. 2016;11:e0163932.
39. Doolittle RF, McNamara K, Lin K. Correlating structure and function during the evolution of fibrinogen-related domains. *Protein Sci*. 2012;21:1808–23.
40. Wu C, Soderhall K, Soderhall I. Two novel ficolin-like proteins act as pattern recognition receptors for invading pathogens in the freshwater crayfish *Pacifastacus leniusculus*. *Proteomics*. 2011;11:2249–64.
41. Palmer JC, Kehoe PG, Love S. Endothelin-converting enzyme-1 in Alzheimer's disease and vascular dementia. *Neuropathol Appl Neurobiol*. 2010;36:487–97.
42. Langlois VS, Zhang D, Cooke GM, Trudeau VL. Evolution of steroid-5alpha-reductases and comparison of their function with 5beta-reductase. *Gen Comp Endocrinol*. 2010;166:489–97.
43. Auteri M, Zizzo MG, Serio R. GABA and GABA receptors in the gastrointestinal tract: from motility to inflammation. *Pharmacol Res*. 2015;93:11–21.
44. Erickson AR, Cantarel BL, Lamendella R, Darzi Y, Mongodin EF, Pan C, Shah M, Halfvarson J, Tysk C, Henrissat B, Raes J, Verberkmoes NC, Fraser CM, Hettich RL, Jansson JK. Integrated metagenomics/metaproteomics reveals human host-microbiota signatures of Crohn's disease. *PLoS One*. 2012;7:e49138.
45. Thiennimitr P, Winter SE, Winter MG, Xavier MN, Tolstikov V, Huseby DL, Sterzenbach T, Tsois RM, Roth JR, Baumler AJ. Intestinal inflammation allows *Salmonella* to use ethanolamine to compete with the microbiota. *Proc Natl Acad Sci U S A*. 2011;108:17480–5.
46. Winter SE, Baumler AJ. A breathtaking feat: to compete with the gut microbiota, *Salmonella* drives its host to provide a respiratory electron acceptor. *Gut Microbes*. 2011;2:58–60.
47. Berry D, Schwab C, Milinovich G, Reichert J, Ben Mahfoudh K, Decker T, Engel M, Hai B, Hainzl E, Heider S, Kenner L, Muller M, Rauch I, Strobl B, Wagner M, Schleper C, Urich T, Loy A. Phylotype-level 16S rRNA analysis reveals new bacterial indicators of health state in acute murine colitis. *ISME J*. 2012;6:2091–106.
48. Hill DA, Hoffmann C, Abt MC, Du Y, Kobuley D, Kirn TJ, Bushman FD, Artis D. Metagenomic analyses reveal antibiotic-induced temporal and spatial changes in intestinal microbiota with associated alterations in immune cell homeostasis. *Mucosal Immunol*. 2010;3:148–58.
49. Danzeisen JL, Kim HB, Isaacson RE, Tu ZJ, Johnson TJ. Modulations of the chicken cecal microbiome and metagenome in response to anticoccidial and growth promoter treatment. *PLoS One*. 2011;6:e27949.
50. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30:2114–20.
51. Peng Y, Leung HC, Yiu SM, Chin FY. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*. 2012;28:1420–8.

52. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics*. 2011;27:578–9.
53. Hunt M, Newbold C, Berriman M, Otto TD. A comprehensive evaluation of assembly scaffolding tools. *Genome Biol*. 2014;15:R42.
54. Jolley KA, Bliss CM, Bennett JS, Bratcher HB, Brehony C, Colles FM, Wimalaratna H, Harrison OB, Sheppard SK, Cody AJ, Maiden MC. Ribosomal multilocus sequence typing: universal characterization of bacteria from domain to strain. *Microbiology*. 2012;158:1005–15.
55. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res*. 2014;42:D206–14.
56. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome assemblies. *Bioinformatics*. 2013;29:1072–5.
57. Kopylova E, Noe L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics*. 2012;28:3211–7.
58. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*. 2012;19:455–77.
59. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. Clustal W and Clustal X version 2.0. *Bioinformatics*. 2007;23:2947–8.
60. Ronquist F, Huelsenbeck JP. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*. 2003;19:1572–4.
61. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res*. 2016;44:W242–5.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

