



# Whole Genome Sequencing of Extended-Spectrum- and AmpC- $\beta$ -Lactamase-Positive Enterobacterales Isolated From Spinach Production in Gauteng Province, South Africa

Loandi Richter<sup>1,2</sup>, Erika M. du Plessis<sup>1,2\*</sup>, Stacey Duvenage<sup>1,2</sup>, Mushal Allam<sup>3</sup>, Arshad Ismail<sup>3</sup> and Lise Korsten<sup>1,2</sup>

<sup>1</sup>Department of Plant and Soil Sciences, University of Pretoria, Pretoria, South Africa, <sup>2</sup>Department of Science and Innovation, National Research Foundation Centre of Excellence in Food Security, Pretoria, South Africa, <sup>3</sup>Sequencing Core Facility, National Institute for Communicable Diseases, National Health Laboratory Service, Johannesburg, South Africa

## OPEN ACCESS

### Edited by:

Aloysius Wong,  
Kean University-Wenzhou, China

### Reviewed by:

Ralf Lopes,  
University of São Paulo, Brazil  
Masaki Yamamoto,  
Kyoto University Hospital, Japan

### \*Correspondence:

Erika M. du Plessis  
Erika.duplessis@up.ac.za

### Specialty section:

This article was submitted to  
Food Microbiology,  
a section of the journal  
Frontiers in Microbiology

Received: 06 July 2021

Accepted: 01 September 2021

Published: 01 October 2021

### Citation:

Richter L, du Plessis EM, Duvenage S, Allam M, Ismail A and Korsten L (2021) Whole Genome Sequencing of Extended-Spectrum- and AmpC-  $\beta$ -Lactamase-Positive Enterobacterales Isolated From Spinach Production in Gauteng Province, South Africa. *Front. Microbiol.* 12:734649. doi: 10.3389/fmicb.2021.734649

The increasing occurrence of multidrug-resistant (MDR) extended-spectrum  $\beta$ -lactamase- (ESBL) and/or AmpC  $\beta$ -lactamase- (AmpC) producing Enterobacterales in irrigation water and associated irrigated fresh produce represents risks related to the environment, food safety, and public health. In South Africa, information about the presence of ESBL/AmpC-producing Enterobacterales from non-clinical sources is limited, particularly in the water-plant-food interface. This study aimed to characterize 19 selected MDR ESBL/AmpC-producing *Escherichia coli* ( $n = 3$ ), *Klebsiella pneumoniae* ( $n = 5$ ), *Serratia fonticola* ( $n = 10$ ), and *Salmonella enterica* ( $n = 1$ ) isolates from spinach and associated irrigation water samples from two commercial spinach production systems within South Africa, using whole genome sequencing (WGS). Antibiotic resistance genes potentially encoding resistance to eight different classes were present, with *bla*<sub>CTX-M-15</sub> being the dominant ESBL encoding gene and *bla*<sub>ACT</sub>-types being the dominant AmpC encoding gene detected. A greater number of resistance genes across more antibiotic classes were seen in all the *K. pneumoniae* strains, compared to the other genera tested. From one farm, *bla*<sub>CTX-M-15</sub>-positive *K. pneumoniae* strains of the same sequence type 985 (ST 985) were present in spinach at harvest and retail samples after processing, suggesting successful persistence of these MDR strains. In addition, ESBL-producing *K. pneumoniae* ST15, an emerging high-risk clone causing nosocomial outbreaks worldwide, was isolated from irrigation water. Known resistance plasmid replicon types of Enterobacterales including IncFIB, IncFIA, IncFII, IncB/O, and IncHI1B were observed in all strains following analysis with PlasmidFinder. However, *bla*<sub>CTX-M-15</sub> was the only  $\beta$ -lactamase resistance gene associated with plasmids (IncFII and IncFIB) in *K. pneumoniae* ( $n = 4$ ) strains. In one *E. coli* and five *K. pneumoniae* strains, integron In191 was observed. Relevant similarities to human pathogens were predicted with PathogenFinder for all 19 strains, with a confidence of

0.635–0.721 in *S. fonticola*, 0.852–0.931 in *E. coli*, 0.796–0.899 in *K. pneumoniae*, and 0.939 in the *S. enterica* strain. The presence of MDR ESBL/AmpC-producing *E. coli*, *K. pneumoniae*, *S. fonticola*, and *S. enterica* with similarities to human pathogens in the agricultural production systems reflects environmental and food contamination mediated by anthropogenic activities, contributing to the spread of antibiotic resistance genes.

**Keywords:** WGS, food safety, leafy greens, multidrug resistance, foodborne bacterial pathogens

## INTRODUCTION

The discovery of antibiotics in the 1940s led to a new age in medical care. However, the global increase in antimicrobial resistance (AMR) is reducing the effectiveness of clinically important antibiotics (Lobanovska and Pilla, 2017; Dandachi et al., 2019). Examples of shifting resistance profiles in bacteria within the  $\beta$ -lactam class of antibiotics, including penicillins and third generation cephalosporins, which are the most widely used in human and veterinary medicine and widely expressed AMR are being reported (Finton et al., 2020). Persistent exposure to these antibiotics has resulted in bacteria becoming resistant by evolving extended-spectrum  $\beta$ -lactamases (ESBLs), which hydrolyze the  $\beta$ -lactam ring within the antibiotic, thus rendering it inactive (Bush and Jacoby, 2010). Consequently, production of ESBLs is regarded as one of the most clinically significant resistance mechanisms (Bush and Jacoby, 2010), with ESBL-producing Enterobacterales (*Escherichia coli*, *Klebsiella pneumoniae*, and *Serratia* spp., among others) listed as priority pathogens for research and development in the new frontier of antibiotics (WHO, 2017).

Classified into several groups according to their amino acid sequence homology, the CTX-M, TEM, and SHV ESBL variants are the most common  $\beta$ -lactamases identified in Enterobacterales (van Duin and Doi, 2017). In addition, AmpC  $\beta$ -lactamases (AmpCs) are chromosomally encoded by several Enterobacterales species and play a key role in resistance development (van Duin and Doi, 2017). Plasmid encoded AmpC genes have been known since 1989 (Jacoby, 2009) and are now regularly reported in clinical and environmental strains (Khari et al., 2016; Colosi et al., 2020; Tekele et al., 2020). Both chromosomally encoded and plasmid-mediated AmpC  $\beta$ -lactamases confer resistance to a broad spectrum of  $\beta$ -lactams such as penicillins, oxyimino-cephalosporins (including cefotaxime and ceftazidime), cephamycins, and aztreonam at variable levels (Jacoby, 2009; Palzkill, 2018; Furlan and Stehling, 2021; Lopes et al., 2021b).

The increase in antimicrobial resistant strains and effective resistance mechanisms among Enterobacterales has led to numerous global reports of ESBLs, AmpC-, and more recently carbapenemase-producing Enterobacterales not only in clinical settings, but also in the agricultural environment (Ye et al., 2017; Al-Kharousi et al., 2019; Dandachi et al., 2019; Hassen et al., 2020; Richter et al., 2020). Although members of the Enterobacterales family occur naturally in human and animals' gastrointestinal tracts as well as in the environment (water, soil, and plants; Blaak et al., 2014; Ye et al., 2017), the occurrence of multidrug-resistant (MDR) strains in the different habitats

is concerning. Inadequately treated or untreated effluents from industries, households, and zootechnical farms are reported as one of the main contamination causes of South African surface- and ground water resources (Verlicchi and Grillini, 2020). It is also well-documented that the three principal antibiotic contamination channels in the environment are animal-, human-, and manufacturing waste (O'Neill, 2016). Consequently, contamination of soil, irrigation-, and drinking water as well as crops can occur, adding additional exposure routes to humans (Finton et al., 2020; Lopes et al., 2021a).

Previous surveillance studies have shown prevalence of MDR ESBL/AmpC-producing Enterobacterales in fresh vegetables sold in South Africa (Richter et al., 2019) and in other countries e.g., the Netherlands, Switzerland, and Germany (Reuland et al., 2014; Zurfluh et al., 2015; Reid et al., 2020). Occurrence of ESBL-producing Enterobacterales has also been reported in corresponding irrigation water sources and cultivated crops (Blaak et al., 2014; Njage and Buys, 2014; Ye et al., 2017). Furthermore, Richter et al. (2020) reported the occurrence of ESBL/AmpC-producing Enterobacterales in different spinach supply chains from irrigation water and produce at harvest, throughout processing and at retail in the Gauteng Province of South Africa.

The high discriminatory power of whole genome sequencing (WGS) has led to an increase in use of this method for detecting points of contamination, source tracking, pathogen surveillance, and outbreak investigations (Oniciuc et al., 2018; CDC, 2019). WGS provides information regarding multiple AMR genes, genomic mutations, mobile genetic elements, and association with resistance genes, as well as molecular typing like multi-locus sequence typing (MLST; Oniciuc et al., 2018; CDC, 2019; Kim et al., 2020). Consequently, the WGS results can aid in elucidating the genetic relationship among isolates from different environments and along the food chain (Adator et al., 2020). Surveillance of antimicrobial resistant strains through WGS is increasingly being used due to increasing accessibility and affordability (Adator et al., 2020). In South Africa, WGS has been used for characterization of clinical ESBL-producing *K. pneumoniae* strains among others (Founou et al., 2019), as well as typing of *Listeria monocytogenes* from environmental and clinical settings during the 2017 listeriosis outbreak (Thomas et al., 2020). However, the use of WGS for surveillance of antimicrobial resistant potential pathogenic Enterobacterales in retailed fresh produce and the production environment has not been reported locally.

The World Health Organization (WHO) developed the Global AMR Surveillance System (GLASS) in 2015 supporting research

and surveillance as well as a global data sharing through a standardized analysis approach (WHO, 2020). Initially, the GLASS focus was mainly on surveillance of human priority pathogens, but has since shifted to include AMR in foodborne pathogens (WHO, 2020). Moreover, the One Health framework for understanding AMR in pathogenic gram-negative bacteria is increasingly attracting attention (Collignon and McEwen, 2019). In South Africa, information regarding AMR in fresh produce production systems and specifically focusing on the Enterobacteriales is lacking. The aim of this study was thus to use WGS for analysis of AMR genes, associated mobile genetic elements, virulence factors, serotypes, multi-locus sequence types, and pathogenicity of selected, partially characterized, ESBL/AmpC-producing environmental Enterobacteriales from commercial spinach production systems (Richter et al., 2020). These isolates included four different species (*E. coli*, *K. pneumoniae*, *S. fonticola*, and *S. enterica*) listed by the WHO as a particular threat for gram-negative bacteria that are resistant to multiple antibiotics (WHO, 2017), while isolates harbouring integrons as described in Richter et al. (2020) were preferentially selected. The results of this study will contribute to address the problem of antimicrobial drug resistance at the water–plant–food interface and how it might impact human health and disease.

## MATERIALS AND METHODS

### Isolation and DNA Extraction of ESBL/AmpC-Producing Enterobacteriales

Irrigation water and fresh produce samples from spinach production systems were collected and ESBL-producing Enterobacteriales were isolated as described (Richter et al., 2020). A selection of 19 isolates were further characterized (Table 1). The genomic DNA of each isolate was extracted with the DNeasy PowerSoil kit (Qiagen, South Africa) according to the manufacturer's instructions. Following gDNA extraction, the concentrations were determined using the Qubit dsDNA Broad Range Assay and a Qubit 2.0 fluorometer (Life Technologies, Johannesburg) and quantification was determined on a Nanodrop 2000 (ThermoScientific, Johannesburg).

### DNA Sequencing and Whole Genome Analysis

Sequencing was performed on an Illumina MiSeq instrument (2 × 300 bp) with 100× coverage by the National Institute for Communicable Diseases Sequencing Core Facility, South Africa, following preparation of multiplexed paired-end libraries with the Nextera XT DNA sample preparation kit (Illumina, San Diego, CA, United States). The resultant reads were quality trimmed using CLC version 20<sup>1</sup> and *de novo* assembled. The contiguous sequences were then submitted to the National Centre for Biotechnology Information (NCBI) Prokaryotic Genome Annotation Pipeline.<sup>2</sup> The AMR gene presence was

**TABLE 1** | Isolates selected for whole genome sequence analysis from the agricultural environment in spinach supply chains, Gauteng Province, South Africa.

Strain	Organism identity	Isolation point from spinach production systems	
		Water (W) or spinach (S)	
UPMP2117	<i>Escherichia coli</i>	W	Water reservoir
UPMP2120	<i>Escherichia coli</i>	S	Unwashed spinach bunches at retailer
UPMP2130	<i>Escherichia coli</i>	W	Holding dam water (source water)
UPMP2112	<i>Klebsiella pneumoniae</i>	W	Irrigation pivot point water
UPMP2114	<i>Klebsiella pneumoniae</i>	S	Spinach at harvest
UPMP2118	<i>Klebsiella pneumoniae</i>	W	Irrigation pivot point water
UPMP2121	<i>Klebsiella pneumoniae</i>	S	Unwashed spinach bunches at retailer
UPMP2122	<i>Klebsiella pneumoniae</i>	S	Spinach at retailer
UPMP2115	<i>Salmonella spp.</i>	W	River water
UPMP2116	<i>Serratia fonticola</i>	W	River water
UPMP2119	<i>Serratia fonticola</i>	W	Irrigation pivot point water
UPMP2123	<i>Serratia fonticola</i>	S	Unwashed spinach punnet at retailer
UPMP2124	<i>Serratia fonticola</i>	S	Spinach at receipt
UPMP2125	<i>Serratia fonticola</i>	S	Spinach after pack
UPMP2126	<i>Serratia fonticola</i>	S	Spinach at receipt
UPMP2127	<i>Serratia fonticola</i>	S	Unwashed spinach at retailer
UPMP2128	<i>Serratia fonticola</i>	S	Unwashed spinach at retailer
UPMP2129	<i>Serratia fonticola</i>	S	Spinach at receipt
UPMP2131	<i>Serratia fonticola</i>	S	Unwashed spinach at retailer

corroborated using ABRicate<sup>3</sup> that included the Comprehensive Antibiotic Resistance Database (CARD), ARG-ANNOT, ResFinder, NCBI AMRFinder Plus, and MEGAREs databases (Zankari et al., 2012; Gupta et al., 2014; Jia et al., 2017; Feldgarden et al., 2019; Doster et al., 2020). Plasmid replicon types were determined with PlasmidFinder (version 2.1; Carattoli et al., 2014). Using the Centre for Genomic Epidemiology (CGE) platform<sup>4</sup>, mobile genetic elements for all four species, sequence types of *E. coli*, *K. pneumoniae*, and *S. enterica* as well as the *E. coli* serotypes based on lipopolysaccharide (O-antigen) and capsular flagella (protein; H-antigen), and virulence genes of *E. coli* were determined with MGEFinder, Multilocus Sequence Typing (MLST; version 2.2), SeroTypeFinder (version 2.0), and VirulenceFinder (version 2.0), respectively (Larsen et al., 2012; Joensen et al., 2014, 2015; Johansson et al., 2021). The following parameters were used in the Serotype Finder Web-based tool: 85% threshold for %ID and 60% minimum length (the number of nucleotides in a sequence of interest that must overlap a serotype gene to count as a hit for that gene; Joensen et al., 2015). The *in silico* serotyping based on the capsule polysaccharide (K-antigen) of *K. pneumoniae* strains was conducted using Kaptive Web (Wick et al., 2018), while the presence of virulence genes for *K. pneumoniae* was

<sup>1</sup><https://digitalinsights.qiagen.com>

<sup>2</sup><https://pubmed.ncbi.nlm.nih.gov/27342282/>

<sup>3</sup><https://github.com/tseemann/abricate>

<sup>4</sup><https://cge.cbs.dtu.dk/services/>

identified by using the Institut Pasteur's *Klebsiella* database.<sup>5</sup> Additionally, paired reads of the WGS raw data files for the *S. enterica* strain were uploaded to the online SeroSeq tool version 1.0 which predicted the *Salmonella* serotype of the requested isolate (Zhang et al., 2015; Thompson et al., 2018). The *Salmonella* Pathogenicity Islands (SPI) were identified with SPIFinder 2.0 (Roer et al., 2016). Next, the existence of virulence factors in each SPI was analyzed by performing BLAST analysis on the predicted SPIs against the virulence factor database (VFDB; Chen et al., 2016; Ashari et al., 2019). The virulence factors of *S. fonticola* were determined using the VFDB with ABRicate (Chen et al., 2016). All sequences were submitted to the INTEGRALL database<sup>6</sup> for annotation and integron number assignment. Using PathogenFinder (version 1.1) on the CGE platform, the strains' pathogenicity towards humans was predicted (Cosentino et al., 2013).

## Data Availability

The nucleotide sequences of the 19 Enterobacterales strains described in this paper were deposited in the National Center for Biotechnology Information GenBank database in the BioProject number: PRJNA642017, accession numbers NZ\_JACAAL010000000, NZ\_JACBIV000000000-NZ\_JACBJE000000000, and NZ\_JACNY M000000000-NZ\_JACNYT000000000 (Table 2).

## RESULTS

### Detection of Antimicrobial Resistance Genes

The selected 19 ESBL/AmpC producing Enterobacterales isolates all harboured at least one  $\beta$ -lactamase encoding gene in addition to the ESBL/AmpC genetic determinants, accompanied by resistance genes from different antibiotic classes including fluoroquinolone, sulfonamide, fosfomycin, aminoglycoside, trimethoprim, phenicol, and/or tetracycline (Figure 1). The  $\beta$ -lactamase resistance genes included chromosomally encoded AmpC in the *S. enterica* strain as well as all three *E. coli* strains. Plasmid-mediated AmpC genes (*bla*<sub>CMY-113</sub> and *bla*<sub>CMY-101</sub>) were present in two *E. coli* strains from irrigation water and *bla*<sub>ACT-13</sub>, *bla*<sub>ACT-38</sub>, *bla*<sub>ACT-6</sub>, and/or *bla*<sub>ACT-58</sub> were present in 10 *S. fonticola* strains from irrigation water ( $n=2$ ) and spinach ( $n=8$ ) samples (Figure 1). Additionally, *bla*<sub>FONA-5</sub> ( $n=8$ ) from irrigation water and spinach and *bla*<sub>FONA-6</sub> ( $n=2$ ) from spinach were present in *S. fonticola* strains. The ESBL genes included *bla*<sub>SFO-1</sub> in all 10 *S. fonticola* strains, *bla*<sub>CTX-M-15</sub> in five *K. pneumoniae* strains from irrigation water and spinach, and one *E. coli* strain from spinach. It also included *bla*<sub>CTX-M-14</sub> in an *E. coli* strain from irrigation water, while *bla*<sub>SHV-187</sub> ( $n=3$ ), *bla*<sub>SHV-106</sub> ( $n=1$ ), and *bla*<sub>SHV-178</sub> ( $n=1$ ) were present in *K. pneumoniae* strains (Figure 1).

Interestingly, a greater number of resistance genes across more classes were seen in all the *K. pneumoniae* strains ( $n=5$ ), compared to the other genera tested. All five *K. pneumoniae* strains had

chloramphenicol (*catB3*), aminoglycosides [*aac*(6')-Ib-cr, *aph*(6)-Ia and *aph*(3'')-Ib], fosfomycin (*fosA6*), and sulfonamide (*sul2*) resistance genes present (Figure 1). Other resistance genes included fluoroquinolone *oqxA* ( $n=4$ ), *oqxB* ( $n=4$ ), and *qnrB1* ( $n=4$ ) in *K. pneumoniae* from spinach and water, *qnrS1* ( $n=1$ ) in *E. coli* from spinach and *qnrB6* ( $n=3$ ), *qnrB37* ( $n=5$ ), *qnrE1* ( $n=10$ ) in *S. fonticola* from spinach and water, while *mdtK* ( $n=4$ ), and *mdtH* ( $n=3$ ) were present in *S. fonticola* from water only. The *qnrB17* resistance gene was present in *K. pneumoniae* ( $n=4$ ) and *S. fonticola* ( $n=2$ ) strains from spinach and water (Figure 1). The *S. enterica* strain isolated from irrigation water also harboured *aac*(6')-Iaa and *aac*(6')-Iy aminoglycoside resistance genes (Figure 1) and a *S. fonticola* strain from irrigation water harboured an aminoglycoside [*aph*(3'')-Ib] and sulfonamide (*sul2*) resistance gene (Figure 1).

### Detection of Mobile Genetic Elements and Association to Antimicrobial Resistance Genes

Known resistance plasmid replicon types of Enterobacterales including IncFIB, IncFIA, IncFII, IncB/O, and IncHI1B were observed in all strains following analysis with PlasmidFinder (data not shown). The  $\beta$ -lactamase gene, *bla*<sub>CTX-M-15</sub>, was the only resistance gene associated with plasmids (IncFII\_pKP91 and/or IncFIB(K)\_1\_Kpn3) in four *K. pneumoniae* strains upon further analysis (Table 3). The IS6 family elements (IS6100) have been reported to play a pivotal role in the dissemination of resistance determinants in gram-negative bacteria (Partridge et al., 2018), and were observed in relation to the *dfrA14b* resistance gene in all five *K. pneumoniae* strains (Table 3). The *bla*<sub>CTX-M-14</sub> and *sul2* resistance genes were related to the ISEcp1 element within the IS1380 family in one *E. coli* and three *K. pneumoniae* strains, respectively, while one *S. fonticola* strain carried a *sul2* gene that was related to IS110 (Table 3). One *E. coli* strain carried the *qnrS1* resistance gene that was related to ISKra4. Other insertion sequences detected belonged predominantly to the IS3 and IS110 families (data not shown), with one *K. pneumoniae* strain carrying the *bla*<sub>SHV-80</sub> broad spectrum  $\beta$ -lactamase that was related to IS3 (Table 3). In all *K. pneumoniae* strains ( $n=5$ ) where the *qnrB1* resistance gene was present, association to Tn5403 was seen (Table 1). In one *E. coli* and five *K. pneumoniae* strains, integron In191 was observed, with *dfrA14* in the cassette array (Table 3).

### In silico Analysis of Serotypes, Multi-Locus Sequence Types, and Virulence Factors

The *in silico* MLST analysis, predicted serotypes, and pathogenicity probability of all 19 strains are shown in Table 2. Three different sequence types (ST58, ST117, and ST10) and three different serotypes (O75:H9, O11:H4, and O8:H17) were observed in the three *E. coli* strains. The five *K. pneumoniae* strains belonged to three different sequence types and three different serotypes (KL27, KL24, and KL39) which were observed based on the K-antigen, while the O-serotype included O4 and O1 (Table 2). The predicted antigenic profile of the *S. enterica* strain was O11:k:1,2. Furthermore, the *S. enterica* strain contained 11 *Salmonella* SPI, namely SPI-1,

<sup>5</sup><https://bigsd.bpasteur.fr/klebsiella/klebsiella.html>

<sup>6</sup><http://integrall.bio.ua.pt>





**TABLE 3** | Extended-spectrum  $\beta$ -lactamase (ESBL)/AmpC  $\beta$ -lactamase (AmpC)-producing Enterobacterales with resistance genes related to mobile genetic elements.

Isolate information			Resistance genes associated with mobile genetic elements					
Source	Strain	Species	Genes			Mobile genetic elements		
			$\beta$ -lactamase	Other	Plasmids	Insertion sequence families	Transposons	Integron
W	UPMP2130	<i>Escherichia coli</i>	CTX-M-14			IS1380		
S	UPMP2120	<i>Escherichia coli</i>		qnrS1 dfrA14b		ISKra4		In191
W	UPMP2112	<i>Klebsiella pneumoniae</i>	SHV-80 CTX-M-15	sul2 qnrB1 dfrA14b	IncFIB(K)_1_ Kpn3	IS3 IS1380		Tn5403 In191
W	UPMP2118	<i>Klebsiella pneumoniae</i>	TEM-1B	dfrA14b qnrB1		IS1380 IS6		In191 Tn5403
S	UPMP2114	<i>Klebsiella pneumoniae</i>	CTX-M-15	sul2 qnrB1 dfrA14b	IncFII_pKP91 IncFIB(K)_1_ Kpn3	IS1380 IS6		Tn5403 In191
S	UPMP2121	<i>Klebsiella pneumoniae</i>	CTX-M-15 TEM-1B	qnrB1 dfrA14b	IncFII_pKP91	IS1380 IS6		Tn5403 In191
S	UPMP2122	<i>Klebsiella pneumoniae</i>	CTX-M-15	sul 2 qnrB1 dfrA14b	IncFII_pKP91 IncFIB(K)_1_ Kpn3	IS1380 IS6		Tn5403 In191
W	UPMP2116	<i>Serratia fonticola</i>		sul2		IS110		

W, water; S, spinach.

strains (**Supplementary Table S4**). This included 25, 18, 16, and 6 of the virulence factors present in 100% ( $n=10$ ), 90, 80, and 70% of the selected *S. fonticola* strains, respectively, while the remaining 24 virulence factors were present in varying numbers in 1–6 of the strains (**Supplementary Table S4**). The *iroN* salmochelin siderophore receptor which plays a role in disease establishment was present in three *S. fonticola* strains (two from unwashed baby spinach samples at the retailer and one from the irrigation pivot point water), one *E. coli* strain from the ground water, as well as in the SPI-13 in the *S. enterica* strain from river irrigation water. Relevant similarities to human pathogens were predicted for all 19 strains with a confidence of 0.635–0.721 in the *S. fonticola* strains ( $n=10$ ), 0.852–0.931 in the *E. coli* strains ( $n=3$ ), 0.796–0.899 in the *K. pneumoniae* strains ( $n=5$ ), and 0.939 in the *S. enterica* strain (**Table 2**).

## DISCUSSION

To the authors knowledge, this is the first study to use WGS for in-depth molecular characterization of ESBL/AmpC-producing

*E. coli*, *K. pneumoniae*, *S. enterica*, and *S. fonticola* isolates, previously identified and partially characterized, from spinach and irrigation water samples in commercial production chains (Richter et al., 2020). Characterization included AMR, mobile genetic elements (e.g., insertion sequences, plasmids, and integrons), serotypes, and determining the pathogenicity. All these factors are crucial in defining and attributing infection sources of food-related outbreaks caused by resistant microorganisms (Oniciuc et al., 2018). Overall, the results corresponded with main global findings where AMR genes and associated mobile genetic elements have been reported in Enterobacterales from fresh produce and irrigation water, with the potential to pose a health risk to humans upon exposure (Jones-Dias et al., 2016; Finton et al., 2020). Previously, the presence of *intI3* was reported in a high percentage of isolates from the current study following conventional PCR and sequencing (Richter et al., 2020). However, in-depth WGS analysis showed that no *attI* fragment preceded the *IntI3* genes; consequently, the *IntI3* genes detected did not form part of complete integrons, which typically include an integrase *intI* gene encoding a site-specific recombinase, a recombination site *attI* as well as a promoter ( $P_{\int}$ ; Kaushik et al., 2018). Overall, six isolates in the current study

were positive for Class 1 integrons (In191), similar to In191 positive clinical ESBL-producing Enterobacterales from an academic teaching hospital in Pretoria, SA (Sekyere et al., 2020). Additionally, these MDR environmental isolates harbored various virulence factors central to pathogenicity, including genes associated with urinary tract infections and iron sequestering systems crucial for disease establishment. All isolates had relevant similarity to human pathogens and form part of the WHO 3rd generation cephalosporin resistant critical priority pathogens (WHO, 2017).

Two of the *E. coli* strains from the current study harboured plasmid-mediated AmpC *bla*<sub>CMY-2-like</sub> genes (*bla*<sub>CMY-113</sub> and *bla*<sub>CMY-101</sub>), which correspond to the phenotypic profile of resistance to expanded-spectrum cephalosporins previously reported for these isolates using traditional PCR analysis (Richter et al., 2020). The *bla*<sub>CMY-2</sub> pAmpC genes are the most commonly reported in *E. coli* and other Enterobacterales species and have clinical relevance, as it inactivates third generation cephalosporins and mediates resistance to carbapenems (Jacoby, 2009; Bortolaia et al., 2014). Three different multi-locus sequence types, namely ST58, ST10, and ST117, were identified in the *E. coli* isolates. Isolated from the retail unwashed spinach samples in the current study, ST58 *E. coli* have previously also been associated with human extra-intestinal infections including sepsis, and have emerged worldwide in wild and food-production animals (Reid et al., 2020). As an example, ST58 *E. coli* with serotype O75:H9 corresponded to an *E. coli* strain of bovine origin from Pakistan and also carried the IncFIB plasmid (Ali et al., 2020). Although the strain from the current study had less AMR genes than reported in ST58 *E. coli* with serotype O75:H9 by Ali et al. (2020), the trimethoprim (*dhfrA14*), fluoroquinolone (*qnrS1*), and  $\beta$ -lactam (*bla*<sub>CTX-M-15</sub>) genes corresponded. Similarly, uropathogenic ST58 *E. coli* with resistance to fluoroquinolone and trimethoprim have previously been isolated from hospital patients in Australia (McKinnon et al., 2018). The *bla*<sub>CTX-M-15</sub> gene identified in the ST58 *E. coli* strain from the current study was associated with the ISKra4 insertion sequence, previously identified in *K. pneumoniae* harbouring *bla*<sub>CTX-M-15</sub>, and was responsible for the movement to different parts of the genome through a replicative transposition mechanism (Razavi et al., 2020). In contrast to Hauser et al. (2013) who identified food-associated shiga-toxin producing *E. coli* ST58, no *stx* genes were present in the strains. The *E. coli* ST58 from the current study harboured the *gad* (glutamate decarboxylase) virulence gene, similar to *E. coli* ST58 strains isolated from arugula (rocket; Reid et al., 2020). However, the presence of *lpfA* (long polar fimbriae) and *terC* (tellurium ion resistance protein) virulence factors in the strain from the current study contrasted the virulence gene profiles reported by Reid et al. (2020). *Escherichia coli* ST10 have previously been associated with human clinical infections and has been isolated from different sources including recreational and/or wastewater samples (Falgenhauer et al., 2019). From the current study, the *E. coli* ST10 with serotype O8:H17 was isolated from borehole water used for irrigation. Although this sequence type has previously been associated with shiga-toxin-producing *E. coli* (STEC; Gonzalez-Escalona and Kase, 2018), no *stx* genes were detected in the current study. The virulence factors present

were *terC* (tellurium ion resistance protein), *astA* (EAST-1 heat-stable toxin), *fyuA* (ferric yersiniabactin uptake receptor), *irp2* (nonribosomal peptide synthetases), *iss* (increased serum survival), and *sitA* (iron transport protein). Previously, *E. coli* ST10 with similar virulence gene profiles were isolated from human blood cultures and reported as extra-intestinal pathogenic *E. coli* (ExPEC; Maluta et al., 2017). Additionally, ESBL-producing *E. coli* ST10 of the same serotype have been isolated from wastewater and are depicted as a probable environmental reservoir of *bla*<sub>CTX-M</sub> genetic determinants (Tanaka et al., 2019). In the current study, the ST58 *E. coli* strain harboured the *bla*<sub>CTX-M-15</sub> genetic determinant, while *bla*<sub>CTX-M-14</sub> was present in the ST10 *E. coli* strain. Globally, the CTX-M type ESBLs (especially *bla*<sub>CTX-M-14</sub> and *bla*<sub>CTX-M-15</sub>) have become the dominant genotype and the most widely distributed (Cantón et al., 2012; Adamski et al., 2015). *Escherichia coli bla*<sub>CTX-M-14</sub> positive strains have previously been isolated from store-bought produce in Germany and South Africa (Richter et al., 2019; Reid et al., 2020), food producing animals in China (Liao et al., 2015) and clinical settings in Brazil and South Africa (Cergole-Novella et al., 2010; Peirano et al., 2011).

The third *E. coli* sequence type (ST117) detected from irrigation source water in the current study has previously been reported as part of a group of multi-serotype extra-intestinal pathogenic *E. coli* (ExPEC) and avian pathogenic *E. coli* (APEC) strains (Kim et al., 2017). The *E. coli* ST117 strain from the current study harboured 20 virulence factors including the ExPEC *hlyF* (Hemolysin F) virulence gene. In previous studies, *stx* genes were identified in *E. coli* strains with the same STs detected in the current study, yet the virulence gene content and serotypes differ from the strains in the current study (Gonzalez-Escalona and Kase, 2018). However, the three non-STEC *E. coli* strains (ST58, ST10, and ST117) from the current study had a 93, 89, and 85% probability of being human pathogens, based on the pathogenic protein families.

In addition to *E. coli*, other Enterobacterales isolates harbouring *bla*<sub>CTX-M-15</sub> have also been detected in different environments. In the current study, all five *K. pneumoniae* strains harboured the *bla*<sub>CTX-M-15</sub> genetic determinant. The prevalence and dissemination of *bla*<sub>CTX-M</sub> throughout various environments globally underlines the different contamination routes through which fresh produce may also become contaminated with these MDR organisms. For instance, Gekenidis et al. (2020) have demonstrated the long-term persistence of *E. coli* harbouring *bla*<sub>CTX-M-15</sub> in soil and lettuce after its introduction via irrigation water. Similarly, *bla*<sub>CTX-M-15</sub> positive ST985 *K. pneumoniae* strains were present in spinach at harvest on the farm as well as retail samples after processing in the current study, suggesting successful persistence of these MDR strains. In four *K. pneumoniae* strains (ST3559,  $n=1$  and ST985,  $n=3$ ), the *bla*<sub>CTX-M-15</sub> genes were associated with IncF replicons (IncFII<sub>K</sub> and IncFIB) which have previously been linked to diverse *K. pneumoniae* outbreak strains (Dolejska et al., 2012, 2013; Löhr et al., 2015). Moreover, in *K. pneumoniae* ST3559, *bla*<sub>CTX-M-15</sub> was also associated with *ISEcp1* (also called *ISEc9*), a member of the widely reported IS1380 family, and can enable the independent transposition with insertion mutation and

genetic relocations (Partridge, 2011). The *K. pneumoniae* strains in the current study also harboured *bla*<sub>SHV</sub> ESBL encoding genes (*bla*<sub>SHV-187</sub>, *bla*<sub>SHV-106</sub>, and *bla*<sub>SHV-178</sub>). Previously, SHV genetic determinants were reported in *K. pneumoniae* from hospitals and receiving wastewater treatment plants in Romania (Surleac et al., 2020) as well as irrigation water and agricultural soil in South Africa (Iwu et al., 2020; Richter et al., 2020). Interestingly, the *K. pneumoniae* ST15 strain isolated from water in the current study harboured *bla*<sub>SHV-106</sub> which Liakopoulos et al. (2016) previously reported to be geographically constrained and have only been described in *K. pneumoniae* isolates from Portugal together with *bla*<sub>TEM-1</sub>. Similarly, the *K. pneumoniae* ST15 strain from the current study also harboured *bla*<sub>SHV-106</sub> together with *bla*<sub>TEM-1</sub>. *Klebsiella pneumoniae* ST15 is regarded as an emerging international high-risk clone causing nosocomial outbreaks worldwide with high-levels of antibiotic resistance including production of ESBLs, mainly CTX-M-15 (Han et al., 2021).

The *K. pneumoniae* ST3559 strain isolated from irrigation water in the current study was capsular type 27 and serotype O4, which is similar to an O4 serotype MDR *K. pneumoniae* outbreak strain from a neonatal care unit in sub-Saharan Africa (Cornick et al., 2020). In addition, *K. pneumoniae* ST3559 harboured the *bla*<sub>SHV-178</sub> gene which, to the best of our knowledge, have previously only been reported in clinical *Enterobacter hormaechei* strains from the First Affiliated Hospital of Zhejiang University in Hangzhou (Gou et al., 2020). Apart from  $\beta$ -lactamase genes, the *K. pneumoniae* strains also harboured aminoglycoside, fosfomycin, fluoroquinolone, tetracycline, phenicol, trimethoprim, and sulfonamide resistance genes, which is a greater diversity of resistance genes than previously reported in Enterobacterales isolates from German surface waters (Falgenhauer et al., 2019). Similar to the results of clinical *K. pneumoniae* strains reported by Mbelle et al. (2020), In191, harbouring *dfrA14* was identified in the three different *K. pneumoniae* sequence types of the current study, reiterating that it is not a narrow spectrum integron. In addition, *dfrA14b* was associated with *IS6* that has previously been reported as having a vital role in the rearrangement and dissemination of antibiotic resistance (Varani et al., 2021). The presence of *fosA* and *sul2* in all the *K. pneumoniae* strains of the current study also corresponds to the results reported by Mbelle et al. (2020) from clinical *K. pneumoniae* strains in Pretoria. The high-level of trimethoprim resistance globally has however led to trimethoprim-sulfamethoxazole no longer being recommended for outpatient treatment of urinary tract infections and similarly, the use of fosfomycin might not be efficacious anymore (Mbelle et al., 2020). Four MDR *K. pneumoniae* isolates from irrigation water (ST15, *n*=1) and spinach (ST985, *n*=3) had O1 serotypes, previously reported as the most commonly isolated serotypes from human hosts and dominant in human disease (Follador et al., 2016). However, it is noteworthy that no genes encoding carbapenamases nor resistance to colistin were identified in the current study. All five characterized *K. pneumoniae* strains also harbored several virulence factors including those that coded for an iron uptake system (*kfu*) and type 3 fimbrial adhesins (*mrk*) that play an

important role in adhesion to medical devices such as catheters (Albasha et al., 2020; Finton et al., 2020).

*Serratia* spp. are opportunistic pathogens that may pose a health threat to immunocompromised and hospitalized patients (Petersen and Tisa, 2013). The *S. marcescens* species is most often associated with nosocomial infections; however, *S. fonticola* has been reported to function as a human pathogen when detected alone or may be a bystander and act as a carrier of resistance genes when discovered with other organisms (Petersen and Tisa, 2013; Aljorayid et al., 2016). Characterizing virulence genes of the MDR environmental strains therefore becomes important within the plant-food producing environment. In the current study, all *S. fonticola* strains harboured *bla*<sub>SFO-1</sub> and numerous plasmid incompatibility (Inc) groups were identified in these *S. fonticola* strains (data not shown). However more in-depth plasmid typing and analysis will be required to fully understand the risk/probability of *bla*<sub>SFO-1</sub> dissemination in the environment where *S. fonticola* naturally occurs. In certain Enterobacterales species, ESBL genes are inherently carried on chromosomes (Naas et al., 2008). This includes the *bla*<sub>SFO-1</sub> ESBL gene from *S. fonticola* that differs from most class A ESBLs, as the  $\beta$ -lactamases' production can be induced by a high level of imipenem (Naas et al., 2008). The *bla*<sub>SFO-1</sub> ESBL does not form part of the most clinically relevant ESBLs and are therefore rarely reported. Zhou et al. (2020) reported in contrast an increasing trend of the co-existence of plasmid-borne *bla*<sub>SFO-1</sub> and carbapenemase genes in clinical *Enterobacter* spp. in China. All the *S. fonticola* strains also harboured numerous fluoroquinolone resistance genes, raising a health concern for treatment options, as fluoroquinolones are often used for management of conditions including typhoid fever and MDR tuberculosis (Richards et al., 2019). Interestingly, one *S. fonticola* strain harboured an acquired trimethoprim (*sul2*) resistance gene associated with IS110, corresponding to *K. pneumoniae* from a German university hospital (Schwanbeck et al., 2021). The *Serratia* genus naturally lacks resistance genes for trimethoprim and sulfonamides (Sandner-Miranda et al., 2018). Previous reports of potential pathogenic *S. fonticola* primarily focused on the antibiotic resistance profiles (Tasić et al., 2013; Aljorayid et al., 2016; Hai et al., 2020). The strains from the current study additionally harboured various virulence factors. This included flagellar biosynthesis- and chemotaxis-related genes as well as genes encoding iron uptake systems corresponding to those previously reported in important MDR nosocomial pathogenic *S. marcescens* (Iguchi et al., 2014).

Only one *S. enterica* strain isolated from river irrigation water was characterized in the current study. Irrigation water is well documented as a source for fresh produce contamination of foodborne pathogens including *Salmonella* spp. (Liu et al., 2018). The strain harboured an AmpC resistance gene, similar to *S. enterica* characterized from surface water in the United States (Li et al., 2014). In addition, the *S. enterica* from the current study carried aminoglycoside resistance genes [*aac*(6')-Iaa and *aac*(6')-Iy], similar to results reported by Nair et al. (2016) for non-typhoidal *Salmonella* spp. isolated from a United Kingdom population. Of the 23 known *Salmonella* SPIs previously described (Mansour et al., 2020), the isolate from



the current study carried 11 SPIs. This included SPIs that are commonly reported in *S. enterica* and encode genes responsible for enabling invasion of epithelial cells (SPI1), facilitating the replication of intracellular bacteria (SPI2), adhesion to epithelial cells (SPI3, 4, 5, and 9; Waterman and Holden, 2003; Velásquez et al., 2016; Mansour et al., 2020), as well as SPI13 and 14 which corresponds to being part of the core genome of invasive non-typhoidal *Salmonella* spp. (Suez et al., 2013). Additionally, pathogenicity islands C63PI and CS54 were present in the *S. enterica* strain in this study, which has previously been found in the *S. Typhimurium* and *S. Typhi* genomes (Sabbagh et al., 2010; Jibril et al., 2021). Since no phenotypic indication of virulence was investigated, the prediction of virulence genes using *in silico* tools should be regarded with care; however, using PathogenFinder, the *S. enterica* strain from the current study showed 94% probability of being a human pathogen.

## CONCLUSION

This is the first WGS analysis study of MDR ESBL/AmpC-producing *E. coli*, *K. pneumoniae*, *S. fonticola*, and *S. enterica* isolates from spinach production systems within South Africa. The selected isolates represent potential pathogenic genera listed by the WHO as a priority for surveillance of AMR screening. Numerous clinically relevant resistance genes were detected in the screened samples. This study showed the potential of using WGS in metadata studies for detailed molecular characterization of potential pathogenic Enterobacterales. Furthermore, the study highlighted the importance of the agricultural production environment as a source of antibiotic resistance genes within Enterobacterales at the water-plant-food interface. A more in-depth and controlled analysis, with a greater number of sequenced isolates from the farm-to-retail supply chain is required to better understand the prevalence and resistance gene transmission through the supply chain. The results from this study further highlight the need for expanded surveillance in agricultural systems.

## DATA AVAILABILITY STATEMENT

The nucleotide sequences of the 19 Enterobacteriaceae strains described in this paper were deposited in the National Center for Biotechnology Information GenBank database in the BioProject number: PRJNA642017, accession numbers NZ\_JAC AAL010000000, NZ\_JACBIV000000000-NZ\_JACBJE000000000, and NZ\_JACNYM000000000-NZ\_JACNYT000000000 (Table 2).

## REFERENCES

- Adamski, C. J., Cardenas, A. M., Brown, N. G., Horton, L. B., Sankaran, B., Prasad, B. V. V., et al. (2015). Molecular basis for the catalytic specificity of the CTX-M extended-spectrum  $\beta$ -lactamases. *Biochemistry* 54, 447–457. doi: 10.1021/bi501195g
- Adator, E. H., Walker, M., Narvaez-Bravo, C., Zaheer, R., Goji, N., Cook, S. R., et al. (2020). Whole genome sequencing differentiates presumptive extended spectrum beta-lactamase producing *Escherichia coli* along segments of

## AUTHOR CONTRIBUTIONS

EP, SD, LR, and LK contributed to the conception and design of the study. LR performed the experiments. LR, SD, MA, and AI analyzed the data. LR, EP, and SD contributed to interpretation and presentation. SD, EP, and LK were involved in funding acquisition. All authors contributed to the article and approved the submitted version.

## FUNDING

This research was funded by the Water Research Commission (WRC) funded project “Measurement of water pollution determining the sources and changes of microbial contamination and impact on food safety from farming to retail level for fresh vegetables” (WRC Project No K5/2706/4 and WRC Knowledge Review 2017/18) and the Partnerships for Enhanced Engagement in Research (PEER) a USAID/DST funded project “Characterizing and tracking of antimicrobial resistance in the water-plant-food public health interface” (Grant no. 48). Furthermore, the financial assistance of the Department of Science and Innovation (DSI) – National Research Foundation (NRF) Centre of Excellence in Food Security funded this research under the Food Safety Program’s “Safe Food for the Food Insecure” project (Project 160301 and 160302). This work is based on the research supported in part by the NRF of South Africa [Grant specific unique reference number (UID): 74426 and Grant Number: 120319]. Opinions expressed and conclusions arrived at, are those of the authors and are not necessarily to be attributed to the NRF.

## ACKNOWLEDGMENTS

The authors acknowledge the University of Pretoria, DSI-NRF Centre of Excellence in Food Security, PEER, and the WRC for funding and bursary support. The opinions, findings, conclusions, and/or recommendations expressed in this publication are the author(s) alone.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2021.734649/full#supplementary-material>

the one health continuum. *Microorganisms* 8:448. doi: 10.3390/microorganisms8030448

- Albasha, A. M., Osman, E. H., Abd-Alhalim, S., Alshuib, E. F., Al-Hassan, L., and Altayb, H. N. (2020). Detection of several carbapenems resistant and virulence genes in classical and hyper-virulent strains of *Klebsiella pneumoniae* isolated from hospitalized neonates and adults in Khartoum. *BMC. Res. Notes* 13:312. doi: 10.1186/s13104-020-05157-4
- Ali, A., Ali, Q., Ali, R., and Mohsin, M. (2020). Draft genome sequence of an extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* ST58 isolate

- from cattle in Pakistan. *J. Glob. Antimicrob. Resist.* 21, 303–305. doi: 10.1016/j.jgar.2020.04.020
- Aljorayid, A., Viau, R., Castellino, L., and Jump, R. L. P. (2016). *Serratia fonticola*, pathogen or bystander? A case series and review of the literature. *IDCases* 5, 6–8. doi: 10.1016/j.idcr.2016.05.003
- Al-Kharousi, Z. S., Guizani, N., Al-Sadi, A. M., and Al-Bulushi, I. M. (2019). Antibiotic resistance of Enterobacteriaceae isolated from fresh fruits and vegetables and characterization of their AmpC  $\beta$ -lactamases. *J. Food Prot.* 82, 1857–1863. doi: 10.4315/0362-028X.JFP-19-089
- Ashari, K. S., Roslan, N. S., Omar, A. R., Bejo, M. H., Ideris, A., and Isa, N. M. (2019). Genome sequencing and analysis of *Salmonella enterica* subsp. *enterica* serovar Stanley UPM 517: insights on its virulence-associated elements and their potentials as vaccine candidates. *PeerJ* 7:e6948. doi: 10.7717/peerj.6948
- Blaak, H., van Hoek, A. H. A. M., Veenman, C., Docters van Leeuwen, A. E., Lynch, G., van Overbeek, W. M., et al. (2014). Extended spectrum  $\beta$ -lactamase and constitutively AmpC-producing Enterobacteriaceae on fresh produce and in the agricultural environment. *Int. J. Food Microbiol.* 168–169, 8–16. doi: 10.1016/j.ijfoodmicro.2013.10.006
- Bortolaia, V., Hansen, K. H., Nielsen, C. A., Fritsche, T. R., and Guardabassi, L. (2014). High diversity of plasmids harbouring blaCMY-2 among clinical *Escherichia coli* isolates from humans and companion animals in the upper Midwestern USA. *J. Antimicrob. Chemother.* 69, 1492–1496. doi: 10.1093/jac/dku011
- Bush, K., and Jacoby, G. A. (2010). Updated functional classification of  $\beta$ -lactamases. *Antimicrob. Agents Chemother.* 54, 969–976. doi: 10.1128/AAC.01009-09
- Cantón, R., González-Alba, J. M., and Galán, J. C. (2012). CTX-M enzymes: origin and diffusion. *Front. Microbiol.* 3:110. doi: 10.3389/fmicb.2012.00110
- Carattoli, A., Zankari, E., García-Fernández, A., Larsen, M. V., Lund, O., Villa, L., et al. (2014). In silico detection and typing of plasmids using plasmidfinder and plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* 58, 3895–3903. doi: 10.1128/AAC.02412-14
- CDC (2019). Antibiotic resistance threats in the United States. Atlanta, GA. Available at: [https://www.cdc.gov/drugresistance/biggest\\_threats.html](https://www.cdc.gov/drugresistance/biggest_threats.html) (Accessed June 18, 2021).
- Cergole-Novella, M. C., Guth, B. E. C., Castanheira, M., Carmo, M. S., and Pignatari, A. C. C. (2010). First description of blaCTX-M-14-and blaCTX-M-15-producing *Escherichia coli* isolates in Brazil. *Microb. Drug Resist.* 16, 177–184. doi: 10.1089/mdr.2010.0008
- Chen, L., Zheng, D., Liu, B., Yang, J., and Jin, Q. (2016). VFDB 2016: hierarchical and refined dataset for big data analysis—10 years on. *Nucleic Acids Res.* 44, D694–D697. doi: 10.1093/nar/gkv1239
- Collignon, P. J., and McEwen, S. A. (2019). One health—its importance in helping to better control antimicrobial resistance. *Trop. Med. Infect. Dis.* 4:22. doi: 10.3390/tropicalmed4010022
- Colosi, I. A., Baci, A. M., Oprea, R. V., Peca, L., Gudat, T., Simon, L. M., et al. (2020). Prevalence of ESBL, AmpC and carbapenemase-producing enterobacteriales isolated from raw vegetables retailed in Romania. *Foods* 9:1726. doi: 10.3390/foods9121726
- Cornick, J., Musicha, P., Peno, C., Saeger, E., Iroh Toh, P. Y., Bennett, A., et al. (2020). Genomic investigation of a suspected multi-drug resistant *Klebsiella pneumoniae* outbreak in a neonatal care unit in sub-Saharan Africa. bioRxiv [Preprint]. doi: 10.1101/2020.08.06.236117
- Cosentino, S., Voldby Larsen, M., Møller Aarestrup, F., and Lund, O. (2013). PathogenFinder—distinguishing friend from foe using bacterial whole genome sequence data. *PLoS One* 8:e77302. doi: 10.1371/journal.pone.0077302
- Dandachi, L., Chaddad, A., Hanna, J., Matta, J., and Daoud, Z. (2019). Understanding the epidemiology of multi-drug resistant gram-negative bacilli in the middle east using a one health approach. *Front. Microbiol.* 10:1941. doi: 10.3389/fmicb.2019.01941
- Dolejska, M., Brhelova, E., Dobiasova, H., Krivdova, J., Jurankova, J., Sevcikova, A., et al. (2012). Dissemination of IncFIIK-type plasmids in multiresistant CTX-M-15-producing Enterobacteriaceae isolates from children in hospital paediatric oncology wards. *Int. J. Antimicrob. Agents* 40, 510–515. doi: 10.1016/j.ijantimicag.2012.07.016
- Dolejska, M., Vill, L., Dobiasova, H., Fortini, D., Feudi, C., and Carattoli, A. (2013). Plasmid content of a clinically relevant *Klebsiella pneumoniae* clone from the Czech Republic producing CTX-M-15 and QnrB1. *Antimicrob. Agents Chemother.* 57, 1073–1076. doi: 10.1128/AAC.01886-12
- Doster, E., Lakin, S. M., Dean, C. J., Wolfe, C., Young, J. G., Boucher, C., et al. (2020). MEGARes 2.0: A database for classification of antimicrobial drug, biocide and metal resistance determinants in metagenomic sequence data. *Nucleic Acids Res.* 48, D561–D569. doi: 10.1093/nar/gkz1010
- Falgenhauer, L., Schwengers, O., Schmiedel, J., Baars, C., Lambrecht, O., Heß, S., et al. (2019). Multidrug-resistant and clinically relevant gram-negative bacteria are present in german surface waters. *Front. Microbiol.* 10:2779. doi: 10.3389/fmicb.2019.02779
- Feldgarden, M., Brover, V., Haft, D. H., Prasad, A. B., Slotta, D. J., Tolstoy, I., et al. (2019). Validating the AMRFinder tool and resistance gene database by using antimicrobial resistance genotype-phenotype correlations in a collection of isolates. *Antimicrob. Agents Chemother.* 63, e00483–e00419. doi: 10.1128/AAC.00483-19
- Finton, M. D., Meisal, R., Porcellato, D., Brandal, L. T., and Lindstedt, B. A. (2020). Whole genome sequencing and characterization of multidrug-resistant (MDR) bacterial strains isolated from a Norwegian university campus pond. *Front. Microbiol.* 11:1273. doi: 10.3389/fmicb.2020.01273
- Follador, R., Heinz, E., Wyres, K. L., Ellington, M. J., Kowarik, M., Holt, K. E., et al. (2016). The diversity of *Klebsiella pneumoniae* surface polysaccharides. *Microb. Genomics* 2:e000073. doi: 10.1099/mgen.0.000073
- Founou, R. C., Founou, L. L., Allam, M., Ismail, A., and Essack, S. Y. (2019). Whole genome sequencing of extended spectrum  $\beta$ -lactamase (ESBL)-producing *Klebsiella pneumoniae* isolated from hospitalized patients in KwaZulu-Natal, South Africa. *Sci. Rep.* 9:6266. doi: 10.1038/s41598-019-42672-2
- Furlan, J. P. R., and Stehling, E. G. (2021). Multiple sequence types, virulence determinants and antimicrobial resistance genes in multidrug- and colistin-resistant *Escherichia coli* from agricultural and non-agricultural soils. *Environ. Pollut.* 288:117804. doi: 10.1016/j.envpol.2021.117804
- Gekenidis, M. T., Rigotti, S., Hummerjohann, J., Walsh, F., and Drissner, D. (2020). Long-term persistence of blaCTX-m-15 in soil and lettuce after introducing extended-spectrum  $\beta$ -lactamase (EsbI)-producing *Escherichia coli* via manure or water. *Microorganisms* 8:1646. doi: 10.3390/microorganisms8111646
- Gonzalez-Escalona, N., and Kase, J. A. (2018). Virulence gene profiles and phylogeny of Shiga toxin-positive *Escherichia coli* strains isolated from FDA regulated foods during 2010–2017. bioRxiv [Preprint], 1–26. doi: 10.1101/461327
- Gou, J. J., Liu, N., Guo, L. H., Xu, H., Lv, T., Yu, X., et al. (2020). Carbapenem-resistant Enterobacter hormaechei ST1103 with IMP-26 carbapenemase and ESBL gene blaShv-178. *Infect. Drug Resist.* 13, 597–605. doi: 10.2147/IDR.S232514
- Gupta, S. K., Padmanabhan, B. R., Diene, S. M., Lopez-Rojas, R., Kempf, M., Landraud, L., et al. (2014). ARG-annot, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrob. Agents Chemother.* 58, 212–220. doi: 10.1128/AAC.01310-13
- Hai, P. D., Hoa, L. T. V., Tot, N. H., Phuong, L. L., Quang, V. V., Thuyet, B. T., et al. (2020). First report of biliary tract infection caused by multidrug-resistant *Serratia fonticola*. *New Microb. New Infect.* 36:100692. doi: 10.1016/j.nmni.2020.100692
- Han, Y., Huang, L., Liu, C., Huang, X., Zheng, R., Lu, Y., et al. (2021). Characterization of carbapenem-resistant *Klebsiella pneumoniae* st15 clone coproducing kpc-2, ctx-m-15 and shv-28 spread in an intensive care unit of a tertiary hospital. *Infect. Drug Resist.* 14, 767–773. doi: 10.2147/IDR.S298515
- Hassen, B., Abbassi, M. S., Benlabidi, S., Ruiz-Ripa, L., Mama, O. M., Ibrahim, C., et al. (2020). Genetic characterization of ESBL-producing *Escherichia coli* and *Klebsiella pneumoniae* isolated from wastewater and river water in Tunisia: predominance of CTX-M-15 and high genetic diversity. *Environ. Sci. Pollut. Res.* 27, 44368–44377. doi: 10.1007/s11356-020-10326-w
- Hauser, E., Mellmann, A., Semmler, T., Stoeber, H., Wieler, L. H., Karch, H., et al. (2013). Phylogenetic and molecular analysis of food-borne Shiga toxin-producing *Escherichia coli*. *Appl. Environ. Microbiol.* 79, 2731–2740. doi: 10.1128/AEM.03552-12
- Iguchi, A., Nagaya, Y., Pradel, E., Ooka, T., Ogura, Y., Katsura, K., et al. (2014). Genome evolution and plasticity of *Serratia marcescens*, an important multidrug-resistant nosocomial pathogen. *Genome Biol. Evol.* 6, 2096–2110. doi: 10.1093/gbe/evu160
- Iwu, C. D., de Plessis, E. M., Korsten, L., Nontongana, N., and Okoh, A. I. (2020). Antibigram signatures of some enterobacteria recovered from irrigation water and agricultural soil in two district municipalities of South Africa. *Microorganisms* 8:1206. doi: 10.3390/microorganisms8081206
- Jacoby, G. A. (2009). AmpC  $\beta$ -Lactamases. *Clin. Microbiol. Rev.* 22, 161–182. doi: 10.1128/CMR.00036-08

- Jia, B., Raphenya, A. R., Alcock, B., Wagglechner, N., Guo, P., Tsang, K. K., et al. (2017). CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res.* 45, 566–573. doi: 10.1093/nar/gkw1004
- Jibril, A. H., Okeke, I. N., Dalsgaard, A., Menéndez, V. G., and Olsen, J. E. (2021). Genomic analysis of antimicrobial resistance and resistance plasmids in salmonella serovars from poultry in Nigeria. *Antibiotics* 10:99. doi: 10.3390/antibiotics10020099
- Joensen, K. G., Scheutz, F., Lund, O., Hasman, H., Kaas, R. S., Nielsen, E. M., et al. (2014). Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J. Clin. Microbiol.* 52, 1501–1510. doi: 10.1128/JCM.03617-13
- Joensen, K. G., Tetzschner, A. M. M., Iguchi, A., Aarestrup, F. M., and Scheutz, F. (2015). Rapid and easy in silico serotyping of *Escherichia coli* isolates by use of whole-genome sequencing data. *J. Clin. Microbiol.* 53, 2410–2426. doi: 10.1128/JCM.00008-15
- Johansson, M. H. K., Bortolaia, V., Tansirichaiya, S., Aarestrup, F. M., Roberts, A. P., and Petersen, T. N. (2021). Detection of mobile genetic elements associated with antibiotic resistance in *Salmonella enterica* using a newly developed web tool: MobileElementFinder. *J. Antimicrob. Chemother.* 76, 101–109. doi: 10.1093/jac/dkaa390
- Jones-Dias, D., Manageiro, V., Ferreira, E., Barreiro, P., Vieira, L., Moura, I. B., et al. (2016). Architecture of class 1, 2, and 3 integrons from gram negative bacteria recovered among fruits and vegetables. *Front. Microbiol.* 7:1400. doi: 10.3389/fmicb.2016.01400
- Kaushik, M., Kumar, S., Kapoor, R. K., Virdi, J. S., and Gulati, P. (2018). Integrons in Enterobacteriaceae: diversity, distribution and epidemiology. *Int. J. Antimicrob. Agents* 51, 167–176. doi: 10.1016/j.ijantimicag.2017.10.004
- Khari, F. I. M., Karunakaran, R., Rosli, R., and Tay, S. T. (2016). Genotypic and phenotypic detection of AmpC  $\beta$ -lactamases in Enterobacter spp. isolated from a teaching hospital in Malaysia. *PLoS One* 11:e0150643. doi: 10.1371/journal.pone.0150643
- Kim, S., Karns, J. S., Van Kessel, J. A. S., and Haley, B. J. (2017). Genome sequences of five multidrug-resistant *Escherichia coli* sequence type 117 isolates recovered from dairy calves. *Genome Announc.* 5, e00732–e00717. doi: 10.1128/genomeA.00732-17
- Kim, S., Kim, H., Kim, Y., Kim, M., Kwak, H., and Ryu, S. (2020). Whole-genome sequencing-based characteristics in extended-spectrum beta-lactamase-producing *Escherichia coli* isolated from retail meats in Korea. *Microorganisms* 8:508. doi: 10.3390/microorganisms8040508
- Larsen, M. V., Cosentino, S., Rasmussen, S., Friis, C., Hasman, H., Marvig, R. L., et al. (2012). Multilocus sequence typing of total-genome-sequenced bacteria. *J. Clin. Microbiol.* 50, 1355–1361. doi: 10.1128/JCM.06094-11
- Li, B., Vellidis, G., Liu, H., Jay-Russell, M., Zhao, S., Hu, Z., et al. (2014). Diversity and antimicrobial resistance of *salmonella enterica* isolates from surface water in southeastern United States. *Appl. Environ. Microbiol.* 80, 6355–6365. doi: 10.1128/AEM.02063-14
- Liakopoulos, A., Mevius, D., and Ceccarelli, D. (2016). A review of SHV extended-spectrum  $\beta$ -lactamases: neglected yet ubiquitous. *Front. Microbiol.* 7:1374. doi: 10.3389/fmicb.2016.01374
- Liao, X. P., Xia, J., Yang, L., Li, L., Sun, J., Liu, Y. H., et al. (2015). Characterization of CTX-M-14-producing *Escherichia coli* from food-producing animals. *Front. Microbiol.* 6:1136. doi: 10.3389/fmicb.2015.01136
- Liu, H., Whitehouse, C. A., and Li, B. (2018). Presence and persistence of salmonella in water: the impact on microbial quality of water and food safety. *Front. Public Health* 6:159. doi: 10.3389/fpubh.2018.00159
- Lobanovska, M., and Pilla, G. (2017). Penicillin's discovery and antibiotic resistance: lessons for the future? *Yale J. Biol. Med.* 90, 135–145. doi: 10.1103/PhysRevA.32.435
- Löhr, I. H., Hülter, N., Bernhoff, E., Johnsen, P. J., Sundsfjord, A., and Naseer, U. (2015). Persistence of a pKPN3-like CTX-M-15-encoding IncFIIK plasmid in a *Klebsiella pneumoniae* ST17 host during two years of intestinal colonization. *PLoS One* 10:e0116516. doi: 10.1371/journal.pone.0116516
- Lopes, R., Fuentes-Castillo, D., Fontana, H., Rodrigues, L., Dantas, K., Cerdeira, L., et al. (2021a). Endophytic lifestyle of global clones of extended-spectrum beta-lactamase-producing priority pathogens in fresh vegetables: a trojan horse strategy favoring human colonization? *mSystems* 6, e01125–e01120. doi: 10.1128/mSystems.01125-20
- Lopes, R., Furlan, J. P. R., Dos Santos, L. D. R., Gallo, I. F. L., and Stehling, E. G. (2021b). Colistin-resistant mcr-1-positive *Escherichia coli* ST131-H22 carrying Bla CTX-M-15 and qnrB19 in agricultural soil. *Front. Microbiol.* 12:659900. doi: 10.3389/fmicb.2021.659900
- Maluta, R. P., Leite, J. L., Rojas, T. C. G., Scaletsky, I. C. A., Guastalli, E. A. L., de Carvalho Ramos, M., et al. (2017). Variants of astA gene among extra-intestinal *Escherichia coli* of human and avian origin. *FEMS Microbiol. Lett.* 364, 1–5. doi: 10.1093/femsle/fnw285
- Mansour, M. N., Yaghi, J., El Khoury, A., Felten, A., Mistou, M. Y., Atoui, A., et al. (2020). Prediction of salmonella serovars isolated from clinical and food matrices in Lebanon and genomic-based investigation focusing on Enteritidis serovar. *Int. J. Food Microbiol.* 333:108831. doi: 10.1016/j.ijfoodmicro.2020.108831
- Mbelle, N. M., Feldman, C., Sekyere, J. O., Maningi, N. E., Modipane, L., and Essack, S. Y. (2020). Pathogenomics and evolutionary epidemiology of multidrug resistant clinical *Klebsiella pneumoniae* isolated from Pretoria, South Africa. *Sci. Rep.* 10:1232. doi: 10.1038/s41598-020-58012-8
- McKinnon, J., Roy Chowdhury, P., and Djordjevic, S. P. (2018). Genomic analysis of multidrug-resistant *Escherichia coli* ST58 causing urosepsis. *Int. J. Antimicrob. Agents* 52, 430–435. doi: 10.1016/j.ijantimicag.2018.06.017
- Naas, T., Poirel, L., and Nordmann, P. (2008). Minor extended-spectrum  $\beta$ -lactamases. *Clin. Microbiol. Infect.* 14, 42–52. doi: 10.1111/j.1469-0691.2007.01861.x
- Nair, S., Ashton, P., Doumith, M., Connell, S., Painsent, A., Mwaigwisya, S., et al. (2016). WGS for surveillance of antimicrobial resistance: A pilot study to detect the prevalence and mechanism of resistance to azithromycin in a UK population of non-typhoidal salmonella. *J. Antimicrob. Chemother.* 71, 3400–3408. doi: 10.1093/jac/dkw318
- Njage, P. M. K., and Buys, E. M. (2014). Pathogenic and commensal *Escherichia coli* from irrigation water show potential in transmission of extended spectrum and AmpC  $\beta$ -lactamases determinants to isolates from lettuce. *Microb. Biotechnol.* 8, 462–473. doi: 10.1111/1751-7915.12234
- O'Neill, J. (2016). Tackling Drug-Resistant Infections Globally: Final Report and Recommendations the Review on Antimicrobial Resistance.
- Oniciuc, E. A., Likotrafti, E., Alvarez-Molina, A., Prieto, M., Santos, J. A., and Alvarez-Ordóñez, A. (2018). The present and future of whole genome sequencing (WGS) and whole metagenome sequencing (WMS) for surveillance of antimicrobial resistant microorganisms and antimicrobial resistance genes across the food chain. *Gene* 9:268. doi: 10.3390/genes9050268
- Palzkill, T. (2018). Structural and mechanistic basis for extended-spectrum drug-resistance mutations in altering the specificity of TEM, CTX-M, and KPC  $\beta$ -lactamases. *Front. Mol. Biosci.* 5:16. doi: 10.3389/fmolb.2018.00016
- Partridge, S. R. (2011). Analysis of antibiotic resistance regions in gram-negative bacteria. *FEMS Microbiol. Rev.* 35, 820–855. doi: 10.1111/j.1574-6976.2011.00277.x
- Partridge, S. R., Kwong, S. M., Firth, N., and Jensen, S. O. (2018). Mobile genetic elements associated with antimicrobial resistance. *Clin. Microbiol. Rev.* 31, e00088–e00017. doi: 10.1128/CMR.00088-17
- Peirano, G., van Greune, C. H. J., and Pitout, J. D. D. (2011). Characteristics of infections caused by extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* from community hospitals in South Africa. *Diagn. Microbiol. Infect. Dis.* 69, 449–453. doi: 10.1016/j.diagmicrobio.2010.11.011
- Petersen, L. M., and Tisa, L. S. (2013). Friend or foe? A review of the mechanisms that drive serratia towards diverse lifestyles. *Can. J. Microbiol.* 59, 627–640. doi: 10.1139/cjm-2013-0343
- Razavi, M., Kristiansson, E., Flach, C.-F., and Larsson, D. G. J. (2020). The association between insertion sequences and antibiotic resistance genes. *mSphere* 5, 418–420. doi: 10.1128/mSphere.00418-20
- Reid, C. J., Blau, K., Jechalke, S., Smalla, K., Djordjevic, S. P., and Campo, R. Del (2020). Whole genome sequencing of *Escherichia coli* from store-bought produce. *Front. Microbiol.* 10:3050. doi:10.3389/fmicb.2019.03050
- Reuland, E. A., Al Naiemi, N., Raadsen, S. A., Savelkoul, P. H. M., Kluytmans, J. A. J. W., and Vandenbroucke-Grauls, C. M. J. E. (2014). Prevalence of ESBL-producing Enterobacteriaceae in raw vegetables. *Eur. J. Clin. Microbiol. Infect. Dis.* 33, 1843–1846. doi: 10.1007/s10096-014-2142-7
- Richards, G. A., Brink, A. J., and Feldman, C. (2019). Rational use of the fluoroquinolones. *South Afr. Med. J.* 109, 378–381. doi: 10.7196/SAMJ.2019.v109i6.14002
- Richter, L., Du Plessis, E. M., Duvenage, S., and Korsten, L. (2019). Occurrence, identification, and antimicrobial resistance profiles of extended-spectrum and AmpC  $\beta$ -lactamase-producing enterobacteriaceae from fresh vegetables retailed in Gauteng province. *South Afr. Foodborne Pathog. Dis.* 16, 421–427. doi: 10.1089/fpd.2018.2558

- Richter, L., du Plessis, E. M., Duvenage, S., and Korsten, L. (2020). Occurrence, phenotypic and molecular characterization of extended-spectrum- and AmpC- $\beta$ -lactamase producing enterobacteriaceae isolated from selected commercial spinach supply chains in South Africa. *Front. Microbiol.* 11:638. doi: 10.3389/fmicb.2020.00638
- Roer, L., Hendriksen, R. S., Leekitcharoenphon, P., Lukjancenko, O., Kaas, R. S., Hasman, H., et al. (2016). Is the evolution of *Salmonella enterica* subsp. *enterica* linked to restriction-modification systems? *mSystems* 1, e00009–e00016. doi: 10.1128/mSystems.00009-16
- Sabbagh, S. C., Forest, C. G., Lepage, C., Leclerc, J. M., and Daigle, F. (2010). So similar, yet so different: uncovering distinctive features in the genomes of *Salmonella enterica* serovars typhimurium and Typhi. *FEMS Microbiol. Lett.* 305, 1–13. doi: 10.1111/j.1574-6968.2010.01904.x
- Sandner-Miranda, L., Vinuesa, P., Cravioto, A., and Morales-Espinosa, R. (2018). The genomic basis of intrinsic and acquired antibiotic resistance in the genus *Serratia*. *Front. Microbiol.* 9:828. doi: 10.3389/fmicb.2018.00828
- Schwanbeck, J., Bohne, W., Hasdemir, U., Groß, U., Pfeifer, Y., Bunk, B., et al. (2021). Detection of a new resistance-mediating plasmid chimera in a blaOXA-48-positive *Klebsiella pneumoniae* strain at a German university hospital. *Microorganisms* 9:720. doi: 10.3390/microorganisms9040720
- Sekyere, J. O., Maningi, N. E., Modipane, L., and Mbelle, N. M. (2020). Emergence of mcr-9.1 in extended-spectrum-beta-lactamase-producing clinical Enterobacteriaceae in Pretoria, South Africa: global evolutionary Phylogenomics, Resistome, and Mobilome. *mSystems* 5, e00148–e00120. doi: 10.1128/mSystems.00148-20
- Suez, J., Porwollik, S., Dagan, A., Marzel, A., Schorr, Y. I., Desai, P. T., et al. (2013). Virulence gene profiling and pathogenicity characterization of non-typhoidal salmonella accounted for invasive disease in humans. *PLoS One* 8:e58449. doi: 10.1371/journal.pone.0058449
- Surleac, M., Barbu, I. C., Paraschiv, S., Popa, L. I., Gheorghe, I., Marutescu, L., et al. (2020). Whole genome sequencing snapshot of multidrug resistant *Klebsiella pneumoniae* strains from hospitals and receiving wastewater treatment plants in southern Romania. *PLoS One* 15:e0228079. doi: 10.1371/journal.pone.0228079
- Tanaka, H., Hayashi, W., Iimura, M., Taniguchi, Y., Soga, E., Matsuo, N., et al. (2019). Wastewater as a probable environmental reservoir of extended-spectrum-beta-lactamase genes: detection of chimeric beta-lactamases CTX-M-64 and CTX-M-123. *Appl. Environ. Microbiol.* 85, e01740–e01719. doi: 10.1128/AEM.01740-19
- Tasić, S., Obradović, D., and Tasić, I. (2013). Characterization of *Serratia fonticola*, an opportunistic pathogen isolated from drinking water. *Arch. Biol. Sci.* 65, 899–904. doi: 10.2298/ABS1303899T
- Tekele, S. G., Teklu, D. S., Tullu, K. D., Birru, S. K., and Legese, M. H. (2020). Extended-spectrum Beta-lactamase and AmpC beta-lactamases producing gram negative bacilli isolated from clinical specimens at international clinical laboratories, Addis Ababa, Ethiopia. *PLoS One* 15:e0241984. doi: 10.1371/journal.pone.0241984
- Thomas, J., Govender, N., McCarthy, K. M., Erasmus, L. K., Doyle, T. J., Allam, M., et al. (2020). Outbreak of listeriosis in South Africa associated with processed meat. *N. Engl. J. Med.* 382, 632–643. doi: 10.1056/NEJMoa1907462
- Thompson, C. P., Doak, A. N., Amirani, N., Schroeder, E. A., Wright, J., Kariyawasam, S., et al. (2018). High-resolution identification of multiple salmonella serovars in a single sample by using CRISPR-SeroSeq. *Appl. Environ. Microbiol.* 84, e01859–e01818. doi: 10.1128/AEM.01859-18
- van Duin, D., and Doi, Y. (2017). The global epidemiology of carbapenemase-producing Enterobacteriaceae. *Virulence* 8, 460–469. doi: 10.1080/21505594.2016.1222343
- Varani, A., He, S., Siguier, P., Ross, K., and Chandler, M. (2021). The IS6 family, a clinically important group of insertion sequences including IS26. *Mob. DNA* 12:11. doi: 10.1186/s13100-021-00239-x
- Velásquez, J. C., Hidalgo, A. A., Villagra, N., Santiviago, C. A., Mora, G. C., and Fuentes, J. A. (2016). SPI-9 of *Salmonella enterica* serovar typhi is constituted by an operon positively regulated by RpoS and contributes to adherence to epithelial cells in culture. *Microbiology* 162, 1367–1378. doi: 10.1099/mic.0.000319
- Verlicchi, P., and Grillini, V. (2020). Surface water and ground water quality in South Africa and Mozambique-analysis of the most critical pollutants for drinking purposes and challenges in water treatment selection. *WaterSA* 12:305. doi: 10.3390/w12010305
- Waterman, S. R., and Holden, D. W. (2003). Functions of the salmonella pathogenicity island 2 (SPI-2) type III secretion system. *Cell. Microbiol.* 5, 501–511. doi: 10.1046/j.1462-5822.2003.00294.x
- WHO (2017). Global Priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics.
- WHO (2020). Global Antimicrobial Resistance and Use Surveillance System (GLASS) Report. Available at: <https://www.who.int/glass/resources/publications/early-implementation-report-2020/en/> (Accessed June 15, 2021).
- Wick, R. R., Heinz, E., Holt, K. E., and Wyres, K. L. (2018). Kaptive web: user-friendly capsule and lipopolysaccharide serotype prediction for *Klebsiella* genomes. *bioRxiv* [Preprint]. doi: 10.1101/260125
- Ye, Q., Wu, Q., Zhang, S., Zhang, J., Yang, G., Wang, J., et al. (2017). Characterization of extended-spectrum  $\beta$ -lactamase-producing enterobacteriaceae from retail food in China. *Front. Microbiol.* 9:1709. doi: 10.3389/fmicb.2018.01709
- Zankari, E., Hasman, H., Cosentino, S., Vestergaard, M., Rasmussen, S., Lund, O., et al. (2012). Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.* 67, 2640–2644. doi: 10.1093/jac/dks261
- Zhang, S., Yin, Y., Jones, M. B., Zhang, Z., Kaiser, B. L. D., Dinsmore, B. A., et al. (2015). Salmonella serotype determination utilizing high-throughput genome sequencing data. *J. Clin. Microbiol.* 53, 1685–1692. doi: 10.1128/JCM.00323-15
- Zhou, K., Zhou, Y., Zhang, C., Song, J., Cao, X., Yu, X., et al. (2020). Dissemination of a 'rare' extended-spectrum  $\beta$ -lactamase gene blaSFO-1 mediated by epidemic clones of carbapenemase-producing Enterobacter hormaechei in China. *Int. J. Antimicrob. Agents* 56:106079. doi: 10.1016/j.ijantimicag.2020.106079
- Zurfluh, K., Nuesch-Inderbinen, M., Morach, M., Berner, A. Z., Hachler, H., and Stephan, R. (2015). Extended-spectrum-beta-lactamase-producing Enterobacteriaceae isolated from vegetables imported from the Dominican Republic, India, Thailand, and Vietnam. *Appl. Environ. Microbiol.* 81, 3115–3120. doi: 10.1128/AEM.00258-15

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Richter, du Plessis, Duvenage, Allam, Ismail and Korsten. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.