

Wild Plants, Mushrooms and Nuts

Wild Plants, Mushrooms and Nuts: Functional Food Properties and Applications

Edited by Isabel C. F. R. Ferreira, Patricia Morales, and Lillian Barros

Mountain Research Centre (CIMO), School of Agriculture, Polytechnic Institute of Bragança, Portugal

Department of Nutrition and Bromatology II, Faculty of Pharmacy, Complutense University of Madrid, Spain

Mountain Research Centre (CIMO), School of Agriculture, Polytechnic Institute of Bragança, Portugal

WILEY Blackwell

This edition first published 2017

© 2017 John Wiley & Sons, Ltd

Registered Office

John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, United Kingdom

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com.

The right of the author to be identified as the author of this work has been asserted in accordance with the Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. It is sold on the understanding that the publisher is not engaged in rendering professional services and neither the publisher nor the author shall be liable for damages arising herefrom. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data

Names: Ferreira, Isabel C. F. R., 1979– editor. | Barros, Lillian, editor. | Patricia Morales, editor.

Title: Wild plants, mushrooms and nuts : functional food properties and applications /

[edited by] Isabel Ferreira, Patricia Morales, Lillian Barros

Description: Chichester, UK ; Hoboken, NJ : John Wiley & Sons, 2017. |

Includes bibliographical references and index.

Identifiers: LCCN 2016036173 (print) | LCCN 2016045177 (ebook) | ISBN 9781118944622 (cloth) |

ISBN 9781118944639 (pdf) | ISBN 9781118944646 (epub)

Subjects: LCSH: Functional foods. | Mushrooms. | Nuts. | Wild plants, Edible.

Classification: LCC QP144.F85 W54 2016 (print) | LCC QP144.F85 (ebook) | DDC 581.6/32–dc23

LC record available at <https://lcn.loc.gov/2016036173>

A catalogue record for this book is available from the British Library.

Set in 10/12pt Warnock by SPi Global, Pondicherry, India

Contents

List of Contributors *xi*

Preface *xv*

| | | |
|----------|--|-----------|
| 1 | Introduction: The Increasing Demand for Functional Foods | 1 |
| | <i>Natália Martins, Patricia Morales, Lillian Barros, and Isabel C. F. R. Ferreira</i> | |
| 1.1 | Food Patterns: A Cross-sectional Approach and Brief Overview | 1 |
| 1.2 | Nutrition and Health: Facts and Tendencies | 2 |
| 1.2.1 | Evidence-based Medicine: Past to Present | 2 |
| 1.2.2 | Modern Food Patterns: An (Un)Healthy Yield | 3 |
| 1.3 | Functional Foods Diversity and Related Applications: A World of (Un)Explored Biofunctionalities | 4 |
| 1.3.1 | Food and Dietary Supplements, Botanicals, and Nutraceuticals: Clarifying Misinterpreted Concepts | 5 |
| 1.4 | Functional Foods Versus Bioactive Molecules: Hierarchies and Regulatory Practices | 6 |
| 1.5 | Challenges and Opportunities: A Multidimensional Perspective | 8 |
| 1.6 | Conclusion | 9 |
| | References | 10 |
| | | |
| 2 | The Numbers Behind Mushroom Biodiversity | 15 |
| | <i>Anabela Martins</i> | |
| 2.1 | Origin and Diversity of Fungi | 15 |
| 2.2 | Ecological Diversity | 18 |
| 2.2.1 | Freshwater Fungi | 21 |
| 2.2.2 | Marine Fungi | 21 |
| 2.2.3 | Endophytes of Plant Leaves and Stems | 22 |
| 2.2.4 | Fungi from Arthropod and Invertebrate Animals | 22 |
| 2.3 | Global Diversity of Soil Fungi | 22 |
| 2.4 | Wild Edible Fungi | 24 |
| 2.4.1 | Diversity of Wild Edible Mushrooms | 30 |
| 2.4.2 | Medicinal Mushrooms | 32 |
| 2.5 | Cultivation of Edible Fungi | 38 |
| 2.6 | Social and Economic Interest in Edible Mushrooms | 41 |
| 2.7 | Edible Mushroom World Production and Commercialization | 42 |
| 2.8 | Conclusion | 49 |
| | References | 50 |

3 The Nutritional Benefits of Mushrooms 65*Carolina Barroetaveña and Carolina V. Toledo*

- 3.1 Introduction 65
- 3.2 Nutritional Properties of Mushrooms 66
 - 3.2.1 Proteins and Amino Acids 66
 - 3.2.2 Carbohydrates: Available Carbohydrates and Dietary Fiber 69
 - 3.2.3 Lipids 71
 - 3.2.4 Energetic Value/Caloric Content 72
 - 3.2.5 Ash and Mineral Elements 73
- 3.3 Vitamins 73
- 3.4 Conclusion 75
- References 76

4 The Bioactive Properties of Mushrooms 83*Marina Soković, Ana Ćirić, Jasmina Glamočlija, and Dejan Stojković*

- 4.1 Introduction 83
- 4.2 Antimicrobial Activity of Edible and Medicinal Fungi 84
 - 4.2.1 Antibacterial Activity of Mushroom Extracts 84
 - 4.2.2 Compounds Isolated from Mushrooms as Bacterial Growth Inhibitors 84
 - 4.2.3 Antifungal Activity of Crude Mushroom Extracts 92
 - 4.2.4 Isolated Compounds from Mushrooms Express Antifungal Potency 93
- 4.3 Mushrooms as a Reliable Source of Antioxidants for Disease Prevention 95
- 4.4 Could Mushrooms Be Used as Cytotoxic and Antitumor Agents? 100
 - 4.4.1 Cytotoxic Features of Wild Mushroom Extracts 100
 - 4.4.2 Mushroom Polysaccharides, β -, and α -Glucans as Antitumor Agents 100
 - 4.4.3 Cytotoxic Potency of Terpenoids and Related Compounds from Mushrooms 104
 - 4.4.4 Mushroom Sterols Inhibit the Growth of Carcinoma Cell Lines 104
- 4.5 Controlling Obesity, Metabolic Syndrome, and Diabetes Mellitus with Mushrooms 108
- 4.6 Conclusion 111
- References 111

5 The Use of Mushrooms in the Development of Functional Foods, Drugs, and Nutraceuticals 123*Humberto J. Morris, Gabriel Llauradó, Yaixa Beltrán, Yamila Lebeque, Rosa C. Bermúdez, Nora García, Isabelle Gaime-Perraud, and Serge Moukha*

- 5.1 Introduction 123
- 5.2 A Window into the “Garden” of a Novel Class of Products 125
- 5.3 Main Uses of Edible Medicinal Mushrooms in the Age of Human Health Crises 127
 - 5.3.1 Mushrooms as Functional Foods: A Paradigm of Integrating Tradition and Novelty 128
 - 5.3.1.1 Proven Functional Properties 129

| | | |
|-------|---|-----|
| 5.3.2 | Mushroom Nutraceuticals | 135 |
| 5.3.3 | Mushrooms as a Significant Source of Drugs: Lessons from Wasser's Discovery Pathway | 139 |
| 5.4 | Conclusion | 146 |
| | References | 149 |

| | | |
|----------|--|------------|
| 6 | The Consumption of Wild Edible Plants | 159 |
| | <i>Ana Maria Carvalho and Ana Maria Barata</i> | |
| 6.1 | Wild Edible Plants | 159 |
| 6.1.1 | Contribution of Wild Edible Plants to People's Diets and Daily Lives | 160 |
| 6.1.1.1 | Famine Foods | 161 |
| 6.1.1.2 | Weeds | 162 |
| 6.1.2 | New Trends in Edible Wild Plant Consumption | 163 |
| 6.1.3 | Wild Edible Plants, Food Security, and Research Approaches | 164 |
| 6.2 | Foraging and Wild Edible Plant Resources | 165 |
| 6.2.1 | Wild Plant Resources Worldwide | 165 |
| 6.2.1.1 | Africa | 166 |
| 6.2.1.2 | Americas | 171 |
| 6.2.1.3 | Asia | 173 |
| 6.2.1.4 | Europe | 174 |
| 6.2.1.5 | Oceania | 175 |
| 6.3 | Wild Relatives of Crop Plants | 177 |
| 6.3.1 | CWR Inventories and Checklists | 179 |
| 6.4 | Enhancing Biodiversity and Plant Genetic Resources Conservation | 181 |
| 6.4.1 | Conservation Strategies | 181 |
| 6.4.2 | Promoting and Strengthening Biocultural Heritage | 183 |
| 6.5 | Culturally Significant Wild Edible Plants | 185 |
| 6.6 | Conclusion | 187 |
| | References | 188 |

| | | |
|----------|---|------------|
| 7 | Wild Greens as Source of Nutritive and Bioactive Compounds Over the World | 199 |
| | <i>Patricia Morales, Patricia García Herrera, Maria Cruz Matallana González, Montaña Cámara Hurtado, and Maria de Cortes Sánchez Mata</i> | |
| 7.1 | Introduction | 199 |
| 7.2 | Wild Greens as a Source of Nutritive and Bioactive Compounds in Different Geographical Areas | 200 |
| 7.2.1 | Traditional Wild Greens from Africa | 200 |
| 7.2.2 | Wild Vegetables Consumed in the Americas | 213 |
| 7.2.3 | Asian Wild Edible Greens | 222 |
| 7.2.4 | Vegetables Traditionally Consumed in Europe | 226 |
| 7.3 | Implications of Wild Greens Consumption for Human Health: Safely Gathering Wild Edible Plants | 243 |
| 7.4 | Conclusion | 248 |
| | References | 249 |

| | | |
|-----------|--|------------|
| 8 | Nutrients and Bioactive Compounds in Wild Fruits Through Different Continents | 263 |
| | <i>Virginia Fernández-Ruiz, Patricia Morales, Brígida María Ruiz-Rodríguez, and Esperanza Torija Isasa</i> | |
| 8.1 | Introduction | 263 |
| 8.2 | African Wild Fruits as a Source of Nutrients and Bioactive Compounds | 264 |
| 8.3 | American Wild Fruits as a Source of Nutrients and Bioactive Compounds | 273 |
| 8.4 | Asian Wild Fruits as a Source of Nutrients and Bioactive Compounds | 287 |
| 8.5 | European Wild Fruits as a Source of Nutrients and Bioactive Compounds | 291 |
| 8.6 | Conclusion | 306 |
| | References | 306 |
| 9 | Wild Plant-Based Functional Foods, Drugs, and Nutraceuticals | 315 |
| | <i>José Pinela, Márcio Carochó, Maria Inês Dias, Cristina Caleja, Lillian Barros, and Isabel C. F. R. Ferreira</i> | |
| 9.1 | Introduction | 315 |
| 9.2 | Wild Plants and Functional Foods | 316 |
| 9.2.1 | The Concept and Recent Trends in Functional Foods | 316 |
| 9.2.2 | Classification and Development of Functional Foods | 318 |
| 9.2.3 | Wild Plants Used as Functional Foods | 319 |
| 9.3 | Wild Plant-Based Nutraceuticals | 326 |
| 9.3.1 | The Emerging Concept and Applications of Nutraceuticals | 326 |
| 9.3.2 | Recent Advances in Formulations for Nutraceuticals | 328 |
| 9.3.3 | Examples of Nutraceuticals Based on Wild Plants | 329 |
| 9.4 | Wild Plant-Based Drugs | 335 |
| 9.4.1 | From the Bioactive Phytochemical to the Active Principle | 335 |
| 9.4.2 | Common Formulations in Drugs from Plant Origin | 337 |
| 9.4.3 | Wild Plant-Based Drugs for Different Therapeutic Targets | 340 |
| 9.5 | Conclusion | 341 |
| | References | 342 |
| 10 | Nuts: Agricultural and Economic Importance Worldwide | 353 |
| | <i>Albino Bento, Paula Cabo, and Ricardo Malheiro</i> | |
| 10.1 | Introduction | 353 |
| 10.2 | Almond | 354 |
| 10.2.1 | Evolution of Almond Production and Trade Facts | 356 |
| 10.2.2 | Consumption of Almonds Worldwide | 357 |
| 10.3 | Chestnut | 359 |
| 10.3.1 | Evolution of Chestnut Production and Trade Facts | 360 |
| 10.3.2 | Consumption of Chestnuts Worldwide | 362 |
| 10.4 | Hazelnut | 362 |
| 10.4.1 | Evolution of Hazelnut Production and Trade Facts | 365 |
| 10.4.2 | Consumption of Hazelnuts Worldwide | 367 |
| 10.5 | Walnut | 367 |
| 10.5.1 | Evolution of Walnut Production and Trade Facts | 369 |
| 10.5.2 | Consumption of Walnuts Worldwide | 374 |
| 10.6 | Conclusion | 374 |
| | References | 374 |

| | | |
|-----------|---|------------|
| 11 | Recent Advances in Our Knowledge of the Biological Properties of Nuts | 377 |
| | <i>Ryszard Amarowicz, Yi Gong, and Ronald B. Pegg</i> | |
| 11.1 | Introduction | 377 |
| 11.2 | Nuts as a Source of Nutrients, Phytosterols, and Natural Antioxidants | 378 |
| 11.2.1 | Nuts as a Source of Energy and Macronutrients | 378 |
| 11.2.2 | Biological Value of Nut Proteins | 378 |
| 11.2.3 | Nuts as a Source of Vitamins and Minerals | 378 |
| 11.2.4 | Nuts as a Source of Essential Fatty Acids and Phytosterols | 382 |
| 11.2.5 | Phenolic Compounds Originating from Tree Nuts as Natural Antioxidants | 382 |
| 11.3 | Health Benefits of Nuts | 389 |
| 11.3.1 | Health-Promoting Properties of Nuts | 389 |
| 11.3.2 | Nuts and Body Weight Control | 389 |
| 11.3.3 | Nuts and Cardiovascular Disease | 389 |
| 11.3.4 | Nuts and Diabetes | 393 |
| 11.3.5 | Nuts and Cancer | 395 |
| 11.3.6 | Application of Nuts in the Functional Food Industry | 398 |
| 11.4 | Tree Nuts and Allergy | 399 |
| 11.5 | Conclusion | 401 |
| | References | 401 |
| | | |
| 12 | Nuts as Sources of Nutrients | 411 |
| | <i>João C. M. Barreira, M. Beatriz P. P. Oliveira, and Isabel C. F. R. Ferreira</i> | |
| 12.1 | <i>Prunus dulcis</i> (Miller) D. A. Webb (almond) | 411 |
| 12.1.1 | Botanical Aspects and Geographical Distribution | 411 |
| 12.1.2 | Main Applications and Nutritional Overview | 411 |
| 12.1.3 | Major Components | 412 |
| 12.1.4 | Minor Components | 417 |
| 12.2 | <i>Castanea sativa</i> Miller (Chestnut) | 418 |
| 12.2.1 | Botanical Aspects and Geographical Distribution | 418 |
| 12.2.2 | Main Applications and Nutritional Overview | 418 |
| 12.2.3 | Major Components | 418 |
| 12.2.4 | Minor Components | 419 |
| 12.3 | <i>Corylus avellana</i> L. (Hazelnut) | 420 |
| 12.3.1 | Botanical Aspects and Geographical Distribution | 420 |
| 12.3.2 | Main Applications and Nutritional Overview | 420 |
| 12.3.3 | Major Components | 420 |
| 12.3.4 | Minor Components | 421 |
| 12.4 | <i>Juglans regia</i> L. (Walnut) | 422 |
| 12.4.1 | Botanical Aspects and Geographical Distribution | 422 |
| 12.4.2 | Main Applications and Nutritional Overview | 422 |
| 12.4.3 | Major Components | 422 |
| 12.4.4 | Minor Components | 423 |
| 12.5 | Conclusion | 423 |
| | References | 424 |

13 The Contribution of Chestnuts to the Design and Development of Functional Foods 431

Ariane Mendonça Kluczkovski

- 13.1 Introduction 431
- 13.2 Chestnut Composition 431
- 13.3 Biotechnology and Safety 435
 - 13.3.1 Functional Properties and Health Effects 435
 - 13.3.1.1 Antioxidants 436
 - 13.3.1.2 Prebiotics 437
 - 13.3.1.3 Gluten-Free Products 438
 - 13.3.2 Functionality of Chestnut Products 439
- 13.4 Conclusion 440
- References 441

14 Emerging Functional Foods Derived from Almonds 445

Isabela Mateus Martins, Qianru Chen, and C. Y. Oliver Chen

- 14.1 Introduction 445
- 14.2 Overview of Almond Nutrients 446
- 14.3 Health Benefits and Bioactions of Almonds 447
 - 14.3.1 Cholesterol Reduction 447
 - 14.3.2 Glucose Regulation 451
 - 14.3.3 Antiinflammation 451
 - 14.3.4 Antioxidation 453
 - 14.3.5 Body Weight Control 456
 - 14.3.6 Prebiotics 458
- 14.4 Development of Functional Foods with Almonds 459
- 14.5 Conclusion 462
- References 462

Index 471

List of Contributors

Ryszard Amarowicz

Division of Food Science,
Institute of Animal Reproduction and
Food Research of the Polish Academy of
Sciences, ul,
Poland

Ana Maria Barata

Instituto Nacional de Investigação
Agrária (INIAV),
Banco Português de Germoplasma
Vegetal (BPGV),
Portugal

João C. M. Barreira

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Carolina Barroetaveña

Centro de Investigación y Extensión
Forestal Andino Patagónico CIEFAP,
Argentina

Lillian Barros

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Yaixa Beltrán

Center for Studies on Industrial
Biotechnology (CEBI),
University of Oriente,
Cuba

Albino Bento

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Rosa C. Bermúdez

Center for Studies on Industrial
Biotechnology (CEBI),
University of Oriente,
Cuba

Paula Cabo

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Cristina Caleja

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Montaña Cámara Hurtado

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Márcio Carochó

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Ana Maria Carvalho

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

C. Y. Oliver Chen

Antioxidants Research Laboratory,
Jean Mayer USDA Human Nutrition
Research Center on Aging,
Tufts University, USA

Qianru Chen

Antioxidants Research Laboratory,
Jean Mayer USDA Human Nutrition
Research Center on Aging,
Tufts University, USA

Ana Ćirić

University of Belgrade, Institute for
Biological Research “Siniša Stanković”,
Serbia

Maria de Cortes Sánchez Mata

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Maria Inês Dias

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Virginia Fernández-Ruiz

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Isabel C. F. R. Ferreira

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Isabelle Gaime-Perraud

IMBE Biotechnologies et Bioremediation
(IMBE-EBB),
Faculte St Jerome,
France

Nora García

Center for Studies on Industrial
Biotechnology (CEBI),
University of Oriente,
Cuba

Patricia García Herrera

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Jasmina Glamočlija

University of Belgrade, Institute for
Biological Research “Siniša Stanković”,
Serbia

Yi Gong

Department of Food Science and
Technology,
College of Agricultural and
Environmental Sciences,
University of Georgia,
USA

Yamila Lebeque

Center for Studies on Industrial
Biotechnology (CEBI),
University of Oriente,
Cuba

Gabriel Llauradó

Center for Studies on Industrial
Biotechnology (CEBI),
University of Oriente,
Cuba

Ricardo Malheiro

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Anabela Martins

Polytechnic Institute of Bragança,
School of Agriculture (IPB-ESA),
Portugal

Natália Martins

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Maria Cruz Matallana González

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Isabela Mateus Martins

Antioxidants Research Laboratory,
Jean Mayer USDA Human Nutrition
Research Center on Aging,
Tufts University,
USA

Ariane Mendonça Kluczkowski

Faculty of Pharmaceutical Sciences,
Federal University of Amazonas,
Brazil

Patricia Morales

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Humberto J. Morris

Center for Studies on Industrial
Biotechnology (CEBI),
University of Oriente,
Cuba

Serge Moukha

Department of Toxicology, UFR des
Sciences,
Pharmaceutiques-Université Bordeaux
Segalen,
France

M. Beatriz P. P. Oliveira

REQUIMTE/LAQV, Faculty of Pharmacy,
University of Porto,
Portugal

Ronald B. Pegg

Department of Food Science and Technology,
College of Agricultural and
Environmental Sciences,
University of Georgia,
USA

José Pinela

Mountain Research Centre (CIMO),
School of Agriculture,
Polytechnic Institute of Bragança,
Portugal

Brígida María Ruiz-Rodríguez

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Marina Soković

University of Belgrade, Institute for
Biological Research “Siniša Stanković”,
Serbia

Dejan Stojković

University of Belgrade, Institute for
Biological Research “Siniša Stanković”,
Serbia

Carolina V. Toledo

Centro de Investigación y Extensión
Forestal Andino Patagónico CIEFAP,
Argentina

Esperanza Torija Isasa

Department of Nutrition and
Bromatology II,
Faculty of Pharmacy, Complutense
University of Madrid,
Spain

Preface

The use of healthy ingredients is a natural way of preventing diseases and contributes to the increased use of natural matrices. This book focuses on the nutritional, chemical, and biological properties of natural matrices from the Iberian peninsula, mainly food products such as wild plants, mushrooms, chestnuts, and almond.

Society's attitude to food, as a natural and inevitable necessity, has altered in line with changes in social conditions and development of technology. Current consumers are interested in the composition, properties, safety, and health effects of food products. The desire to consume foods with high biological value from natural origins poses a huge challenge for modern food science and industry. In addition, the recent consumer interest in chemopreventive nutrition has increased the choice of food products (functional foods) with specific components (bioactive compounds). The current increase in the adoption of more active and healthy lifestyles needs to be followed by a concomitant response from all players in the food chain. The knowledge contained in this book will allow scientists and, in the longer term, lay members of society to gain a better understanding of the value that these products exhibit, focusing on their nutritional and chemical composition, bioactivity, and potential as functional foods.

Ongoing research on selected products will lead to a new generation of foods, and will promote their nutritional and medicinal use. Public health authorities consider prevention and treatment with nutraceuticals a powerful instrument in maintaining and promoting health, longevity, and life quality. The beneficial effects of nutraceuticals will undoubtedly have an impact on nutritional therapy; they also represent a growing segment of today's food industry. Therefore wild plants, mushrooms, and nuts have become interesting food products due to the increasing interest in the concept of "functional foods" with "health benefits."

Wild Plants, Mushrooms and Nuts: Functional Food Properties and Applications is a compendium of current and novel research on the chemistry, biochemistry, nutritional and pharmaceutical value of traditional food products, which are becoming more relevant in our current diet, for developing novel health foods and in modern natural food therapies. Topics covered range from their nutritional value, chemical and biochemical characterization, to their multifunctional applications as food with beneficial effects on health, through their biological and pharmacological properties (antioxidant, antibacterial, antifungal, and antitumor capacity, among others).

9

Wild Plant-Based Functional Foods, Drugs, and Nutraceuticals

José Pinela^{1,2}, Márcio Carochó^{1,3}, Maria Inês Dias^{1,2}, Cristina Caleja^{1,2}, Lillian Barros¹, and Isabel C. F. R. Ferreira¹

¹ Mountain Research Centre (CIMO), School of Agriculture, Polytechnic Institute of Bragança, Portugal

² REQUIMTE/LAQV, Faculty of Pharmacy, University of Porto, Rua Jorge Viterbo Ferreira, no 228, 4050-313 Porto, Portugal

³ Department of Nutrition and Bromatology II, Faculty of Pharmacy, Complutense University of Madrid, Plaza Ramón y Cajal, s/n., 28040 Madrid, Spain

9.1 Introduction

Wild plants were originally the main element in the human diet, culminating in the different cultures and societies of today. However, the establishment of agriculture led to the decline of consumption of wild plants in comparison to the cultivars that could be grown every year (Grivetti & Ogle 2000). Nevertheless, consumption of wild plants is still a tradition that remains in many cultures, either for their nutritional and health benefits or for sociocultural behaviors that characterize many societies (Groot *et al.* 2002; Pardo de Santayana *et al.* 2007; Schulp *et al.* 2014). As human health and nutrition are two of the pillars that sustain our survival, it is necessary to find new ways to support medical care, which can be found in the vast wild plant ecosystem (Heywood 2011).

Food with additional functional properties could be the future of health supplies for the world population, and thus food and drugs are increasingly seen as one matrix (Bernal *et al.* 2011). Functional foods, nutraceuticals, and drugs based on wild plants that are still unexplored are emerging as a response to the world market, which has been searching for new, better, and safer products.

Functional foods have a similar appearance to their traditional counterparts, but bring potential beneficial effects when consumed on a regular basis in a varied diet. On the other hand, nutraceuticals are substances that have positive physiological effects on the human body, being consumed in unit dose forms such as tablets, capsules or liquids, allowing the delivery of a concentrated bioactive agent and providing a dose that could not be obtained from a normal food intake (Gulati & Ottaway 2006; Hasler 2000). Both the functional food and nutraceutical sectors have been growing significantly in Europe but in the European Union nutraceuticals are not

considered as a specific food category with a series of rules and guidelines to define the product itself, obeying the general regulations on food safety, traceability, recall, and notification (Coppens *et al.* 2006; Gulati & Ottaway 2006). In terms of the health claims associated with functional foods, relating to Regulation (CE) No. 1924/2006 of the European Parliament on nutritional claims and health properties of food, it is possible to classify a functional food under very strict rules and conditions. In addition to the legislation required for all foodstuffs, scientific evidence of the health claims regarding the relevant food will be mandatory for all new products (Bech-Larsen & Scholderer 2007). In the United States, on the other hand, the Food and Drug Administration (FDA) defines the product's category depending on its characteristics, nutraceuticals being regulated as a food and beverage product and dietary supplement, covered by several safety issues, health claims, labeling, and good manufacturing practices (Milner 2000; Wrick 2005).

Concerning drug development, the market also requires safer products due to increasing worldwide concern about synthetic chemical compounds. In that respect, wild plant-based drugs are now in the forefront of the therapeutic agents used for human health, taking into account their high efficiency and low toxicity (Bhardwaj *et al.* 2014; Carocho & Ferreira 2013a).

In this chapter, wild plants commonly used as functional foods will be reviewed. For nutraceuticals, the emerging concept, their applications and novel formulations will be described, and also some products already available on the market. The relationship between the bioactive phytochemical and the active principle will be explained, listing the common formulations in wild plant-based drugs and the different therapeutic targets that can be explored.

9.2 Wild Plants and Functional Foods

9.2.1 The Concept and Recent Trends in Functional Foods

In the first half of the twentieth century, the focus of nutritional science was on establishing the minimum requirements for essential nutrients that ensure the avoidance of deficiency diseases (MMWR 1999). Nowadays, these concepts are changing significantly in the industrialized world. We are progressing from a concept of “adequate nutrition” to one of “optimal nutrition” (Ashwell 2003); from a matter of survival, satisfying hunger, and ensuring food safety to an emphasis on the potential for foods to promote health, in terms of both preventing nutrition-related diseases and improving physical and mental wellbeing (Nöthlings *et al.* 2007; Takachi *et al.* 2008). In addition, consumers are increasingly better informed about the subject than they were in the past. As a result, their expectations of obtaining health benefits from the food they eat are also increasing (Diplock *et al.* 1999). These changes can be explained by some significant trends in our present society, namely rapid advances in science and technology, the rising costs of healthcare, the increase in the numbers of elderly people and in average life expectancy, changes in food laws affecting label and product claims, and people's desire for a better quality of life (Roberfroid 2007).

The primary role of food is to provide nutrients to meet human metabolic requirements and to give the consumer a feeling of satisfaction and wellbeing through

hedonistic attributes such as taste. In addition to this, food can fulfill specific physiological functions in the human body (Li *et al.* 2014a; Zhang *et al.* 2015). In fact, food can not only help to achieve optimal health and development, but it might also play an important role in reducing or preventing the risk of disease. According to the World Health Organization (WHO) and Food and Agriculture Organization (FAO), several dietary patterns along with lifestyle habits constitute major modifiable risk factors in relation to the development of coronary heart disease, different types of cancer, diabetes, obesity, osteoporosis, and periodontal disease (WHO 2003). Foods with these properties were first regulated in Japan in 1981 as Foods for Specified Health Use (FOSHU) (Hasler 2002; Ohama *et al.* 2006). Later, in Europe, the project Functional Food Science in Europe (FUFOSE) was created to assess critically the science base required to provide evidence that specific nutrients and food components beneficially affect target functions in the human body (Tijhuis *et al.* 2012). Currently, this kind of food is generally referred to as “functional food,” if in accordance with the definition given below.

Although there is no universally accepted definition for functional foods (Hasler 2002), and because functional foods are more of a concept than a well-defined group of food products, here we present the definition described previously by Diplock *et al.* (1999). According to these authors, a food can be regarded as “functional” if it is satisfactorily demonstrated to affect beneficially one or more target functions in the human body, beyond adequate nutritional effects, in a way relevant to either an improved state of health and wellbeing and/or disease risk reduction. These foods must remain foods in appearance and they must demonstrate their effects in amounts that can normally be expected to be consumed in the usual diet, i.e. they are not pills or capsules, but part of a normal food pattern. Additionally, a functional food can be a natural or unmodified food, or one to which a component has been added or removed by technological or biotechnological means. It can also be a food where the nature of one or more components has been modified, the bioavailability altered, or any combination of these possibilities. Additionally, a functional food might be functional for all members of a population or for particular groups only. It is also important to note that, along with the nonuniversal definition, global markets also do not have the same regulatory systems for these foods (Bagchi 2014).

Functional food science is still at an early stage in its development. However, since knowledge about the functional effects of foods is increasing and the functionality of particular foods and food components is more extensively recognized, technology will have a continuing role to play in making those foods and food components more widely available and accessible (Howlett 2008). On the other hand, it is now known that genetic factors influence the relationship between diet and disease, and the ways in which different protective and risk factors can act. Furthermore, it is possible to visualize differences between genetic profiles of individuals at the molecular level and understand how they relate to differences between those individuals’ responses to physiological factors. Thus, in the near future, knowledge gained in the fields of genomics, proteomics, and metabolomics (collectively known as “omics”) will be of great importance for the development of functional foods and to create customized diet programs, as well as verifying the influence of dietary factors on human health and disease, which can lead to the identification of new food functionality routes (Howlett 2008).

9.2.2 Classification and Development of Functional Foods

Functional foods represent one of the most interesting and active areas of research and innovation in the food industry (Annunziata & Vecchio 2011). Their design and development, besides being an expensive process (Betoret *et al.* 2011), is a key issue, as well as a scientific challenge, which should rely on basic scientific knowledge relevant to target functions and their possible modulation by food components (Diplock *et al.* 1999). It is possible to separate them into natural (or nonaltered) and modified functional foods. But whether modified or not, they should always be safe, without any consideration of a trade-off between health benefit and health risk. More specifically, and according to the definition of functional foods presented before, they can be classified as:

- *nonaltered products*: foods naturally containing increased content of nutrients and/or health-promoting compounds
- *fortified products*: foods wherein the content of the existing components is increased
- *enriched products*: foods to which a component not normally found is added to provide benefits
- *altered products*: foods in which a component is removed or replaced by an alternative component with favorable properties
- *enhanced commodities*: the food composition is altered by changing the raw commodity, i.e. one of the components is enhanced through special growing conditions, breeding, or biotechnological means.

Although the functional food industry is growing steadily worldwide, the successful commercialization of new functional foods remains a challenge, especially due to the need for a strategic approach to their production processes (Howlett 2008). For this reason, during the development or reengineering of modified functional foods, it is necessary to take into account many variables, such as sensory acceptance, convenience, stability, chemical and functional properties, and price (Betoret *et al.* 2011; Granato *et al.* 2010). In fact, the relationship “structure-property” needs to be noted, once the functional effect depends on the active component gaining access to the functional target site. However, foods are mostly complex mixtures that can trap active compounds, modulate their release, or inhibit their activity. Thus, the food matrix in its raw state, after culinary preparation, or storage can have a significant influence on the activity or release of the key components. According to Betoret *et al.* (2011), the design of appropriate food vehicles to maintain the active form until the time of consumption, and to deliver this form to the desired target site within the organism, is vital to the success of functional foods.

Betoret *et al.* (2011) grouped the available technologies for functional foods development into three main categories. The first group is formed by the most commonly used technologies for functional foods development, including technologies traditionally used in food processing, formulation, and blending as well as for cultivation and breeding. The second group, constituted by methodologies that form a structure to try to prevent the deterioration of physiologically active compounds, includes microencapsulation, edible films and coatings, and vacuum impregnation technologies. The third group, formed by recent technologies that are intended to design functional foods aimed at personalized nutrition, is the one that has grown significantly in recent years.

9.2.3 Wild Plants Used as Functional Foods

Plants are irreplaceable food resources for humans. Their interchangeable use as foods and as medicines, or healthy foods, has been part of human heritage since prehistoric times. Despite only a small number of existing plant foods having substantial clinical documentation of their health benefits, an even smaller number (and including only cultivated plants) have surpassed the rigorous standard of “significant scientific agreement” required by the FDA and EFSA for authorization of a health claim (Hasler 2002). Oat soluble fiber, soluble fiber from psyllium seed husk, soy protein and sterol and stanol ester-fortified margarine are plant-based foods currently eligible to bear an FDA-approved health claim (Hasler 2002). However, there is growing clinical research supporting the potential health benefits of various plant foods (including wild plants) or food constituents that currently do not have approved health claims, and thus are described as having “moderately strong evidence.” Examples include berries, leafy vegetables, garlic, grapes and chocolate, among others listed in Table 9.1.

Table 9.1 presents wild edible plants that have been investigated due to their claimed functional properties. These plants are interesting sources of physiologically active ingredients which are linked to various beneficial health effects. Various berries, including elderberry, bilberry, cranberry, blackberry, raspberry, and wild strawberry, stand out as a source of anthocyanins, proanthocyanidins, flavonols, phenolic acids, and vitamins, among other bioactive compounds. These molecules, isolated or in combined extracts, have antioxidant, antiinflammatory, anticarcinogenic, cardioprotective, and antibacterial properties (Barros *et al.* 2011a; Bowen-Forbes *et al.* 2010; Madhavi *et al.* 1998; Najda *et al.* 2014; Sidor & Gramza-Michalowska 2014; Singh *et al.* 2009). Wild strawberry fruits harvested from natural habitats were highlighted by Najda *et al.* (2014) as containing more anthocyanins and higher antioxidant activity than those from cultivation. Likewise, Lv *et al.* (2014) showed that the wild litchi cultivar Hemaoli has high total phenolic and flavonoid content in comparison to one of the main market cultivars. This fruit also has high levels of carotenoids and vitamin C, which contribute to its antioxidant, antiapoptotic, and hepatoprotective effects (Bhoopat *et al.* 2011; Huang *et al.* 2010a; Lv *et al.* 2014). *Physalis* (*Physalis* spp.) is another berry with claimed functional properties. Physalins, withanolides, sterols, polysaccharides, and flavones are compounds present in this golden berry. According to Li *et al.* (2014c), it has antiinflammatory, antioxidant, antitumor, hypoglycemic, and analgesic properties.

Other plants, like the root of beet (*Beta* spp.), have antioxidant, hepatoprotective, anticancer, and antiproliferative activity in MRC5 and MCF-7 cell lines, antihypertensive, and hypoglycemic effects. These health benefits are conferred by the high content of phenolic acids, flavonoids, betalains, minerals (P, Mg, Fe, Zn, Ca, and Na), folic acid, biotin, and soluble fiber (Ninfali & Angelino 2013; Vulić *et al.* 2014; Wootton-Beard & Ryan 2011). In turn, ginger (*Zingiber officinale* Roscoe) has been described as a source of gingerols (6-gingerol), shogaols (6-shogaol), fiber, and flavonoids, as well as having antioxidant, antiinflammatory, antithrombotic, cholesterol-lowering, analgesic, antipyretic, and hypotensive effects (Mojani *et al.* 2014; Thomson *et al.* 2002).

Regarding leafy vegetables, the aerial parts of water blinks (*Montia fontana* L.) have high amounts of tocopherols and vitamin C, compounds that provide antioxidant benefits (Morales *et al.* 2012; Pereira *et al.* 2011), while watercress (*Nasturtium officinale* W.T. Aiton) is a rich source of phenolic compounds and minerals (P, Mg, Ca, and Mn)

Table 9.1 Some wild edible plant foods claimed to have functional properties.

| Plant species | Common name and used part | Evaluated extract | Functional compounds | Potential health benefits | Reference |
|--|---------------------------|---|---|--|--|
| <i>Allium ampeloprasum</i> L. | Wild leek (bulb) | Aqueous and ethanolic extracts | Fiber, zinc, polyunsaturated fatty acids (mainly palmitic acid), polysaccharides (glucofructan), and steroidal saponins | Antioxidant, antiinflammatory, antiulcerogenic, and gastroprotective | Adão <i>et al.</i> 2011; García-Herrera <i>et al.</i> 2014; Malafaia <i>et al.</i> 2015 |
| <i>Arbutus unedo</i> L. | Strawberry tree (fruit) | Methanolic, hydromethanolic (80%), and phenolic extracts | Tocopherols, ascorbic acid, flavan-3-ols, and galloyl derivatives | Antioxidant activity and antitumor potential against NCI-H460 human cell line | Barros <i>et al.</i> 2010; Guimarães <i>et al.</i> 2013, 2014 |
| <i>Asparagus acutifolius</i> L. | Wild asparagus (shoots) | | Tocopherols, vitamin C, and glycosides of flavonols | Antioxidant | Barros <i>et al.</i> 2011b; Martins <i>et al.</i> 2011 |
| <i>Beta</i> spp. | Beet (root) | Aqueous, hydroethanolic, methanolic, and betelain-rich extracts and juice | Phenolic acids (ferulic, vanillic, <i>p</i> -hydroxybenzoic, caffeic and protocatechuic acids), flavonoids (catechin, epicatechin, rutin and vitexin), betalains (betanin, isobetanin and vulgaxanthin I), minerals (potassium, magnesium, iron, zinc, calcium, sodium), folic acid, biotin and soluble fiber | Antioxidant, hepatoprotective, anticancer, antiproliferative activity in MRC5 and MCF-7 cell lines, antihypertensive and hypoglycemic | Wootton-Beard & Ryan 2011; Ninfali & Angelino 2013; Vulić <i>et al.</i> 2014 |
| <i>Capparis decidua</i> (Forssk.) Edgew. | Caper or kair (fruit) | Aqueous, methanolic, acidified methanolic, hydroalcoholic, and ethanolic extracts | N-pentacosane, β -sitosterol, β -carotene, alkaloids, phenolic compounds including flavonoids and minerals (manganese, copper, and iron) | Antioxidant, antidiabetic, diuretic, hypercholesterolemic, antihypertensive, antiatherosclerotic, hypolipidemic, antimicrobial, and anthelmintic | Rathee <i>et al.</i> 2010; Sharma <i>et al.</i> 2010; Zia-Ul-Haq <i>et al.</i> 2011; Shad <i>et al.</i> 2014 |

| | | | | | |
|--------------------------------|--|---|--|---|---|
| <i>Dimocarpus longan</i> Lour. | Longan (fruit) | Aqueous, hydromethanolic (80%), acetone:ethanol (1:1, v/v), hydroacetone (70%) and polyphenol-rich extracts | Phenolic compounds (corilagin, gallic and ellagic acids, flavone glycosides, glycosides of quercetin and kaempferol and epicatechin), vitamin C, fiber, and minerals | Antioxidant, antiinflammatory, antityrosinase, antilycated, anticancer, and memory-enhancing effects | Huang <i>et al.</i> 2010a; Yang <i>et al.</i> 2011; Huang <i>et al.</i> 2012 |
| <i>Eugenia uniflora</i> L. | Pitanga or Brazilian cherry (fruit) | Ethyl acetate and ethanolic extracts | Anthocyanins, flavonols, and carotenoids | Antidiarrheic, diuretic, antirheumatic, antifebrile, antidiabetic, antimicrobial, and antitrypanosoma | Costa <i>et al.</i> 2013 |
| <i>Euterpe oleracea</i> Mart. | Acai, assai or açai (fruit) | Ethyl acetate, <i>n</i> -buthanolic, hydromethanolic (50%), and hydroacetonic (70%) extracts | Flavonoids (anthocyanins and procyanidin), phenolic acids, lignans, and stilbenes | Antioxidant, antiallergic, anticancer, antiinflammatory, atheroprotective, improves the endothelial function and platelet aggregation, vasodilation, and prevents cardiovascular disease | Rufino <i>et al.</i> 2010; Kang <i>et al.</i> 2011; Costa <i>et al.</i> 2013 |
| <i>Ficus carica</i> L. | Fig (fruit) | Hexane, methanolic, and hydromethanolic extracts | Phenolic acids (chlorogenic acid), anthocyanins, flavonols, flavones (luteolin), minerals (iron, potassium, sodium and calcium), fiber, sugars, and vitamin A | Antioxidant, anticholinesterase, anticarcinogenic, antiproliferative activity in several cancer cell lines, digestive, antifungal, and anthelmintic | Huang <i>et al.</i> 2010a; Barolo <i>et al.</i> 2014; Shad <i>et al.</i> 2014 |
| <i>Fragaria vesca</i> L. | Wild strawberry or European strawberry (fruit) | Aqueous extracts and combined extract of <i>n</i> -buthanolic and to HCl (1 mol/dm ³) | Flavonoids (e.g. anthocyanins), phenolic acids, and salicylic acid | Antioxidant | Najda <i>et al.</i> 2014 |
| <i>Garcinia mangostana</i> L. | Mangosteen or purple mangosteen (fruit) | Aqueous, methanolic, ethanolic, hydroethanolic (40 and 50%), and juice extracts | Xanthones (α -, β -, and γ -mangostins, garcinone E, 8-deoxygartanin and gartanin) | Antioxidant, antitumor, antiproliferative, proapoptotic, antiinflammatory, antiallergic, antibacterial, antifungal, antiviral, antimalarial, antidiabetic, antihyperlipidemic and antiatherogenic, cardioprotective, hepatoprotective, immunomodulator, and antiulcer | Pedraza-Chaverri <i>et al.</i> 2008; Gutierrez-Orozco & Failla 2013 |

(Continued)

Table 9.1 (Continued)

| Plant species | Common name and used part | Evaluated extract | Functional compounds | Potential health benefits | Reference |
|--------------------------------------|--------------------------------------|--|--|--|--|
| <i>Gardenia jasminoides</i> J. Ellis | Gardenia (fruit) | Hydromethanolic (80%), hydroethanolic (60%), and acetone:ethanol (1:1, v/v) extracts | Caffeoylquinic acid derivatives (chlorogenic acid, dicaffeoyl-quinic acid and other caffeoyl-conjugate quinic acid derivatives), flavonoids (rutin), iridoids (geniposide), and carotenoids (crocin) | Antioxidant and antiinflammatory | Huang <i>et al.</i> 2010a; Peng <i>et al.</i> 2013 |
| <i>Litchi chinensis</i> Sonn. | Litchi or lychee (fruit) | Methanolic (70% and 80%), acetone:ethanol, (1:1) and juice extracts | Phenolic compounds (cinnamic acid and procyanidins), carotenoids, and vitamin C | Antioxidant, antiapoptotic, and hepatoprotective | Huang <i>et al.</i> 2010a; Bhoopat <i>et al.</i> 2011; Lv <i>et al.</i> 2014 |
| <i>Lycium barbarum</i> L. | Goji berry (fruit) | Aqueous, methanolic, and crude and purified polysaccharide extracts | Polysaccharides, carotenoids (zeaxanthin), betaine, cerebroside, β -sitosterol, <i>p</i> -coumaric acid, and vitamin C | Antioxidant, antiaging, antiinflammatory, anticancer, cytoprotective, neuroprotective, metabolism stimulator; glucose regulator in diabetics, glaucoma (eye health benefits), immunomodulatory, antibacterial and cardioprotective | Amagase & Farnsworth 2011 |
| <i>Malpighia emarginata</i> D.C. | Acerola or wild crepe myrtle (fruit) | Methanolic, hydromethanolic (50%), hydroacetonic (70%), and aqueous extracts and juice | Vitamin C, carotenoids (β -carotene), riboflavin, thiamine, fiber, minerals (phosphorus, calcium, and iron), anthocyanins (cyanidin-3-rhamnoside and pelargonidin-3-rhamnoside) and flavonols (quercetin) | Antioxidant, antiaging, antiinflammatory and prevents weight gain and dyslipidemia | Mezadri <i>et al.</i> 2008; Rufino <i>et al.</i> 2010 ; Delva & Goodrich-Schneider 2013; Dias <i>et al.</i> 2014 |
| <i>Montia fontana</i> L. | Water blinks (aerial parts) | Methanolic extract | Tocopherols and vitamin C | Antioxidant | Pereira <i>et al.</i> 2011; Morales <i>et al.</i> 2012 |

| | | | | | |
|---|---|---|---|--|---|
| <i>Myrciaria cauliflora</i> (Mart.) O. Berg | Jaboticaba or guapuru (fruit) | Methanol:formic acid (9:1, v/v), methanol:water:acetic acid (85:15:0.5, v/v/v/), methanolic, hydromethanolic (50%), ethanolic, acetonic, and hydroacetonic (70%) extracts | Anthocyanins, ellagic and gallic acid, carotenoids, depsides, tannins, rutin, and vitamin C | Antioxidant, antiinflammatory, inhibits IL-8 production, antiproliferative effects against tumor cells, protective effect in cardiovascular disease and type 2 diabetes mellitus | Rufino <i>et al.</i> 2010; Leite <i>et al.</i> 2011; Costa <i>et al.</i> 2013 |
| <i>Myrciaria dubia</i> (Kunth.) McVaugh | Camu-camu, cacari or camocamo (fruit) | Hydromethanolic (50%) and hydroacetonic (70%) extracts | Anthocyanins, myricetin and conjugates, ellagic acid and conjugates, ellagittannins, flavan-3-ols, proanthocyanidins, and vitamin C | Antioxidant, antiinflammatory and inhibits the LPS-induced NO release in RAW 264.7 cells | Rufino <i>et al.</i> 2010; Costa <i>et al.</i> 2013; Fracassetti <i>et al.</i> 2013 |
| <i>Nasturtium officinale</i> W. T. Alton | Watercress (aerial parts) | Methanolic and hydromethanolic (70%) extracts | Phenolic compounds and minerals (phosphorus, potassium, calcium, and manganese) | Antioxidant, anticarcinogenic, and chemopreventive | Pereira <i>et al.</i> 2011; Manchali <i>et al.</i> 2012 |
| <i>Origanum vulgare</i> L. | Oregano (aerial parts) | Aqueous (infusions and decoctions) and hydromethanolic extracts (80%) | Flavonoids and phenolic acids | Antioxidant and antimicrobial potential | Martins <i>et al.</i> 2014 |
| <i>Physalis</i> spp. | Physalis or golden berry (fruit) | Hydroethanolic (70%) extracts | Physalins, withanolides, sterols, polysaccharides, and flavones | Antiinflammatory, antioxidant, antitumor, hypoglycemic, and analgesic | Li <i>et al.</i> 2014c |
| <i>Prosopis cineraria</i> (L.) Druce | Ghaf, khejri, sami or golden tree of Indian deserts (pod) | Aqueous and methanolic extracts | Triterpenoids (3-benzyl-2-hydroxy-urs-12-en-28-oic acid and maslinic acid-3 glucoside), fatty acid (linoleic acid), piperidine alkaloid (prosophylline) and polyphenols (5,5'-oxybis-1,3-benzenediol, 3,4,5-trihydroxycinnamic acid 2-hydroxyethyl ester and 5,3',4'-trihydroxyflavanone 7-glycoside) | Antioxidant and antiinflammatory | Liu <i>et al.</i> 2012 |

(Continued)

Table 9.1 (Continued)

| Plant species | Common name and used part | Evaluated extract | Functional compounds | Potential health benefits | Reference |
|---|----------------------------|--|--|---|---|
| <i>Prunus spinosa</i> L. | Blackthorn or sloe (fruit) | Methanolic and phenolic enriched extracts | Ascorbic acid, phenolic acids and flavonoids (anthocyanins, flavonols and flavones) | Antioxidant and antitumor potential | Barros <i>et al.</i> 2010; Guimarães <i>et al.</i> 2013, 2014 |
| <i>Psidium cattleianum</i> Sabine | Strawberry guava (fruit) | Hexane, ethyl acetate, acetonic, aqueous, ethanolic, and methanolic extracts | Phenolic compounds (ellagic acid, ellagic acid deoxyhexoside and epicatechin gallate), carotenoids, vitamin C, and fiber | Antioxidant, antiinflammatory, and antimicrobial | McCook-Russell <i>et al.</i> 2012; Ribeiro <i>et al.</i> 2014 |
| <i>Psidium guajava</i> L. | Guava (fruit) | Methanolic, hydromethanolic (80%), acetone:ethanol (1:1, v/v), hexane, ethyl acetate and ethanol/water/formic acid (70:25:5, v/v/v) extracts | Phenolic acids (chlorogenic acid), flavan-3-ols (catechin), anthocyanins (delphinidin-3-O-glucoside and cyanidin-3-O-glucoside) | Antioxidant, antiinflammatory, and antimicrobial | Huang <i>et al.</i> 2010a; McCook-Russell <i>et al.</i> 2012; Flores <i>et al.</i> 2015 |
| <i>Punica granatum</i> L. | Pomegranate (fruit) | Aqueous, ethyl acetate, acetonic, and methanolic extracts | Anthocyanins, gallotannins, ellagitannins (ellagic acid, gallic acid, and punicalagin), gallagyl esters, hydroxybenzoic and hydroxycinnamic acids and dihydroflavonol | Antioxidant, antiinflammatory, antiallergic, chemopreventive, anticancer, cardioprotective, gastroprotective, antimicrobial, and anthelmintic | Ismail <i>et al.</i> 2012 |
| <i>Rhodomyrtus tomentosa</i> (Aiton) Hassk. | Rose myrtle (fruit) | Hexane, methanolic, hydromethanolic (80%), acetone:ethanol (1:1, v/v), acetone:water:acetic acid (50:49:1, v/v/v), and flavonoid-rich extracts | Flavonoids (galocatechin gallate, dihydromyricetin, quercetin, kaempferol, anthocyanins and vitexin), organic acids, polysaccharides, fiber, vitamin E (α -tocopherol), minerals (manganese and copper) and essential fatty acids (mainly linoleic acid) | Antioxidant | Huang <i>et al.</i> 2010a; Lai <i>et al.</i> 2015; Wu <i>et al.</i> 2015 |

| | | | | | |
|-------------------------------------|---|---|--|--|--|
| <i>Rubus</i> spp. | Blackberry and raspberry (fruits) | Hexane, ethyl acetate, and methanolic extracts | Anthocyanins, flavonols, phenolic acids (ellagic acid), vitamins C and E, folic acid, and β -sitosterol | Antioxidant, antiinflammatory, and chemopreventive | Bowen-Forbes <i>et al.</i> 2010 |
| <i>Sambucus nigra</i> L. | Elderberry, black elder or European elder (fruit) | Methanolic, acidified methanolic, ethanolic, and hydroethanolic (80%) extracts | Polyphenols (anthocyanins, flavonols, phenolic acids and proanthocyanidins), terpenes, lectins, unsaturated fatty acids, fiber, vitamins A, B, C and E, and minerals | Antioxidant, cardiovascular protection, antidiabetic and antiobesity, reinforces the immune system, antiviral, antibacterial, and UV radiation protector | Barros <i>et al.</i> 2011a; Sidor & Gramza-Michałowska 2014 |
| <i>Syzygium cumini</i> (L.) Skeels. | Jambul or jambolan (fruit) | Methanolic, hydromethanolic (50%), hydroacetic (70%), and hexane extracts | Anthocyanins, phenolic acids (ellagic acid), flavonols (quercetin and rutin), carotenoids, vitamin C, and manganese | Antioxidant, antiscorbutic, diuretic, and antidiabetic | Rufino <i>et al.</i> 2010; Costa <i>et al.</i> 2013; Shad <i>et al.</i> 2014 |
| <i>Vaccinium myrtillus</i> L. | Bilberry (fruit) | Acidified methanolic, ethyl acetate, hexane, anthocyanins, and proanthocyanidin-rich extracts | Flavonoids (proanthocyanidins and anthocyanins), carotenoids (lutein and zeaxanthin), and sterols | Antioxidant, antibacterial (inhibition of urinary tract infections), anticarcinogenic and antiproliferative activity in two human breast cancer cell lines MCF-7 and BT-20 | Madhavi <i>et al.</i> 1998 |
| <i>Vaccinium</i> spp. | Cranberry (fruit) | Hydroacetic (80%), ethyl acetate, and phenolic extracts | Phenolic acids and flavonoids (anthocyanins, proanthocyanidins, and flavonols) | Antioxidant, antiinflammatory and cardiovascular and urinary tract protection | Singh <i>et al.</i> 2009; Khoo & Falk 2014 |
| <i>Zingiber officinale</i> Roscoe | Ginger (rhizome) | Aqueous and methanolic extracts | Gingerols (6-gingerol), shogaols (6-shogaol), fiber, and flavonoids | Antioxidant, antiinflammatory, antithrombotic and cholesterol lowering, analgesic, antipyretic, and hypotensive | Thomson <i>et al.</i> 2002; Mojani <i>et al.</i> 2014 |
| <i>Ziziphus jujuba</i> Mill. | Jujube or red date (fruit) | Aqueous, hexane, methanolic, and hydromethanolic extracts | Saponins, tannins, terpenoids, flavonoids, and iron | Antioxidant, antiinflammatory, and gastrointestinal protector | Yu <i>et al.</i> 2012; Shad <i>et al.</i> 2014 |

which confer its claimed antioxidant, anticarcinogenic, and chemopreventive effects (Manchali *et al.* 2012; Pereira *et al.* 2011). The aerial parts of oregano (*Origanum vulgare* L.), prepared in infusions, decoctions or hydromethanolic extracts (80%), have antioxidant and antimicrobial potential probably related to flavonoids and phenolic acids (Martins *et al.* 2014).

Today, aggressive marketing highlighting the health-promoting benefits of mangosteen, acai, acerola or goji berry, among other fruits, bulbs, roots, seeds or leafy vegetables presented in Table 9.1, has resulted in their classification as “superfruits” or “superfoods.” Scientific research carried out in recent years proves their effectiveness as healthy foods, and due to high profits, the food and pharmaceutical industries are increasingly interested in developing new products based on these plants.

However, in addition to edible plant parts, wild nonedible parts or plants can also be used as a source of health-promoting ingredients. Thus, medicinal and aromatic plants play an important role in the development of new or improved functional foods, as well as nutraceuticals. At the research level, some wild plant extracts are being incorporated into food products to increase their health-promoting properties. Martins *et al.* (2014) formulated new yogurts based on phenolic extracts of wild blackberry (*Rubus ulmifolius* Schott) flowers. The authors microencapsulated the hydroalcoholic extract in an alginate-based matrix and incorporated this into a yogurt to achieve antioxidant benefits. Recently, Caleja *et al.* (2015) improved the antioxidant properties of cottage cheese by the incorporation of fennel (*Foeniculum vulgare* Mill.) decoction (phenolic-enriched extract), improving not only functionality of the final product but also preservation effectiveness due to the antimicrobial potential of fennel.

Carocho *et al.* (2015a) transformed the Portuguese “Serra da Estrela” cheese into a functional food by incorporating dried chestnut (*Castanea sativa* Mill.) flowers or lemon balm (*Melissa officinalis* L.) plants, as well as their decocted extracts. The functionalized cheeses showed higher antioxidant activity, especially lipid peroxidation inhibition, bringing benefits both for consumers (healthier product) and producers (added-value product). The same authors also functionalized the Portuguese traditional cakes “económicos” by incorporation of dried chestnut (*C. sativa*) flowers or decoctions prepared from them (Carocho *et al.* 2015b). The final product showed increased antioxidant activity and phenolic content, without causing visible changes in inner and outer appearance.

9.3 Wild Plant-Based Nutraceuticals

9.3.1 The Emerging Concept and Applications of Nutraceuticals

A new generation of processed food is coming, which is a controversial subject for many people. Nutraceutical products represent a fast-growing sector within the food industry, aiming to increasingly attract the buyer to consume these novel dietary supplements and phytotherapeutic products. It is expected that in the near future, “food for special dietary needs,” such as soups, smoothies, processed meat, bread and sausages, among others, will be enriched with nutraceutical formulations (Andlauer & Furst 2002; Regulation (EC) No. 2002/46).

Nutraceuticals can be defined as diet supplements that contain bioactive compounds or extracts, prepared from raw natural matrices that will provide a higher dosage that

could not be obtained from normal food products and functional foods (DeFelice 1992; Espín *et al.*, 2007; Zeisel 1999). Directive 2002/46/EC of the European Parliament and Council, on the approximation of laws of Member States relating to food supplements, defines “food supplements” as foodstuffs with the purpose of supplementing the normal diet and which are concentrated sources of nutrients or other substances with a nutritional or physiological effect, alone or in combination, marketed in dose form, such as capsules, pastilles, tablets, pills, sachets of powder, ampoules of liquids, drop dispensing bottles, and other similar forms of liquids and powders designed to be taken in measured small unit quantities (Regulation (EC) No. 2002/46).

The health industry is using nutraceutical formulations as complements to prevent some diseases. Some authors have stated that any food or parts of foods can be considered nutraceutical compounds, as long as their beneficial health and nutritional claims are proved scientifically (Braithwaite *et al.*, 2014; McNamara, 1997; Ross, 2000). On the other hand, Gulati and Ottaway (2006) and Espín *et al.* (2007) distinguished nutraceuticals as components that are often consumed in unit dose forms such as tablets, capsules or liquids. They can be isolated nutrients or herbal products presented in pharmaceutical forms or processed products like cereals, smoothies, and soups for special diet requirements (Andlauer & Furst 2002; Braithwaite *et al.* 2014; Regulation (EC) No. 2002/46).

The concept of nutraceuticals is relatively recent, only appearing in the 1990s with the first publications and patents related to the subject (Figure 9.1). However, the increasing number of publications from academics (through articles and reviews) and industry (through patents) is notable. This can be explained by the fact that there is increasingly market demand for new, better, and safer food products. However, regarding plant-based nutraceuticals, the number of articles (and reviews) and patents is very low (see Figure 9.1) although it is growing. Many of the primary studies on nutraceuticals were made with individual compounds with known beneficial effects, but there is now interest in exploring the synergisms existing within plant extracts and incorporating them into nutraceuticals or modified functional food.




Figure 9.1 Number of research articles and reviews (■ and - - -), and patents (■ and.....) published in the period from 1990 to 2015 regarding nutraceuticals and nutraceuticals formulated with plant material, respectively (obtained on Web of Science, January 2015; keyword: nutraceutical; nutraceutical + plant).

Therefore, the development of legislation that regulates the production of nutraceutical formulations, their labeling and market supply is crucial. In January 2002, the European Food Safety Authority (EFSA) established Regulation (EC) No. 178/2002, setting down the general principles and requirements for food law. This Regulation also contains procedures for food safety, increasing the health protection of consumers. These guidelines are applied to all food products, including those with added functional properties, such as nutraceuticals and functional foods. From the perspective of the global pharmaceutical and medical industry, nutraceutical products are dietary supplements (Kwak & Jukes 2001). Specific regulation of nutraceutical products is still very patchy; in European law they have no specific category, being considered under the same parameters used for dietetic foods, dietetic supplements, and food supplements (Coppens *et al.* 2006; Regulation (EC) No. 2002/46) or even under medicinal classification (Gulati & Ottaway 2006). In the USA, nutraceuticals are considered only as dietary supplements (Bernal *et al.* 2011; Espín *et al.* 2007). The differences between European and USA regulation may be due to cultural, historical, and traditional backgrounds (Gulati & Ottaway 2006). However, the development of specific legislation in Europe is necessary to ensure food safety for consumers and to prove that nutraceuticals are safe and scientifically accepted, and this may dictate the future success of these products (Braithwaite *et al.* 2014; Byrne 2003).

Recent research has shown very promising prospects for different natural ingredients added to food products, creating benefits for consumers' health and added value for manufacturers (Coppens *et al.* 2006). Many of the published papers on nutraceuticals are focused on their beneficial health properties (Bernal *et al.* 2011), for instance their ability to decrease the development of heart disease (Garcia-Rios *et al.* 2013; Giordano *et al.* 2012; Izzo *et al.* 2010; Scicchitano *et al.* 2014) such as hypercholesterolemia (Mannarino *et al.* 2014), and also for the prevention and treatment of prostate cancer (Li *et al.* 2014b). Nutraceutical formulations have been proved to be safe and well tolerated, but further studies are required to assess the decreasing of secondary effects of nutraceuticals when compared to analogue commercial drugs for the treatment of certain diseases (Bernal *et al.* 2011; McAlindon 2006).

9.3.2 Recent Advances in Formulations for Nutraceuticals

Due to the difficulty in the classification of nutraceuticals, we are faced with two types of products: nutraceuticals in the form of dietary supplements (tablets, capsules, solutions, syrups, powders, chewing tablets, among others) and those in the form of free or encapsulated extracts/compounds to be inserted into a food matrix (i.e. used to develop functional foods). For that reason, the formulation of nutraceuticals involves a wide range of methodologies and techniques, from the most used (tableting) to the newest and most advanced, such as microencapsulation complemented with nanotechnology.

First, it is necessary to ensure the safety and quality of the nutraceutical product. The chemical, nutritional, and bioactive characterization of the compound/extract that will be part of the formulation is required, as well as control of the dosage. For this, some advanced analytical techniques are used such as mass spectrometry (MS), nuclear magnetic resonance (NMR), high-performance liquid chromatography (HPLC), capillary electrophoresis (CE), and gas chromatography (GC), among others (Bernal *et al.* 2011; Sener & Orhan 2005).

The vast majority of nutraceutical formulations are designed for oral administration. Braithwaite *et al.* (2014) reported a description of some new nutraceutical formulation strategies to improve dosage, design, and delivery of the bioactives. From liposomal carriers, electrospun fiber mats, microsponges and nanosponges, cyclodextrin complexations to biodegradable hydrogels, all these technologies prove the importance of nutraceuticals in today's economy, with a growing investment by the industry in new formulations that respond to market demand. Second-generation nanocrystals, another new formulation, are an emerging technology for the delivery of poorly soluble bioactives. They are mostly used for drug delivery to solve poor solubility and bioavailability. However, they also represent a reliable response for the delivery of many nutraceutical compounds already on the market, such as antioxidants. The main advantage of nanocrystal systems is the capacity to be applied via oral, intravenous, dermal, mucosal, ocular and even pulmonary routes (Shegokar & Müller 2010).

It is important to realize that nutraceutical formulations go far beyond diet products or products enriched with a certain bioactive compound. Formulations are already on the “micro” and “nano” scales, which can be incorporated in food matrices but also in pharmaceutical formulations, serving as a complement to traditional medicine. Microencapsulation complemented with nanotechnology appears to overcome problems related to the use of free bioactives but also to provide controlled target delivery release (Braithwaite *et al.* 2014; Dias *et al.* 2015 Ezhilarisi *et al.* 2013; Huang *et al.* 2010b). Nanoscale delivery systems have the advantages of improving solubility, masking undesirable flavors and smells, and preventing the degradation of the bioactive compounds; they provide a triggered controlled release and, most important of all, increased bioavailability by prolonging contact within the gastrointestinal tract (Cerqueira *et al.* 2014). Microemulsions, for instance, are one of the most used techniques for the solubilization and transport of water and oil-insoluble compounds, presenting easier formulation and manufacture and also high stability during storage (Spernath & Aserin 2006).

Food protein-based materials can also be used at “micro” and “nano” scales, depending on the type of encapsulation methodology used to produce the capsules and also the objective of the work. Proteins present the ability to form gels and emulsions due to their functional properties, which makes them appealing to the industry and academia for the encapsulation of nutraceuticals (Chen *et al.* 2006). Hydrocolloids fibers are being proposed to encapsulate nutraceutical compounds and extracts; they are nontoxic, inexpensive, and generally recognized as safe (GRAS). Furthermore, since they are complex carbohydrates, consumption on a regular basis showed health benefits for cardiovascular disease and diabetes (Janaswamy & Youngren 2012). Researchers are also developing formulations linking nutraceuticals with drugs to enhance efficacy and reduce dosage and side-effects of chemical compounds (Braithwaite *et al.* 2014).

9.3.3 Examples of Nutraceuticals Based on Wild Plants

For economic and ecological sustainable reasons, the FAO recommends the cultivation of medicinal and aromatic plants that represent a genetic pool of raw material with better control of biotic and abiotic factors, allowing the standardization of the final product (Schippmann *et al.* 2002). For that reason all the listed examples in Table 9.2 are

Table 9.2 Nutraceutical formulations based on plants with traditional wild use.

| Plant species | Used part | Extract | Formulation | Application | Available commercial product | Reference |
|--|-----------------|-----------------------------|--------------|--|------------------------------|-----------------------------------|
| <i>Abelmoschus manihot</i> L. | – | Standardized extract | Pills | Renal inflammation, glomerular; chronic kidney disease | – | Tu <i>et al.</i> 2013 |
| <i>Artemisia annua</i> L. | Leaves | Dried leaves | Tablet | Antimalarial drug | – | Weathers & Towler 2014 |
| <i>Berberis aristata</i> D.C. | Root | Standardized extract | Tablet | Glycemia and lipid alteration | Berbenol® | Pierro <i>et al.</i> 2013 |
| <i>Silybum marianum</i> (L.) Gaertn. | Fruit | | | | | |
| <i>Boswellia serrate</i> Triana & Planch. | – | Gum extract | Pills | Osteoarthritis | 5-Loxin® | Sengupta <i>et al.</i> 2010 |
| | – | Gum extract + essential oil | Pills | Osteoarthritis | Aflapin® | Sengupta <i>et al.</i> 2010 |
| <i>Cajanus cajan</i> L. | Seed | | Syrup | Sickle cell anemia | Ciklavit® | Imaga <i>et al.</i> 2013 |
| <i>Commiphora angolensis</i> Welw. | Roots | Aqueous extract | Syrup, Pills | Antihepatocellular carcinoma | – | Pereira <i>et al.</i> 2013b, 2014 |
| <i>Cucumis melo cantalupensis</i> | Fruit | Juice | Pills | Stress and fatigue | Extramel® | Milesi <i>et al.</i> 2009 |
| <i>Cynara scolymus</i> L. | Leaves | Aqueous extract | Syrup, Pills | Antihepatocellular carcinoma | – | Pereira <i>et al.</i> 2013b, 2014 |
| <i>Echinacea angustifolia</i> (D.C.) Hell. | Root | Polysaccharide extract | Syrup | Immunomodulatory effect | Polinacea® | Dapas <i>et al.</i> 2014 |
| <i>Echinacea purpurea</i> (L.) Moench | Leaves and root | Dried leaves and roots | Pills | Cold | Echinaforce® | Brinkeborn <i>et al.</i> 1989 |

| | | | | | | |
|---|------------|----------------------|--------------|--|------------|---------------------------------|
| <i>Echinacea purpurea</i> (L.) Moench | Root | Ethanol extract | Tablet | Bioavailability of bioactive compounds | – | Matthias <i>et al.</i> 2007 |
| <i>Echinacea angustifolia</i> (D.C.) Hell. | | | | | | |
| <i>Echinacea purpurea</i> (L.) Moench | Root | Standardized extract | Tablet | Immunological effects | Revitonil® | Wagner & Jurcic 2002 |
| <i>Glycyrrhiza glabra</i> L. | | | | | | |
| <i>Ficus carica</i> L. | Fruits | – | Syrup | Bioactive compounds | – | Puoci <i>et al.</i> 2011 |
| <i>Ginkgo biloba</i> L. | Leaves | Aqueous extract | Syrup, Pills | Antioxidant activity | – | Pereira <i>et al.</i> 2013a |
| | Leaves | – | Tablet | Mild cognitive impairment | – | Bäurle <i>et al.</i> 2009 |
| | Leaves | Standardized extract | Pills | Mild cognitive impairment | Memo® | Yakoot <i>et al.</i> 2013 |
| <i>Ginkgo biloba</i> L. | | | | | | |
| <i>Panax ginseng</i> sp. | Root | | | | | |
| <i>Ginseng panax</i> sp. | Root | Standardized extract | Tablet | Antianging | Eufortyn® | Xu <i>et al.</i> 2010 |
| <i>Hedera helix</i> Linné L. | Leaves | Dry leaves | Tablet | Cough | Prospan® | Stauss-Grabo <i>et al.</i> 2011 |
| <i>Hypericum perforatum</i> L. | Shoot tips | – | Tablet | Depression | – | Lenoir <i>et al.</i> 1999 |
| | | | | | | |
| <i>Juglans regia</i> L. | Leaves | Ethanol extract | Pills | Hyperglycemia | – | Hosseini <i>et al.</i> 2014 |
| <i>Magnolia officinalis</i> Rehder & Wilson | Bark | Standardized extract | Pills | Reducing stress and anxiety | Relora® | Talbott <i>et al.</i> 2013 |
| <i>Phellodendron amurense</i> Rupr. | | | | | | |
| <i>Mikania laevigata</i> Willd. | – | – | Syrup | Antispasmodic and respiratory diseases | – | Graca <i>et al.</i> 2007 |
| | | | | | | |
| <i>Murraya koenigii</i> (L.) Sprengel | Leaves | Standardized extract | Pills | Benign prostatic hyperplasia | – | Sengupta <i>et al.</i> 2011 |
| <i>Tribulus terrestris</i> L. | | | | | | |

(Continued)

Table 9.2 (Continued)

| Plant species | Used part | Extract | Formulation | Application | Available commercial product | Reference |
|---|------------------|-------------------------|--------------|--|------------------------------|-----------------------------------|
| <i>Peumus boldus</i> L. | Leaves | Dried leaves | Tablet | Choleretic, diuretic, stomachic, cholagogic properties | – | Palma <i>et al.</i> 2002 |
| <i>Phellodendron amurense</i> Rupr. | Bark | Standardized extract | Pills | Joint pain and movements | – | Oben <i>et al.</i> 2009 |
| <i>Citrus sinensis</i> (L.) Osbeck | Peel | – | – | – | – | – |
| <i>Phoenix canariensis</i> Chabaud | Sap | – | Syrup | Sugar and nutritional source | – | Luis <i>et al.</i> 2012 |
| <i>Phyllanthus amarus</i> Schum. & Thonn | Leaves and stems | – | Syrup | Antitussive | – | Avbunudiogba <i>et al.</i> 2013 |
| <i>Proposis</i> pods L. | Fruit | – | Syrup | Bioactive compounds | – | Quispe <i>et al.</i> 2014 |
| <i>Salvia officinalis</i> L. | Leaves | Ethanolic/water extract | Pills | Glycemia | – | Kianbakht & Dabaghian 2013 |
| <i>Silybum marianum</i> (L.) Gaertn. | Bark | Aqueous extract | Syrup, Pills | Antihepatocellular carcinoma | – | Pereira <i>et al.</i> 2013b, 2014 |
| <i>Smallanthus sonchifolius</i> Poepp. & Endl | Root | – | Syrup | Obesity and insulin resistance | – | Genta <i>et al.</i> 2009 |
| <i>Uncaria tomentosa</i> (Willd.) D.C. | Bark | Aqueous extract | Pills | Macrophage resistance | – | Lenzi <i>et al.</i> 2013 |
| <i>Vitis vinifera</i> L. cv. <i>Cabernet sauvignon</i> ; <i>Vitis vinifera</i> L. cv. <i>Merlot</i> | Grape | – | Tablet | Bioavailability of resveratrol | – | Ortuño <i>et al.</i> 2010 |

plants that are normally consumed as wild and that present some bioactive properties, allowing the development of nutraceutical formulations. In this chapter, we only discuss nutraceutical formulations in the form of dietetic supplements (capsules, tablets, syrup). A detailed description of microencapsulated nutraceuticals based on plants has been previously provided by Dias *et al.* (2015), where the most frequently used microencapsulation techniques and materials are described, and also the most common extracts and bioactive compounds, including also some applicability studies for the developed microcapsules (e.g. milk, cheese, yogurt, ice cream, pasta, meat, bread, and chewing gum enhanced with bioactive extracts of plant origin).

Currently, some trademarked plant-based nutraceuticals are used as adjuvants in several illness processes. For instance, Ciklavit® is a syrup which has effects against sickle cell anemia, prepared with an aqueous extract of *Cajanus cajan* L. seed, a plant found in semiarid tropical regions, commonly consumed in soups and rice dishes (Imaga *et al.* 2013). Xu *et al.* (2010) studied the effect of a commercial product, Eufortyn®, that comprises chemical compounds including terclatrated coenzyme Q₁₀ and creatine, but also an extract of *Ginseng panax* sp. roots, which has shown effects in the antiaging process of rats. The roots of *Echinacea purpurea* (L.) Moench and *Glycyrrhiza glabra* L. are the major constituents of a commercial tablet, Revitonil®, used mainly for its immunological effects, while the leaves of *Hedera helix* L. are used in Prospan®, marketed for cough symptoms (Stauss-Grabo *et al.* 2011; Wagner & Jurcic 2002). Extramel® consists of small pills of melon juice (*Cucumis melo* var. *cantalupensis* Naudin) used to treat stress and fatigue symptoms (Milesi *et al.* 2009), while 5-Loxin® and Aflapin® contain gum extract and gum extract plus oil, respectively, of *Boswellia serrata* Triana & Planch, being used for osteoarthritis (Sengupta *et al.* 2010).

The genus *Echinacea* is well known for its medicinal properties. Brinkeborn *et al.* (1989) reported the effects of pills (Echinaforce®) prepared from *E. purpurea* roots and leaves in the treatment of the common cold, while Dapas *et al.* (2014) studied the root syrup (Polinacea®) obtained using *E. angustifolia* (D.C.) Hell. for its immunomodulatory effects. Berbenol®, a tablet formulation made from an extract of *Berberis aristata* D.C. and *Silybum marianum* (L.) Gaertn., taking advantage of the synergistic effects of both plants, is used for the treatment of glycemia and lipid value alterations in patients with type 2 diabetes (Pierro *et al.* 2013).

However, most of the formulations reported in the literature as nutraceuticals do not reach the market due to a lack of more in-depth studies, including clinical trials, or for legal or technical reasons. Lenoir *et al.* (1999) tested the effects of tablets containing three different concentrations of shoot tips from *Hypericum perforatum* L. on symptoms in patients with mild to moderate depression. The bark of *Phellodendron amurense* Rupr., traditionally used in Chinese medicine, and the peel of *Citrus sinensis* (L.) Osbeck were inserted into pills in order to evaluate their beneficial effects in joint pain; both species contributed to weight loss in tested patients and also an improvement in their health status (Oben *et al.* 2009). The leaves of *Murraya koenigii* (L.) Sprengel and *Tribulus terrestris* L. are traditionally used in India for curry and to treat infertility and impotence, respectively. Sengupta *et al.* (2011) studied the effects of both plant pills in benign prostatic hyperplasia, obtaining satisfactory results.

Type 2 diabetes is increasing worldwide; moreover, the additional health problems related to this disease are also an important concern. Kianbakht and Dabaghian (2013) reported the effects of pills prepared from *Salvia officinalis* L. leaves in patients with

type 2 diabetes and hyperlipidemia, describing good results in contrast to the placebo group, and without adverse side-effects. Furthermore, Hosseini *et al.* (2014) proved the effectiveness of pills obtained from *Juglans regia* L. leaves in patients with type 2 diabetes. Genta *et al.* (2009) studied humans given “yacon” syrup obtained from the roots of *Smallanthus sonchifolius* Poepp. & Endl with a high fructooligosaccharide content, demonstrating beneficial health effects in insulin-resistant patients.

Age-related cognitive changes and dementia are also a worldwide concern. *Ginkgo biloba* L. leaves have been described as being able to affect some neurological properties. Bäurle *et al.* (2009) studied the effects of tablets made from the extract of this plant in mild cognitive impairment; the authors reported the nutraceutical as safe, effective, and acting as an adjuvant to patients who suffer from this illness. There is already on the market a product, Memo®, prepared from *G. biloba* leaves and *Panax ginseng* sp. roots, used against mild cognitive impairment, by slowing the cognitive decline that occurs during the aging process (Yakoot *et al.* 2013). Cognitive wellbeing is also related to stress, depression, and fatigue, and the methods used to treat stress conditions range from a balanced nutritional plan to powerful drugs such as benzodiazepines. Relora® is a pill formulation consisting of a blend of bark extracts of *Magnolia officinalis* Rehder & Wilson and *P. amurense* standardized to honokiol and berberine, respectively, used in the treatment of stress and anxiety; the results achieved in a clinical trial performed by Talbott *et al.* (2013) showed that the combination of these two plants improved a variety of mood state parameters, lowering fatigue and increasing vigor.

Current stress-related diseases are a direct consequence of our modern lifestyle; the human organism produces reactive oxygen species, which are related to higher incidences of cardiovascular, brain, and immune system diseases (Carocho & Ferreira 2013b). Therefore, nutraceutical formulations are also being studied for their antioxidant properties, such as the syrup obtained from the fruits of *Ficus carica* L. and *Prosopis* pods, both widely used in traditional cuisine to prepare desserts and sweets (Puoci *et al.* 2011; Quispe *et al.* 2014). Furthermore, the syrup and pills prepared from *G. biloba* leaves, known for their action against degenerative neurological diseases, as previously mentioned, but also for their action in the cardiovascular system and cerebral vascular activity, were studied for their antioxidant capacity, showing higher activity than the corresponding infusion and extract; this higher activity was attributed to the highest content in phenolic compounds (Pereira *et al.* 2013a). Pereira *et al.* (2013b, 2014) also tested different nutraceutical formulations (pills and syrup) prepared using *Cynara scolymus* L., *Cochlospermum angolensis* Welw., and *S. marianum*, known for their capacity to prevent oxidative stress and liver disease, in terms of antioxidant and antihepatocellular carcinoma activities; the synergistic effects between these nutraceuticals (mixtures) were also assessed, showing many advantages over individual components.

The bioavailability of nutraceutical formulations is also a hot research topic as metabolic reactions can decrease their bioactive properties. There are already some studies in this direction, such as the one conducted by Matthias *et al.* (2007) on liquid (alcoholic solution) and tablet formulations prepared with *E. purpurea* and *E. angustifolia* roots. Alkylamides, found in both species, were used as target compounds to evaluate the nutraceuticals' bioavailability; these compounds were rapidly and easily absorbed in both formulations. A similar study was performed with tablets prepared with red wine grape extracts made from *Vitis vinifera* L. cv. *Cabernet sauvignon* and *Vitis vinifera*

L. cv. Merlot, in order to assess the bioavailability of resveratrol; however, in this case the bioavailability was higher in the natural matrix than in the nutraceutical formulation (Ortuño *et al.* 2010).

Nutraceuticals can also combine plant-based principles with other natural matrices such as mushrooms. A good example is ASHMITM, a pill formulation use in asthma treatment, containing the plants *Sophora flavescens* Aiton and *glycyrrhiza uralensis* Fisch. (root aqueous extracts) and the mushroom *Ganoderma lucidum* (Curtis) P. Karst. (fruiting body aqueous extracts) (Kelly-Pieper *et al.* 2009). Wong *et al.* (2004) also studied the effects of *Coriolus versicolor* (L.: Fr.) Quél. and *Salvia miltiorrhiza* Bunge pills (polysaccharides extract) on the improvement of cellular immunity in healthy subjects, which proved to be effective and without adverse effects.

9.4 Wild Plant-Based Drugs

9.4.1 From the Bioactive Phytochemical to the Active Principle

Plants have been used as medicine by humans for thousands of years, since their first use as teas, tinctures, poultices, etc. to the isolation of morphine from opium in the early nineteenth century. Since then, administration methods have changed drastically (Balunas & Kinghorn 2005; Newman *et al.* 2000). Today, there are many sources of new bioactive compounds, including plants, bacteria, fungi, and marine organisms; in fact, from 1981 to 2002, 61% of the 877 new small molecule chemical compounds were derived from natural products, and in specific therapeutic areas (antibacterial, antifungal, antiparasitic, and antiviral treatments), these compounds have provided 70% of total drugs (Cechinel-Filho 2012). There are six classes of compounds that result from botanical sources:

- bioactive compounds that are used directly as drugs, as in the case of digoxin, used for heart conditions
- bioactive compounds with structures that may act as lead compounds to more potent drugs, for instance, paclitaxel, a mitotic inhibitor used in cancer chemotherapy
- chemophores, which are cells that transduce energy, and may be converted into druggable compounds
- pure phytochemicals that can be used as markers to standardize crude plant material
- phytochemicals that can be used as pharmacological tools
- herbal extracts as botanical drugs or green tea extracts (Katiyar *et al.* 2012).

Although there are numerous classes of compounds and methods of obtaining them, the pharmaceutical industry faces unprecedented challenges, with fewer compounds being found, tested, and released to the public. Typically, after *in vitro* assays showing bioactivity of a specific compound, it may start preclinical studies on animal models followed by a “New Drug Application” addressed to the FDA (USA) and EFSA (EU). If approved, the human studies take place, divided into three phases with escalating numbers of participants to determine the toxicity, side-effects, and other effects not detectable in animal models. The ideal approval process of a new drug is hardly ever linear, and several drawbacks ensue, meaning that several years to some decades may elapse before a compound is marketed as a drug (FDA 2014; Paul *et al.* 2010). Compounds leading to hypothetical

drugs must achieve suitable solubility and chemical stability, demonstrate effectiveness in animals (adequate pharmacological profile) and satisfactory bioavailability (with a good half-life), the interactions with cytochrome p450 (CYP450) must be clarified and finally, there must be no obvious toxicity (Cechinel-Filho 2012).

With the reduction of new compounds appearing as potential drugs, humans have once again turned to Nature in order to mitigate the relative void of combinatorial chemistry to find new compounds (Phillipson 2007). The quest for compounds in plants can be carried out in many ways.

- Random selection followed by chemical screening (simple tests that may lead to false positives and false negatives, rendering conclusions difficult to assess and the class of compounds responsible for the activity impossible to specify).
- Random selection followed by one or more biological assays (carried out in animals or *in vitro* assays that screen high volumes of plant species in order to find new drugs).
- Follow-up of biological activity reports (reports of plant extracts with interesting biological activity, which were not studied for their active principles).
- Follow-up of ethnomedical (traditional medicine) uses of plants – plants used in traditional systems like Ayurveda, Unani, Kampo, and traditional Chinese medicine which are not seen as credible by Western scientific methods and are harder to assess, but their undeniable results in many illnesses are impossible to overlook. Herbalism, folklore, and shamanism, which are also viewed with scepticism, are also considered due to their strong reliance on endemic plants.
- Use of databases (large literature sources systematically organized that allow correlation of ethnomedical practices with experimental biochemical and pharmacological activities or to identify plants with multiple effects) (Fabricant & Farnsworth 2001).

To achieve the final compound, a large number of molecules must be extracted from the medicinal plant through various methods.

- Percolation, used for poorly soluble plants or when the price of the plant is relevant. The matrix is placed in a container with solvent flowing through it.
- Countercurrent extraction is obtained by moving solvent through the raw plant in countercurrent.
- Supercritical fluid extraction is carried out by placing the raw plant in a container and filling it with supercritical fluid until the pressure and temperature rise by a considerable amount. These conditions help the fluid to achieve a very high solubility capacity, extracting the compounds of interest.
- Microwave-assisted extraction relies on microwaves that extract compounds more selectively and rapidly while depending less on solvents.
- Maceration is the process of placing the raw plant in a container for different periods of time, while kinetic maceration uses the same process but the mixture is maintained under constant stirring.
- Turbo-extraction uses a cold solvent at high sheer forces, which leads to particle reduction, cell disintegration, and temperature increase.
- Decoctions and infusions rely on hot water as the extractor. Infusions are prepared by adding the plant to boiling water, and maintaining it for 5–10 minutes, while decoctions are prepared by adding the plant to cold water and heating it until it boils, maintaining it for 5–10 minutes.

- Soxhlet extraction relies on cycles of extraction within a glass chamber in which the solvent boils and condenses back into contact with the plant. After filling the chamber it is unloaded into a glass recipient that is heated, evaporating the solvent, only to condense back into the chamber, in a cyclical way.
- Sublimation extraction sublimates the compounds of interest leaving behind impurities which then condense in another chamber.
- Steam distillation relies on steam to carry the compounds from the boiling mixture containing the plant which then condenses.
- Ultrasonic-assisted extraction is used to increase mass transfer between the plant material and a solution by inducing liquid circulation and turbulence (Cechinel-Filho 2012; Sarker & Nahar 2012; Sticher 2008).

After extraction, the solutions have to be screened to determine their constituents and dereplication (which is the process that recognizes previously studied components that are not important for a screening of new ones) to then prepare for separation and isolation. To separate and isolate the mixtures into their constituents, several methods are used; HPLC is the simplest and can yield results in a short time without needing derivatization steps, although the results can be poor in resolution, and confusing. Ultra high-pressure liquid chromatography (UHPLC) is an improvement on HPLC by enhancing the resolution and throughput for rapid fingerprinting of crude extracts. Liquid chromatography coupled to a photo diode array (LC-PDA) detector is another add-on to a HPLC by allowing a view of the UV spectra, which is useful for detecting compounds with characteristic chromophores. HPLC-MS is HPLC that is coupled to a mass spectrometer, aiding detection, quantification, and identification by providing at the same time a chromatographic (retention times) and a mass spectrometric (m/z) dimension. HPLC-NMR is one of the strongest HPLC methods used to separate compounds. It has the advantage of not relying on commercial databases for spectral comparison, like HPLC-MS. HPLC-NMR provides structural information or even stereochemical information, as well as detection of any hydrogen-containing compounds. LC-SPE-NMR uses a solid-phase extraction coupled to a HPLC and finally a NMR detector, and allows the NMR detection after HPLC separation by either trapping the peaks on SPE or by HPLC microfractionation, drying, and reinjection of the concentrated peak in a microflow capillary LC-NMR probe. Microflow NMR and cryogenized probes are derivations of this technique (Cechinel-Filho 2012).

It is incontestable that medicinal plants provide unlimited opportunities for new drug discovery because of the unmatched availability of chemical diversity. Nevertheless, since bioactive phytochemicals occurring in plant materials consist of multicomponent mixtures, their extraction, separation, and isolation still create problems. In fact, extraction techniques can negatively affect the integrity of active principles, and practically all of them have to be purified by the combination of several chromatographic techniques or various other purification methods. Thus, it is expected that improvements in these methods will allow us to overcome some of the current limitations, as well as driving the development and introduction of new technologies.

9.4.2 Common Formulations in Drugs from Plant Origin

Drug development has evolved steadily since it first began as part of traditional medicine, and today more and more plant compounds are used as precursors, prototypes, and probes in drug production (Ramawat & Mérillon 2008). Depicted in Table 9.3 are

Table 9.3 Drugs derived from natural products.

| Plant of origin before modification | Used part | Active principle | Application | Reference |
|--|---------------|-----------------------------|--|--|
| <i>Artemisia annua</i> L. | Aerial parts | Artemisinin | Antimalarial | Phillipson 2007 |
| <i>Atropa belladonna</i> L. | Aerial parts | Tiotropium | Chronic obstructive pulmonary disease | Balunas & Kinghorn 2005 |
| | Aerial parts | Atropine | Mydriatic agent, antispasmodic | Ramawat & Mérillon 2008 |
| <i>Betula</i> spp. L. | Bark | Betulinic acid | Melanoma, anticancer, antimalarial, anti-HIV, anthelmintic, antiinflammatory, antiretroviral | Balunas & Kinghorn 2005; Ramawat & Mérillon 2008 |
| <i>Callistemon citrinus</i> Curtis | Aerial parts | Nitisinone | Tyrosinemia | Balunas & Kinghorn 2005 |
| <i>Calophyllum lanigerum</i> W. | Aerial parts | Calanolide | Anti-HIV | – |
| <i>Camptotheca acuminata</i> Decne | Bark and stem | Camptothecin | Anticancer | Phillipson 2007 |
| | Bark and stem | Topotecan | Metastatic ovarian cancer | – |
| | Bark and stem | Irinotecan | Colorectal cancer | – |
| | Bark and stem | Exatecan | Anticancer agent | Balunas & Kinghorn 2005 |
| <i>Capsicum</i> spp. L. | Fruit | Capsaicin | Osteoarthritis, psoriasis, diabetic, neuropathy | Ramawat & Mérillon 2008 |
| <i>Catharanthus roseus</i> L. | Aerial parts | Vindesine | Leukemia and lung cancer | Phillipson 2007 |
| | Aerial parts | Vinorelbine | Breast cancer | – |
| | Aerial parts | Vinflunine | Anticancer agent | Balunas & Kinghorn 2005 |
| <i>Chondrodendron tomentosum</i> Ruiz & Pavón | Aerial parts | Tubocurarine | Neuromuscular blocking agent | Phillipson 2007 |
| <i>Combretum cafrum</i> (Eckl. & Zeyh.) Kuntze | Aerial parts | Combretastatin A4 phosphate | Anaplastic thyroid cancer | Ramawat & Mérillon 2008 |
| <i>Dioscorea</i> genus | Tubers | Diosgenin | Contraceptive | – |
| <i>Erythroxylum pervillei</i> Bail | Stem bark | Pervilleine A | Epidermoid cancer | Balunas & Kinghorn 2005 |

| | | | | |
|--|-------------------|---------------------------------|--|-------------------------|
| <i>Euphorbia peplos</i> L. | Sap | Ingenol 3-angelate | Skin conditions | Ramawat & Mérillon 2008 |
| <i>Galanthus woronowii</i> Losinsk. | Bulbs and flowers | Galantamine | Alzheimer's | Balunas & Kinghorn 2005 |
| <i>Galega officinalis</i> L. | Aerial parts | Guanidine derivatives | Type 2 diabetes | Ramawat & Mérillon 2008 |
| <i>Glycine max</i> L. Merrill | Aerial parts | Phenoxadiol | Cervical, ovarian, prostate, renal, and vaginal cancer | – |
| <i>Huperzia serrata</i> Thunb. (Ex Murray) | Aerial parts | Huperizine | Alzheimer's | – |
| <i>Illicium verum</i> Hoof f. | Fruit | Oseltamivir phosphate (Tamiflu) | Influenza | – |
| <i>Panax ginseng</i> L. | Aerial parts | Protopanaxadiol | Apoptotic effect in cancer cells | – |
| <i>Papaver somniferum</i> L. | Seed pods | M6G | Pain medication | Balunas & Kinghorn 2005 |
| | Seed pods | Apokyn | Parkinson's | Ramawat & Mérillon 2008 |
| <i>Podophyllum peltatum</i> L. | Root | Etoposide | Small cell lung cancer, lymphomas, testicular cancer | Phillipson 2007 |
| | Root | Teniposide | Brain tumors | – |
| <i>Physostigma venenosum</i> Balf. | Seed | Physostigmine | Parkinson's | – |
| <i>Plectranthus barbatus</i> Andrews | Aerial parts | Colforsin daropate | Anticancer | Butler 2005 |
| <i>Taxus brevifolia</i> Nutt. | Bark | Taxol | Anticancer chemotherapy | Phillipson 2007 |
| | Bark | Taxotere | Breast cancer and nonsmall cell lung cancer (adjuvant) | – |

some of the most important drugs either developed using compounds derived from plants, or synthetic ones that were inspired by them, along with the plant from which they were first isolated and the illnesses they are used for. The recent change of attitude from big pharmaceutical companies, which are starting to look for natural compounds, has been a major tonic in the industry, helping to develop new drugs. The applications of natural compounds for human health are endless, and considering that currently only one-quarter of flowering plants is used, there is hope of finding treatments and solutions for many patients around the world (Lange 2004).

The WHO reports that over 21 000 plant taxa are used for medicinal purposes, although this number does not include cosmetics, spirits, and aromas (FAO 2002; Lange 2004). Roughly 80% of developing countries depend on plant-based drugs, although the WHO suggests that in the near future a similar percentage of the entire world population will depend on them. Furthermore, 30% of the drugs sold worldwide contain products derived from plants (FAO 2005).

Of the global trade in medicinal plants, it is hard to know how much is represented by wild or cultivated ones. Although the pharmaceutical industry has isolated a large number of bioactive compounds from wild plants (edible and medicinal), there are considerable disadvantages in harvesting wild medicinal plants rather than cultivating them for industrial drug development. The pharmaceutical industry mainly uses cultivated plants as primary material, despite the expensive domestication and cultivation process, in order to obtain a standard and well-known source of the active principle, in the necessary amounts for industrial-level processing. Moreover, there are some disadvantages related to wild plant gathering, including uncontrolled harvest that leads to extinction of the plant and erosion of the ecosystem. Other problems include poor knowledge about the biology of the plants, little or no inventory, ownership conflicts of the harvest zones, and scarce income due to overharvesting. Cultivation in small farms and households or in large and extensive production facilities could be an alternative, although the disadvantages are still great, due to the large investments needed, the reduction of incentive to conserve native ecosystems, devaluation of wild plants, reduction of genetic diversity and the risk of the introduced plant becoming an invasive species (FAO 2002).

9.4.3 Wild Plant-Based Drugs for Different Therapeutic Targets

Wild plant-based drugs are everywhere; the definition of a drug is quite vague, encompassing all “chemical substances used in the treatment, cure, prevention, or diagnosis of disease or used to otherwise enhance physical or mental well-being.” In this way, all molecules used by any type of medicine, modern or traditional, could be classed as drugs. To narrow down the results, only drugs used and approved in Western modern medicine are considered here, otherwise the list would be endless, although alternative medicines are quite well documented (Ahmad *et al.* 2006; Hawkins 2008; Osbourn & Lanzotti 2009; Trivedi 2009).

Medicinal plants represent 25% of prescription drugs in modern medicine. Of the 3000 plants traded for medicinal purposes, only 900 are cultivated, which means that 70–80% of the whole market depends on wild collection (Hawkins 2008). The conservation of habitats of these plants is the responsibility of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES), which

is the most important source of information on wild medicinal plants in use. *Galanthus* spp. L., an herbaceous plant endemic to the northern hemisphere, is the source of galanthamine, an approved drug used against Alzheimer's disease (Heinrich & Teoh 2004). *Taxus brevifolia* Nutt., the conifer that is the source of the anticancer agent Taxol®, also known as paclitaxel, is another success story of the medicinal power of plants, although it has endangered some cultivars of the tree. The alkaloid colchicine, derived from *Colchicum autumnale* L. corms, is used for the treatment of gout, under the name Colcrys (Romano 2013). The treatment of cardiac diseases also depends on compounds derived from wild medicinal plants, including digoxin, a cardiac glycoside extracted from the herb *Digitalis lanata* Ehrh. It is also marketed under the names Lanoxin®, Lanoxicaps®, Cardoxin® and Digitek®, among others (Hawkins 2008). The cinchona tree, *Cinchona officinalis* L., endemic to South America, is a natural source of quinine, a known antimalarial alkaloid that is used against this disease in modern medicine. There are reports of other uses of this molecule, which have recently been investigated (Christoforidis 2014). *Camptotheca acuminata* Decne is a tree native to China and Tibet which is rich in an alkaloid called camptothecin, used as an anticancer agent (Gaur *et al.* 2014). These examples illustrate some of the illnesses that can be cured or attenuated with wild plant compounds.

The endless combination of compounds found in nature that may have application in medicine provides hope for treatments of illnesses that have not yet been controlled or cured. The search to find new compounds continues at a steady pace and technology keeps lending precious help to this quest. Wild medicinal plants are today still as valuable as they were in the pre-modern medicine era. However, the pursuit of bioactive compounds should never overlook the habitats and wellbeing of the species. Research should continue to try and cultivate the plants that are not yet fit to be intensively grown, therefore reducing dependency on wild plants. But while there is no alternative, mankind should harvest them from nature, but always ensuring their continuity for generations to come.

9.5 Conclusion

Functional foods and nutraceuticals have been reported as one of the top trends of today's food industry. Apart from the naturally occurring functional foods, the development of new functional foods, nutraceuticals, and drugs based on plants is an active and very promising area of research, indispensable for the substantiation of health claims and benefits. The characterization of plant ingredients by advanced technologies, standardization of human clinical trials, and the use of emerging methodologies are crucial strategies for the development of new functional products and drugs. Additionally, the degree of acceptance and awareness of functional foods and nutraceuticals by consumers, the association between manufacturers and academic researchers, and the effects of new regulations for nutrition and health claims are crucial factors for future market evolution. Despite all the potential of these products to prevent diseases and promote human health, health professionals, nutritionists, and regulatory toxicologists should work together to plan appropriate regulation to provide the ultimate health and therapeutic benefit to humans.

However, due to the rising demand for plant-based functional foods, nutraceuticals, and drugs in higher quantities to promote health, longevity, and quality of life, wild harvested medicinal plants are taking on an increasing role and many of them have become endangered due to irresponsible collection, associated with economic interests. Therefore, the cultivation of these species is an alternative that needs to be taken into account. Furthermore, the next phase of market growth depends on valid scientific research for new product technologies, patents, more effective branding, and trade-mark strategies in product manufacture and international regulatory compliance.

References

- Adão, C. R., Silva, B. P. & Parente, J. P. (2011) A new steroidal saponin with antiinflammatory and antiulcerogenic properties from the bulbs of *Allium ampeloprasum* var. *porrum*. *Fitoterapia* **82**, 1175–1180.
- Ahmad, I., Aqil, F., Owais, M., eds. (2006) *Modern Phytomedicine: Turning Plants into Drugs*. Weinheim: Wiley-Vch.
- Amagase, H. & Farnsworth, N. R. (2011) A review of botanical characteristics, phytochemistry, clinical relevance in efficacy and safety of *Lycium barbarum* fruit (Goji). *Food Research International* **44**, 1702–1717.
- Andlauer, W. & Furst, P. (2002) Nutraceuticals: a piece of history, present status and outlook. *Food Research International* **35**, 171–176.
- Annunziata, A. & Vecchio, R. (2011) Functional foods development in the European market: a consumer perspective. *Journal of Functional Foods* **3**, 223–228.
- Ashwell, M. (2003) *Concepts of Functional Foods*. ILSI Europe Concise Monograph Series. Brussels: ILSI Europe.
- Avbunudiogba, J. A., Alalor, C. A., Builders, P. F., *et al.* (2013) Development and evaluation of liquid oral phytoformulation of *Phyllanthus amarus*. *Journal of Pharmacy Research* **6**, 908–912.
- Bagchi, D., ed. (2014) *Nutraceutical and Functional Food Regulations in the United States and Around the World*, 2nd edn. Houston: Academic Press.
- Balunas, M. J. & Kinghorn, A. D. (2005) Drug discovery from medicinal plants. *Life Sciences* **78**, 431–441.
- Barolo, M. I., Mostacero, N. R. & López, S. N. (2014) *Ficus carica* L. (Moraceae): an ancient source of food and health. *Food Chemistry* **164**, 119–127.
- Barros, L., Carvalho, A. M., Sa Morais, J., *et al.* (2010) Strawberry-tree, blackthorn and rose fruits: detailed characterisation in nutrients and phytochemicals with antioxidant properties. *Food Chemistry* **120**, 247–254.
- Barros, L., Cabrita, L., Vilas Boas, M., *et al.* (2011a) Chemical, biochemical and electrochemical assays to evaluate phytochemicals and antioxidant activity of wild plants. *Food Chemistry* **127**, 1600–1608.
- Barros, L., Dueñas, M., Ferreira, I. C. F. R., *et al.* (2011b) Use of HPLC–DAD–ESI/MS to profile phenolic compounds in edible wild greens from Portugal. *Food Chemistry* **127**, 169–173.
- Bäurle, P., Suter, A. & Wormstall, H. (2009) Safety and effectiveness of a traditional ginkgo fresh plant extract – results from a clinical trial. *Forschende Komplementärmedizin* **16**, 156–161.

- Bech-Larsen, T. & Scholderer, J. (2007) Functional foods in Europe: consumer research, market experiences and regulatory aspects. *Trends in Food Science and Technology* **18**, 231–234.
- Bernal, J., Mendiola, J. A., Ibáñez, E., *et al.* (2011) Advanced analysis of nutraceuticals. *Journal of Pharmaceutical and Biomedical Analysis* **55**, 758–774.
- Betoret, E., Betoret, N., Vidal, D. *et al.* (2011) Functional foods development: trends and technologies. *Trends in Food Science and Technology* **22**, 498–508.
- Bhardwaj, Y. R., Pareek, A., Jain, V., *et al.* (2014) Chemical delivery systems and soft drugs: retrometabolic approaches to drug design. *Saudi Pharmaceutical Journal* **22**, 290–302.
- Bhoopat, L., Srichairatanakool, S., Kanjanapothi, D., *et al.* (2011) Hepatoprotective effects of lychee (*Litchi chinensis* Sonn.): a combination of antioxidant and anti-apoptotic activities. *Journal of Ethnopharmacology* **136**, 55–66.
- Bowen-Forbes, C. S., Zhang, Y. & Nair, M. G. (2010) Anthocyanin content, antioxidant, anti-inflammatory and anticancer properties of blackberry and raspberry fruits. *Journal of Food Composition and Analysis* **23**, 554–560.
- Braithwaite, M. C., Tyagi, C., Tomar, L. K., *et al.* (2014) Nutraceutical-based therapeutics and formulation strategies augmenting their efficiency to complement modern medicine: an overview. *Journal of Functional Foods* **6**, 82–99.
- Brinkeborn, R. M., Shah, D. V. & Degenring F. H. (1989) Echinaforce® and other *Echinacea* fresh plant preparations in the treatment of the common cold. A randomized, placebo controlled, double-blind clinical trial. *Phytomedicine* **6**, 1–5.
- Butler, M. S. (2005) Natural products to drugs: natural product derived compounds in clinical trials. *Natural Products Reports* **22**, 162–195.
- Byrne, D. (2003) Health nutrition and labeling. *Food Science and Technology* **17**, 26–28.
- Caleja, C., Barros, L., Antonio, A. L., *et al.* (2015) *Foeniculum vulgare* Mill. as natural conservation enhancer and health promoter by incorporation in cottage cheese. *Journal of Functional Foods* **12**, 428–438.
- Carocho, M. & Ferreira, I. C. F. R. (2013a) The role of phenolic compounds in the fight against cancer – a review. *Anti-Cancer Agents in Medicinal Chemistry* **13**, 1236–1238.
- Carocho, M. & Ferreira, I. C. F. R. (2013b) A review on antioxidants, prooxidants and related controversy: natural and synthetic compounds, screening and analysis methodologies and future perspectives. *Food and Chemical Toxicology* **51**, 15–25.
- Carocho, M., Barreira, J. C. M., Antonio, A. L., *et al.* (2015a) The incorporation of plant materials in “Serra da Estrela” cheese improves antioxidant activity without changing the fatty acid profile and visual appearance. *European Journal of Lipid Science and Technology* **117**, 1607–1614.
- Carocho, M., Barreira, J. C. M., Barros, L., *et al.* (2015b) Traditional pastry with chestnut flowers as natural ingredients: an approach of the effects on nutritional value and chemical composition. *Journal of Food Composition and Analysis* **44**, 93–101.
- Cechinel-Filho, V., ed. (2012) *Plant Bioactivities and Drug Discovery: Principle, Practice, and Perspectives*. New Jersey: John Wiley.
- Cerqueira, M. A., Pinheiro, A. C., Silva, H. D., *et al.* (2014) Design of bio-nanosystems for oral delivery of functional compounds. *Food Engineering Reviews* **6**, 1–19.
- Chen, L., Remondetto, G. E. & Subirade, M. (2006) Food protein-based materials as nutraceutical delivery systems. *Trends in Food Science and Technology* **17**, 272–283.
- Christoforidis, J. (2014) Quinine. Reference module in biomedical sciences. In: P. Wexler, ed. *Encyclopedia of Toxicology*, 3rd edn. Philadelphia: Elsevier, pp 19–22.

- Coppens, P., Silva, M. F. & Pettman, S. (2006) European regulations on nutraceuticals, dietary supplements and functional foods: a framework based on safety. *Toxicology* **221**, 59–74.
- Costa, A. G. V., Garcia-Diaz, D. F., Jimenez, P. *et al.* (2013) Bioactive compounds and health benefits of exotic tropical red-black berries. *Journal of Functional Foods* **5**, 539–549.
- Dapas, B., Dall'Acqua, S., Bullac, R., *et al.* (2014) Immunomodulation mediated by a herbal syrup containing a standardized Echinacea root extract: a pilot study in healthy human subjects on cytokine gene expression. *Phytomedicine* **21**, 1406–1410.
- DeFelice, S. L. (1992) The nutraceutical initiative: a recommendation for U.S. economic and regulatory reforms. *Genetic Engineering News* **12**, 13–15.
- Delva, L. & Goodrich-Schneider, R. (2013) Antioxidant activity and antimicrobial properties of phenolic extracts from acerola (*Malpighia emarginata* DC) fruit. *International Journal of Food Science and Technology* **48**, 1048–1056.
- Dias, F. M., Leffa, D. D., Daumann, F., *et al.* (2014) Acerola (*Malpighia emarginata* DC.) juice intake protects against alterations to proteins involved in inflammatory and lipolysis pathways in the adipose tissue of obese mice fed a cafeteria diet. *Lipids in Health and Disease* **13**, 1–9.
- Dias, M. I., Ferreira, I. C. F. R. & Barreiro, M. F. (2015) Microencapsulation of bioactives for food applications. *Food and Function* **6**, 1035–1052.
- Diplock, A., Aggett, P. J., Ashwell, M., *et al.* (1999) Scientific concepts of functional foods in Europe: consensus document. *British Journal of Nutrition* **81**, 1–27.
- Espín, J. C., García-Conesa, M. & Tomás-Barberán, F. (2007) Nutraceuticals: facts and fiction. *Phytochemistry* **68**, 2986–3008.
- Ezhilarisi, P. N., Karthik, P., Chhanwal, N., *et al.* (2013) Nanoencapsulation techniques for food bioactive components: a review. *Food Bioprocess Technology* **6**, 628–647.
- Fabricant, D. S. & Farnsworth, N. R. (2001) The value of plants used in traditional medicine for drug discovery. *Environmental Health Perspectives* **109**, 69–75.
- FAO (2002) *Impact of Cultivation and Gathering of Medicinal Plants on Biodiversity: Global Trends and Issues*. Available at: <ftp://ftp.fao.org/docrep/fao/005/aa010e/AA010E00.pdf> (accessed 27 June 2016).
- FAO (2005) *Trade in Medicinal Plants*. Raw Materials, Tropical and Horticultural Products Service Commodities and Trade Division. Available at: <ftp://ftp.fao.org/docrep/fao/008/af285e/af285e00.pdf> (accessed 27 June 2016).
- FDA (2014) FDA's Drug Review Process: Continued. Available at: www.fda.gov/Drugs/ResourcesForYou/Consumers/ucm289601.htm (accessed 27 June 2016).
- Flores, G., Wu, S. B., Negrin, A., *et al.* (2015) Chemical composition and antioxidant activity of seven cultivars of guava (*Psidium guajava*) fruits. *Food Chemistry* **170**, 327–335.
- Fracassetti, D., Costa, C., Moulay, L., *et al.* (2013) Ellagic acid derivatives, ellagitannins, proanthocyanidins and other phenolics, vitamin C and antioxidant capacity of two powder products from camu-camu fruit (*Myrciaria dubia*). *Food Chemistry* **139**, 578–588.
- García-Herrera, P., Morales, P., Fernández-Ruiz, V., *et al.* (2014) Nutrients, phytochemicals and antioxidant activity in wild populations of *Allium ampeloprasum* L., a valuable underutilized vegetable. *Food Research International* **62**, 272–279.
- Garcia-Rios, A., Delgado-Lista, J., Alcala-Diaz, J. F., *et al.* (2013) Nutraceuticals and coronary heart disease. *Current Opinion in Cardiology* **28**, 475–482.

- Gaur, S., Wang, Y., Kretzner, L., *et al.* (2014) Pharmacodynamic and pharmacogenomics study of the nanoparticle conjugate of camptothecin CRLX10 for the treatment of cancer. *Nanomedicine: Nanotechnology, Biology, and Medicine* **10**, 1477–1486.
- Genta, S., Cabrera, W., Habib, N., *et al.* (2009) Yacon syrup: beneficial effects on obesity and insulin resistance in humans. *Clinical Nutrition* **28**, 182–187.
- Giordano, P., Scicchitano, P., Locorotondo, M., *et al.* (2012) Carotenoids and cardiovascular risk. *Current Pharmaceutical Design* **18**, 5577–5589.
- Graça, C., Freitas, C. S., Baggio, C.H., *et al.* (2007) *Mikania laevigata* syrup does not induce side effects on reproductive system of male Wistar rats. *Journal of Ethnopharmacology* **111**, 29–32.
- Granato, D., Branco, G. F. & Nazzaro, F. (2010) Functional foods and nondairy probiotic food development: trends, concepts and products. *Comprehensive Reviews in Food Science and Food Safety* **9**, 292–302.
- Grivetti, L. E. & Ogle, B. M. (2000) Value of traditional foods in meeting macro- and micronutrient needs: the wild plant connection. *Nutrition Research Reviews* **13**, 31–46.
- Groot, R. S., Wilson, M. A. & Boumans, R. M. J. (2002) A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecological Economics* **41**, 393–408.
- Guimarães, R., Barros, L., Dueñas, M., *et al.* (2013) Characterisation of phenolic compounds in wild fruits from Northeastern Portugal. *Food Chemistry* **141**, 3721–3730.
- Guimarães, R., Barros, L., Calhelha, R. C., *et al.* (2014) Bioactivity of different enriched phenolic extracts of wild fruits from Northeastern Portugal: a comparative study. *Plant Foods for Human Nutrition* **69**, 37–42.
- Gulati, O. P. & Ottaway, P. B. (2006) Legislation relating to nutraceuticals in the European Union with a particular focus on botanical-sourced products. *Toxicology* **221**, 75–87.
- Gutierrez-Orozco, F. & Failla, M. L. (2013) Biological activities and bioavailability of mangosteen xanthones: a critical review of the current evidence. *Nutrients* **5**, 3163–3183.
- Hasler, C. M. (2000) The changing face of functional foods. *Journal of the American College of Nutrition* **19**, 499–506.
- Hasler, C. M. (2002) Functional foods: benefits, concerns and challenges – a position paper from the American Council on Science and Health. *Journal of Nutrition* **132**, 3772–3781.
- Hawkins, B., ed. (2008) *Plants for Life: Medicinal Plant Conservation and Botanic Gardens*. London: Botanic Gardens Conservation International.
- Heinrich, M. & Teoh, H. L. (2004) Galanthamine from snowdrop – the development of a modern drug against Alzheimer's disease from local Caucasian knowledge. *Journal of Ethnopharmacology* **92**, 147–162.
- Heywood, V. H. (2011) Ethnopharmacology, food production, nutrition and biodiversity conservation: towards a sustainable future for indigenous peoples. *Journal of Ethnopharmacology* **137**, 1–15.
- Hosseini, S., Jamshidi, L., Mehrzadi, S., *et al.* (2014) Effects of *Juglans regia* L. leaf extract on hyperglycemia and lipid profiles in type two diabetic patients: a randomized double-blind, placebo-controlled clinical trial. *Journal of Ethnopharmacology* **152**, 451–456.
- Howlett, J. (2008) *Functional Foods: From Science to Health and Claims*. Brussels: ILSI Europe.
- Huang, G. J., Wang, B. S., Lin, W. C., *et al.* (2012) Antioxidant and anti-inflammatory properties of longan (*Dimocarpus longan* Lour.) pericarp. *Evidence-Based Complementary and Alternative Medicine* **2012**, 1–10.

- Huang, Q., Yu, H. & Ru, Q. (2010b) Bioavailability and delivery of nutraceuticals using nanotechnology. *Journal of Food Science* **75**, 50–57.
- Huang, W. Y., Cai, Y. Z., Corke, H., *et al.* (2010a) Survey of antioxidant capacity and nutritional quality of selected edible and medicinal fruit plants in Hong Kong. *Journal of Food Composition and Analysis* **23**, 510–517.
- Imaga, N. A., Chukwu, C. E., Blankson, A., *et al.* (2013) Biochemical assessment of Ciklavit®, a nutraceutical used in sickle cell anemia management. *Journal of Herbal Medicine* **3**, 137–148.
- Ismail, T., Sestili, P. & Akhtar, S. (2012) Pomegranate peel and fruit extracts: a review of potential anti-inflammatory and anti-infective effects. *Journal of Ethnopharmacology* **143**, 397–405.
- Izzo, R., Simone, G., Giudice, R., *et al.* (2010) Effects of nutraceuticals on prevalence of metabolic syndrome and on calculated Framingham risk score in individuals with dyslipidaemia. *Journal of Hypertension* **28**, 1482–1487.
- Janaswamy, S. & Youngren, S. R. (2012) Hydrocolloid-based nutraceutical delivery systems. *Food and Function* **3**, 503–507.
- Kang, J., Xie, C., Li, Z., *et al.* (2011) Flavonoids from acai (*Euterpe oleracea* Mart.) pulp and their antioxidant and anti-inflammatory activities. *Food Chemistry* **128**, 152–157.
- Katiyar, C., Gupta, A., Kanjilal, S., *et al.* (2012) Drug discovery from plant sources: an integrated approach. *AYU* **33**, 10–19.
- Kelly-Pieper, K., Patil, S. P., Busse, P., *et al.* (2009) Safety and tolerability of an antiasthma herbal formula (ASHMITM) in adult subjects with asthma: a randomized, double-blinded, placebo-controlled, dose-escalation phase I study. *Journal of Alternative and Complementary Medicine* **15**, 735–743.
- Khoo, C. & Falk, M. (2014) Cranberry polyphenols: effects on cardiovascular risk factors. In: R. R. Watson, V. R. Preedy & S. Zibadi, eds. *Polyphenols in Human Health and Disease*. San Diego: Academic Press, pp 1049–1065.
- Kianbakht, S. & Dabaghian, F. H. (2013) Improved glycemic control and lipid profile in hyperlipidemic type 2 diabetic patients consuming *Salvia officinalis* L. leaf extract: a randomized placebo controlled clinical trial. *Complementary Therapies in Medicine* **21**, 441–446.
- Kim, J., Lee, K. W. & Lee, H. J. (2011) Cocoa (*Theobroma cacao*) seeds and phytochemicals in human health. In: V. R. Preedy, R. R. Watson & V. B. Patel, eds. *Nuts and Seeds in Health and Disease Prevention*. Philadelphia: Elsevier, pp 351–360.
- Kwak, N. S. & Jukes, D. J. (2001) Functional foods. Part 2: The impact of current regulatory terminology. *Food Control* **12**, 109–117.
- Lai, T. N. H., Andre, C., Rogez, H., *et al.* (2015) Nutritional composition and antioxidant properties of the sim fruit (*Rhodomyrtus tomentosa*). *Food Chemistry* **168**, 410–416.
- Lange, D. (2004) Medicinal and aromatic plants: trade, production, and management of botanical resources. *Acta Horticulturae* **629**, 177–197.
- Leite, A. V., Malta, L. G., Riccio, M. F., *et al.* (2011) Antioxidant potential of rat plasma by administration of freeze-dried jaboticaba peel (*Myrciaria jaboticaba* Vell Berg). *Journal of Agricultural and Food Chemistry* **59**, 2277–2283.
- Lenoir, S., Degenring, F. H. & Saller, R. (1999) A double-blind randomised trial to investigate three different concentrations of a standardised fresh plant extract obtained from the shoot tips of *Hypericum perforatum* L. *Phytomedicine* **6**, 141–146.

- Lenzi, R. M., Campestrini, L. H., Okumura, L. M., *et al.* (2013) Effects of aqueous fractions of *Uncaria tomentosa* (Willd.) D.C. on macrophage modulatory activities. *Food Research International* **53**, 767–779.
- Li, A. N., Li, S., Li, H. B., *et al.* (2014a) Total phenolic contents and antioxidant capacities of 51 edible and wild flowers. *Journal of Functional Foods* **6**, 319–330.
- Li, X., Zhao, J., Yang, M., *et al.* (2014c) Physalins and withanolides from the fruits of *Physalis alkekengi* L. var. *franchetii* (Mast.) Makino and the inhibitory activities against human tumor cells. *Phytochemistry Letters* **10**, 95–100.
- Li, Y., Ahmad, A., Kong, D., *et al.* (2014b) Recent progress on nutraceutical research in prostate cancer. *Cancer and Metastasis Review* **33**, 629–640.
- Liu, Y., Singh, D. & Nair, M. G. (2012) Pods of Khejri (*Prosopis cineraria*) consumed as a vegetable showed functional food properties. *Journal of Functional Foods* **4**, 116–121.
- Luis, G., Rubio, C., Gutiérrez, A. J., *et al.* (2012) Palm tree syrup: nutritional composition of a natural edulcorant. *Nutrición Hospitalaria* **27**, 548–552.
- Lv, Q., Si, M., Yan, Y., *et al.* (2014) Effects of phenolic-rich litchi (*Litchi chinensis* Sonn.) pulp extracts on glucose consumption in human HepG2 cells. *Journal of Functional Foods* **7**, 621–629.
- Madhavi, D. L., Bomser, J., Smith, M. A. L., *et al.* (1998) Isolation of bioactive constituents from *Vaccinium myrtillus* (bilberry) fruits and cell cultures. *Plant Science* **131**, 95–103.
- Malafaia, C. R. A., Silva, B. P., Tinoco, L. W., *et al.* (2015) Structural characterization and gastroprotective property of a novel glucofructan from *Allium ampeloprasum* var. *porrum*. *Carbohydrate Research* **402**, 44–49.
- Manchali, S., Murthy, K. N. C. & Patil, B. S. (2012) Crucial facts about health benefits of popular cruciferous vegetables. *Journal of Functional Foods* **4**, 94–106.
- Mannarino, M. R., Ministrini, S. & Pirro, M. (2014) Nutraceuticals for the treatment of hypercholesterolemia. *European Journal of Internal Medicine* **25**, 592–599.
- Martins, A., Barros, L., Carvalho, A. M., *et al.* (2014) Phenolic extracts of *Rubus ulmifolius* Schott flowers: characterization, microencapsulation and incorporation into yogurts as nutraceutical sources. *Food and Function* **5**, 1091–1100.
- Martins, D., Barros, L., Carvalho, A. M., *et al.* (2011) Nutritional and *in vitro* antioxidant properties of edible wild greens in Iberian Peninsula traditional diet. *Food Chemistry* **125**, 488–494.
- Matthias, A., Addison, R. S., Agnew, L. L., *et al.* (2007) Comparison of Echinacea alkylamide pharmacokinetics between liquid and tablet preparations. *Phytomedicine* **14**, 587–590.
- McAlindon, T.E. (2006) Nutraceuticals: do they work and when should we use them? *Clinical Rheumatology* **20**, 99–115.
- McCook-Russell, K. P., Nair, M. G., Facey, P. C., *et al.* (2012) Nutritional and nutraceutical comparison of Jamaican *Psidium cattleianum* (strawberry guava) and *Psidium guajava* (common guava) fruits. *Food Chemistry* **134**, 1069–1073.
- McNamara, S. H. (1997) Dietary supplement legislation enhances opportunities to market nutraceutical-type products. *Journal of Nutraceuticals, Functional and Medical Foods* **1**, 47–59.
- Mezadri, T., Villaño, D., Fernández-Pachón, M. S., *et al.* (2008) Antioxidant compounds and antioxidant activity in acerola (*Malpighia emarginata* DC.) fruits and derivatives. *Journal of Food Composition and Analysis* **21**, 282–290.

- Milesi, M., Lacan, D., Brosse, H., *et al.* (2009) Effect of an oral supplementation with a proprietary melon juice concentrate (Extramel®) on stress and fatigue in healthy people: a pilot, double-blind, placebo-controlled clinical trial. *Nutrition Journal* **8**, 1–7.
- Milner, J. A. (2000) Functional foods: the US perspective. *American Journal of Clinical Nutrition*, **71**, 1654–1659.
- MMWR (1999) Achievements in public health, 1900–1999: safer and healthier foods. *Morbidity and Mortality Weekly Report*, **48**, 905–913.
- Mojani, M. S., Ghasemzadeh, A., Rahmat, A., *et al.* (2014) Assessment of bioactive compounds, nutritional composition and antioxidant activity of Malaysian young ginger (*Zingiber officinale* Roscoe). *International Food Research Journal* **21**, 1931–1935.
- Morales, P., Carvalho, A. M., Sánchez-Mata, M. C., *et al.* (2012) Tocopherol composition and antioxidant activity of Spanish wild vegetables. *Genetic Resources and Crop Evolution* **59**, 851–863.
- Najda, A., Dyduch-Siemńska, M., Dyduch, J., *et al.* (2014) Comparative analysis of secondary metabolites contents in *Fragaria vesca* L. fruits. *Annals of Agricultural and Environmental Medicine* **21**, 339–343.
- Newman, D. J., Cragg, G. M. & Snader, K. M. (2000) The influence of natural products upon drug discovery. *Natural Product Reports* **17**, 215–234.
- Ninfali, P. & Angelino, D. (2013) Nutritional and functional potential of *Beta vulgaris cicla* and *rubra*. *Fitoterapia* **89**, 188–199.
- Nöthlings, U., Murphy, S. P., Wilkens, L. R., *et al.* (2007) Flavonols and pancreatic cancer risk – the multiethnic cohort study. *American Journal of Epidemiology* **166**, 924–931.
- Oben, J., Enonchong, E., Kothari, S., *et al.* (2009) *Phellodendron* and *Citrus* extracts benefit joint health in osteoarthritis patients: a pilot, double-blind, placebo-controlled study. *Nutrition Journal* **8**, 1–9.
- Ohama, H., Ikeda, H. & Moriyama, H. (2006) Health foods and foods with health claims in Japan. *Toxicology* **221**, 95–111.
- Ortuño, J., Covas, M. I., Farre, M., *et al.* (2010) Matrix effects on the bioavailability of resveratrol in humans. *Food Chemistry* **120**, 1123–1130.
- Osbourne, A. E. & Lanzotti, V., eds. (2009) *Plant-Derived Natural Products: Synthesis, Function and Application*. Berlin: Springer Science + Business Media, LLC.
- Palma, S., Luján, C., Llabot, J. M., *et al.* (2002) Design of *Peumus boldus* tablets by direct compression using a novel dry plant extract. *International Journal of Pharmaceutics* **233**, 191–198.
- Pardo de Santayana, M., Tardío, J., Blanco, E., *et al.* (2007) Traditional knowledge of wild edible plants used in the northwest of the Iberian Peninsula (Spain and Portugal): a comparative study. *Journal of Ethnobiology and Ethnomedicine* **3**, 1–11.
- Paul, S. M., Mytelka, D. S., Dunwiddie, C. T., *et al.* (2010) How to improve R&D productivity: the pharmaceutical industry's grand challenge. *Nature Reviews* **9**, 203–214.
- Pedraza-Chaverri, J., Cárdenas-Rodríguez, N., Orozco-Ibarra, M., *et al.* (2008) Medicinal properties of mangosteen (*Garcinia mangostana*). *Food and Chemical Toxicology* **46**, 3227–3239.
- Peng, K., Yang, L., Zhao, S., *et al.* (2013) Chemical constituents from the fruit of *Gardenia jasminoides* and their inhibitory effects on nitric oxide production. *Bioorganic and Medicinal Chemistry Letters* **23**, 1127–1131.

- Pereira, C., Barros, L., Carvalho, A. M., *et al.* (2011) Nutritional composition and bioactive properties of commonly consumed wild greens: potential sources for new trends in modern diets. *Food Research International* **44**, 2634–2640.
- Pereira, C., Calhelha, R. C., Barros, L., *et al.* (2013b) Antioxidant properties, anti-hepatocellular carcinoma activity and hepatotoxicity of artichoke, milkthistle and borututu. *Industrial Crops Production* **49**, 61–65.
- Pereira, C., Calhelha, R. C., Barros, L., *et al.* (2014) Synergisms in antioxidant and anti-hepatocellular carcinoma activities of artichoke, milk thistle and borututu syrups. *Industrial Crops and Products* **52**, 709–713.
- Pereira, E., Barros, L. & Ferreira, I. C. F. R. (2013a) Chemical characterization of *Ginkgo biloba* L. and antioxidant properties of its extracts and dietary supplements. *Industrial Crops and Products* **51**, 244–248.
- Phillipson, J. D. (2007) Phytochemistry and pharmacognosy. *Phytochemistry* **68**, 2960–2972.
- Pierro, F. D., Putignano, P., Villanova, N., *et al.* (2013) Preliminary study about the possible glycemic clinical advantage in using a fixed combination of *Berberis aristata* and *Silybum marianum* standardized extracts versus only *Berberis aristata* in patients with type 2 diabetes. *Clinical Pharmacology: Advances and Applications* **5**, 167–174.
- Puoci, F., Iemma, F., Spizzirri, U. G., *et al.* (2011) Antioxidant activity of a Mediterranean food product: “fig syrup”. *Nutrients* **3**, 317–329.
- Quispe, C., Petroll, K., Theoduloz, C., *et al.* (2014) Antioxidant effect and characterization of South American *Prosopis pods* syrup. *Food Research International* **56**, 174–181.
- Ramawat, K. G. & Mérillon, J., eds. (2008) *Bioactive Molecules and Medicinal Plants*. Berlin: Springer-Verlag.
- Rathee, S., Rathee, P., Rathee, D., *et al.* (2010) Phytochemical and pharmacological potential of Kair (*Capparis decidua*). *International Journal of Phytomedicine* **2**, 10–17.
- Ribeiro, A. B., Chisté, R. C., Freitas, M., *et al.* (2014) *Psidium cattleianum* fruit extracts are efficient *in vitro* scavengers of physiologically relevant reactive oxygen and nitrogen species. *Food Chemistry* **165**, 140–148.
- Roberfroid, M. B. (2007) Concepts and strategy of functional food science: the European perspective. *American Journal of Clinical Nutrition* **71**, 1660–1664.
- Romano, J. (2013) Therapeutic review: colchicine. *Journal of Exotic Pet Medicine* **22**, 405–408.
- Ross, S. (2000) Functional foods: the Food and Drug Administration perspective. *American Journal of Clinical Nutrition* **71**, 1735–1738.
- Rufino, M. S. M., Alves, R. E., Brito, E. S., *et al.* (2010) Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. *Food Chemistry* **121**, 996–1002.
- Sarker S. D. & Nahar, L., eds. (2012) *Natural Products Isolation*. New York: Humana Press, Springer.
- Schippmann, U., Leaman, D. J. & Cunningham, A. B. (2002) *Impact of Cultivation and Gathering of Medicinal Plants on Biodiversity: Global Trends and Issues*. FAO Biodiversity and the Ecosystem Approach in Agriculture, Forestry and Fisheries. Satellite event on the occasion of the Ninth Regular Session of the Commission on Genetic Resources for Food and Agriculture. Inter-Departmental Working Group on Biological Diversity for Food and Agriculture. Rome: FAO.
- Schulp, C. J. E., Thuiller, W. & Verburg, P. H. (2014) Wild food in Europe: a synthesis of knowledge and data of terrestrial wild food as an ecosystem service. *Ecological Economics* **105**, 292–305.

- Scicchitano, P., Cameli, M., Maiello, M., *et al.* (2014) Nutraceuticals and dyslipidaemia: beyond the common therapeutics. *Journal of Functional Food* **6**, 11–32.
- Sener, B. & Orhan, L. (2005) Discovery of drug candidates from some Turkish plants and conservation of biodiversity. *Pure and Applied Chemistry* **77**, 53–64.
- Sengupta, G., Hazra, A., Kundu, A., *et al.* (2011) Comparison of *Murraya koenigii*- and *Tribulus terrestris*-based oral formulation versus tamsulosin in the treatment of benign prostatic hyperplasia in men aged >50 years: a double-blind, double-dummy, randomized controlled trial. *Clinical Therapeutics* **33**, 1943–1952.
- Sengupta, K., Krishnaraju, A. V., Vishal, A. A., *et al.* (2010) Comparative efficacy and tolerability of 5-Loxin® and Aflapin® against osteoarthritis of the knee: a double blind, randomized, placebo controlled clinical study. *International Journal of Medicinal Sciences* **7**, 366–377.
- Shad, A. A., Ahmad, S., Ullah, R., *et al.* (2014) Phytochemical and biological activities of four wild medicinal plants. *Scientific World Journal* **2014**, 1–7.
- Sharma, B., Salunke, R., Balomajumder, C., *et al.* (2010) Anti-diabetic potential of alkaloid rich fraction from *Capparis decidua* on diabetic mice. *Journal of Ethnopharmacology*, **127**, 457–462.
- Shegokar, R. & Müller, R. H. (2010) Nanocrystals: industrially feasible multifunctional formulation technology for poorly soluble actives. *International Journal of Pharmaceutics* **399**, 129–139.
- Sidor, A. & Gramza-Michałowska, A. (2014) Advanced research on the antioxidant and health benefit of elderberry (*Sambucus nigra*) in food – a review. *Journal of Functional Foods* **18**, 941–958.
- Singh, A. P., Wilson, T., Kalk, A. J., *et al.* (2009) Isolation of specific cranberry flavonoids for biological activity assessment. *Food Chemistry* **116**, 963–968.
- Singh, J., & Sinha, S. (2012) Classification, regulatory acts and applications of nutraceuticals for health. *International Journal of Pharma and Bio Sciences* **2**, 177–187.
- Spernath, A. & Aserin, A. (2006) Microemulsions as carriers for drugs and nutraceuticals. *Advances in Colloid and Interface Science* **128-130**, 47–64.
- Stauss-Grabo, M., Atiyea, S., Warnke, A., *et al.* (2011) Observational study on the tolerability and safety of film-coated tablets containing ivy extract (Prospan® Cough Tablets) in the treatment of colds accompanied by coughing. *Phytomedicine* **18**, 433–436.
- Sticher, O. (2008) Natural product isolation. *Natural Products Reports* **25**, 517–554.
- Takachi, R., Inoue, M., Ishihara, J., *et al.* (2008) Fruit and vegetable intake and risk of total cancer and cardiovascular disease Japan public health center-based prospective study. *American Journal of Epidemiology* **167**, 59–70.
- Talbott, S. M., Talbott, J. A. & Pugh, M. (2013) Effect of *Magnolia officinalis* and *Phellodendron amurense* (Relora®) on cortisol and psychological mood state in moderately stressed subjects. *Journal of the International Society of Sports Nutrition* **10**, 2–6.
- Thomson, M., Al-Qattan, K. K., Al-Sawan, S. M., *et al.* (2002) The use of ginger (*Zingiber officinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. *Prostaglandins, Leukotrienes and Essential Fatty Acids* **67**, 475–478.
- Tijhuis, M. J., Jong, N., Pohjola, M. V., *et al.* (2012) State of the art in benefit-risk analysis: food and nutrition. *Food and Chemical Toxicology* **50**, 5–25.
- Trivedi, P. C., ed. (2009) *Medicinal Plants: Utilisation and Conservation*. Chaura Rasta: Aaviskar Publishers

- Tu, Y., Sun, W., Wan, Y., *et al.* (2013) Huangkui capsule, an extract from *Abelmoschus manihot* (L.) medic, ameliorates adriamycin-induced renal inflammation and glomerular injury via inhibiting p38MAPK signaling pathway activity in rats. *Journal of Ethnopharmacology* **147**, 311–320.
- Vulić, J. J., Čebović, T. N., Čanadanović-Brunet, J. M., *et al.* (2014) In vivo and in vitro antioxidant effects of beetroot pomace extracts. *Journal of Functional Foods* **6**, 168–175.
- Wagner, H. & Jurcic, K. (2002) Immunological studies of Revitonil®, a phytopharmaceutical containing *Echinacea purpurea* and *Glycyrrhiza glabra* root extract. *Phytomedicine* **9**, 390–397.
- Weathers, P. J. & Towler, M. J. (2014) Changes in key constituents of clonally propagated *Artemisia annua* L. during preparation of compressed leaf tablets for possible therapeutic use. *Industrial Crops and Products* **62**, 173–178.
- WHO (2003) *Diet, Nutrition and the Prevention of Chronic Diseases*. WHO Technical Report Series 916. Geneva: World Health Organization.
- Wong, C. K., Tse, P. S., Wong, E. L. Y., *et al.* (2004) Immunomodulatory effects of Yun Zhi and Danshen capsules in health subjects – a randomized, double-blind, placebo-controlled, crossover study. *International Immunopharmacology* **4**, 201–211.
- Wootton-Beard, P. C. & Ryan, L. (2011) A beetroot juice shot is a significant and convenient source of bioaccessible antioxidants. *Journal of Functional Foods* **3**, 329–334.
- Wrick, K. L. (2005) The impact of regulation on the business of nutraceuticals in the United States: yesterday, today, and tomorrow. In: C. M. Hasler, ed. *Regulation of Functional Foods and Nutraceuticals: A Global Perspective*. Ames: Wiley-Blackwell, pp 3–36.
- Wu, P., Ma, G., Li, N., *et al.* (2015) Investigation of *in vitro* and *in vivo* antioxidant activities of flavonoids rich extract from the berries of *Rhodomyrtus tomentosa* (Ait.) Hassk. *Food Chemistry* **173**, 194–202.
- Xu, J., Seo, A. Y., Vorobyeva, D. A., *et al.* (2010) Beneficial effects of a Q-ter® based nutritional mixture on functional performance, mitochondrial function, and oxidative stress in rats. *PLoS One* **5**, 1–10.
- Yakoot, M., Salem, A. & Helmy, S. (2013) Effect of Memo®, a natural formula combination, on Mini-Mental State Examination scores in patients with mild cognitive impairment. *Clinical Interventions in Aging* **8**, 975–981.
- Yang, B., Jiang, Y., Shi, J., *et al.* (2011) Extraction and pharmacological properties of bioactive compounds from longan (*Dimocarpus longan* Lour.) fruit – a review. *Food Research International* **44**, 1837–1842.
- Yu, L., Jiang, B. P., Luo, D., *et al.* (2012) Bioactive components in the fruits of *Ziziphus jujuba* Mill. against the inflammatory irritant action of *Euphorbia* plants. *Phytomedicine* **19**, 239–244.
- Zeisel, S. H. (1999) Regulation of nutraceuticals. *Science* **285**, 1853–1855.
- Zhang, C. R., Dissanayake, A. A., Kevseroğlu, K., *et al.* (2015) Evaluation of coriander spice as a functional food by using *in vitro* bioassays. *Food Chemistry* **167**, 24–29.
- Zia-Ul-Haq, M., Čavar, S., Qayum, M., *et al.* (2011) Compositional studies: antioxidant and antidiabetic activities of *Capparis decidua* (Forsk.) Edgew. *International Journal of Molecular Sciences* **12**, 8846–8861.