

Worldwide Prevalence and Incidence of Dementia

Laura Fratiglioni,^{1,2} Diana De Ronchi^{2,3} and Hedda Agüero-Torres^{1,2}

- 1 Stockholm Gerontology Research Center, Stockholm, Sweden
- 2 Department of Clinical Neuroscience, Occupational Therapy and Elderly Care Research, Division of Geriatric Medicine, Karolinska Institute, Stockholm, Sweden
- 3 Institute of Psychiatry, University of Bologna, Bologna, Italy

Contents

Abstract	365
1. Occurrence of Dementia	366
2. Prevalence of Dementia and Different Dementing Disorders	367
2.1 Geographical Distribution of Dementia	367
2.2 Geographical Distribution of Dementia Types	369
3. Incidence of Dementia and Different Dementing Disorders	370
3.1 Geographical Variation of Dementia	371
3.2 Dementia and Aging	372
4. Conclusions	372

Abstract

Dementia is a common and disabling disorder in the elderly. Because of the worldwide aging phenomenon, existing in both developed and developing countries, dementia has a growing public health relevance. This article reviews the prevalence and incidence data for dementia reported in the international literature in the last 10 years. Results from 36 prevalence and 15 incidence studies have been examined. Prevalence is equal to 0.3 to 1.0 per 100 people in individuals aged 60 to 64 years, and increases to 42.3 to 68.3 per 100 people in individuals 95 years and older. The incidence varies from 0.8 to 4.0 per 1000 person years in people aged 60 to 64 years, and increases to 49.8 to 135.7 per 1000 person years when the population was older than 95 years. The international comparison allows the following conclusions: (i) both prevalence and incidence show little geographical variation, as differences between countries seem to reflect methodological rather than real differences [the low prevalence of dementia in Africa needs to be confirmed by incidence data]; (ii) both incidence and prevalence figures increase with age even in the advanced ages; (iii) regarding dementia types, most of the inconsistency in results from different studies is due to vascular dementia rather than to Alzheimer’s disease (AD); (iv) it is still unclear if the reported higher frequency of vascular dementia in Asian populations is due to differential distribution of genetic and/or environmental factors, or due to methodological differences; (v) different dementia types might have different age distributions.

Dementia is an emerging public health problem as it is one of the most common diseases in the elderly and a major cause of disability and mortality.^[1,2] It is becoming more relevant from a public health perspective because of the increasing number of elderly in the world. In fact, during the past few decades, the aging of the population has become a worldwide phenomenon, no longer confined to Western societies.^[3,4] The unprecedented declines in mortality and fertility have resulted in a rapid population aging process in most of the developing countries. In 1990, 26 nations had more than 2 million elderly citizens (≥ 65 years); by 2025, an additional 33 countries are likely to join the list. Between 1985 and 2025, the rate of increase of the elderly population in some developing countries will be up to 15 times higher than the increase in, for example, Sweden. On the other hand, the developed countries, which have already experienced a dramatic increase in the population aged 65 years and older, will face the progressive aging of the elderly population itself. In most of these countries, the oldest (those 80 years and older) are the fastest growing part of the elderly population.

Consequently, both developed and developing countries will face the challenge of coping with a high frequency of chronic diseases, such as dementia, which are characteristic of aging societies. In response to the worldwide increase in an older population and increased number of individuals affected by disabling disorders, many initiatives have been taken to promote attention to this topic. In April 1995, the World Health Organization (WHO) launched a new programme on aging and health in which research was one of the main objectives. In December 1996, the editor of the *Journal of the American Medical Association (JAMA)* announced that 80 major medical journals through more than 200 proposals had identified aging to be the most important topic to be addressed in the second global theme issue.^[5] 11 months later this Global Theme Issue on Aging was published with the participation of 97 journals in 31 countries.^[6] The European Commission has proposed, with the approval of the European Parliament in December 1997, that the

aging population be one of the key initiatives for the programme for research and technological development in 1998 to 2002 (the 5th Framework Program, FP5).^[3] This initiative is devoted to promoting healthy aging and independence in old age by preventing and treating age-related diseases and disability.^[7]

All of these initiatives reflect the attention of the scientific world on the third age and related diseases. Epidemiology is one of the leading research areas, studying the aging process from an epidemiological point of view is a priority from a public health perspective. A large part of such studies have been devoted to dementia. The first epidemiological studies concerning dementia and different dementia types were carried out during the 1960s in the Scandinavian countries; however, it was during the 1980s that epidemiological methods first began to be widely used in the field of dementia research. These methods were used to describe different patterns of the dementias, and to identify risk factors for different dementing disorders. Finally, in the 1990s, follow-up studies regarding incidence, aetiology and natural history of the dementias have been initiated.

In spite of this large amount of research, several questions concerning the occurrence of dementia in the world are still open. This review aims to explore the following questions:

- Are there geographical differences in the occurrence of dementia?
- Does dementia incidence vary in different ethnic groups?
- Are the degenerative and vascular forms of dementia differently distributed in different countries?

1. Occurrence of Dementia

The occurrence of a disease can be measured as the proportion of individuals affected by the disease in a defined population at a specific instant (prevalence), or as the number of new cases that develop in a population of individuals at risk during a specific time interval (incidence). Whereas the prevalence provides an estimate of the risk of an individ-

ual to be ill at a certain point in time, the incidence estimates the risk of becoming ill.

The prevalence is determined both by incidence and disease duration. Thus, differences in prevalence between groups may be due to differences in treatment/care of the disease and/or general health conditions that can affect disease duration. Most of our knowledge about the occurrence of the dementias is based on prevalence rather than incidence data,^[8,9] as the detection of incident cases is expensive and time consuming. International comparison based on prevalence figures needs to be examined with caution because of the uncertainties of interpretation.

Unfortunately, even the incidence data have limitations as they are often hampered by methodological problems such as small sample sizes, detection of disease onset, attrition rate and diagnostic accuracy. Worldwide comparison of incidence data needs to take into account methodological differences between studies.

Several attempts to review the prevalence of dementia in the world have already been reported,^[10-18] either in the form of meta-analysis or review. In contrast, only 2 meta-analyses of published incidence data in the world^[19,20] and a comparison of incidence studies in Europe^[21] are present in the literature.

In this review, we will compare separately prevalence and incidence data in the last decade among different countries in the world, and among different ethnic groups in the same nation. Published prevalence and incidence studies were identified through: a Medline search from 1990 to 1998, existing reviews of the field and reference lists of identified studies. Population-based studies, in which dementia diagnosis was made according to comparable diagnostic criteria, were included. When the same study was replicated in the same population at different times, only the most recent figures were taken into account. We did not include articles in which age-stratified prevalence or incidence figures were not reported, or in which both the number of cases and the size of the study population were missing. Data from studies carried out in Europe, North America,

Asia and multi-ethnic communities were pooled after stratification in 5-year age groups. When a study did not stratify the oldest ages, only data from the youngest groups were pooled. Prevalence and incidence studies included in this review are reported in tables I to III.

2. Prevalence of Dementia and Different Dementing Disorders

Table I shows the European prevalence studies on dementia. There is a good representation of western European countries, with wide north-south distribution: 2 studies were performed in Finland,^[23,35] 1 in Norway,^[34] 5 in Sweden,^[22,32,36,40,46] 1 in Denmark,^[42] 4 in the UK,^[24,26,29,30] 2 in The Netherlands,^[25,44] 1 in Belgium,^[37] 3 in Germany,^[27,38,39] 4 in Italy,^[31,41,43,45] and 1 in Spain.^[28] We could include only 3 North American studies that investigated Caucasian populations;^[45-49] 3 other studies examined Asian-Americans^[55,56] and African-Americans^[57] (table I). In Africa, only 1 study was performed in Nigeria.^[57] From Asia, we included 2 studies from Japan^[50,51] and 1 each from China,^[54] Korea^[52] and India^[53] (table I).

2.1 Geographical Distribution of Dementia

Pooled age-specific prevalence figures from different continents and multi-ethnic groups are presented in figure 1. In all countries, dementia prevalence increases almost exponentially with age. This finding confirms previous reports that the disease prevalence doubles almost every 5 years.^[11,12,19] Prevalence is very low in individuals under the age of 60 years in all countries (0.3 to 0.7 per 100 people) and increases even in the most advanced ages (33.3 to 68.3 per 100 people). In the age groups up to 85 years, if we exclude the data from Africa,^[57] the prevalence figures are surprisingly similar, leading to a figure per 100 people equal to 0.5 cases in the age group 60 to 65 years, 1.5 in the age group 65 to 69 years, 3 in the age group 70 to 74 years, 6 in the age group 75 to 79 years, and 12 in the age group 80 to 84 years. The differences in the oldest ages are likely to be as a result of unstable prevalences caused by small sample sizes, especially in

Table I. Prevalence studies included in this review

Reference	Site of study	Age (y)	Dementia diagnostic criteria
Europe			
Hagnell et al. ^[22]	Lund, Sweden	30+	DSM-III
Sulkava et al. ^[23]	Finland	65+	DSM-III
Copeland et al. ^[24]	Liverpool, UK	65+	GMS-AGECAT, DSM-III
Heeren et al. ^[25]	Leiden, Netherlands	85+	DSM-III, GMS
Brayne & Calloway ^[26]	Cambridgeshire, UK	70+, women	CAMDEX
Cooper & Bickel ^[27]	Mannheim, Germany	65+	CAMDEX, ICD-9
Lobo et al. ^[28]	Zaragoza, Spain	65+	DSM-III, ICD-10, GMS
O'Connor et al. ^[29]	Cambridge, UK	75+	CAMDEX
Livingston et al. ^[30]	London, UK	65+	DSM-III, GMS
Rocca et al. ^[31]	Appignano, Italy	60+	DSM-III, NINCDS-ADRDA
Fratiglioni et al. ^[32]	Stockholm, Sweden	75+	DSM-III-R
Breteler et al. ^[33]	Amsterdam/Rotterdam, Netherlands	65+	CAMDEX, DSM-III-R
Engedal et al. ^[34]	Oslo, Norway	75+	DSM-III
Juva et al. ^[35]	Helsinki, Finland	75+	CDR
Skoog et al. ^[36]	Gothenburg, Sweden	85+	DSM-III-R, NINCDS-ADRDA, CT-scan
Roelands et al. ^[37]	Heist-op-den-Berg, Belgium	65+	DSM-III-R
Wernicke & Reischies ^[38]	Berlin, Germany	70+	DSM-III-R
Fichter et al. ^[39]	Munich, Germany	85+	DSM-III-R, GMS, ICD-10
Johansson & Zarit ^[40]	Jonkoping, Sweden	85+	DSM-III-R
Prencipe et al. ^[41]	L'Aquila, Italy	65+	DSM-III, NINCDS-ADRDA, NINCDS-AIREN
Andersen et al. ^[42]	Odense, Denmark	65+	CAMDEX, DSM-III-R, NINCDS-ADRDA
Ferini-Strambi et al. ^[43]	Vescovato, Italy	60+	NINCDS-ADRDA, NINCDS-AIREN
Boersma et al. ^[44]	Zwolle, Netherlands	65+	CAMDEX, DSM-III-R
De Ronchi et al. ^[45]	Granarolo, Italy	60+	DSM-III-R
von Strauss et al. ^[46]	Stockholm, Sweden	77+	DSM-III-R
North America			
Kokmen et al. ^[47]	Rochester, Minnesota, US	55+	DSM-III, NINCDS-ADRDA
Bachman et al. ^[48]	Framingham, Massachusetts, US	60+	DSM-III-R, NINCDS-ADRDA
Ebly et al. ^[49]	Canadian Study of Health and Aging (CSHA), Canada	85+	DSM-III-R, NINCDS-ADRDA
Asia			
Hasegawa et al. ^[50]	Kanagawa, Japan	65+	DSM-III
Ueda et al. ^[51]	Hisayama, Japan	65+	DSM-III, CT-scan
Park et al. ^[52]	Myun, Korea	65+	DSM-III-R, CAMDEX
Shaji et al. ^[53]	Kerala, India	60+	CAMDEX, DSM-III-R, ICD-10
Chiu et al. ^[54]	Hong Kong, China	65+	CAMDEX, DSM-IV, ICD-10, NINCDS-AIREN
Asian-Americans			
Graves et al. ^[55]	King County, Washington, US	65+	DSM-III-R, NINCDS-ADRDA
White et al. ^[56]	Honolulu, Hawaii, US	60+	DSM-III-R, NINCDS-ADRDA
African and African-Americans			
Hendrie et al. ^[57]	Ibada, Nigeria and Copiah, Indianapolis, US	65+ 65+	DSM-III-R, ICD-10 DSM-III-R, ICD-10

CAMDEX = Cambridge Mental Disorders of the Elderly Examination; **CDR** = Clinical Dementia Rating; **CT-scan** = computed tomography scan; **DSM-III** = Diagnostic and Statistical Manual of Mental Disorders (third edition); **DSM-III-R** = DSM-III revised; **DSM-IV** = DSM (fourth edition); **GMS** = Geriatric Mental State Interview; **GMS-AGECAT** = GMS–Automated Geriatric Examination for Computer Assisted Taxonomy; **ICD-9** = International Classification of Diseases (ninth edition); **ICD-10** = ICD (tenth edition); **NINCDS-ADRDA** = National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association; **NINCDS-AIREN** = NINCDS and the Association Internationale pour la Recherche et L'Enseignement en Neurosciences.




Fig. 1. Prevalence rates, per 100 people, for dementia on different continents. Pooled data from Europe, Asia and North America.^[22-57]

the oldest European studies.^[22,23,28,30,33] In fact, a recent report from Stockholm, Sweden,^[46] found a prevalence of 37.3 and 47.9 per 100 people in the age groups 90 to 94 years and 95 years and older. These data are similar to those recently reported from North America.^[47-49]

The prevalences in the only study from Africa are low in all ages.^[57] Lower survival or methodological differences can explain this discrepancy,^[16] but possible ethnic variation cannot be excluded. Environmental or genetic factors acting as prognostic and/or risk factors might determine this variation. For example, it has been reported that the apolipoprotein E e4 allele, which is a well known risk factor for dementia in Caucasian^[73] and Asian populations, is not associated with a higher risk of dementia among Africans.^[74]

2.2 Geographical Distribution of Dementia Types

Despite similar overall prevalence of dementia in North America, Europe and Asia, the relative pro-

portions attributed to Alzheimer’s disease (AD) and vascular dementia seems to differ among continents and multi-ethnic communities of Western countries (fig. 2). In most Western countries, 50 to 70% of the total dementia prevalence is attributed to AD, and 20 to 30% to vascular dementia. Only a few reports differ from this pattern.^[31,36] In these 2 European studies^[31,36] carried out in Italy and Sweden, respectively, there was a lower proportion of AD than vascular dementia, and age-specific prevalences were similar. In comparison with other European studies, Rocca et al.^[31] and Skoog et al.^[36] found higher vascular dementia prevalences, but similar AD figures, as shown in table II, in which age-specific prevalences from these studies are compared with our data from the Kungsholmen Project.^[32]

In most Asian studies, higher percentages are found for vascular dementia, as vascular dementia represents 38% of the total dementia cases. Never-




Fig. 2. Prevalent dementia cases: proportion of different dementing disorders in different continents. Pooled data from Europe, Asia and North America. AD = Alzheimer’s disease; VaD = vascular dementia.^[22-57]

Table II. Comparison of studies performed in Stockholm, Appignano and Gothenburg

Study [age (y)]	Prevalence per 100 population (95%confidence interval)		
	dementia	Alzheimer's disease	vascular disease
Stockholm, Sweden^[32]			
75-79	5.7 (4.1-7.7)	3.4 (2.0-4.7)	1.4 (1.0-2.6)
80-89	13.3 (11.1-15.5)	7.0 (5.4-8.7)	3.1 (2.0-4.2)
85-89	20.4 (16.2-25.2)	11.8 (8.5-15.8)	4.9 (2.9-7.9)
Appignano, Italy^{a[31]}			
70-79	5.6 (3.0-8.2)	2.0 (0.4-3.5)	2.6 (0.8-4.4)
80-89	25.9 (17.7-34.2)	10.2 (4.5-15.9)	8.3 (3.1-13.5)
90+	14.3 (0.4-57.9)	14.3 (0.4-57.9)	0.0 (0.0-41.0)
Gothenburg, Sweden^[35]			
85	29.8 (25.8-34.0)	13.0 (10.0-15.9)	14.0 (10.9-17.0)

a Mixed-dementia cases are not included in vascular disease.

theless, if we take into account only recent studies,^[52-54] vascular dementia accounts for 31% and AD for 61%. These figures are similar to the findings reported in Europe, North America and Africa. However, the proportion of vascular dementia is 3 times higher among Asian-Americans than among the other ethnic groups.

In conclusion, it seems that most of the inconsistency present in the literature is due to the estimate of vascular dementia occurrence, rather than AD. The interpretation of the geographical differences in vascular dementia prevalence, if they exist, needs to take into account the following points:

- Variation in diagnostic criteria and procedure may occur. AD and vascular dementia diagnoses are hampered by the lack of diagnostic markers and by the controversial conceptual definition of vascular dementia.^[75] Most epidemiological studies focused on multi-infarct dementia, whereas others used the broader definition of vascular dementia. Very few studies^[33,36] used neuroimaging in the diagnosis of dementia type.
- Dementia duration may vary in different countries because of differential survival and care. This may explain different dementia prevalences between different countries.
- Different age structure of the study populations can affect the relative proportions of dementia types, when not stratified by age. It has been suggested^[36] that vascular dementia may be the ma-

ajor cause of dementia in the highest ages, whereas other researchers have reached an opposite conclusion, suggesting that this type of dementing disorder may be more frequent in the youngest old ages.^[76]

- Differential geographical distribution of vascular risk factors and stroke may cause differential distribution of vascular dementia. High rates of vascular diseases are present among Asian populations, which would inevitably predict a higher proportion of vascular dementias among this ethnic group.^[77] However, the study on African and African-Americans did not show any significantly higher proportion of vascular dementia compared with Caucasian populations, although a higher frequency of hypertension and stroke has been reported in this group.^[77]

3. Incidence of Dementia and Different Dementing Disorders

A limited number of incidence studies are present in the literature, especially regarding different dementing disorders (table III). Only 2 incidence studies were carried out in Asian countries, specifically in Beijing, China, and southern Taiwan^[71,72] and only 1 assessed multi-ethnic American communities.^[78]

3.1 Geographical Variation of Dementia

In figures 3 and 4, the findings from single studies are reported separately for Europe, North America and Asia. Incidence figures are surprisingly similar in the youngest old, whereas a wide variation is present in the 3 oldest groups, probably because of imprecise estimates. During 1 year, there is 1 person per 1000 people aged 60 to 65 years who becomes demented; when the population is older than 95 years, this number increases to an average of 90 new cases in 1 year per 1000 people. One study from Germany^[65] (not reported in fig. 3), showed very high incidence rates in the age range 85 to 89 years (113.6 new cases per 1000 person years), 90 to 94 years (112.5 new cases per 1000 person years) and 95 years and older (235.7 new cases per 1000 person

years), but the case ascertainment was performed during a very short time interval (1 year).

North American studies show a tendency towards lower incidence rates than European studies. Some methodological differences could account for these lower results: (i) in the Framingham study,^[69] the inception of the dementia-free cohort was based on psychological tests, therefore early dementia cases, and maybe preclinical cases, were removed from the study population; (ii) in the Rochester study,^[70] only those people who sought medical attention could be identified as demented cases, because the study was based on a record linkage system.

The incidence studies confirm that AD is the most common dementia disorder, perhaps more common in Caucasian populations than in Asian (fig. 5). Only 1 study reported similar rates for AD and

Table III. Incidence studies included in this review

Reference	Site of study	Age (y)	Dementia diagnostic criteria
Europe			
Copeland et al. ^[68]	Liverpool, UK	65+	GMS-AGECAT
Bickel & Cooper ^[59]	Mannheim, Germany	65+	ICD-9, CAMDEX
Boothby et al. ^[60]	London, UK	65+	CAMCOG, NINCDS-ADRDA
Letenneur et al. ^[61]	Bordeaux, France	65+	DSM-III-R, NINCDS-ADRDA
Paykel et al. ^[62]	Cambridge, UK	75+	CAMDEX
Gussekloo et al. ^[63]	Leiden, Netherlands	85+	DSM-III, GMS-AGECAT
Aevarsson & Skoog ^[64]	Gothenburg, Sweden	85+	ICD-10, NINCDS-ADRDA, CT-scan
Fichter et al. ^[65]	Munich, Germany	85+	DSM-III-R
Brayne et al. ^[66]	Cambridge, UK	70+, women	CAMDEX
Fratiglioni et al. ^[67]	Stockholm, Sweden	75+	DSM-III-R
Ott et al. ^[68]	Rotterdam, Netherlands	55+	CAMDEX, DSM-III-R, NINCDS-ADRDA, NINCDS-AIREN
North America			
Bachman et al. ^[69]	Framingham, Massachusetts, US	60+	NINCDS-ADRDA
Rocca et al. ^[70]	Rochester, Minnesota, US	50+	DSM-III-R, NINCDS-ADRDA
Asia			
Li et al. ^[71]	Beijing, China	60+	DSM-III
Liu et al. ^[72]	Kaohsiung, Taiwan	60+	ICD-10, NINCDS-ADRDA, NINCDS-AIREN

CAMCOG = Cambridge Cognitive Examination; **CAMDEX** = Cambridge Mental Disorders of the Elderly Examination; **CT-scan** = computed tomography scan; **DSM-III** = Diagnostic and Statistical Manual of Mental Disorders (third edition); **DSM-III-R** = DSM-III revised; **GMS-AGECAT** = GMS-Automated Geriatric Examination for Computer Assisted Taxonomy; **ICD-9** = International Classification of Diseases (ninth edition); **ICD-10** = ICD (tenth edition); **NINCDS-ADRDA** = National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association; **NINCDS-AIREN** = NINCDS and the Association Internationale pour la Recherche et L'Enseignement en Neurosciences.




Fig. 3. Age-specific incidence rates for dementia per 1000 person-years in European studies.^[58-64,66-68]

vascular dementia in the ages of 85 to 88 years.^[64] Further investigations are needed in this area.

3.2 Dementia and Aging

In recent years, several studies have addressed the issue of whether dementia is an ‘age- or aging-related disorder’.^[14,19,38] In spite of the current availability of incidence data, it is still unclear if the exponential increase in dementia frequency with age continues even in the advanced ages as an inevitable feature of aging, or if the occurrence reaches a plateau in people aged 85 years and older. This uncertainty derives mainly from the difficulty in studying samples of individuals aged 90 years and older that are large enough to allow reliable estimations of dementia incidence. Moreover, the degree of diagnostic difficulty increases with age. In very old people, it may be problematic to distinguish between

early manifestations of dementia and changes associated with ‘normal aging’.^[46]

Both dementia and AD incidence increase steeply with age, while the relationship between vascular dementia and age is not so clear. In very old ages, it is likely that both vascular and degenerative mechanisms contribute together to the development of dementia, which can be attributed to one or other subtype of dementia depending on the diagnostic tools used.^[8,79]

4. Conclusions

Dementia is a growing worldwide medical, social and economic problem. In all countries, both the prevalence and incidence of dementia increase with increasing age, perhaps even in the most advanced ages. Both prevalence and incidence show little geographical variation, as differences between countries seem to reflect methodological rather than




Fig. 4. Age-specific incidence rates for dementia per 1000 person-years in American and Asian studies.^[69-72]




Fig. 5. Incident dementia cases: proportion of different dementing disorders in different countries, pooled data from Europe, Asia and the US. AD = Alzheimer's disease; VaD = vascular dementia.^[58-72]

real differences. The low prevalence of dementia in Africa needs to be replicated by other prevalence studies and confirmed by incidence data.^[80,81] Regarding dementia types, it is still unclear if the reported higher frequency of vascular dementia in Asian populations is due to differential distribution of genetic and/or environmental factors, or if it is just a result of methodological differences. Finally, it is still controversial whether different dementia types have different age distributions.

Most of the inconsistency is with vascular dementia rather than AD. In fact, both AD and vascular dementia diagnoses are hampered by the lack of diagnostic markers. In addition, there is controversy regarding the definition of vascular dementia. Most likely, vascular and degenerative mechanisms contribute together to the development of dementia especially in very old ages. More incidence studies with comparable methodology are needed for a better understanding of the worldwide occurrence of dementia, and in order to generate new aetiological hypotheses.

References

1. Agüero-Torres H, Fratiglioni L, Guo Z, et al. Prognostic factors in very old demented adults: a seven-year follow-up from a population-based survey in Stockholm. *J Am Geriatr Soc* 1998 Apr; 46 (4): 444-52
2. Agüero-Torres H, Fratiglioni L, Guo Z, et al. Dementia is the major cause of functional dependence in the elderly: 3-year follow-up data from a population-based study. *Am J Public Health* 1998 Oct; 88 (10): 1452-6
3. Kalache A. Ageing worldwide. In: Ebrahim S, Kalche A, editors. *Epidemiology in old age*. London: BMJ, 1996: 22-31
4. Kinsella K. Demographic aspects. In: Ebrahim S, Kalche A, editors. *Epidemiology in old age*. London: BMJ, 1996: 32-40
5. Winker MA, Glass RM. The aging global population: a call for papers. *JAMA* 1996; 276: 1758
6. Winker MA. Aging: a global issue. *JAMA* 1997; 278: 1377
7. Anonymous. Reducing disability among the elderly in Europe [editorial]. *Lancet* 1997; 350: 1261
8. Fratiglioni L. Epidemiology. In: Winblad B, Wimo A, Jonsson B, et al., editors. *Health economics of dementia*. Chichester: John Wiley & Son Inc., 1998: 13-31
9. Fratiglioni L, Agüero-Torres H, Winblad B. Impact of dementia in our societies and current possibilities of prevention. *Int J Geriatr Psychopharmacol* 1998; 1: 179-83
10. Jorm AF, Korten AE, Henderson AS. The prevalence of dementia: a quantitative integration of the literature. *Acta Psychiatr Scand* 1987 Nov; 76 (5): 465-79
11. Hofman A, Rocca WA, Brayne C, et al. The prevalence of dementia in Europe: a collaborative study of 1980-1990 findings: Eurodem Prevalence Research Group. *Int J Epidemiol* 1991 Sep; 20 (3): 736-48
12. Rocca WA, Hofman A, Brayne C, et al. Frequency and distribution of Alzheimer's disease in Europe: a collaborative study of 1980-1990 prevalence findings: the EURODEM-Prevalence Research Group. *Ann Neurol* 1991 Sep; 30 (3): 381-90
13. Rocca WA, Hofman A, Brayne C, et al. The prevalence of vascular dementia in Europe: facts and fragments from 1980-1990 studies: EURODEM-Prevalence Research Group. *Ann Neurol* 1991 Dec; 30 (6): 817-24
14. Ritchie K, Kildea D. Is senile dementia 'age-related' or 'ageing-related'? Evidence from meta-analysis of dementia prevalence in the oldest old. *Lancet* 1995 Oct 7; 346 (8980): 931-4
15. Ritchie K, Kildea D, Robine JM. The relationship between age and the prevalence of senile dementia: a meta-analysis of recent data. *Int J Epidemiol* 1992 Aug; 21 (4): 763-9
16. Corrada M, Brookmeyer R, Kawas C. Sources of variability in prevalence rates of Alzheimer's disease. *Int J Epidemiol* 1995 Oct; 24 (5): 1000-5
17. Jorm AF. Cross-national comparisons of the occurrence of Alzheimer's and vascular dementias. *Eur Arch Psychiatry Clin Neurosci* 1991; 240 (4-5): 218-22
18. Rockwood K, Stadnyk K. The prevalence of dementia in the elderly: a review. *Can J Psychiatry* 1994 Jun; 39 (5): 253-7
19. Jorm AF, Jolley D. The incidence of dementia: a meta-analysis. *Neurology* 1998 Sep; 51 (3): 728-33
20. Launer LJ, Fratiglioni L, Andersen K, et al. Regional differences in the incidence of dementia in Europe: Eurodem Collaborative analysis. In: Iqbal K, Swaab DF, Winblad B, et al. *Alzheimer's disease and related disorders: etiology, pathogenesis and therapeutics*. Chichester: John Wiley & Son Inc., 1999: 9-17
21. Gao S, Hendrie HC, Hall KS, et al. The relationships between age, sex and the incidence of dementia and Alzheimer's disease: a meta-analysis. *Arch Gen Psychiatry* 1998; Sep; 55: 809-15

22. Hagnell O, Lanke J, Rorsman B, et al. Does the incidence of age psychosis decrease? A prospective, longitudinal study of a complete population investigated during the 25-year period 1947-1972: the Lundby study. *Neuropsychobiology* 1981; 7 (4): 201-11
23. Sulkava R, Wikstrom J, Aromaa A, et al. Prevalence of severe dementia in Finland. *Neurology* 1985 Jul; 35 (7): 1025-9
24. Copeland JR, Dewey ME, Wood N, et al. Range of mental illness among the elderly in the community: prevalence in Liverpool using the GMS-AGECAT package. *Br J Psychiatry* 1987 Jun; 150: 815-23
25. Heeren TJ, Lagaav AM, Hilmans W, et al. Prevalence of dementia in the 'oldest old' in a Dutch population. *J Am Geriatr Soc* 1991 Aug; 39 (8): 755-9
26. Brayne C, Calloway P. An epidemiological study of dementia in a rural population of elderly women. *Br J Psychiatry* 1989 Aug; 155: 214-9
27. Cooper B, Bickel HE. Prevalence and incidence of dementia diseases in the elderly population: results of a population-related longitudinal study in Mannheim [in German]. *Nervenarzt* 1989 Aug; 60 (8): 472-82
28. Lobo A, Dewey M, Copeland J, et al. The prevalence of dementia among elderly people living in Zaragoza and Liverpool. *Psychol Med* 1992 Feb; 22 (1): 239-43
29. O'Connor DW, Pollitt PA, Hyde JB, et al. The prevalence of dementia as measured by the Cambridge Mental Disorders of the Elderly Examination. *Acta Psychiatr Scand* 1989 Feb; 79 (2): 190-8
30. Livingston G, Sax K, Willison J, et al. The Gospel Oak Study stage II: the diagnosis of dementia in the community. *Psychol Med* 1990 Nov; 20 (4): 881-91
31. Rocca WA, Bonaiuto S, Lippi A, et al. Prevalence of clinically diagnosed Alzheimer's disease and other dementing disorders: a door-to-door survey in Appignano, Macerata Province, Italy. *Neurology* 1990 Apr; 40 (4): 626-31
32. Fratiglioni L, Grut M, Forsell Y, et al. Prevalence of Alzheimer's disease and other dementias in an elderly urban population: relationship with age, sex, and education. *Neurology* 1991 Dec; 41 (12): 1886-92
33. Breteler MM, van den Ouweland FA, Grobbee DE, et al. A community-based study of dementia: the Rotterdam Elderly Study. *Neuroepidemiology* 1992; 11 Suppl. 1: 23-8
34. Engedal K, Gilje K, Laake K. Prevalence of dementia in a Norwegian sample aged 75 years and over and living at home. *Compr Gerontol [A]* 1988 Oct; 2 (3): 102-6
35. Juva K, Sulkava R, Erkinjuntti T, et al. Prevalence of dementia in the city of Helsinki. *Acta Neurol Scand* 1993 Feb; 87 (2): 106-10
36. Skoog I, Nilsson L, Palmertz B, et al. A population-based study of dementia in 85-year-olds. *N Engl J Med* 1993 Jan 21; 328 (3): 153-8
37. Roelands M, Wostyn P, Dom H, et al. The prevalence of dementia in Belgium: a population-based door-to-door survey in a rural community. *Neuroepidemiology* 1994; 13 (4): 155-61
38. Wernicke TF, Reischies FM. Prevalence of dementia in old age: clinical diagnoses in subjects aged 95 years and older. *Neurology* 1994 Feb; 44 (2): 250-3
39. Fichter MM, Meller I, Schroppe H, et al. Dementia and cognitive impairment in the oldest old in the community: prevalence and comorbidity. *Br J Psychiatry* 1995 May; 166 (5): 621-9
40. Johansson B, Zarit SH. Prevalence and incidence of dementia in the oldest old: a longitudinal study of a population-based sample of 84-90-year-olds in Sweden. *Int J Geriatr Psychiatry* 1995; 10: 359-66
41. Prencipe M, Casini AR, Ferretti C, et al. Prevalence of dementia in an elderly rural population: effects of age, sex, and education. *J Neurol Neurosurg Psychiatry* 1996 Jun; 60 (6): 628-33
42. Andersen K, Lolk A, Nielsen H, et al. Prevalence of very mild to severe dementia in Denmark. *Acta Neurol Scand* 1997 Aug; 96 (2): 82-7
43. Ferini-Strambi L, Marcone A, Garancini P, et al. Dementing disorders in north Italy: prevalence study in Vescovato, Cremona Province. *Eur J Epidemiol* 1997 Feb; 13 (2): 201-4
44. Boersma F, Eefsting JA, van den Brink W, et al. Prevalence of dementia in a rural Netherlands population and the influence of DSM-III-R and CAMDEX criteria for the prevalence of mild and more severe forms. *J Clin Epidemiol* 1998 Mar; 51 (3): 189-97
45. De Ronchi D, Fratiglioni L, Rucci P, et al. The effect of education on dementia occurrence in an Italian population with middle to high socioeconomic status. *Neurology* 1998 May; 50 (5): 1231-8
46. von Strauss E, Viitanen M, De Ronchi D, et al. Dementia is an ageing related disorder even in the oldest old: evidence from a population aged 90 years and over. *Arch Neurol* 1999; 56: 587-92
47. Kokmen E, Beard CM, Offord KP, et al. Prevalence of medically diagnosed dementia in a defined United States population: Rochester, Minnesota, January 1, 1975. *Neurology* 1989 Jun; 39 (6): 773-6
48. Bachman DL, Wolf PA, Linn R, et al. Prevalence of dementia and probable senile dementia of the Alzheimer type in the Framingham Study. *Neurology* 1992 Jan; 42 (1): 115-9
49. Eibly EM, Parhad IM, Hogan DB, et al. Prevalence and types of dementia in the very old: results from the Canadian Study of Health and Aging. *Neurology* 1994 Sep; 44 (9): 1593-600
50. Hasegawa K, Homma A, Yukimichi I. An epidemiological study of age-related dementia in the community. *Int J Geriatr Psychiatry* 1986; 1: 45-55
51. Ueda K, Kawano H, Hasuo Y, et al. Prevalence and etiology of dementia in a Japanese community. *Stroke* 1992 Jun; 23 (6): 798-803
52. Park J, Ko HJ, Park YN, et al. Dementia among the elderly in a rural Korean community. *Br J Psychiatry* 1994 Jun; 164 (6): 796-801
53. Shaji S, Promodu K, Abraham T, et al. An epidemiological study of dementia in a rural community in Kerala, India. *Br J Psychiatry* 1996 Jun; 168 (6): 745-9
54. Chiu HF, Lam LC, Chi I, et al. Prevalence of dementia in Chinese elderly in Hong Kong. *Neurology* 1998 Apr; 50 (4): 1002-9
55. Graves AB, Larson EB, Edland SD, et al. Prevalence of dementia and its subtypes in the Japanese American population of King County, Washington state: the Kame Project. *Am J Epidemiol* 1996 Oct 15; 144 (8): 760-71
56. White L, Petrovitch H, Ross GW, et al. Prevalence of dementia in older Japanese-American men in Hawaii: the Honolulu-Asia Aging Study (see comments). *MA* 1996 Sept 25; 276 (12): 955-60
57. Hendrie HC, Osuntokun BO, Hall KS, et al. Prevalence of Alzheimer's disease and dementia in two communities: Nigerian Africans and African Americans. *Am J Psychiatry* 1995 Oct; 152 (10): 1485-92
58. Copeland JR, Davidson IA, Dewey ME, et al. Alzheimer's disease, other dementias, depression and pseudodementia: prevalence, incidence and three-year outcome in Liverpool. *Br J Psychiatry* 1992 Aug; 161: 230-9

59. Bickel H, Cooper B. Incidence and relative risk of dementia in an urban elderly population: findings of a prospective field study. *Psychol Med* 1994 Feb; 24 (1): 179-92
60. Boothby H, Blizard R, Livingston G, et al. The Gospel Oak Study stage III: the incidence of dementia. *Psychol Med* 1994 Feb; 24 (1): 89-95
61. Letenneur L, Commenges D, Dartigues JF, et al. Incidence of dementia and Alzheimer's disease in elderly community residents of south-western France. *Int J Epidemiol* 1994 Dec; 23 (6): 1256-61
62. Paykel ES, Brayne C, Huppert FA, et al. Incidence of dementia in a population older than 75 years in the United Kingdom. *Arch Gen Psychiatry* 1994 Apr; 51 (4): 325-32
63. Gussekloo J, Heeren TJ, Izaks GJ, et al. A community based study of the incidence of dementia in subjects aged 85 years and over. *J Neurol Neurosurg Psychiatry* 1995 Nov; 59 (5): 507-10
64. Aearvsson O, Skoog I. A population-based study on the incidence of dementia disorders between 85 and 88 years of age. *J Am Geriatr Soc* 1996 Dec; 44 (12): 1455-60
65. Fichter MM, Schroppe H, Meller I. Incidence of dementia in a Munich community sample of the oldest old. *Eur Arch Psychiatry Clin Neurosci* 1996; 246 (6): 320-8
66. Brayne C, Best N, Muir M, et al. Five-year incidence and prediction of dementia and cognitive decline in a population sample of women aged 70-79 at baseline. *Int J Geriatr Psychiatry* 1997 Nov; 12 (11): 1107-18
67. Fratiglioni L, Viitanen M, von Strauss E, et al. Very old women at highest risk of dementia and Alzheimer's disease: incidence data from the Kungsholmen Project, Stockholm. *Neurology* 1997 Jan; 48 (1): 132-8
68. Ott A, Breteler MM, van Harskamp F, et al. Incidence and risk of dementia: the Rotterdam Study. *Am J Epidemiol* 1998 Mar 15; 147 (6): 574-80
69. Bachman DL, Wolf PA, Linn RT, et al. Risk Incidence of dementia and probable Alzheimer's disease in a general population: the Framingham Study. *Neurology* 1993 Mar; 43 (3 Pt 1): 515-9
70. Rocca WA, Cha RH, Waring SC, et al. Incidence of dementia and Alzheimer's disease: a reanalysis of data from Rochester, Minnesota, 1975-1984. *Am J Epidemiol* 1998 Jul 1; 148 (1): 51-62
71. Li G, Shen YC, Chen CH, et al. A three-year follow-up study of age-related dementia in an urban area of Beijing. *Acta Psychiatr Scand* 1991 Feb; 83 (2): 99-104
72. Liu CK, Lai CL, Tai CT, et al. Incidence and subtypes of dementia in southern Taiwan: impact of socio-demographic factors. *Neurology* 1998 Jun; 50 (6): 1572-9
73. Corder EH, Lannfelt L, Bogdanovic N, et al. The role of APOE polymorphisms in late-onset dementias. *Cell Mol Life Sci* 1998 Sep; 54 (9): 928-34
74. Kalaria RN, Ogeng'o JA, Patel NB, et al. Evaluation of risk factors for Alzheimer's disease in elderly east Africans. *Brain Res Bull* 1997; 44 (5): 573-7
75. Fratiglioni L. Classifications and diagnostics. In: Winblad B, Wimo A, Jonsson B, et al., editors. *Health economics of dementia*. Chichester: John Wiley & Son Inc., 1998: 3-11
76. Hebert R, Brayne C. Epidemiology of vascular dementia. *Neuroepidemiology* 1995; 14 (5): 240-57
77. van Duijn CM. Epidemiology of the dementias: recent developments and new approaches. *J Neurol Neurosurg Psychiatry* 1996 May; 60 (5): 478-88
78. Perkins P, Annegers JF, Doody RS, et al. Incidence and prevalence of dementia in a multiethnic cohort of municipal retirees. *Neurology* 1997 Jul; 49 (1): 44-50
79. Pasquier F, Leys D. Why are stroke patients prone to develop dementia? *J Neurol* 1997 Mar; 244 (3): 135-42
80. Graham C, Howard R, Ha Y. Dementia and ethnicity. *Int Psychogeriatr* 1998 Jun; 10 (2): 183-91
81. Prince M. The need for research on dementia in developing countries. *Trop Med Int Health* 1997 Oct; 2 (10): 993-1000

Correspondence and reprints: Dr L. Fratiglioni, Stockholm Gerontology Research Center, Box 6401, S-113 82 Stockholm, Sweden.
E-mail: Laura.Fratiglioni@cnsf.ki.se