

X-LINKED B-LYMPHOCYTE IMMUNE DEFECT IN CBA/N MICE

II. Studies of the Mechanisms Underlying the Immune Defect*

BY IRWIN SCHER, ALFRED D. STEINBERG, ALICE K. BERNING, AND WILLIAM E. PAUL

(From the Department of Clinical and Experimental Immunology, Naval Medical Research Institute; the Division of Clinical Immunology, Department of Medicine, National Naval Medical Center; the Arthritis, and Rheumatism Branch, National Institute of Arthritis, Metabolism and Digestive Diseases, and the Laboratory of Immunology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20014)

CBA/N (CN)¹ mice are a subline of CBA mice with a marked defect in the function and a diminution in the number of thymus-independent (B) lymphocytes. These animals are unable to form specific antibody to Type III pneumococcal polysaccharide (SIII), bacterial lipopolysaccharide (LPS) (1), or polyriboinosinic-polyribocytidylic acid (poly I·C) (2). Each of these antigens is known to evoke an antibody response in all conventional mouse strains studied, even if these mice are deprived of thymus-dependent (T) lymphocytes (3-5). The inability of the CN mice to respond to these "T-independent" antigens is inherited as an X-linked recessive trait (1, 2). We have shown that CN mice and F₁ male mice of the CN × DBA/2N (DN) cross (CN × DN F₁ males) have a diminished number of immunoglobulin (Ig)-bearing spleen cells, an impaired response to agents mitogenic for B lymphocytes (B mitogens) and a diminished ability to participate in antibody-dependent cell-mediated cytotoxicity (6).

Although the X-linked immune defect in the CN mice is expressed as a functional abnormality of B lymphocytes, the mechanism underlying the defect has not been studied. In particular, it is not known whether the abnormality of CN B-lymphocyte function represents a defect intrinsic to the B-lymphocyte line or results either from an abnormality in the microenvironment in which CN B lymphocytes differentiate or from abnormal function of CN T lymphocytes. Because of the X-linked nature of the immune defect of CN mice, lymphoid cell

* This work was supported in part by the Bureau of Medicine and Surgery, Navy Department, Work Unit nos. MR041.02.01.0020B2GI and CIC 3-06-132. The opinions or assertions contained herein are the private ones of the authors and are not to be construed as official or reflecting the views of the Navy Department or the naval service at large. The animals used in this study were handled in accordance with the provisions of Public Law 89-54 as amended by Public Law 91-579, "Animal Welfare Act of 1970," and the principles outlined in the "Guide for the Care and Use of Laboratory Animals," U. S. Department of Health, Education and Welfare publication no. (NIH) 73-23.

¹ *Abbreviations used in this paper:* anti- θ , anti-Thy 1.2; B, thymus independent; BRBC, burro erythrocytes; Con A, concanavalin A; CN, CBA/N mice; DN, DBA/2N mice; DNP-lys-Ficoll, 2,4-dinitrophenyl-lysyl-derivative of Ficoll; LPS, bacterial lipopolysaccharide; PFC, plaque-forming cell; poly I·C, polyriboinosinic-polyribocytidylic acid; SIII, type III pneumococcal polysaccharide; T, thymus dependent; TNP, trinitrophenyl.

transfers between phenotypically normal F₁ female mice and their abnormal male littermates can be performed. By immunization of recipients of transferred cells with either of two T-independent antigens, poly I-C and the 2,4-dinitrophenyl-lysyl-derivative of Ficoll (DNP-lys-Ficoll) (7, 8), we have studied the cellular basis of the immune defect of CN mice. Our data indicate that the failure of these mice to respond to T-independent antigens is a result of a deficiency or an intrinsic abnormality of B lymphocytes and/or their progenitors rather than a microenvironmental or T-lymphocyte abnormality.

Materials and Methods

Animals. CN,² DN mice, and F₁ animals derived from these strains, (CN × DN) F₁, were obtained from the Rodent and Rabbit Production Section of the National Institutes of Health, Bethesda, Md. All mice were 6–12wk of age at the time of study. The CN mice are a distinct subline of CBA mice and the history of their establishment has been described elsewhere (1, 2). F₁ mice of both sexes were produced by breeding CN females with DN males (CN × DN F₁ males or females).

Lethally irradiated mice (1,000 rads, ⁶⁰Co source, dose rate 40 rads/min) were reconstituted on the day after irradiation by the administration of spleen or bone marrow cells intravenously. CN × DN F₁ males or females were thymectomized at 6 wk of age by aspiration through a sternum-splitting incision. These mice were lethally irradiated 14 days after surgery and reconstituted with 10 × 10⁶ F₁ male or female bone marrow cells which had been treated with AKR anti-Thy 1.2 (anti-θ) plus rabbit complement (C) as has been previously described (9).

Cell Suspensions. Mice were killed by cervical dislocation and their thymuses and/or spleens removed. Spleen and thymus cells were obtained by gentle teasing with a rubber policeman and forceps into RPMI 1640 (Grand Island Biological Co, Grand Island, N. Y.). Bone marrow cells were flushed from the femur and tibial bones using a 25 gauge needle. Cell aggregates were disrupted by passing the cell suspensions through a 26 gauge needle. The single cell suspensions were then washed twice with RPMI 1640 and the number of cells counted.

Antigens. Poly I-C, prepared by annealing the homopolymers polyriboinosinic acid and polyribocytidylic acid, (P-L Biochemicals, Inc., Milwaukee, Wis.) was dissolved in borate buffer, pH 8, and emulsified with an equal volume of complete Freund's adjuvant (Difco Laboratories, Detroit, Mich.) so that 0.3 ml of the final emulsion contained 100 μg of antigen. Mice were immunized with 0.3 ml of this material by intraperitoneal injection.

DNP-lys-Ficoll was prepared according to the method of Sharon et al. (8). Briefly, Ficoll (Pharmacia Fine Chemicals, Inc., Piscataway, N. J.) was reacted with cyanuric chloride at 4°C, followed by the addition of ε-DNP-L-lysine and subsequent reaction at room temperature. The DNP-lys-Ficoll used in these studies had a molar ratio of DNP to Ficoll of 32:1.

Sheep erythrocytes (SRBC) and burro erythrocytes (BRBC) were obtained from a single sheep or burro and were collected sterily in citric acid-dextrose solution (ACD solution; Abbott Laboratories, Chemical Marketing Div., North Chicago, Ill.). These were washed three times in Hanks' balanced salt solution before use. SRBC were heavily conjugated with the trinitrophenyl (TNP) hapten by reacting sodium 2,4,6-trinitrobenzenesulfonate with SRBC according to the method of Kettman and Dutton (10). Mice were immunized by the intravenous injection of 2 × 10⁸ of these heavily conjugated TNP-SRBC immediately after their preparation.

Antibody and Plaque-Forming Cell Assays. Antibody directed against poly I-C was assayed in sera obtained by orbital sinus puncture of immunized mice by an ammonium sulfate precipitation assay using [¹⁴C]poly I-C (Miles Laboratories Inc., Miles Research Div., Elkhart, Ind.) as ligand (11). Data are expressed as the percent binding of 80 ng of [¹⁴C]poly I-C (4,000 dpm/μg) by 25 μl of mouse serum. Binding of greater than 20% was considered a positive response; in previous studies we have shown that serum from unimmunized mice do not exceed this degree of binding (2) as is the case for mice immunized with noncross-reacting antigens or adjuvant alone.

Spleen cells from mice immunized with SRBC, heavily substituted TNP-SRBC or DNP-lys-

² CBA/N mice were formerly designated CBA/HN.

Ficoll were assayed for cells releasing specific antibody by a modification of the Jerne hemolytic plaque technique (12), using SRBC or lightly conjugated TNP-BRBC or TNP-SRBC as indicator cells, respectively. TNP-SRBC and TNP-BRBC were prepared by the method of Rittenberg and Pratt (13). There was less than 1% cross-reactivity between SRBC and BRBC in our system. Plaque-forming cells (PFC) releasing IgG antibodies were detected according to the method of Pierce et al. (14) by inhibition of IgM PFC with goat antimouse μ -chain antibody incorporated in the agar and development of IgG PFC with rabbit polyvalent antimouse γ -chain antibody (both antisera were the gift of Dr. R. Asofsky, NIAID, Bethesda, Md.). Data are expressed as the mean PFC per spleen or the PFC per 10^6 cells \pm SE unless otherwise indicated.

Results

Response of CN \times DN F₁ Male and Female Mice to SRBC and TNP-SRBC. As a basis for comparison with subsequent studies of T-independent antibody responses of male and female CN \times DN F₁ mice, we initially evaluated the response of these animals to two highly thymus-dependent antigens, SRBC and TNP-SRBC. In each of these experiments, the IgM response of F₁ male mice at 4 days after immunization with SRBC or TNP-SRBC was less than those of female mice when considered on the basis of PFC per spleen (Table I). However, since male mice have substantially fewer nucleated cells per spleen (6), it is more instructive to consider the antibody response on the basis of the number of PFC per 10^6 spleen cells. In the first two experiments shown in Table I, the male response, considered on this basis, was approximately half that of the female response and in the third experiment, only slightly less than that of the female. These results are particularly significant in view of our recent finding that the percent of B lymphocytes in the spleen of F₁ male mice is substantially less than that of F₁ female mice (6), and they indicate that F₁ male mice make considerable responses to thymus-dependent antigens. Or on other hand, the IgG anti-SRBC response of F₁ male mice was substantially less than that of F₁ females

TABLE I
IgM and IgG Responses of CN \times DN F₁ Male and Female Mice to TNP-SRBC

Exp.	Sex (no. of mice)	Immuno-globulin	TNP-BRBC*		SRBC*	
			PFC/spleen	PFC/ 10^6 cells	PFC/spleen	PFC/ 10^6 cells
1	Male (6)	IgM	—	—	2,000 \pm 488	44 \pm 6.1
	Female (6)		—	—	7,583 \pm 1,450	77 \pm 10.6
2	Male (3)	IgM	4,037 \pm 678	83.4 \pm 10.7	4,175 \pm 724	86 \pm 11.4
	Female (3)		17,443 \pm 2,362	208.2 \pm 23.4	15,400 \pm 1,584	166 \pm 30.0
3	Male (3)	IgM	28,900 \pm 8,636	586.0 \pm 158.0	9,925 \pm 1,615	204 \pm 22.4
	Female (3)		59,417 \pm 17,774	700.0 \pm 76.0	29,933 \pm 11,855	287 \pm 95.4
3	Male (4)	IgG	—	—	2,858 \pm 414	36 \pm 8.2
	Female (4)		—	—	18,427 \pm 3,080	216 \pm 19.2

* The mean number of IgM or IgG PFC measured at 4 or 11 days, respectively, in response to 2×10^8 heavily conjugated TNP-SRBC. Data are expressed as the net mean number of PFC \pm SE.

even when considered on a per B-cell basis (Table I, exp. 3).

Response of CN and CN × DN F₁ Mice to DNP-lys-Ficoll. DNP-lys-Ficoll has been recently demonstrated to be a potent T-independent antigen which elicits both IgM and IgG anti-DNP antibody in many strains of mice (8). CN × DN F₁ male and female mice were immunized with varying doses of DNP-lys-Ficoll and their direct (IgM) anti-DNP PFC response was measured at 4 days (Table II). All of the F₁ female mice formed large numbers of PFC after immunization with from 0.1–500 μg of DNP-lys-Ficoll. In contrast to the vigorous responses of the F₁ females, 10 of 12 F₁ males immunized with DNP-lys-Ficoll had fewer PFC than the number seen in unimmunized F₁ males. Two of the F₁ male mice had responses of 800 and 1,600 PFC/spleen after immunization with 0.1 and 1.0 μg of DNP-lys-Ficoll, respectively. These values were the highest responses seen with CN × DN F₁ male mice in our experience and are in

TABLE II
Response of CN × DN F₁ Male and Female Mice to Varying Doses of DNP-Lys-Ficoll

DNP-lys-Ficoll	TNP-SRBC at day 4*			
	Male		Female	
	PFC/spleen	PFC/10 ⁶ cells	PFC/spleen	PFC/10 ⁶ cells
μg				
0	250 ± 72	6.9 ± 1.4	955 ± 325	12 ± 3.6
0.1	330 ± 235	9.4 ± 7.6	13,600 ± 3,372	156 ± 24.0
1.0	641 ± 480	11.8 ± 9.0	22,108 ± 9,125	357 ± 113.1
20.0	150 ± 68	3.3 ± 0.5	37,000 ± 4,485	340 ± 60.4
500.0	100 ± 14	2.1 ± 0.6	13,616 ± 3,156	169 ± 28.4

* The mean number (three animals per group) of IgM PFC (±SE) measured on day 4 in response to varying doses of DNP-lys-Ficoll.

the same range as the number of background TNP-SRBC plaques seen in CN × DN F₁ females.

Studies of the 4- and 8-day IgM and IgG anti-DNP PFC responses of CN and CN × DN F₁ mice to DNP-lys-Ficoll are shown in Table III. The CN males and females and CN × DN F₁ males had no response (less than background) while the CN × DN F₁ females made large numbers of PFC at both 4 and 8 days. These experiments demonstrate the CBA/N mice have an X-linked defect in responsiveness to another T-independent antigen, DNP-lys-Ficoll. Moreover, since CN × DN F₁ male mice are able to form antibody to the highly cross-reactive hapten (TNP) coupled to erythrocytes, their X-linked immune defect is not a simple absence of the genetic information required to synthesize anti-DNP antibody.

Reconstitution of poly I·C Response in Lethally Irradiated CN × DN F₁ Mice by CN × DN Spleen Cells. In order to study the influence of the CN × DN F₁ male and female environment on the ability of splenic B lymphocytes to respond to T-independent antigens, lethally irradiated F₁ male and female mice were reconstituted with either F₁ male or F₁ female spleen cells and immunized immediately with poly I·C. At 14 days, antibody to poly I·C was found in 9 of 11

TABLE III
IgM and IgG Responses of Male and Female CN and CN × DN F₁ Mice to DNP-Lys-Ficoll (100 μg)

Strain	Sex	TNP-SRBC*			
		IgM (day 4)		IgG (day 7)	
		PFC/spleen	PFC/10 ⁶ cells	PFC/spleen	PFC/10 ⁶ cells
CN	Male	325 ± 160	5.8 ± 3.1	125 ± 11.1	2.6 ± 0.2
	Female	125 ± 32	2.9 ± 1.6	13 ± 2.0	0.3 ± 0.1
CN × DN	Male	150 ± 25	3.0 ± 0.2	25 ± 5.0	0.8 ± 0.3
	Female	11,775 ± 1,041	201.0 ± 24.1	3,695 ± 150.0	53.0 ± 10.2

* The mean number (four animals per group) of IgM or IgG PFC ± SE measured at 4 or 7 days, respectively, in response to 100 μg or DNP-lys-Ficoll.

TABLE IV
Response of Lethally Irradiated CN × DN F₁ Mice to Poly I·C After Reconstitution with CN × DN F₁ Spleen Cells

Sex of recipient	[¹⁴ C]poly I·C binding*	
	Female donor	Male donor
Female	%	%
	56.1	4.7
	80.6	3.2
	76.9	0.2
	28.5	4.7
Male	21.0	1.7
	0.5	2.4
	14.4	0.2
	71.7	0.0
	21.1	0.0
	71.7	2.4
	50.9	0.0

* Recipient mice were lethally irradiated (1,000 R) and reconstituted with 50 × 10⁶ F₁ male or female spleen cells given intravenously on the day of irradiation. They were immediately challenged with 100 μg of poly I·C in CFA and the percent [¹⁴C]poly I·C binding was assayed on 25 μl of serum obtained 14 days after immunization.

recipients of F₁ female spleen cells. This included four of four F₁ female recipients and five of seven F₁ male recipients (Table IV). In contrast, none of the 11 F₁ mice (either male or female) which had received male F₁ spleen cells formed antibody to poly I·C.

Reconstitution of DNP-Lys-Ficoll Response in Lethally Irradiated CN × DN Mice with CN × DN Bone Marrow. The previous experiment demonstrated

that a population of female cells containing splenic B lymphocytes was able to transfer to irradiated F_1 males the ability to form antibody to poly I-C upon immediate immunization. In order to study the influence of environment on the maturation of these responsive cells, we reconstituted lethally irradiated male and female $CN \times DN F_1$ mice with 10×10^6 bone marrow cells derived from either male or female $CN \times DN$ donors and waited 8 wk before immunizing with DNP-lys-Ficoll (Table V). The mice were sacrificed 4 days later and the numbers of nucleated spleen cells and direct TNP-SRBC PFC determined. Both male and female recipients of female bone marrow cells had substantially greater numbers of nucleated spleen cells than male and female recipients of male bone marrow cells. Indeed, the number of nucleated spleen cells in lethally irradiated recipients which received F_1 male cells resembled that of nonirradiated F_1 male animals, while the numbers of nucleated spleen cells in recipients of F_1 female cells resembled that of normal F_1 females (6). Furthermore, male and female

TABLE V
Response of Lethally Irradiated $CN \times DN F_1$ Male and Female Mice to DNP-Lys-Ficoll after Reconstitution with $10 \times 10^6 F_1$ Male or Female Bone Marrow Cells

Sex of recipient	Sex of donor	No. of nucleated cells per spleen \ddagger	TNP-SRBC*	
			PFC/spleen	PFC/ 10^6 cells
Male	Male	37.8 ± 4.2	50 ± 25	1.5 ± 1.2
	Female	90.3 ± 4.8	$21,875 \pm 709$	243.7 ± 24.0
Female	Male	58.0 ± 7.5	$1,575 \pm 368$	29.2 ± 9.8
	Female	112.0 ± 8.5	$26,217 \pm 3,397$	210.3 ± 7.9

* Recipient mice were lethally irradiated (1,000 R) at 8 wk of age and reconstituted with 10×10^6 bone marrow cells. 8 wk after reconstitution, they were immunized with 100 μ g of DNP-lys-Ficoll. IgM PFC were measured at 4 days.

\ddagger The number of nucleated cells per spleen \pm SE (three animals per group).

recipients of F_1 female bone marrow cells made a vigorous response to DNP-lys-Ficoll while male and female recipients of male cells had very few TNP-SRBC PFC 4 days after immunization. Male recipients of male cells made essentially no response; female recipients of male cells mounted a small response, which can probably be ascribed to the contribution of F_1 female bone marrow cells that escaped or recovered from the effects of irradiation.

These experiments demonstrate that F_1 female spleen and bone marrow cells can transfer to lethally irradiated F_1 male recipients the ability to respond to T-independent antigens and imply that the maturation and function of F_1 female B cells can proceed normally in the environment of the irradiated abnormal host. Similarly, they show that cells from the abnormal donor do not develop normally even when placed in a normal (although irradiated) environment.

Reconstitution of Poly I-C and DNP-Lys-Ficoll Response in Nonirradiated $CN \times DN F_1$ Male Mice. Transfers of spleen cells into irradiated recipients as described above do not rule out the possibility that an abnormal radiation-

sensitive regulatory mechanism operates in the F_1 male animals nor do they provide information about the cell type critical for restoring responsiveness upon transfer.

To approach these problems, we transferred cells from F_1 male and female donors to nonirradiated F_1 male and female recipients. In the experiment presented in Table VI, 50×10^6 male or female spleen cells were transferred to nonirradiated F_1 male recipients which were immunized on the same day with 100 μg of poly I-C. Recipients of F_1 female cells made substantial responses to poly I-C, indicating that female cells could function in the environment of an intact F_1 male mouse. When F_1 male cells were transferred to intact F_1 female recipients, no suppression of the anti-poly I-C response was obtained (data not shown) further indicating that abnormal T-cell regulation of B-cell activation was not responsible for the defect of the F_1 male.

TABLE VI
Response of CN \times DN F_1 Male Mice to Poly I-C after the Administration of CN \times DN F_1 Spleen Cells

Sex of recipient	$[^{14}\text{C}]$ poly I-C binding*	
	Female donor	Male donor
Male	%	%
	38.5	0.0
	41.3	0.0
	58.7	13.0
	44.0	0.0
	72.3	0.0
	63.5	5.0
51.6	2.4	

* Recipient CN \times DN F_1 male mice received 50×10^6 F_1 male or female spleen cells intravenously. They were immediately challenged with 100 μg of poly I-C in CFA and 14 days later the percent $[^{14}\text{C}]$ poly I-C binding by 25 μl of serum was determined.

These results were confirmed and extended by transferring varying numbers of F_1 female spleen cells to F_1 male recipients which were immunized with 100 μg of DNP-lys-Ficoll immediately. As few as 1×10^6 female spleen cells allowed males to make a detectable response; 50×10^6 cells transferred a response comparable, in terms of PFC per 10^6 spleen cells, to that of intact F_1 female mice (Fig. 1). Moreover, treatment of F_1 female spleen cells with anti- θ and C before transfer had no effect on the ability of these cells to reconstitute the response of F_1 male recipients (Fig. 2). The effectiveness of treatment with anti- θ and C in removing T lymphocytes was demonstrated by the marked impairment of the responsiveness of such cells to concanavalin A (Con A). Thus, anti- θ - and C-treated cells gave a net incorporation of $[^3\text{H}]$ thymidine of 3,081 cpm while cells treated with normal mouse serum and C incorporated 86,869 cpm in response to Con A. This result indicates that F_1 female T lymphocytes are probably not required to reconstitute the responsiveness of F_1 male mice. On the other hand, *in vitro* irradiation (1,000 R) of F_1 female spleen cells completely abolished the




FIG. 1. The number of DNP-PFC/10⁶ spleen cells in CN × DN F₁ males which were given different numbers of CN × DN F₁ female spleen cells intravenously and immunized with 100 μg of DNP-lys-Ficoll. The response to DNP-lys-Ficoll was measured on the 6th day after the administration of the female cells and immunization.

ability of these cells to transfer responsiveness to DNP-lys-Ficoll to intact F₁ male recipients. Thus, no PFC (less than background) were seen in recipients of 50×10^6 irradiated F₁ female cells while three F₁ male recipients of nonirradiated F₁ female cells had 275 ± 18 PFC/10⁶ spleen cells after immunization with 100 μg of DNP-lys-Ficoll. Further investigation of this transfer model demonstrated that 3×10^6 F₁ female lymph node cells could also transfer responsiveness to DNP-lys-Ficoll to nonirradiated F₁ male recipients, whereas 50×10^6 F₁ female thymocytes were ineffective in initiating a response (Table VII).

Reconstitution of F₁ male mice with spleen cells from F₁ female donors not only allowed responses of the recipients to immediate challenge with DNP-lys-Ficoll but, as shown in separate experiments (Table VIII), allowed responses by the males to primary challenges administered as late as 42 days after cell transfer. This indicates that transferred F₁ female B lymphocytes survive in the recipient or that B lymphocytes develop from F₁ female stem cells present in the spleen cells used for transfer.

Lack of Thymic Influence on the Reconstitution of Responsiveness in Lethally Irradiated CN × DN F₁ Male Mice. The previous studies provide convincing




FIG. 2. The number of DNP-PFC/10⁶ spleen cells in CN × DN F₁ males which were given different numbers of CN × DN F₁ female spleen cells which had been previously treated with anti- θ and C (Δ), or NMS and C (\circ). The response to DNP-lys-Ficoll was measured on the 6th day after the administration of the female cells and 100 μ g of the antigen.

evidence that F₁ female cells can transfer to both intact and irradiated F₁ male recipients the capability to respond to T-independent antigens. Moreover, since anti- θ -treated spleen cells are effective, they provide strong evidence that the defect of the F₁ male is at the B-lymphocyte level. However, it could be argued that F₁ male B lymphocytes develop abnormally in the F₁ male because of an abnormality in the F₁ male T lymphocytes which, in some way, regulate B-cell development or, alternatively, because of an abnormal sensitivity of the F₁ male B cells to normal regulatory effects of F₁ male thymus or T lymphocytes. In order to investigate this point, F₁ male mice were thymectomized at 6 wk of age, irradiated (1,000 R) at 8 wk, and reconstituted with 10×10^6 anti- θ -treated male or female bone marrow cells 1 day later. They were held for 10 wk to allow repopulation of their lymphoid system and were then immunized with 100 μ g of DNP-lys-Ficoll. Animals reconstituted with anti- θ -treated F₁ female bone marrow made excellent responses to DNP-lys-Ficoll, but those repopulated with anti- θ -treated F₁ male bone marrow were unresponsive to DNP-lys-Ficoll (Table IX). Spleen cells isolated from these F₁ male mice were cultured in the presence of Con A in order to determine if functional T lymphocytes were present. The proliferative response of these spleen cells were all less than 4% of normal F₁

TABLE VII
Response of CN × DN F₁ Male Mice to DNP-Lys-Ficoll after the Administration of CN × DN F₁ Female Spleen, Lymph Node, or Thymus Cells

F ₁ male recipient	F ₁ female donors		TNP-SRBC*	
	Cell source	No. of Cells	PFC/spleen	PFC/10 ⁶ cells
1	—	—	75	1.5
2	—	—	100	3.2
3	Spleen	3 × 10 ⁶	10,950	267.0
4	Spleen	3 × 10 ⁶	6,450	140.0
5	Spleen	10 × 10 ⁶	11,050	254.0
6	Spleen	10 × 10 ⁶	8,450	275.0
7	Lymph node	3 × 10 ⁶	5,500	149.0
8	Lymph node	3 × 10 ⁶	5,075	108.0
9	Lymph node	10 × 10 ⁶	6,650	155.0
10	Lymph node	10 × 10 ⁶	8,275	194.0
11	Thymus	50 × 10 ⁶	75	1.2
12	Thymus	50 × 10 ⁶	200	3.1
13	Thymus	50 × 10 ⁶	150	2.2

* Recipient F₁ male mice were given varying numbers of F₁ female spleen, lymph node, or thymus cells intravenously. They were then challenged with 100 μg of DNP-lys-ficoll and IgM TNP-SRBC PFC were assayed on day 6.

TABLE VIII
Response of CN × DN F₁ Male Mice to DNP-Lys-Ficoll at Different Times after the Administration of F₁ Female Spleen Cells

Days after F ₁ female cells	TNP-SRBC*	
	PFC/spleen	PFC/10 ⁶ spleen
18	1,316 ± 719	35.2 ± 13.5
25	1,550 ± 123	38.5 ± 1.9
39	3,000 ± 177	57.4 ± 1.7
46	1,900 ± 50	45.8 ± 0.5

* Recipient F₁ male mice were given 50 × 10⁶ F₁ female cells intravenously. At 14, 21, 35, and 42 days after the administration of these cells, three F₁ male mice were immunized with 100 μg of DNP-lys-Ficoll and their IgM TNP-SRBC PFC response was measured 4 days later. Data are expressed as the mean number of PFC of three mice ± SE.

male mice. Thus, it seems clear that the B-lymphocyte defect exhibited by F₁ male mice cannot be ascribed to development under the influence of abnormal T-dependent stimuli or to an abnormal response to normal T-dependent stimuli.

TABLE IX
*Response of Thymectomized, Lethally Irradiated, Bone Marrow
 Reconstituted CN × DN F₁ Male Mice to DNP-Lys-Ficoll*

Sex of recipient	Sex of donor	TNP-SRBC*	
		PFC/spleen	PFC/10 ⁶ spleen
Female‡	—	60,450	530
Male	Female	51,000	362
Male	Female	27,927	338
Male	Male	25	0.9
Male	Male	75	1.1

* Recipient F₁ mice were thymectomized at 6 wk of age, lethally irradiated (1,000 R) at 8 wk and reconstituted with 10 × 10⁶ anti- θ -treated bone marrow cells on the day after irradiation. At 18 wk of age, these mice were immunized with 100 μ g of DNP-lys-Ficoll and their IgM TNP-SRBC PFC response was assayed 4 days later.

‡ Nonirradiated normal female immunized with 100 μ g of DNP-lys-Ficoll.

Discussion

CN mice and male CN × DN F₁ mice have been previously shown to have a profound defect in immune responses to SIII, LPS, and poly I-C (1, 2). We show here that they are essentially unresponsive to DNP-lys-Ficoll. Moreover, *in vitro* studies confirm the unresponsiveness of F₁ male mice to DNP-lys-Ficoll while F₁ female mice mount vigorous responses.³ Thus, CN mice fail to respond to four different T-independent antigens. Our results with DNP-lys-Ficoll are particularly impressive as this T-independent antigen yields very large numbers of IgM and IgG PFC in a variety of normal strains (8).

On the other hand, F₁ male CN × DN mice mount substantial responses to SRBC and TNP-SRBC. When considered in terms of PFC per unit of B lymphocytes (rather than as PFC per spleen), the IgM response of F₁ males is similar to that of F₁ females. This is so because of F₁ males generally have half as many nucleated cells per spleen as F₁ females (6). In addition, F₁ males have only 26% Ig-bearing cells among spleen lymphocytes, whereas 40% of spleen lymphocytes from F₁ females bear surface Ig (6). It should be noted that considering responses of F₁ males to DNP-lys-Ficoll on this basis does not result in such a normalization, as F₁ males make no net TNP-specific PFC as a result of immunization with this T-independent antigen. These results confirm our previous conclusion that the X-linked functional immune defect expressed by CN and male CN × DN F₁ mice is principally an inability to mount T-independent responses rather than a global defect in all humoral immune responses (6).

The finding that CN and CN × DN F₁ male mice fail to respond to T-independent antigens does not, by itself, establish that the defect of these

³ Cohen, P., I. Scher, and D. Mosier. Manuscript submitted for publication.

animals resides in the B-lymphocyte pool. In order to study this problem, we evaluated the capacity of lymphoid cells obtained from F₁ female donors to transfer responsiveness to F₁ male recipients. Spleen cells from F₁ female mice were able to reconstitute an easily detectable antibody response to poly I-C or DNP-lys-Ficoll in F₁ males after immediate challenge. Moreover, both anti- θ -treated spleen cells and normal lymph node cells from F₁ female donors transferred responsiveness to F₁ males, while irradiated F₁ female spleen cells did not transfer responsiveness. The former indicates that mature T lymphocytes from the F₁ female are not required for the DNP-lys-Ficoll response; the latter experiments suggest that macrophages are not the limiting cell type, since macrophages are less frequent in lymph node cell populations than in spleen cell suspensions and are quite radiation resistant. Indeed, Mosier et al. (7) have shown that the in vitro response to DNP-lys-Ficoll is much less dependent on adherent cells than is the response to SRBC, further suggesting that macrophages are not likely to be the defective cell type in CN and male CN \times DN F₁ mice.

Since cell populations rich in mature B lymphocytes from F₁ female mice will reconstitute the response of F₁ male mice, it is likely that the X-linked functional defect in these mice resides in the B-cell pool. Nonetheless, it could be proposed that the B-cell defect results from a defect in control of B-cell development or function based upon abnormal regulatory cells or abnormal sensitivity of B cells to external regulatory influences. Several of our experiments bear on these points. Firstly, mature F₁ female spleen cells function normally in intact or irradiated males, indicating that an excess of an acutely acting regulatory factor is not present in the F₁ male. Furthermore, nonirradiated F₁ males which have received F₁ female spleen cells mount anti-DNP-responses to DNP-lys-Ficoll administered up to 42 days after transfer, making it unlikely that a suppressive factor active only over a prolonged period limits responsiveness of mature B cells in the F₁ male. Also supporting this contention is the failure of F₁ male spleen cells to inhibit the response of intact F₁ females.

These data strongly indicate that mature B lymphocytes capable of responding to T-independent antigens are absent in CN and CN \times DN F₁ mice. Such an absence could represent a defect in the differentiative potential of B-lymphocyte progenitors or, alternatively, a defect in the microenvironment in which B lymphocytes develop. Our data strongly points to the former rather than the latter. Thus, F₁ female bone marrow cells, when transferred to lethally irradiated F₁ male mice, allow the latter to respond to challenge with DNP-lys-Ficoll administered 8 wk later. This implies that female B-lymphocyte progenitors can develop normally in the environment of the F₁ male. On the other hand, F₁ male bone marrow cells transferred to lethally irradiated F₁ female mice do not reconstitute the response to DNP-lys-Ficoll of the recipients. The small response which these animals mount is equivalent to the number of background TNP-specific PFC in normal females and may represent a response on the part of surviving female B lymphocytes. Similarly, failure in development of male B cells responsive to DNP-lys-Ficoll cannot be ascribed to a regulatory influence of the male thymus or T lymphocytes since anti- θ -treated F₁ male bone marrow cells could not reconstitute responsiveness in thymectomized, lethally irradiated F₁ male recipients.

A final objection to the concept that a functional class of B lymphocytes is absent in CN and F₁ male mice could be raised, if it could be shown that the cells responding to poly I-C or DNP-lys-Ficoll in F₁ males which had received F₁ female cells were the defective F₁ male cells. According to this hypothesis, the presence of the F₁ female cells would allow the F₁ male cells to form antibody against these T-independent antigens. In order to study this problem, we have reconstituted F₁ males with T-lymphocyte-depleted DN spleen cells. In preliminary experiments, we have shown that the cell responding to DNP-lys-Ficoll was derived from the DN strain rather than the F₁ male.

Our data present convincing evidence that the X-linked defect in responsiveness to T-independent antigens by CN mice represents an intrinsic defect in the B-lymphocyte line rather than an abnormal extrinsic regulation of B-lymphocyte development and/or function. We cannot, as yet, determine whether these animals have a deletion in a single line of B lymphocytes (i.e., B cells responsive to T-independent antigens) or, alternatively, a defect in all lymphocytes which affects T-independent responses more profoundly than T-dependent responses. These mice provide a model system in which an analysis of the molecular mechanism of B-lymphocyte activation and a determination of the nature of the functional heterogeneity of such cells should be possible. In addition, they provide an excellent model for understanding the role of X-linked genes in the control of immune responses.

Summary

The mechanisms underlying the X-linked thymus-independent (B) lymphocyte functional defect in the CBA/N (CN) mice and their F₁ progeny were studied. Immune defective mice were unable to respond to the T-independent antigen 2,4-dinitrophenyl-lysyl-derivative of Ficoll (DNP-lys-Ficoll) but were able to form antibody against the highly cross-reactive hapten (trinitrophenyl) when it was coupled to an erythrocyte carrier. Immune defective CN × DBA/2N (DN) F₁ male mice, which do not normally respond to T-independent antigens, were able to respond to both polyribosinic-polyribocytidylic acid and DNP-lys-Ficoll after the administration of CN × DN F₁ female spleen cells even if these cells had been depleted of T lymphocytes.

In addition, it was shown that the inability of the CN mice and their F₁ progeny to respond to T-independent antigens was not due to an intrinsic abnormality of their microenvironment or the suppressive actions of a T lymphocyte. Our data present evidence that the X-linked defect in the CN mice is due to an intrinsic defect in B-lymphocyte development.

We would like to acknowledge the excellent editorial assistance of Mrs. Betty J. Sylvester.

Received for publication 27 May 1975.

References

1. Amsbaugh, D. F., C. T. Hansen, B. Prescott, P. W. Stashak, D. R. Barthold, and P. J. Baker. 1972. Genetic control of the antibody response to Type III pneumococcal polysaccharide in mice. I. Evidence that an X-linked gene plays a decisive role in determining responsiveness. *J. Exp. Med.* 136:931.

2. Scher, I., M. Frantz, and A. D. Steinberg. 1973. The genetics of the immune response to a synthetic double-stranded RNA in a mutant CBA mouse strain. *J. Immunol.* 110:1396.
3. Andersson, B., and H. Blomgren. 1971. Evidence for thymus-dependent humoral antibody production in mice against polyvinylpyrrolidone and *E. coli* lipopolysaccharide. *Cell. Immunol.* 2:411.
4. Davies, A. J. S., R. L. Carter, E. Leuchars, V. Wallis, and F. M. Dietrich. 1970. The morphology of immune reactions in normal thymectomized and reconstituted mice. III. Response to bacterial antigens: salmonella flagellar antigen and pneumococcal polysaccharide. *Immunology.* 19:945.
5. Chused, T. M., A. D. Steinberg, and L. M. Parker. 1973. Enhanced antibody response of mice to polyinosinic-polycytidylic acid by antithymocyte serum and its age-dependent loss in NZB/W mice. *J. Immunol.* 111:52.
6. Scher, I., A. Ahmed, D. M. Strong, A. D. Steinberg, and W. E. Paul. 1975. X-linked B-lymphocyte immune defect in CBA/HN mice. I. Studies of the function and composition of spleen cells. *J. Exp. Med.* 141:788.
7. Mosier, D. E., B. M. Johnson, W. E. Paul, and P. R. B. McMaster. 1974. Cellular requirements of the primary in vitro antibody response to DNP-Ficoll. *J. Exp. Med.* 139:1354.
8. Sharon, R., P. R. B. McMaster, A. M. Kask, J. D. Owens, and W. E. Paul. 1975. DNP-lys-Ficoll: a T-independent antigen which elicits both IgM and IgG anti-DNP antibody secreting cells. *J. Immunol.* 114:1585.
9. Scher, I., D. M. Strong, A. Ahmed, R. Knudsen, and K. W. Sell. 1973. Specific murine B-cell activation by synthetic single- and double-stranded polynucleotides. *J. Exp. Med.* 138:1545.
10. Kettman, J., and R. W. Dutton. 1970. An in vitro primary immune response to 2,4,6-trinitrophenyl substituted erythrocytes: response against carrier and hapten. *J. Immunol.* 104:1558.
11. Steinberg, A. D., T. Pincus, and N. Talal. 1971. The pathogenesis of autoimmunity in New Zealand Mice. III. Factors influencing the formation of anti-nucleic acid antibodies. *Immunology.* 20:523.
12. Mosier, D. E. 1969. Cell interactions in the primary immune response in vitro: a requirement for specific cell clusters. *J. Exp. Med.* 129:351.
13. Rittenberg, M. B., and C. Pratt. 1969. Anti-trinitrophenyl (TNP) plaque assay. Primary response of BALB/c mice to soluble and particulate immunogen. *Proc. Soc. Exp. Biol. Med.* 132:575.
14. Pierce, C. W., B. M. Johnson, H. E. Gershon, and R. Asofsky. 1971. Immune responses in vitro. III. Development of primary γ M, γ G, and γ A plaque-forming cell responses in mouse spleen cell cultures stimulated with heterologous erythrocytes. *J. Exp. Med.* 134:395.