

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

Xylem embolism refilling and resilience against drought-induced mortality in woody plants: processes and trade-offs

### Permalink

<https://escholarship.org/uc/item/5d95j9h1>

### Journal

Ecological Research, 33(5)

### ISSN

0912-3814

### Authors

Klein, T  
Zeppel, MJB  
Anderegg, WRL  
[et al.](#)


### Publication Date

2018-09-01

### DOI

10.1007/s11284-018-1588-y

Peer reviewed

Tamir Klein  · Melanie J. B. Zeppel  
William R. L. Anderegg · Jasper Bloemen  
Martin G. De Kauwe · Patrick Hudson  
Nadine K. Ruehr · Thomas L. Powell  
Georg von Arx · Andrea Nardini

## Xylem embolism refilling and resilience against drought-induced mortality in woody plants: processes and trade-offs

Received: 1 August 2017 / Accepted: 15 February 2018  
© The Ecological Society of Japan 2018

**Abstract** Understanding which species are able to recover from drought, under what conditions, and the mechanistic processes involved, will facilitate predictions of plant mortality in response to global change. In response to drought, some species die because of embolism-induced hydraulic failure, whilst others are able to avoid mortality and recover, following rehydration. Several tree species have evolved strategies to avoid embolism, whereas others tolerate high embolism rates but can recover their hydraulic functioning upon drought relief. Here, we focus on structures and processes that might allow some plants to *recover* from drought stress via embolism reversal. We provide insights into how embolism repair may have evolved, anatomical and physiological features that facilitate this process, and describe possible trade-offs and related costs. Recent controversies

on methods used for estimating embolism formation/repair are also discussed, providing some methodological suggestions. Although controversial, embolism repair processes are apparently based on the activity of phloem and ray/axial parenchyma. The mechanism is energetically demanding, and the costs to plants include metabolism and transport of soluble sugars, water and inorganic ions. We propose that embolism repair should be considered as a possible component of a ‘hydraulic efficiency-safety’ spectrum. We also advance a framework for vegetation models, describing how vulnerability curves may change in hydrodynamic model formulations for plants that recover from embolism.

**Keywords** Hydraulic conductivity · Plant hydraulics · Plant water relations · Recovery · Repair

---

T. Klein and M. J. B. Zeppel contributed equally to this work.

T. Klein (✉)  
Department of Plant and Environmental Sciences, Weizmann  
Institute of Science, 76100 Rehovot, Israel  
E-mail: tamir.klein@weizmann.ac.il  
Tel.: 972-8-9343505

M. J. B. Zeppel  
GenIMPACT National Centre for Genomic Medicine, Faculty of  
Business and Economics, Macquarie University, Sydney, NSW  
2107, Australia

W. R. L. Anderegg  
Department of Biology, University of Utah, Salt Lake City, UT  
84112, USA

J. Bloemen  
Institute of Ecology, University of Innsbruck, Innsbruck, Austria

M. G. De Kauwe  
ARC Centre of Excellence for Climate Extremes, University of  
New South Wales, Sydney, NSW 2052, Australia

P. Hudson  
Department of Biology, University of New Mexico, Albuquerque,  
NM, USA

N. K. Ruehr  
Karlsruhe Institute of Technology (KIT), Institute of Meteorology  
and Climate Research-Atmospheric Environmental Research  
(IMK-IFU), 82467 Garmisch-Partenkirchen, Germany

T. L. Powell  
Department of Organismic and Evolutionary Biology, Harvard  
University, Cambridge, MA, USA

T. L. Powell  
Earth and Environmental Sciences Area, Lawrence Berkeley Na-  
tional Laboratory, Berkeley, CA, USA

G. von Arx  
Swiss Federal Institute for Forest, Snow and Landscape Research  
WSL, Birmensdorf, Switzerland

A. Nardini  
Dipartimento di Scienze della Vita, Università di Trieste, Via L.  
Giorgieri 10, 34127 Trieste, Italy

---

## Introduction

### How plants avoid mortality: resistance and recovery strategies

The frequency, duration and intensity of drought events are expected to increase in the coming decades in many regions of the globe (Kirtman et al. 2014). Why some plants die under drought stress while other co-occurring species resist or recover from stress remains an unresolved topic of recent investigation (e.g. McDowell et al. 2011; Schenk 2014). Plant hydraulic failure (catastrophic loss of hydraulic conductivity) is one of the major and widespread mechanistic causes of drought-induced mortality (Anderegg et al. 2012; Nardini et al. 2014). Much research has investigated how plants avoid hydraulic failure by preventing formation and spread of catastrophic xylem embolism (Brodribb and Cochard 2009; Pittermann et al. 2013). However, recent research suggests the existence of a spectrum of strategies ranging from ‘resistance’ to ‘recovery’, with some plants coming remarkably close to hydraulic failure, but recovering once xylem tension is partially or fully relaxed (Ogasa et al. 2013; Martorell et al. 2014). Indeed, many woody plants commonly operate at xylem pressures close to critical thresholds for runaway embolism (i.e. with narrow safety margins) (Choat et al. 2012). This suggests that many species routinely suffer the risk of reaching high levels of xylem embolism, with angiosperms tending to have smaller safety margins than gymnosperms. A high loss of conductivity has been reported also in drought-surviving individuals (> 50%) of different species, hinting at a possible survival strategy based on post-drought hydraulic recovery, rather than a strategy relying on complete hydraulic safety (Klein et al. 2011; Trifilò et al. 2015).

Woody plants can refill embolised xylem conduits via spontaneous bubble dissolution once xylem pressure nears atmospheric values, or under positive root/stem pressure (Nardini et al. 2018). Trees can also repair the total conducting area following seasonal replacement through regrowth of new xylem (Brodersen and McElrone 2013; Nardini et al. 2014). However, debate remains over how and when woody plants can refill embolised xylem conduits under residual negative water potential, so called “novel refilling” (Hacke and Sperry 2003; Brodersen and McElrone 2013) and which methods are appropriate for studying embolism formation and its reversal in plants (Rockwell et al. 2014). The ‘novel refilling’ phenomenon is very controversial, and some researchers have argued that embolism refilling under any residual negative pressure in still functional xylem is unlikely or impossible due to thermodynamic constraints (Zwieniecki and Holbrook 2000; Rockwell et al. 2014; but see also Gouin 2015). Some researchers have also suggested that methodological uncertainties call for a thorough re-evaluation of current evidence for embolism reversal (Cochard and Delzon 2013; Torres-

Ruiz et al. 2015; Charrier et al. 2016). However, the available evidence around this phenomenon suggests that embolism reversal might be an important vital process essential for plant hydraulic functioning in some or even several plant species (Nardini et al. 2011a). Hence, understanding the process of hydraulic recovery, and why it occurs in some plants and not in others, would improve the robustness of predictions of plant mortality in future climates.

In this paper, we synthesise recent experimental studies on embolism reversal, focusing on currently available methods and techniques to investigate the phenomenon, and related possible pitfalls. Then, we describe structures and processes possibly facilitating embolism reversal, followed by a conceptual framework adding ‘recovery’ to the safety-efficiency spectrum, and finally present a modelling framework to evaluate if, when and where xylem refilling may be an important ecological process.

### Defining recovery

Plants frequently experience physiological stress from disturbance events including heat waves, drought, fire, or freezing. When of extreme duration and/or intensity, such stress factors can result in significant damage eventually leading to death. However, following disturbance, plants can avoid mortality by repairing damage, thus experiencing either partial or complete recovery. Post-stress recovery has been observed at the leaf, tree and landscape scale and can take from days to weeks (Brodribb et al. 2010), months to years (Zeppel et al. 2008), or even decades (Vertessy et al. 2001).

Full recovery may be defined as when ‘pre-disturbance’ levels in defined parameters are reached. From a structural viewpoint, complete recovery of biomass or leaf area index (LAI) after disturbance might represent convenient reference points (Kelley et al. 2014). Alternatively, hydraulic recovery may be defined as the point when plants reach pre-disturbance levels of transpiration, stomatal conductance (Brodribb et al. 2010), or hydraulic conductivity (Ogasa et al. 2013). A trade-off between embolism resistance and refilling ability, mediated by contrasting wood functional traits such as dense and embolism-resistant xylem *versus* less dense but refillable xylem has been reported (Ogasa et al. 2013; Trifilò et al. 2015). Following severe drought, different species may use different strategies and take different times to recover their pre-drought hydraulic status (Brodribb et al. 2010). Some recent studies describe how plants may use recovery from embolism as a mortality avoidance, or resilience strategy. However, the actual trade-offs between wood structural traits, xylem vulnerability and recovery ability among different tree species are not fully understood (Johnson et al. 2012). We now discuss potential trade-offs in mortality avoidance strategies.

---

## Mortality avoidance strategies: resisting drought stress

Across time and spatial scales, short-term responses and long-term adaptations to avoid mortality

Plants can adopt different strategies for avoiding drought-induced mortality at different time-scales and organisational levels (see Nardini et al. 2014). For example, over the short-term, stomata close and plants rely on stored carbohydrates to fuel metabolism (Klein et al. 2014). Some species adjust their hydraulic efficiency both at the apoplastic level ('ionic effect'; Nardini et al. 2011b) and the symplastic level through aquaporins (Javot and Maurel 2002) to reduce resistance of water transport and buffer xylem pressure drops. Over the long-term, an increased root-to-shoot biomass ratio can also accomplish this task (Klein et al. 2011). Alternatively, water-use can be decreased in some species by shedding leaves (Tyree et al. 1993) or branches (Rood et al. 2000). Some species shift activity to periods when sufficient water is available (Nardini et al. 1998). Finally, disrupting the hydraulic path between the leaf and stem (Zufferey et al. 2011) or root and soil interface in some species avoids water loss and very low water potentials (Carminati et al. 2013). Over medium time scales (months–years), adjustments in the xylem anatomy are central to many drought avoidance strategies. For example, a reduction in conduit diameter is commonly observed during periods of acute water stress (von Arx et al. 2012), although opposite trends have also been reported (Bouche et al. 2014). Adjustments in pit characteristics may allow increased resistance to drought-induced embolism (Lens et al. 2011). The thickness of conduit cell walls also apparently adjusts in response to drought stress, with thicker walls providing increased mechanical strength to withstand lower water potential and prevent cell implosion (Hacke et al. 2001).

### The occurrence and significance of embolism recovery

Is recovery from xylem embolism unique to a small set of tree species, from specific biomes, or is it rather ubiquitous? In the past few years research on xylem refilling has been rapidly increasing: the number of scientific papers containing “xylem refilling” has increased from 26 published in 2011 to 80 published in 2016. Since the last review by Brodersen and McElrone (2013) several additional species have been investigated (Table 1; Fig. 1). Overall, xylem refilling emerges as a cross-species, cross-biome phenomenon. In 33 of 43 species surveyed here, the level of native embolism decreased significantly with time following peak drought, suggesting the occurrence of refilling mechanisms. Three additional species showed visually rapid refilling, which was not directly quantified (Table 1). Most of the studied species showing evidence of embolism reversal were angiosperm trees, but observations included key gym-

nosperms such as *Picea abies*, *Picea glauca*, and *Pinus halepensis*, as well as shrubs, lianas, and cushion plants (Table 1). Almost two-thirds of the observations were in mature woody plants, and 78% were done on field-grown plants. Remarkably, 54% of the case studies did not include rewatering of the plants (Table 1), indicating that embolism formation and reversal might be part of a routine physiological process in some species, rather than a post-drought occurrence. In all studied species, the refilling process was accompanied by partial relaxation of xylem tension, observed by concomitant increase in shoot water potential, typically to levels above  $-1.0$  MPa, but still moderately negative (Fig. 1). The literature survey compiled here is still too small to draw conclusions on the partitioning of refilling among species and biomes. Yet, it is intriguing that species from temperate ecosystems had relatively high levels of embolism (37–94%), and typically showed the most efficient recovery (PLC, percent loss of conductivity, decreasing overnight by about 40%). Conversely, many species from Mediterranean ecosystems did not recover or showed relatively mild recovery (PLC decreased overnight by about 20%). This observation might relate to eco-physiological processes described below. More importantly, the 10 cases where no refilling was observed were mostly those where a decrease in PLC was mild or simply not statistically significant. For example, PLC in *Carpinus tschonoskii* stems decreased from 67% at  $-3.9$  MPa to 48% at  $-0.2$  MPa in one experiment (Ogasa et al. 2013), and from 70% at  $-3.3$  MPa to 35% at  $-0.2$  MPa in another experiment (Umebayashi et al. 2016); only the latter observation was significant. Therefore, we cannot exclude the possibility that almost all woody plants are capable of xylem refilling to some extent.

### How do we measure embolism recovery or refilling?

Measurements of xylem embolism and subsequent recovery are often restricted by our limited ability to directly observe in vivo the functional status of xylem conduits. Hence, much evidence comes from indirect observations, using destructive and non-destructive methods. Here, we describe their strengths and weaknesses.

#### Destructive measurements

Destructive measurements of xylem hydraulics can potentially introduce a change to the native status of conduits, thus altering the object of study (Rockwell et al. 2014). However, higher replication possibilities and lower equipment costs make these methods among the most accessible and used in different laboratories. During the past recent decades, measurements of hydraulic conductivity on branch sections became the state-of-the-art methodology for embolism detection and calculation

**Table 1** Study cases on xylem refilling published since 2013, divided by biome and woody plant species. Source references are: (1) Leng et al. (2013); (2) Zwieniecki et al. (2013); (3) Ooeda et al. (2016); (4) Ogasa et al. (2013); (5) Umebayashi et al. (2016); (6) Martorell et al. (2014); (7) Christman et al. (2012); (8) Choat et al. (2015); (9) Laur and Hacke (2014); (10) Mayr et al. (2014); (11) Christensen-Dalsgaard et al. (2014); (12) Rolland et al. (2015); (13) Ganthaler and Mayr (2015); (14) Trifilo et al. (2015); (15) Savi et al. (2016); (16) Klein et al. (2016); (17) Li et al. (2016); (18) Chen et al. (2017)

Species	Setting	Plants	PLC50	Method	Rewatering?	Refilling?
<b>Temperate</b>						
<i>Populus alba</i> × <i>P. glandulosa</i> (1)	G	Seedlings	NA	HHF	Y	Y
<i>Acer rubrum</i> (2)	G	2, 20 years	- 2.70	MRI	N	Y(*)
<i>Morus australis</i> (3)	G	Potted plants	NA	HHF	Y	Y
<i>Salix integra</i> (4)	EF	1 year	- 1.08	HHF	Y	Y
<i>Salix Babylonica</i> (4)	EF	1 year	- 1.04	HHF	Y	Y
<i>Salix Cheilophila</i> (4)	EF	1 year	- 1.70	HHF	Y	Y
<i>Salix Psammophila</i> (4)	EF	1 year	- 1.18	HHF	Y	Y
<i>Betula platyphylla</i> (4)	EF	2-3 years	- 1.96	HHF	Y	Y
<i>Cerasus jamasakura</i> (4)	EF	2-3 years	- 3.62	HHF	Y	N
<i>Carpinus tschonoskii</i> (4)	EF	2-3 years	- 3.88	HHF	Y	N
<i>Carpinus tschonoskii</i> (5)	G	2 year potted plants	NA	C-SEM, MRI	Y	Y
<i>Eucalyptus pauciflora</i> (6)	G	2 year potted plants	- 1.61	HHF	Y	Y
<i>Quercus gambelii</i> (7)	F	5-8 years	- 1.08	HHF	N	Y
<i>Sequoia sempervirens</i> (8)	G	Seedlings	- 1.80	uCT	Y	N(**)
<b>Boreal/Alpine/Frost</b>						
<i>Picea glauca</i> (9)	G	3 years saplings	- 4.30	HHF	N	Y
<i>Picea abies</i> (10)	F	Mature trees	- 3.98	HHF	Y	Y
<i>Malus domestica</i> (11)	F	Mature trees	- 3.95	Cavitron	N	Y
<i>Populus deltoides</i> × <i>P. laurifolia</i> × <i>xP. Nigra</i> (11)	F	Mature trees	- 1.70	Cavitron	N	Y
<i>Salix dasyclados</i> (11)	F	Mature trees	- 1.70	Cavitron	N	Y
<i>Azorella macquariensis</i> (12)	F	Mature cushion plants	NA	BFM	Y	Y(***)
<i>Colobanthus muscooides</i> (12)	F	Mature cushion plants	NA	BFM	Y	Y(***)
<i>Vaccinium myrtillus</i> (13)	F	Mature dwarf shrubs	- 2.08	HHF	Y	Y
<i>Vaccinium vitis-idaea</i> (13)	F	Mature dwarf shrubs	- 1.97	HHF	Y	Y
<b>Mediterranean</b>						
<i>Ceratonia siliqua</i> (14)	F	Adult trees	- 2.60	HHF	N	Y
<i>Laurus nobilis</i> (14)	F	Adult trees	- 2.50	HHF	N	Y
<i>Nerium oleander</i> (14)	F	Adult trees	- 1.50	HHF	N	Y
<i>Pistacia lentiscus</i> (14)	F	Adult trees	- 4.10	HHF	N	N
<i>Quercus ilex</i> (14)	F	Adult trees	- 3.30	HHF	N	N
<i>Eucalyptus camaldulensis</i> (14)	F	Adult trees	- 4.60	HHF	N	Y
<i>Myrtus communis</i> (14)	F	Adult trees	- 3.10	HHF	N	N
<i>Olea europaea</i> (14)	F	Adult trees	- 2.10	HHF	N	Y
<i>Olea oleaster</i> (14)	F	Adult trees	- 4.10	HHF	N	N
<i>Quercus pubescens</i> (15)	G	Potted plants	NA	HHF	Y	Y
<i>Prunus mahaleb</i> (15)	G	Potted plants	NA	HHF	Y	N
<i>Robinia pseudoacacia</i> (15)	G	Potted plants	NA	HHF	Y	N
<i>Ailanthus altissima</i> (15)	G	Potted plants	NA	HHF	Y	Y
<b>Semi-arid</b>						
<i>Pinus halepensis</i> (16)	F	40 years	- 3.10	HHF	N	Y
<i>Salix psammophila</i> (17)	F	Adult shrubs	- 1.70	HHF	N	Y
<i>Caragana korshinskii</i> (17)	F	Adult shrubs	- 2.70	HHF	N	N
<b>Tropical</b>						
<i>Celtis tetrandra</i> (18)	EF	Mature trees	- 1.53	HHF	N	Y
<i>Ficus concinna</i> (18)	EF	Mature trees	- 1.43	HHF	N	Y
<i>Harpullia cupanioides</i> (18)	EF	Mature trees	- 2.51	HHF	N	N
<i>Michelia hypolampra</i> (18)	EF	Mature trees	- 2.25	HHF	N	Y
<i>Streblus asper</i> (18)	EF	Mature trees	- 2.93	HHF	N	Y
<i>Celastrus paniculatus</i> (18)	EF	Mature lianas	- 1.42	HHF	N	Y
<i>Marsdenia sinensis</i> (18)	EF	Mature lianas	- 1.04	HHF	N	Y
<i>Ventilago calyculata</i> (18)	EF	Mature lianas	- 1.57	HHF	N	Y


Numbers in parentheses indicate the reference number of the respective study in the reference list. The observed changes in embolism level for each species are in Fig. 1, except where not quantified: (\*) refilling visually significant after 30 min; (\*\*) no refilling observed; and (\*\*\*) bubbles removed in 50 s

Setting: EF experimental field, F field, G greenhouse. PLC50: The shoot water potential (MPa) at 50% percent loss of xylem conductivity. Method: BFM bright-field microscopy, HHF hydraulic head and flushing, MRI magnetic resonance imaging, C-SEM cryo-scanning electron microscopy, uCT micro computed tomography

of PLC, i.e. the relative change in xylem conductivity between native and water-saturated states, usually taken as a proxy for the embolism level. Although still implicit

and invasive to a certain degree, hydraulic conductivity measures directly the consequence of embolism, and the technique has been the basis for early descriptions of the





**Fig. 1** Embolism dynamics in woody plant species reported in the scientific literature between 2013 and 2017 (for more details see Table 1). Note the logarithmic scale in the x axes. Numbers in parentheses indicate the shoot water potentials (MPa) measured at the start and the end of the observation period, respectively. Embolism dynamics exclude changes related to new xylem growth, even in the longer observation periods (decrease in PLC, percent loss of conductivity) is evident in 33 of the 43 species (Colour figure online)

embolism reversal phenomenon (Salleo et al. 1996). Measurements of PLC can be made on (1) samples collected in the field at different daytime/season, (2) samples from branches air-dehydrated in the lab, (3) stem segments where embolism is induced through air-injection, or (4) by inducing tension in the xylem by centrifugal forces.

#### *Methodological precautions*

When using hydraulic methods to quantify PLC, there are some restrictions set by inherent xylem properties (e.g. resin excretion in some conifers), and by sampling procedures. Three possible pitfalls have been proposed:

- (1) When long-vesselled branches are measured using air-injection or cavitron methods, it is suggested that the ratio of open vessels in the sample must be quantified and removed from the PLC calculation (Cochard and Tyree 1990), as some authors have reported a long-vessel artefact in olives (Torres-Ruiz et al. 2014) and grapevines (Choat et al. 2010). However, we note that the long-vessel artefact has been rebutted by other studies (Sperry et al. 2012; Hacke et al. 2014), and this issue remains unresolved.
- (2) To avoid artefactual embolism induction caused by the cutting of a branch, while the xylem is under tension, rehydration/relaxation of samples has been suggested, followed by re-cutting under degassed water (Wheeler et al. 2013), although this artefact has not been found by others (Venturas et al. 2015). Alternatively, the relaxation procedure has been shown to introduce other potential errors related to rehydration-induced embolism refilling (Trifilò et al. 2014b). Recommendations for standardizing measurements of vulnerability curves have recently been described (Torres-Ruiz et al. 2014). Future research should carefully evaluate the occurrence and magnitude of cutting or relaxation artefact in species investigated for their refilling capability. Trifilò et al. (2014b) suggested the use of refilling inhibitors (e.g. orthovanadate) to investigate the impact of cutting and relaxation on PLC values. A careful re-evaluation of PLC recovery in species already tested in previous studies would also help to assess the eventual impact of experimental artefacts in trends reported in Fig. 2.
- (3) In conifer branches where torus margo sealing occurs in the pits when subjected to high pressure, embolism removal to quantify maximum hydraulic conductivity should be achieved by long (e.g. overnight) vacuum-infiltration (Delzon et al. 2010), and not by high pressure that might lead to pit membrane sealing and prevent embolism displacement. Finally, it is critical that studies report not only the PLC, but also either the sapwood area-specific conductivity or leaf-specific conductivity and ideally

the area under the vulnerability curve, which will enable readers to compare absolute values of branch conductivity across studies and temporal scales.


#### *Early non-destructive methods*

A variety of non-destructive, indirect methods have been used to quantify changes in hydraulic conductivity upon drought stress and recovery. We define them here as non-destructive in the sense that they permit focused analysis of undisturbed part of the xylem. Patterns of sap flow coupled to measurements of water potential gradients, acoustic emissions, and sapwood water content have been used to infer changes in the relative level of embolism (see Brodersen and McElrone 2013 and references below). A caveat to some of the non-destructive methods is that observed changes can be interpreted in relation to a number of environmental factors, including changes in light intensity or temperature, which might result in hydraulic properties that do not necessarily represent the result of embolism formation or refilling.

An exception to the aforementioned methods is the measurement of acoustic emissions, developed to detect embolism. Despite early successes (Pena and Grace 1986), acoustic emissions were not fully consistent with other methods, and were advised to be used as a qualitative, complementary method (Jackson and Grace 1996). Recent advances have solved some of the issues associated with interpretation of signals with promising results (Nolf et al. 2015).

#### *Recent non-destructive methods*

The increasing interest in the physiology of embolism and plant water transport has led to the use of advanced technologies, including Magnetic Resonance Imaging MRI (Holbrook et al. 2001; Nolf et al. 2015), X-ray microtomography (Cochard et al. 2015), and cold neutron radiography (Tötzke et al. 2013). Validation and results from some of these newer techniques remain under debate (Rockwell et al. 2014). High resolution imaging of xylem embolism has been enabled by cryogenic scanning electron microscopy for over a decade (cryo-SEM; Utsumi et al. 1998), but this procedure laborious and not free of artefacts (Cochard et al. 2000). Nuclear MRI has been used to monitor water flow at high spatial and temporal resolutions in vivo (Fukuda et al. 2015). X-ray microcomputed tomography (microCT), likely the most promising technique in terms of resolution and with new desktop devices on the market, has been used to assess embolism dynamics in near real-time using 3D time-lapse imaging (Brodersen 2013). The microCT technique has provided evidence for embolism refilling in the xylem of grapevine (Brodersen and McElrone 2013) but also revealed no novel refilling oc-



**Fig. 2** A schematic diagram of processes involved in the recovery from embolism, for species which have the ability to refill (see “Hydraulic and embolism recovery”). If water is present and the osmotic potential is more negative than the water potential, due to the ionic effect, then water may be drawn into embolised conduits and refill them. Chronic stress and cavitation fatigue may result if water remains limiting. Source references include (but are not

limited to) (1) Secchi and Zwieniecki (2010); (2) Laur and Hacke (2014) Chitarra et al. (2014); (3) Jansen et al. (2011), Nardini, Salleo and Jansen (2011), Trifilo et al. (2014); (4) Secchi and Zwieniecki (2010); (5) Grottsch et al. (2013), Laur and Hacke (2014) and Mayr et al. (2014); (6) Sperry and Love (2015); (7) Brodersen and McElrone (2013); (8) Anderegg (2015) and Hacke et al. (2001) (Colour figure online)

occurred in coastal redwoods, the *Sequoia* (Choat et al. 2015). Interestingly, *Sequoia* is a resprouter (Zeppel et al. 2015), which may be an alternative or complementary recovery strategy to embolism refilling. Indeed, much of our knowledge of embolism refilling has been conducted on only a few woody species, including grapevines (Choat et al. 2010; Zufferey et al. 2011; Chitarra et al. 2014), Bay laurel (Trifilo et al. 2014a) and poplar (Bloemen et al. 2016). Clearly, additional studies on different species and plant functional types are urgently required to understand forest vulnerability and resilience to extreme droughts, probably beyond the capacity of direct imaging techniques. In fact, these methods require considerable time and budget investment and should hence focus on in-depth questions of embolism dynamics in selected woody species. Moreover, many of the non-invasive studies have low replication, measuring one tree in one or two species, and generally on very young trees whose xylem properties may not reflect

mature individuals (McDowell et al. 2013). Still, it is very important that simple and affordable techniques, like the hydraulic ones, are validated with in vivo imaging, to be confident about the output of large-scale hydraulic surveys of embolism formation/reversal.

Three recent studies have cross-validated hydraulic measurements of PLC versus estimates of loss of conductivity based on direct microCT observation of the functional status of xylem conduits in *Laurus nobilis* (Nardini et al. 2017), *Helianthus annuus* (Savi et al. 2017), and *Eucalyptus camaldulensis* (Nolf et al. 2017), reporting substantial agreement between the two techniques. Moreover, Fukuda et al. (2015) have validated hydraulic measurements of PLC in *Cercidiphyllum japonicum* and *Betula platyphylla* var. *japonica* via direct MRI observations of embolism rate in intact seedlings. Further comparisons of different methods, within the same laboratory, using the same plant material, would be a step towards methodological consensus. However,



it is not clear to what extent microCT techniques can be considered as reliable instruments to detect vital processes, like embolism reversal, apparently based on the activity of living parenchyma cells (see below). In fact, the X-ray dose supplied during the microCT scans might be high enough to damage cells (Charrier et al. 2016, Savi et al. 2017) and possibly impede embolism reversal.

---

### Hydraulic and embolism recovery: which structures and processes facilitate conduits refilling?

We categorize hydraulic recovery as occurring under three different circumstances: (1) growth of new xylem (Christensen-Dalsgaard and Tyree 2014); (2) refilling from spontaneous bubble dissolution when xylem pressure nears atmospheric values (Brodersen and McElrone 2013) or becomes positive under root pressure regulation (see Brodersen and McElrone 2013); (3) embolism repair while the bulk of xylem is still under tension (Nardini et al. 2014). While much agreement exists around the first two possibilities, debate remains about the occurrence and mechanisms of the third phenomenon, and around methods used to detect and quantify the refilling process. Here, we discuss structures and processes which are thought to enable embolism reversal (see Fig. 2). We then discuss evidence for and against so-called embolism reversal under tension, here referred to as ‘novel refilling’ (Brodersen and McElrone 2013). Two defining features of novel refilling are that it occurs on short time scales (hours to over-night) and while still functional xylem is under negative pressure. Whether embolism reversal occurs will depend on (1) species-specific ability to refill embolized conduits, (2) the degree of soil rewetting after rain (and the resulting post-drought residual xylem tension), and (3) the severity of drought stress the plant has experienced. It is currently unclear how much xylem tension must relax in each refilling-species before novel refilling commences, and whether there are maximum PLC levels that cannot be reversed at all. However, most studies to date have reported embolism reversal only when xylem tension rises above  $-0.5$  MPa.

Various structures, including roots, phloem, xylem, parenchyma, aquaporins, and xylary chloroplasts have been mechanistically linked with embolism refilling. Current research suggests that embolism refilling is based on osmotic gradients generated by sugars and ions released in refilling conduits, and driving water inflow via intact xylem, phloem and parenchyma (Fig. 2). In this scenario, starch degradation appears to play a key role in providing the osmotica (Salleo et al. 2004; Secchi and Zwieniecki 2012), while aquaporins are required to reduce hydraulic resistance along the symplastic pathway (Mayr et al. 2014). Phloem transport has also been demonstrated to be involved in embolism reversal, as different studies showed that phloem girdling inhibits the process (Salleo et al. 2004; Nardini et al. 2014).

Xylem refilling may vary with stem age and position. Vessels close to the bark have been reported to refill faster than inner ones in grapevine segments (Wang et al. 2013), and current year xylem showed refilling over 12 h in *Betula platyphylla* whereas older xylem remained embolized (Ogasa et al. 2010). Similarly, while rapid refilling was observed in young wood of *Acer rubrum*, older wood showed no signs of recovery (Zwieniecki et al. 2013).

Axial and ray parenchyma have been suggested to influence embolism reversal (see Plavcova and Jansen 2015, for a review of xylem anatomy). Evidence for a role of parenchyma in providing water and carbohydrates have been reported in many studies. In particular, the role of living cells as sources of sugars transported into the embolized vessels has been supported by some studies (Salleo et al. 2004; Trifilò et al. 2014a). While axial parenchyma is more abundant in angiosperm species, ray parenchyma is present in most species and shows a high integration with the xylem conduits that may ensure that each conduit is connected to living cells (Fonti et al. 2015). A review of the amount of ray parenchyma in 38 gymnosperms and 54 angiosperms, found a mean of 7% of sapwood in gymnosperms and 17% in angiosperms (Myer 1922).

Aquaporin channels, which facilitate the transport of water through membranes, are also reported to influence embolism-refilling rates (Chitarra et al. 2014; Laur and Hacke 2014). Specifically, plasma membrane intrinsic proteins (PIP) are often up-regulated before or during refilling. Secchi and Zwieniecki (2010) reported that *Populus trichocarpa* showed considerable up-regulation of the PIP1 subfamily of aquaporin water channels, and down-regulation of the same isoform was detrimental to recovery (Secchi and Zwieniecki 2014). In grapevines, two aquaporin genes were activated on stress or refilling, suggesting a specific role in xylem embolism refilling (Chitarra et al. 2014). Finally, xylary chloroplasts have been observed in close proximity to xylem vessels and were suggested to play a role in refilling of embolism, in Australian mangrove species (Schmitz et al. 2012) and in poplar (Bloemen et al. 2016).

### Processes

Osmotic gradients generated by the release of ions (Tyree et al. 1999) and sugars in embolized conduits are thought to be fundamental for embolism refilling (Nardini et al. 2011a). Refilling conduits have more negative osmotic potential than still functioning ones (Secchi and Zwieniecki 2012). In a hypothetical scenario that found partial experimental confirmation, drought-induced rise in abscisic acid (ABA) concentration would lead to up-regulation of several genes in vessel-associated parenchyma, including those coding for aquaporins and enzymes related to carbohydrate metabolism and starch degradation (Secchi and Zwieniecki 2010). An increased sugar concentration in conduits would create a local positive pressure, promoting phloem unloading to par-

enchyma cells that would become strong sinks for both carbohydrates and water (Nardini et al. 2011b). The described mechanism requires the availability of water, possibly at a sufficient free energy level, and active metabolism and phloem functioning. Related research questions which can shed light on the refilling mechanism are: (1) is there a water potential threshold below which refilling is impossible? (2) Are turgor gradients in the phloem linked to refilling capacity? (3) Is there a threshold concentration of non-structural carbohydrates (NSC) below which refilling is completely impaired? (4) How are NSC concentrations linked to an eventual 'xylem pressure threshold'? Moreover, repeated embolism events have been shown to lead to cavitation fatigue (or air-seeding fatigue, i.e. the inability to repeatedly refill vessels) and may result in death many years after drought (Anderegg 2015), according to the chronic stress hypothesis (Sperry and Love 2015). However, it is not known whether the recovery ability of plants can also undergo significant 'fatigue' over repeated drought events, as a function of tissue damage and progressive depletion of NSC reserves.

Foliar water or fog uptake may also be involved in embolism refilling (Gotsch et al. 2014; McCulloh et al. 2014). Foliar water uptake was reported when relative humidity was high, providing a potential source of water in *Picea abies* (Mayr et al. 2014), *Drimys brasiliensis* (Eller et al. 2013) and *Picea glauca* (Laur and Hacke 2014).

How does xylem refilling relate to xylem vulnerability (P50) among woody plant species?

By contrasting species-specific data on xylem refilling capacity and xylem vulnerability (Brodersen and McElrone 2013; Choat et al. 2012 respectively), we raise the hypothesis that species where xylem refilling has been identified have relatively higher (less negative) values of P50 (water potential at 50% loss of conductivity) than others. A preliminary analysis, using species where data were available to date, for both xylem refilling capacity and P50 values, showed that P50 was on average 0.8 MPa higher in refiller vs. other species ( $P = 0.02$ , 11 refillers and 212 others). Two experimental studies found a positive correlation between xylem recovery and P50 in eight (Ogasa et al. 2013) and nine woody species (Trifilò et al. 2015), supporting the hypothesis. We note however two important caveats: (1) there are very few data on species that do not refill, therefore more data are required to test this hypothesis, and (2) different methods have been used to measure P50 across studies (Brodersen and McElrone 2013). Although no thorough conclusions can be drawn from these data, we present this as a hypothesis to be tested once data on additional species are available. Thus, future research is needed to describe genera/species that do not refill, and compare the structure of these species with species that do refill under comparable conditions (Trifilò et al. 2015).

---

## A new framework: safety–efficiency–recovery


The hydraulic safety–efficiency framework involves a trade-off between the ability to (1) maximise water transport, known as high 'efficiency', and (2) avoiding high levels of xylem embolism, known as 'safety' (McCulloh et al. 2014). High efficiency involves plants transporting water quickly, while high safety entails the ability to withstand very negative water potentials, which may also decrease conductivity (Brodribb and Holbrook 2004). Wood capacitance can also buffer against transient drought and increase hydraulic safety (McCulloh et al. 2014). Here we introduce recovery as an additional dimension, where plants can be both relatively efficient and resilient against hydraulic failure due to their ability to repair embolism (Fig. 3). We hypothesize that species that use high-risk hydraulic strategies would also have higher ability to refill embolism (see Fig. 3). In contrast, species that are more conservative, with less risky hydraulic strategies may not need the ability to refill embolism (Ogasa et al. 2013).

Dry biomes contain coexisting species with contrasting drought resilience strategies, and as such, are likely to represent habitats where some species use embolism refilling. One example is a mixed thermo-Mediterranean forest. *Quercus calliprinos* is a sclerophyllous oak species that maintains high gas-exchange rates throughout the 5–7 month dry season and often reaches water potentials below  $-4.0$  MPa (Klein et al. 2013). This species is, however, supplied with water from deep roots. The co-existing cypress *Cupressus sempervirens* has limited capacity to increase stomatal conductance, but has a high xylem resistance to embolism ( $P50 = -10.4$  MPa), and extends its activity range to lower water potentials. *Pinus halepensis* is another co-existing conifer, characterized by relatively low stomatal conductance and a vulnerable xylem ( $P50 = -3.1$  MPa). Living in this dry environment, *P. halepensis* is inevitably exposed to xylem embolism (Klein et al. 2011). Among the oak, cypress, and pine, embolism is routine only in *P. halepensis*, where both embolism formation and recovery have been observed (Klein et al. 2013). This Mediterranean study provides an example of how embolism refilling may be one of the strategies that one, but not all of the co-occurring species, use to tolerate drought. Further studies like this that examine a range of hydraulic traits will improve our understanding of which species and traits are correlated with embolism refilling.

---

## Modelling hydraulic recovery in vulnerability curves

A consideration of hydrodynamics has a long history of development in stand-scale models (e.g. Williams et al. 1996; Bohrer et al. 2005), ecosystem models (Duursma and Medlyn 2012; Christoffersen et al. 2016), and terrestrial biosphere models (Xu et al. 2016). However, to



**Fig. 3** A hypothesized conceptual framework proposing how embolism refilling may be an additional strategy to the continuum of hydraulic safety and hydraulic efficiency. For example, plants may have low safety, and a high ability to recover from embolism (grey ellipse). Note that capacitance, which acts as a buffer against embolism, may be regarded as one aspect of avoidance, and is presented here as an orthogonal axis, representing an additional

strategy. We hypothesize that species may be able to refill embolism, particularly if they are high water users (upper right box). Species may also be high water users and unable to refill embolism (lower right box), using other drought avoidance/tolerance strategies. Alternatively, species may be able to refill embolism and have conservative hydraulic strategies (upper left box) (Colour figure online)

date, representations of alternative strategies for repairing embolisms are lacking in existing models, even though the impact may be considerable (Sperry and Love, 2015). Although the theoretical challenges outlined in “Introduction” do exist for developing a mechanistic formulation of novel refilling, individual-based dynamic vegetation models with hydrodynamics are ideally structured to explore testable hypotheses.

In hydrodynamic formulations, maximum xylem conductivity ( $K_{x,max}$ ) is proportional to the area of the density distribution of the vessel vulnerability to decreasing xylem pressure ( $\Psi_x$ ), assuming the vessel vulnerability is identical across vessels (red zone in Fig. 4a). Under non water-stressed conditions, xylem conductance ( $k_x$ ) can be computed by multiplying  $K_{x,max}$  by the maximum cross-sectional area of functional-xylem ( $A_{x,max}$ ) and dividing by the path length ( $z$ ). As water-stress increases in the plant,  $k_x$  drops as functional vessels become embolized under decreasing  $\Psi_x$ , thus yielding a so-called xylem vulnerability curve (VC, Fig. 4b). During periods of water-stress, the down-regulation of  $k_x$  can be approximated using a 0 – 1 scaling factor ( $\beta$ ), which represents the fraction of maximum conductivity (or  $1 - PLC/100$ ) as  $\Psi_x$  declines and can be formulated as:


$$k_x = \frac{A_{x,max}K_{x,max}}{z} \beta \quad (1a)$$

$$\beta = \exp \left[ - \left( \frac{-x}{c_1} \right)^{c_2} \right] \quad (1b)$$

Vulnerability curves are typically sigmoidal (Torres-Ruiz et al. 2014) and to achieve this shape,  $\beta$  is expressed here as a Weibull function with  $c_1$  and  $c_2$  as fitting parameters (Bohrer et al. 2005).

During drought events, vessels increasingly lose function as  $\Psi_x$  declines (Fig. 4c). Hydrodynamic formulations (e.g. see references above) have the term of recovery of vessel function (i.e.  $\beta$  returning toward 1 and hence,  $k_x$  returning toward its maximum in Fig. 4d) tightly coupled to  $\Psi_x$ . Therefore,  $k_x$  recovers completely over the same temporal scale as the increase in  $\Psi_x$ , which may occur due to the relaxation of xylem tension at night after stomata close or as  $\Psi_s$  increases following a rain event (Fig. 4d, black curve) irrespective of the degree of embolized vessels in the xylem (dashed black line, Fig. 4c). Hence, such hydrodynamic formulations fail to represent those species where xylem refilling and recovery of  $k_x$  does *not* occur as  $\Psi_x$  increases (Fig. 4d, red curve).

We propose a new framework for evaluating xylem recovery as a strategy for drought tolerance. In real



systems, plants experience highly dynamic  $\Psi_x$  from variation in transpiration demand, soil moisture supply and stem water storage, all of which results in episodic embolisms (Fig. 4e). Therefore,  $k_x$  is governed by two distinct phases for  $A_x$  determined by the sign of  $d\Psi_x/dt$ : (1) a growth phase for when functional xylem area is increasing through repair or growth of new xylem ( $d\Psi_x/dt > 0$ ), and (2) a cavitation phase when functional

xylem is decreasing ( $d\Psi_x/dt \leq 0$ ). Accordingly, Eqs. (1a, 1b) can be reorganized to include a time dimension that represents the recovery of xylem function at different time scales ranging from sub-daily to seasons and is decoupled from both  $\Psi_s$  and  $\Psi_x$ :



◀**Fig. 4** Framework illustrating alternative strategies for xylem repair following embolism events. We hypothesize that the range of potential strategies consists of species with xylem vulnerability curves (VCs) that fully recover, partially recover or do not recover until new xylem is grown. **a** Hypothetical density distribution of vessel vulnerability to embolism along a xylem water potential ( $\Psi_x$ ) continuum. The shape of this density distribution differs between species. **b** Each species' VC is the integral of its vessel density distribution (i.e. panel **a**) and is expressed as the reduction of xylem conductance ( $k_x$ ,  $\text{kg s}^{-1} \text{MPa}^{-1}$ ), as  $\Psi_x$  declines. **c** A single embolism event occurring at a given  $\Psi_x$  (black dashed line), where vulnerable vessels above it embolize while vessels below it remain functional. **d** When vessels refill perfectly following an embolism event, the VC continues to operate along the original VC (solid black curve). When vessels do not refill following embolism, a new VC is established (solid red curve) and remains until new xylem grows. **e** During droughts,  $\Psi_x$  may decrease and increase multiple times leading to several embolism events of varying degrees. **f** Alternative VCs for 3 species after embolism events. Species A completely refills its vessels and  $k_x$  recovers (black curve) on the same time scale as soil moisture re-wetting. Species B recovers  $k_x$  on a time scale that is slower than soil moisture re-wetting. The initial loss of  $k_x$  tracks the original VC (black line), but recoveries are delayed (dashed blue), and subsequent VCs (solid blue) operate in the grey zone beneath the original VC. Species C does not refill; rather, a new VC is established after an embolism event (red). The new maximum  $k_x$  for Species C is established at the point on the original VC where the minimum  $\Psi_x$  is reached and extends horizontally back toward 0 as xylem tension relaxes with soil rewetting. For species C, growth of new xylem is required for  $k_x$  to return to its original maximum (black). In panels **c**–**f**, the dashed vertical black line represents the minimum  $\Psi_x$  reached and hence, the maximum embolism event during a drought period before new xylem is grown (Colour figure online)

cavitation phase  $\left(\frac{d_x}{dt} \leq 0\right)$  :

$$A_x^t = \begin{cases} A_x^{t-1} - f\left(\frac{d_x}{dt}, \tau\right), & A_x^{t-1} < A_{\max}\beta^t - \tau \\ A_{\max}\beta^t, & A_x^{t-1} \geq A_{\max}\beta^t - \tau \end{cases} \quad (2a)$$

growth phase  $\left(\frac{d_x}{dt} > 0\right)$  :

$$A_x^t = \min\left[A_{\max}\beta^t, A_x^{t-1} + f\left(\frac{d_x}{dt}, \tau\right)\right] \quad (2b)$$

$$k_x = \frac{K_{x,\max}}{z} A_x^t \quad (2c)$$

The superscript  $t$  indicates either the current ( $t$ ) or previous ( $t - 1$ ) time step. The parameter  $\tau$  is in units of area and determines how quickly  $A_x$  regenerates over one time step. When  $f(d\Psi_x/dt, \tau) = 0$ , no recovery of  $k_x$  occurs as the xylem tension relaxes; but instead a new VC is established at each minimum  $\Psi_x$  that is reached during a drought period (e.g. red curve, Fig. 4d). In this case as presented in Fig. 4f, during the first event the initial reduction in  $\Psi_x$  also happens to be the maximum embolism event (vertical dashed line, Fig. 4d). Here  $k_x$  first tracks down the original VC to the minimum  $\Psi_x$  that is reached; and then, as xylem tension relaxes during rehydration,  $k_x$  tracks horizontally back toward the origin during rehydration (red horizontal line, Fig. 4f). When subsequent embolism events occur (vertical blue dashed lines, Fig. 4e),  $k_x$  tracks horizontally back to the left along the new VC (red) until it reaches the original

VC (black) where it then resumes its decent along the original VC if the embolism event becomes even more severe (not shown in Fig. 4f). Alternatively, when  $f(d\Psi_x/dt, \tau)$  is relatively high, perfect recovery of  $k_x$  occurs along the original VC as xylem tension relaxes (black curve, Fig. 4f). When  $f(d\Psi_x/dt, \tau)$  is relatively low, but still  $> 0$ , imperfect recovery of  $k_x$  occurs more slowly below the original VC (i.e. blue dashed curves in Fig. 4f). This formulation allows for new VCs to establish after recovery and provides flexibility for them to establish anywhere within the grey zone of Fig. 4f. The right hand portion of Eq. (2a) establishes the conditional logic that constrains  $k_x$  to the grey zone in Fig. 4f during the cavitation phase. The position of any new VC within the grey zone (e.g. solid blue curves, Fig. 4f) is dependent on the species-specific magnitude of  $\tau$  and the minimum  $\Psi_x$  attained during the cavitation phase (dashed blue curves, Fig. 4f). Finally, xylem recovery that occurs through radial growth could be represented either with this model if  $f(d\Psi_x/dt, \tau)$  is parameterized to be relatively low, or through a separate stem-growth function.

Our proposed framework can be readily incorporated into individual-based dynamic vegetation models that represent direct competition for resources between species with alternative growth and survival strategies (e.g. ED2, Xu et al. 2016). We present a general form for our xylem recovery model, rather than suggesting specific functional forms for  $f(d\Psi_x/dt, \tau)$  since mechanisms controlling novel refilling are under debate. This general form can accommodate multiple xylem-refilling hypotheses such that one can be agnostic about the phenomenon of refilling while using it within a dynamic vegetation model to evaluate if, when, and in what ecosystems rapid refilling emerges as ecologically important. For example, one species  $f(d\Psi_x/dt, \tau)$  could be formulated within a given dynamic vegetation model with a curve with relatively steep slope that runs parallel and to the inside of the original VC curve (Fig. 4f), which implies that the larger, more conductive vessels refill first. Another species  $f(d\Psi_x/dt, \tau)$  could be formulated with a curve with relatively shallow slope as  $\Psi_x$  returns toward 0, which implies the smaller and less conductive vessels are the first to refill under tension. A third species could be formulated with no refilling. Competition between these species could then be simulated with the dynamic vegetation model to determine the performance of each strategy under the given meteorological forcing.

It should be noted that the *cost* for xylem recovery must also be accounted for in models that consider plant competition; otherwise, plants that recover xylem function more quickly will eventually dominate. Costs associated with xylem recovery, and drought tolerance in general, are poorly understood; yet research to date suggests that these costs likely involve potassium and carbon (Secchi and Zwieniecki 2011; Trifilò et al. 2014a). This framework is, therefore, also structured to easily accommodate new hypotheses about the economics of

xylem function and recovery. Finally, Eqs. (2a, 2b) provides a testable hypothesis explaining why larger trees are more vulnerable to drought (Nepstad et al. 2007) or have an apparently lower xylem P50 compared to smaller trees of the same species (Rowland et al. 2015).

---

## Future research directions

Plant hydraulics research has rapidly advanced in recent years, and yet many unresolved questions remain, including methodical ones. We suggest that future research focus on the following major unresolved problems:

- Which species, biomes, and plant functional types show embolism refilling, and which do not? Current research shows that the ability to refill is variable across species. Therefore, we need a comprehensive understanding of species that do not refill, as well as refillers, under comparable growth conditions, so we may understand which traits are linked with refilling.
- Do multi-year, or repeated drought events impact the ability of plants to refill previously embolised conduits?
- Is there a continuum for refilling or a threshold, defined by the amount of embolism, xylem pressure reached upon rehydration and/or the availability of stored carbohydrates (and/or partial recovery of photosynthesis)?
- Are changes in vulnerability curves related to adjustments in xylem anatomy, and to what degree do anatomical structures or genetics constrain this plasticity? Quantifying ring-to-ring variability in vulnerability curves especially with regards to drought stress and drought history would be very valuable.

## Conclusions

Some species are reported to show embolism refilling, in certain circumstances, whereas others are not. However, these reports are often based on different methods, each with different strengths and weaknesses. Newly emerging non-destructive methods of measuring embolism and refilling are generally free from cutting artefacts, and should be used to validate hydraulic measurements and to reach consensus on artefact-free sampling and handling protocols. At the same time, some of them might lead to injuries to living cells, thus limiting the possibility to detect vital phenomena like embolism reversal.

Given that some studies report embolism refilling in certain species, while others report no refilling in certain species, we conclude that the synthesis of evidence presented here suggests refilling does occur in some circumstances in some species. This leads to the

question—in what species and circumstances does refilling occur? Embolism refilling has been reported in roots, stems, or petioles, through positive root pressure, under negative pressure (up to a certain water potential), or a combination of these. It appears likely that there must be enough water available, although the water potential threshold remains an unknown key research question. Embolism refilling is influenced by phloem, ray and/or axial parenchyma, xylem age and location. There appears to be a trade-off to embolism refilling that involves costs associated with metabolism and transport of soluble sugars, water and potassium.

Further research is required to test which species can and cannot refill embolism, and under what conditions, which will advance our knowledge of why some plants die, while others are able to survive during drought. This information would be very valuable for future research efforts aimed at breeding and selecting cultivars and genotypes of crops and forest species able to cope with projected increase in frequency/intensity of drought and heat waves. Current research is mainly focused on selecting traits conferring resistance or tolerance to drought stress, like embolism resistance (Cochard et al. 2008; Aranda et al. 2015; Fichot et al. 2015). Evidence accumulating around the role of embolism refilling might indeed suggest shifting the focus to traits conferring resilience to drought. It is also possible that selection strategies aimed at improving plant resilience to drought based on embolism refilling might do so while maintaining or even improving plant productivity, thus avoiding the possible trade-offs between plant hydraulic safety and efficiency/productivity (Gleason et al. 2016). Anatomical (wood parenchyma fraction and spatial pattern) and functional (NSC content) traits might represent useful proxies to estimate the recovery ability of different species/genotypes in screening projects.

Further, understanding which species refill embolism will help constrain model formulations that are important for improving predictions of vegetation dynamics and carbon and water fluxes in hotter, drier climates. Including refilling in functional models of vegetation performance under recurrent drought would facilitate models to capture subtle effects of drought/rewetting cycles on ecosystem productivity, which simpler models would be unable to capture. Considering that future global change-type drought might take multiple shapes in terms of intensity and frequency, the impact of physiological strategies for short-term recovery of plant hydraulic functioning cannot be ignored, if we aim to accurately predict the impact of such droughts on ecosystem functioning.

**Acknowledgements** The paper was inspired by an International Interdisciplinary Workshop on Tree Mortality at the Max-Planck for Biogeochemistry in Jena, Germany (October 2014). Two anonymous reviewers have helped improving the text by providing insightful comments. Funding support was provided in full or in part as follows: MZ by ARC DECRA DE120100518. TK by the Benozio Fund for the Advancement of Science; Mr. and Mrs. Norman Reiser, together with the Weizmann Center for New Sci-

entists; and the Edith and Nathan Goldberg Career Development Chair. WRLA by a NOAA Climate and Global Change Postdoctoral fellowship, administered by the University Corporation of Atmospheric Research. JB by the Austrian Science Fund (FWF): M1757-B22 through the Lise Meitner Program. PJH by the University of New Mexico. NKR by the German Federal Ministry of Education and Research (BMBF), through the Helmholtz Association and its research programme ATMO and by the German Research Foundation through its Emmy Noether Programme (RU 1657/2-1). TLP by student research funding from the OEB department at Harvard University and by the Office of Biological and Environmental Research, US Department of Energy, NGEETropics project. GvA by the Swiss State Secretariat for Education, Research and Innovation SERI (SBFI C14.0104 and C12.0100). The authors declare no conflict of interest. W.R.L.A. acknowledges funding from NSF 1714972 and from the USDA National Institute of Food and Agriculture, Agricultural and Food Research Initiative Competitive Programme, Ecosystem Services and Agroecosystem Management, grant no. 2017-05521.

**Author contributions** The ideas presented here were developed in collaboration by all authors during a workshop held in Jena, Germany. MZ and TK performed the literature review and conceptually organized the manuscript. TLP developed the modelling framework presented in “[Modelling hydraulic recovery in vulnerability curves](#)”; code is available upon request. All authors contributed to writing the manuscript.

## References

- Anderegg WRL (2015) Spatial and temporal variation in plant hydraulic traits and their relevance for climate change impacts on vegetation. *New Phytol* 205:1008–1014
- Anderegg WRL, Berry JA, Smith DD, Sperry JS, Anderegg LDL, Field CB (2012) The roles of hydraulic and carbon stress in a widespread climate-induced forest die-off. *PNAS* 109:233–237
- Aranda I, Cano J, Gascó A, Cochard H, Nardini A, Mancha JA, López R, Sánchez-Gómez D (2015) Variation in photosynthetic performance and hydraulic architecture across European beech (*Fagus sylvatica* L.) populations supports the case for local adaptation to water stress. *Tree Physiol* 35:34–46
- Bloemen J, Vergeynst LL, Overlaet-Michiels L, Steppe K (2016) How important is woody tissue photosynthesis in poplar during drought stress? *Trees Struct Funct* 30:63–72
- Bohrer G, Mourad H, Laursen TA, Drewry D, Avissar R, Poggi D, Oren R, Katul GG (2005) Finite element tree crown hydrodynamics model (FETCH) using porous media flow within branching elements: a new representation of tree hydrodynamics. *Water Resour Res* 41(W11404):1–17
- Bouche PS, Larter M, Domec JC, Burlett R, Gasson P, Jansen S, Delzon S (2014) A broad survey of hydraulic and mechanical safety in the xylem of conifers. *J Exp Bot* 65:4419–4431
- Brodersen CR (2013) Visualizing wood anatomy in three dimensions with high-resolution X-ray micro-tomography (Mct)—a Review. *IAWA J* 34:408–424
- Brodersen CR, McElrone AJ (2013) Maintenance of xylem network transport capacity: a review of embolism repair in vascular plants. *Front Plant Sci* 4
- Brodrribb TJ, Cochard H (2009) Hydraulic failure defines the recovery and point of death in water-stressed conifers. *Plant Physiol* 149:575–584
- Brodrribb T, Holbrook NM (2004) Stomatal protection against hydraulic failure: a comparison of co-existing ferns and angiosperms. *New Phytol* 162:663–670
- Brodrribb TJ, Bowman DJMS, Nichols S, Delzon S, Burlett R (2010) Xylem function and growth rate interact to determine recovery rates after exposure to extreme water deficit. *New Phytol* 188:533–542
- Carminati A, Vetterlein D, Koebernick N, Blaser S, Weller U, Vogel HJ (2013) Do roots mind the gap? *Plant Soil* 367:651–661
- Charrier G, Torres-Ruiz JM, Badel E, Burlett R, Choat B, Cochard H, Delmas CE, Domec JC, Jansen S, King A, Lenoir N (2016) Evidence for hydraulic vulnerability segmentation and lack of xylem refilling under tension. *Plant Physiol* 01079
- Chen YJ, Schnitzer SA, Zhang YJ, Fan ZX, Goldstein G, Tomlinson KW, Lin H, Zhang JL, Cao KF (2017) Physiological regulation and efficient xylem water transport regulate diurnal water and carbon balances of tropical lianas. *Funct Ecol* 31:306–317
- Chitarra W, Balestrini R, Vitali M, Pagliarani C, Perrone I, Schubert A, Lovisolo C (2014) Gene expression in vessel-associated cells upon xylem embolism repair in *Vitis vinifera* L. petioles. *Planta* 239:887–899
- Choat B, Drayton WM, Brodersen C, Matthews MA, Shackel KA, Wada HIR, McElrone AJ (2010) Measurement of vulnerability to water stress-induced cavitation in grapevine: a comparison of four techniques applied to a long-veined species. *Plant Cell Environ* 33:1502–1512
- Choat B, Jansen S, Brodrribb TJ, Cochard H, Delzon S, Bhaskar R, Bucci SJ, Feild TS, Gleason SM, Hacke UG, Jacobsen AL, Lens F, Maherali H, Martínez-Vilalta J, Mayr S, Mencuccini M, Mitchell PJ, Nardini A, Pittermann J, Pratt RB, Sperry JS, Westoby M, Wright IJ, Zanne AE (2012) Global convergence in the vulnerability of forests to drought. *Nature* 491:752–755
- Choat B, Brodersen CR, McElrone AJ (2015) Synchrotron X-ray microtomography of xylem embolism in *Sequoia sempervirens* saplings during cycles of drought and recovery. *New Phytol* 205:1095–1105
- Christensen-Dalsgaard KK, Tyree MT (2014) Frost fatigue and spring recovery of xylem vessels in three diffuse-porous trees in situ. *Plant Cell Environ* 37:1074–1085
- Christman MA, Sperry JS, Smith DD (2012) Rare pits, large vessels and extreme vulnerability to cavitation in a ring-porous tree species. *New Phytol* 193:713–720
- Christoffersen BO, Gloor M, Fauset S, Fyllas NM, Galbraith DR, Baker TR, Sevanto S (2016) Linking hydraulic traits to tropical forest function in a size-structured and trait-driven model (TFS v. 1-Hydro). *Geosci Model Develop* 9:4227
- Cochard H, Delzon S (2013) Hydraulic failure and repair are not routine in trees. *Ann For Sci* 70:659–661
- Cochard H, Tyree MT (1990) Xylem dysfunction in *Quercus*: vessel sizes, tyloses, cavitation and seasonal changes in embolism. *Tree Physiol* 6:393–407
- Cochard H, Bodet C, Ameglio T, Cruiziat P (2000) Cryo-scanning electron microscopy observations of vessel content during transpiration in walnut petioles. Facts or artefacts? *Plant Physiol* 124:1191–1202
- Cochard H, Barigah ST, Kleinhentz M, Eshel A (2008) Is xylem cavitation resistance a relevant criterion for screening drought resistance amongst *Prunus* species? *J Plant Physiol* 165:976–982
- Cochard H, Delzon S, Badel E (2015) X-ray microtomography (micro-CT): a reference technology for high-resolution quantification of xylem embolism in trees. *Plant Cell Environ* 38:201–206
- Delzon S, Douthe C, Sala A, Cochard H (2010) Mechanism of water-stress induced cavitation in conifers: bordered pit structure and function support the hypothesis of seal capillary-seeding. *Plant Cell Environ* 33:2101–2111
- Duursma RA, Medlyn BE (2012) MAESPA: a model to study interactions between water limitation, environmental drivers and vegetation function at tree and stand levels, with an example application to [CO<sub>2</sub>]? drought interactions. *Geosci Model Dev* 5:919
- Eller CB, Lima AL, Oliveira RS (2013) Foliar uptake of fog water and transport belowground alleviates drought effects in the cloud forest tree species, *Drimys brasiliensis* (Winteraceae). *New Phytol* 199:151–162
- Fichot R, Brignolas F, Cochard H, Ceulemans R (2015) Vulnerability to drought-induced cavitation in poplars: synthesis and future opportunities. *Plant Cell Environ* 38:1233–1251



- Fonti P, Tabakova M, Kiryanov A, von Bryukhanova M, Arx G (2015) Variability of radial ray anatomy of *Larix gmelinii* along a forest productivity gradient in Siberia. *Trees* 29:1165–1175
- Fukuda K, Kawaguchi D, Aihara T, Ogasa MY, Miki NH, Haishi T, Umebayashi T (2015) Vulnerability to cavitation differs between current-year and older xylem: non-destructive observation with a compact magnetic resonance imaging system of two deciduous diffuse-porous species. *Plant Cell Environ* 38:2508–2518
- Ganthaler A, Mayr S (2015) Dwarf shrub hydraulics: two *Vaccinium* species (*Vaccinium myrtillus*, *Vaccinium vitis-idaea*) of the European Alps compared. *Physiol planta* 155:424–434
- Gleason S, Westoby M, Jansen S, Choat B, Hacke U, Pratt RB, Bhaskar R, Brodribb T, Bucci S, Cao K, Cochard H, Delzon S, Domec JC, Fan Z, Field T, Jacobsen A, Johnson D, Lens F, Maherali H, Martínez-Vilalta J, Mayr S, McCulloh K, Mencuccini M, Mitchell P, Morris H, Nardini A, Pittermann J, Plavcová L, Schreiber S, Sperry J, Wright I, Zanne A (2016) Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant species. *New Phytol* 209:123–136
- Gotsch SG, Asbjornsen H, Holwerda F, Goldsmith GR, Weintraub AE, Dawson TE (2014) Foggy days and dry nights determine crown-level water balance in a seasonal tropical montane cloud forest. *Plant, Cell Environ* 37:261–272
- Gouin H (2015) The watering of tall trees—embolization and recovery. *J Theor Biol* 369:42–50
- Hacke UG, Sperry JS (2003) Limits to xylem refilling under negative pressure in *Laurus nobilis* and *Acer negundo*. *Plant Cell Environ* 26:303–311
- Hacke UG, Sperry John S, Pockman William T, Davis Stephen D, McCulloh Katherine A (2001) Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* 126:457–461
- Hacke UG, Venturas MD, Mackinnon ED, Jacobsen AL, Sperry JS, Pratt RB (2014) The standard centrifuge method accurately measures vulnerability curves of long-vesselled olive stems. *New Phytol* 205:116–127
- Holbrook NM, Ahrens ET, Burns MJ, Zwieniecki MA (2001) In vivo observation of cavitation and embolism repair using magnetic resonance imaging. *Plant Physiol* 126:27–31
- Jackson GE, Grace J (1996) Field measurements of xylem cavitation: are acoustic emissions useful? *J Exp Bot* 47:1643–1650
- Jansen S, Gortan E, Lens F, Lo Gullo MA, Salleo S, Scholz A, Stein A, Trifilò P, Nardini A (2011) Do quantitative vessel and pit characters account for ion-mediated changes in the hydraulic conductance of angiosperm xylem? *New Phytol* 189:218–228
- Javot H, Maurel C (2002) The role of aquaporins in root water uptake. *Ann Bot* 90:301–313
- Johnson DM, McCulloh KA, Woodruff DR, Meinzer FC (2012) Hydraulic safety margins and embolism reversal in stems and leaves: why are conifers and angiosperms so different? *Plant Sci* 195:48–53
- Kelley DI, Harrison SP, Prentice IC (2014) Improved simulation of fire-vegetation interactions in the land surface processes and exchanges dynamic global vegetation model (LPX-Mv1). *Geosci model develop* 7:931–1000
- Kirtman B, Power S, Adedoyin A, Boer GJ, Bojariu R, Camilloni I, Doblas-Reyes F, Fiore A, Kimoto M, Meehl GA, Prather M, Sarr A, Schar C, Sutton R, van Oldenborgh GJ, Vecchi G, Wang HJ (2014) Working group I contribution to the IPCC fifth assessment report climate change 2013: the physical science basis. (ed. W. g. 1). Stockholm
- Klein T, Cohen S, Yakir D (2011) Hydraulic adjustments underlying drought resistance of *Pinus halepensis*. *Tree Physiol* 31:637–648
- Klein T, Shpringer I, Fikler B, Elbaz G, Cohen S, Yakir D (2013) Relationships between stomatal regulation, water-use, and water-use efficiency of two coexisting key Mediterranean tree species. *For Ecol Manage* 302:34–42
- Klein T, Yakir D, Buchmann N, Grünzweig JM (2014) Towards an advanced assessment of the hydrological vulnerability of forests to climate change-induced drought. *New Phytol* 201:712–716
- Klein T, Cohen S, Paudel I, Preisler Y, Rotenberg E, Yakir D (2016) Diurnal dynamics of water transport, storage and hydraulic conductivity in pine trees under seasonal drought. *iForest* 9:710
- Laur J, Hacke UG (2014) Exploring *Picea glauca* aquaporins in the context of needle water uptake and xylem refilling. *New Phytol* 203:388–400
- Leng H, Lu M, Wan X (2013) Variation in embolism occurrence and repair along the stem in drought-stressed and re-watered seedlings of a poplar clone. *Physiol Planta* 147:329–339
- Lens F, Sperry JS, Christman MA, Choat B, Rabaey D, Jansen S (2011) Testing hypotheses that link wood anatomy to cavitation resistance and hydraulic conductivity in the genus *Acer*. *New Phytol* 190:709–723
- Li Y, Chen W, Chen J, Shi H (2016) Contrasting hydraulic strategies in *Salix psammophila* and *Caragana korshinskii* in the southern Mu Us Desert, China. *Ecol Res* 31:869–881
- Martorell S, Diaz-Espejo A, Medrano H, Ball MC, Choat B (2014) Rapid hydraulic recovery in *Eucalyptus pauciflora* after drought: linkages between stem hydraulics and leaf gas exchange. *Plant Cell Environ* 37:617–626
- Mayr S, Schmid P, Laur J, Rosner S, Charra-Vaskou K, Dämon B, Hacke UG (2014) Uptake of water via branches helps timberline conifers refill embolized xylem in late winter. *Plant Physiol* 164:1731–1740
- McCulloh KA, Johnson DM, Meinzer FC, Woodruff DR (2014) The dynamic pipeline: hydraulic capacitance and xylem hydraulic safety in four tall conifer species. *Plant Cell Environ* 37:1171–1183
- McDowell NG, Beerling DJ, Breashers DD, Fisher RA, Raffa KF, Stitt M (2011) The interdependence of mechanisms underlying climate-driven vegetation mortality. *Trends Ecol Evol* 26:523–532
- McDowell N, Ryan MG, Zeppel MJB, Tissue D (2013) Improving our knowledge of drought-induced forest mortality through experiments, observations, and modeling. *New Phytol* 200:289–293
- Myer JE (1922) Ray volumes of the commercial woods of the United States and their significance. PhD, New York State College of Forestry
- Nardini A, Lo Gullo MA, Salleo S (1998) Seasonal changes of root hydraulic conductance (K-RL) in four forest trees: an ecological interpretation. *Plant Ecol* 139:81–90
- Nardini A, Lo Gullo MA, Salleo S (2011a) Refilling embolized xylem conduits: is it a matter of phloem unloading? *Plant Sci* 180:604–611
- Nardini A, Salleo S, Jansen S (2011b) More than just a vulnerable pipeline: xylem physiology in the light of ion-mediated regulation of plant water transport. *J Experim Bot* 62:4701–4718
- Nardini A, Lo Gullo MA, Trifilò P, Salleo S (2014) The challenge of the Mediterranean climate to plant hydraulics: responses and adaptations. *Environ Experim Bot* 103:68–79
- Nardini A, Savi T, Losso A, Petit G, Pacilè S, Tromba G, Mayr S, Trifilò P, Lo Gullo MA, Salleo S (2017) X-ray micro-tomography observations of xylem embolism in stems of *Laurus nobilis* L. are consistent with hydraulic measurements of percentage loss of conductance. *New Phytol* 213:1068–1075
- Nardini A, Savi T, Trifilò P, Lo Gullo MA (2018) Drought stress and the recovery from xylem embolism in woody plants. *Progress Bot* 79:197–231
- Nepstad DC, Tohver IM, Ray D, Moutinho P, Cardinot G (2007) Mortality of large trees and lianas following experimental drought in an Amazon forest. *Ecology* 88:2259–2269
- Nolf M, Beikircher B, Rosner S, Nolf A, Mayr S (2015) Xylem cavitation resistance can be estimated based on time-dependent rate of acoustic emissions. *New Phytol* 208:625–632
- Nolf M, Lopez R, Peters JMR, Flavel RJ, Koloadin LS, Young IM, Choat B (2017) Visualization of xylem embolism by X-ray microtomography: a direct test against hydraulic measurements. *New Phytol*. <https://doi.org/10.1111/nph.14462>
- Ogasa M, Miki N, Yoshikawa K (2010) Changes of hydraulic conductivity during dehydration and rehydration in *Quercus serrata* Thunb. and *Betula platyphylla* var. *japonica* Hara: the effect of xylem structures. *Tree Physiol* 30:608–617



- Ogasa M, Miki NH, Murakami Y, Yoshikawa K (2013) Recovery performance in xylem hydraulic conductivity is correlated with cavitation resistance for temperate deciduous tree species. *Tree Physiol* 33:335–344
- Ooeda H, Terashima I, Taneda H (2016) Structures of bordered pits potentially contributing to isolation of a refilled vessel from negative xylem pressure in stems of *Morus australis* Poir.: testing of the pit-membrane-osmosis and pit-valve hypotheses. *Plant Cell Physiol* pcw196
- Pena J, Grace J (1986) Water relations and ultrasound emissions of *Pinus sylvestris* L. before, during and after a period of water-stress. *New Phytol* 103:515–524
- Pittermann J, Brodersen C, Watkins JE (2013) The physiological resilience of fern sporophytes and gametophytes: advances in water relations offer new insights into an old lineage. *Front Plant Sci* 4
- Plavcova L, Jansen S (2015) The role of xylem parenchyma in the storage and utilisation of nonstructural carbohydrates. In: Hacke U (ed) *Functional and ecological xylem anatomy*. Springer International Publishing, Switzerland, pp 209–234
- Rockwell FE, Wheeler JK, Holbrook NM (2014) Cavitation and its discontents: opportunities for resolving current controversies. *Plant Physiol* 164:1649–1660
- Rolland V, Dana MB, Lenné T, Bryant G, Chen H, Wolfe J, Holbrook NM, Daniel ES, Marilyn CB (2015) Easy come, easy go: capillary forces enable rapid refilling of embolized primary xylem vessels. *Plant Physiol* 168:1636–1647
- Rood SB, Patiño S, Coombs K, Tyree MT (2000) Branch sacrifice: cavitation-associated drought adaptation of riparian cottonwoods. *Trees* 14:248–257
- Rowland L, da Costa ACL, Galbraith DR, Oliveira RS, Binks OJ, Oliveira AAR, Pullen AM, Doughty CE, Metcalfe DB, Vasconcelos SS, Ferreira LV (2015) Death from drought in tropical forests is triggered by hydraulics not carbon starvation. *Nature* 528:119–122
- Salleo S, Lo Gullo MA, De Paoli D, Zippo M (1996) Xylem recovery from cavitation-induced embolism in young plants of *Laurus nobilis*: a possible mechanism. *New Phytol* 132:47–56
- Salleo S, Lo Gullo MA, Trifilo P, Nardini A (2004) New evidence for a role of vessel-associated cells and phloem in the rapid xylem refilling of cavitated stems of *Laurus nobilis* L. *Plant Cell Environ* 27:1065–1076
- Savi T, Casolo V, Luglio J, Bertuzzi S, Gullo MAL, Nardini A (2016) Species-specific reversal of stem xylem embolism after a prolonged drought correlates to endpoint concentration of soluble sugars. *Plant Physiol Biochem* 106:198–207
- Savi T, Miotto A, Petruzzellis F, Losso A, Pacilè S, Tromba G, Mayr S, Nardini A (2017) Drought-induced embolism in stems of sunflower: a comparison of in vivo micro-CT observations and destructive hydraulic measurements. *Plant Physiol Biochem* 120:24–29
- Schenk HJ (2014) Formation and repair of xylem embolisms: Current state of knowledge and implications for irrigation of horticultural crops. *Acta Hort* 311–326
- Schmitz N, Egerton JGG, Lovelock CE, Ball MC (2012) Light-dependent maintenance of hydraulic function in mangrove branches: do xylary chloroplasts play a role in embolism repair? *New Phytol* 195:40–46
- Secchi F, Zwieniecki MA (2010) Patterns of PIP gene expression in *Populus trichocarpa* during recovery from xylem embolism suggest a major role for the PIP1 aquaporin subfamily as moderators of refilling process. *Plant Cell Environ* 33:1285–1297
- Secchi F, Zwieniecki MA (2011) Sensing embolism in xylem vessels: the role of sucrose as a trigger for refilling. *Plant Cell Environ* 34:514–524
- Secchi F, Zwieniecki MA (2012) Analysis of xylem sap from functional (Nonembolized) and nonfunctional (Embolized) vessels of *Populus nigra*: chemistry of refilling. *Plant Physiol* 160:955–964
- Secchi F, Zwieniecki MA (2014) Down-regulation of plasma intrinsic protein1 aquaporin in poplar trees is detrimental to recovery from embolism. *Plant Physiol* 164:1789–1799
- Sperry JS, Love DM (2015) What plant hydraulics can tell us about responses to climate-change droughts. *New Phytol* 207:14–27
- Sperry JS, Christman MA, Torres-Ruiz JM, Taneda H, Smith DD (2012) Vulnerability curves by centrifugation: is there an open vessel artefact, and are ‘r’ shaped curves necessarily invalid? *Plant Cell Environ* 35:601–610
- Torres-Ruiz JM, Cochard H, Mayr S, Beikircher B, Diaz-Espejo A, Rodriguez-Dominguez CM, Badel E, Fernández JE (2014) Vulnerability to cavitation in *Olea europaea* current-year shoots: further evidence of an open-vessel artefact associated with centrifuge and air-injection techniques. *Physiol Planta* 152:465–474
- Torres-Ruiz JM, Jansen S, Choat B, McElrone AJ, Cochard H, Brodribb TJ, Badel E, Burrell R, Bouche PS, Brodersen CR, Li S, Morris H, Delzon S (2015) Direct X-ray microtomography observation confirms the induction of embolism upon xylem cutting under tension. *Plant Physiol* 167:40–43
- Tötze C, Miranda T, Konrad W, Gout J, Kardjilov N, Dawson M, Manke I, Roth-Nebelsick A (2013) Visualization of embolism formation in the xylem of liana stems using neutron radiography. *Ann Bot* 111:723–730
- Trifilo P, Barbera PM, Raimondo F, Nardini A, Gullo MAL, Meinzer F (2014a) Coping with drought-induced xylem cavitation: coordination of embolism repair and ionic effects in three Mediterranean evergreens. *Tree Physiol* 34:109–122
- Trifilo P, Raimondo F, Lo Gullo MA, Barbera PM, Salleo S, Nardini A (2014b) Relax and refill: xylem rehydration prior to hydraulic measurements favours embolism repair in stems and generates artificially low PLC values. *Plant Cell Environ* 37:2491–2499
- Trifilo P, Nardini A, Gullo MAL, Barbera PM, Savi T, Raimondo F (2015) Diurnal changes in embolism rate in nine dry forest trees: relationships with species-specific xylem vulnerability, hydraulic strategy and wood traits. *Tree Physiol* 35:694–705
- Tyree MT, Cochard H, Cruziat P, Sinclair B, Ameglio T (1993) Drought-induced leaf shedding in walnut: evidence for vulnerability segmentation. *Plant Cell Environ* 16:879–882
- Tyree MT, Salleo S, Nardini A, Lo Gullo MA, Mosca R (1999) Refilling of embolized vessels in young stems of laurel. Do we need a new paradigm? *Plant Physiol* 120:11–21
- Umebayashi T, Ogasa MY, Miki NH, Utsumi Y, Haishi T, Fukuda K (2016) Freezing xylem conduits with liquid nitrogen creates artefactual embolisms in water-stressed broadleaf trees. *Trees* 30:305–316
- Utsumi Y, Sano Y, Fujikawa S, Funada R, Ohtani J (1998) Visualization of cavitated vessels in winter and refilled vessels in spring in diffuse-porous trees by cryo-scanning electron microscopy. *Plant Physiol* 117:1463–1471
- Venturas MD, Mackinnon ED, Jacobsen AL, Pratt RB (2015) Excising stem samples underwater at native tension does not induce xylem cavitation. *Plant Cell Environ* 38:1060–1068
- Vertessy RA, Watson FG, O’Sullivan RSK (2001) Factors determining relations between stand age and catchment water balance in mountain ash forests. *For Ecol Manage* 143:13–26
- von Arx G, Archer SR, Hughes MK (2012) Long-term functional plasticity in plant hydraulic architecture in response to supplemental moisture. *Ann Bot* 109:1091–1100
- Wang MT, Tyree MT, Wasylishen RE (2013) Magnetic resonance imaging of water ascent in embolized xylem vessels of grapevine stem segments. *Can J Plant Sci* 93:879–893
- Wheeler JK, Huggett BA, Tofte AN, Rockwell FE, Holbrook NM (2013) Cutting xylem under tension or supersaturated with gas can generate PLC and the appearance of rapid recovery from embolism. *Plant Cell Environ* 36:1938–1949
- Williams M, Rastetter EB, Fernandes DN, Goulden ML, Wofsy SC, Shaver GR, Melillo JM, Munger JW, Fan S-M, Nadelhoffer KJ (1996) Modelling the soil–plant–atmosphere continuum in a Quercus–Acer stand at Harvard Forest: the regulation of stomatal conductance by light, nitrogen and soil/plant hydraulic properties. *Plant Cell Environ* 19:911–927
- Xu X, Medvigy D, Powers JS, Becknell JM, Guan K (2016) Diversity in plant hydraulic traits explains seasonal and inter-annual variations of vegetation dynamics in seasonally dry tropical forests. *New Phytol* 212:80–95

- Zeppel MJB, Macinnis-Ng CMO, Yunusa IAM, Whitley RJ, Eamus D (2008) Long term trends of stand transpiration in a remnant forest during wet and dry years. *J Hydrol* 349:200–213
- Zeppel MJB, Harrison SP, Adams HD, Kelley DI, Li G, Tissue DT, Dawson TE, Fensham R, Medlyn BE, Palmer A, West AG, McDowell NG (2015) Drought and resprouting plants. *New Phytol* 206:583–589
- Zufferey V, Cochard H, Ameglio T, Spring J-L, Viret O (2011) Diurnal cycles of embolism formation and repair in petioles of grapevine (*Vitis vinifera* cv. Chasselas). *J Exp Bot* 62:3885–3894
- Zwieniecki MA, Holbrook NM (2000) Bordered pit structure and vessel wall surface properties. Implications for embolism repair. *Plant Physiol* 123:1015–1020
- Zwieniecki MA, Melcher PJ, Ahrens ET (2013) Analysis of spatial and temporal dynamics of xylem refilling in *Acer rubrum* L. using magnetic resonance imaging. *Front. Plant Sci* 4:265