



Yin-and-yang bifurcation of opioidergic circuits for descending analgesia at the midbrain of the mouse

Jong-Hyun Kim^{a,b,c}, Gireesh Gangadharan^a, Junweon Byun^{a,d}, Eui-Ju Choi^b, C. Justin Lee^c, and Hee-Sup Shin^{a,d,1}

^aCenter for Cognition and Sociality, Institute for Basic Science, 34141 Daejeon, Korea; ^bDivision of Life Sciences, Korea University, 02841 Seoul, Korea; ^cCenter for Glia-Neuron Interaction, Brain Science Institute, Korea Institute of Science and Technology, 02792 Seoul, Korea; and ^dDepartment of Basic Science, Korea University of Science and Technology, 34141 Daejeon, Korea

Contributed by Hee-Sup Shin, September 12, 2018 (sent for review April 10, 2018; reviewed by Xiaoke Chen, Zhuo Min, and Gerald W. Zamponi)

In the descending analgesia pathway, opioids are known to disinhibit the projections from the periaqueductal gray (PAG) to the rostral ventromedial medulla (RVM), leading to suppression of pain signals at the spinal cord level. The locus coeruleus (LC) has been proposed to engage in the descending pathway through noradrenergic inputs to the spinal cord. Nevertheless, how the LC is integrated in the descending analgesia circuit has remained unknown. Here, we show that the opioidergic analgesia pathway is bifurcated in structure and function at the PAG. A knockout as well as a PAG-specific knockdown of phospholipase C $\beta 4$ (PLC $\beta 4$), a signaling molecule for G protein-coupled receptors, enhanced swim stress-induced and morphine-induced analgesia in mice. Immunostaining after simultaneous retrograde labeling from the RVM and the LC revealed two mutually exclusive neuronal populations at the PAG, each projecting either to the LC or the RVM, with PLC $\beta 4$ expression only in the PAG-LC projecting cells that provide a direct synaptic input to LC-spinal cord (SC) projection neurons. The PAG-LC projection neurons in wild-type mice turned quiescent in response to opiates, but remained active in the PLC $\beta 4$ mutant, suggesting a possibility that an increased adrenergic function induced by the persistent PAG-LC activity underlies the enhanced opioid analgesia in the mutant. Indeed, the enhanced analgesia in the mutant was reversed by blocking $\alpha 2$ -noradrenergic receptors. These findings indicate that opioids suppress descending analgesia through the PAG-LC pathway, while enhancing it through the PAG-RVM pathway, i.e., two distinct pathways with opposing effects on opioid analgesia. These results point to a therapeutic target in pain control.

descending analgesia pathway | opioid | periaqueductal gray | locus coeruleus | phospholipase C

Pain signals are processed by the ascending sensory circuit and modulated by the descending analgesic circuit. The periaqueductal gray (PAG) at the midbrain, a key region in the descending analgesia circuit, is rich in opioid receptors and generates endogenous opioid analgesic signals (1–6). These signals are relayed by the rostral ventromedial medulla (RVM) to the dorsal horn of the spinal cord (SC), where they suppress pain signals (5, 7). Swimming in warm water induces opioid-dependent, swim stress-induced analgesia (SSIA) through this circuit (8). Within the PAG, tonically active GABAergic interneurons inhibit output neurons that project to the RVM (9, 10). Endogenous opioids suppress this inhibitory influence of local GABAergic interneurons through mu opioid receptors (μ ORs), disinhibiting the antinociceptive drive of the neuronal outputs to the RVM to positively control descending analgesia (9, 10). Exogenous opioid analgesics, such as morphine, also act through these μ ORs (11, 12).

Phospholipase C (PLC), the enzyme responsible for calcium mobilization, represents a family of molecules that are coupled to μ ORs and affects protein kinase C (PKC) activation (13, 14). We have previously shown that the PLC β isoform, PLC $\beta 4$, is required for pain sensory transmission at the thalamic level (15, 16). PLC $\beta 4$ -PKC signaling is linked to T-type calcium channels and regulates the firing pattern of thalamocortical neurons to favor either tonic or burst firing, which efficiently sharpens the

transition between open and closed gate status in pain sensory processing (15, 17, 18). In situ hybridization data show that PLC $\beta 4$ is abundantly expressed in the PAG (Allen Brain Atlas, mouse.brain-map.org), but its potential role in the descending pain control circuit has not been explored.

The locus coeruleus (LC) has recently been proposed to play a role in endogenous descending pain control through noradrenergic inputs, which act via $\alpha 2$ -adrenoreceptors to inhibit both primary afferents and second-order projection neurons in the SC (19–28). The LC has an afferent connection from the PAG (29, 30), but how the LC is integrated in the descending analgesia circuits are unresolved.

In this study, motivated by the knowledge of its signaling interactions in the ascending pain pathway (15), we investigated the role of PLC $\beta 4$ in the descending pain system controlled by opioidergic signals. Our results demonstrate the existence of two opioidergic circuits that bifurcate at the PAG and exert opposing effects on descending analgesia.

Results

Opioid-Dependent Analgesia Is Enhanced in PAG-Specific Knockdown and PLC $\beta 4$ -Null Mutant Mice. We previously reported that μ OR-positive GABAergic neurons in the PAG mediate the opioidergic descending analgesia through a mechanism that is dependent on T-type calcium channels (10). In addition, neuronal burst firing, which is mediated by T-type calcium channels, is increased in PLC $\beta 4$ -deleted neurons (15). We thus hypothesized that a deletion of PLC $\beta 4$ at the PAG might enhance opioidergic descending analgesia.

Significance

The midbrain periaqueductal gray (PAG) is a major site in the descending analgesia circuits for the endogenous, opioidergic pain control. Recently, the locus coeruleus (LC) has been proposed to play a role in endogenous pain control. Nevertheless, how the LC is integrated in the descending analgesia circuits remains elusive. Here, we provide evidence that the opioid circuit is bifurcated into two distinct pathways, each with an opposing effect on opioid analgesia at the PAG level: suppressing through the PAG-LC pathway while enhancing through the PAG-rostral ventromedial medulla pathway. Thus, the yin-and-yang model is proposed for the opioidergic descending analgesia. Moreover, our findings suggest an insight for a therapeutic target to control pain.

Author contributions: J.-H.K., E.-J.C., C.J.L., and H.-S.S. designed research; J.-H.K., G.G., and J.B. performed research; J.-H.K., G.G., and J.B. analyzed data; and J.-H.K., J.B., C.J.L., and H.-S.S. wrote the paper.

Reviewers: X.C., Stanford University; Z.M., University of Toronto; and G.W.Z., University of Calgary.

The authors declare no conflict of interest.

Published under the PNAS license.

¹To whom correspondence should be addressed. Email: shin@ibs.re.kr.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1806082115/-DCSupplemental.

Published online October 8, 2018.

To test this hypothesis, we initially examined baseline thermal sensitivity in both PLC β 4-knockout [PLC β 4 $^{-/-}$] mice and PAG-specific PLC β 4-knockdown (PLC β 4-shRNA) mice (Fig. 1A). PLC β 4 knockout significantly increased baseline thermal sensitivity ($t = 4.833$, $P = 0.0003$, unpaired two-tailed Student's t test; Fig. 1B), a result consistent with a previous study showing that PLC β 4 $^{-/-}$ mice exhibit reduced nociception (15, 16). In contrast, PAG-specific knockdown of PLC β 4 did not significantly affect the baseline thermal sensitivity of the mouse ($t = 0.964$, $P = 0.366$, unpaired two-tailed Student's t test; Fig. 1B). We then assessed the performance of these mice in the opioid-dependent swim stress-induced analgesia (SSIA) assay. Interestingly, SSIA was significantly increased in both PLC β 4 $^{-/-}$ ($t = 4.881$, $P = 0.0003$, unpaired two-tailed Student's t test; Fig. 1C) and PAG-specific PLC β 4-knockdown mice (Scrambled-shRNA vs. PLC β 4-shRNA, $t = 8.808$, $P < 0.001$, unpaired two-tailed Student's t test; Fig. 1C). In addition, we carried out the 2% formalin-induced pain test after SSIA in both wild-type and PAG-specific PLC β 4-knockdown group. Correspondingly, the PAG-specific PLC β 4-knockdown mice showed significantly reduced inflammatory pain response compared with the control group [$F_{1,228} = 21.88$, $P < 0.001$, repeated-measures (RM) ANOVA, *SI Appendix*, Fig. S14]. Immunostaining of the brains after SSIA assays confirmed a reduction in PLC β 4 expression in the PAG of knockdown mice (*SI Appendix*, Fig. S2). These results indicate that PLC β 4 in the PAG negatively controls opioid-dependent analgesia without affecting basal thermal sensitization.

Next, we investigated exogenous opioid-induced analgesia in both wild-type and PLC β 4-defective mice after systemic injection (10 mg/kg) or PAG infusion (1 μ g) of morphine through a cannula (*SI Appendix*, Fig. S3), following a previously described procedure (10). In wild-type mice, morphine induced a significant level of analgesia compared with saline-treated controls following both systemic injection ($F_{1,15} = 11.88$, $P = 0.0044$, RM ANOVA) and PAG infusion ($F_{1,32} = 26.57$, $P < 0.001$, RM ANOVA) (Fig. 1D and E). Morphine also induced analgesia compared with saline in PLC β 4 $^{-/-}$ mice, whether injected systemically ($F_{1,14} = 74.84$, $P < 0.001$, RM ANOVA) or infused in the PAG ($F_{1,31} = 69.14$, $P < 0.001$, RM ANOVA) (Fig. 1D

and E). Notably, the extent of the morphine-induced analgesia was significantly augmented in the mutant mice compared with wild-type mice following systemic injection ($t = 3.346$, $P = 0.0065$, unpaired two-tailed Student's t test; *SI Appendix*, Fig. S3A) or infusion into the PAG ($t = 5.533$, $P < 0.001$, unpaired two-tailed Student's t test; *SI Appendix*, Fig. S3B). These findings demonstrate that (i) the PAG is a major site for morphine-induced analgesia and (ii) PLC β 4 deficiency substantially enhances PAG-mediated, morphine-induced analgesia.

PLC β 4 Is Selectively Expressed in PAG \rightarrow LC Circuits, but Not in PAG \rightarrow RVM Circuits. The PAG has extensive connectivity with the brain network that includes not only the RVM (1, 2, 5, 31) but also the LC, which exerts an analgesic effect (19–22, 29, 30, 32). In an effort to define the relationship between PLC β 4 expression and PAG-RVM or PAG-LC projection neurons in the PAG, we injected the retrograde tracers, cholera toxin B (CTB) and fluorospheres, into the LC and RVM, respectively (*SI Appendix*, Fig. S5). Random fluorospheres labeling of PAG-RVM neurons was observed throughout the entire PAG region, consistent with a previous report (10), whereas retrograde labeling by CTB from the LC was predominantly observed in the lateral PAG and ventral PAG. Interestingly, there was no overlap of the two different fluorescent signals in the same cells, indicating that the two projection neurons constitute distinct cell populations.

To investigate the expression of PLC β 4 in these two neuronal populations, we performed immunohistochemistry for PLC β 4 on retrograde-labeled sections of both PAG-LC and PAG-RVM projection neurons. Notably, PLC β 4 was expressed in PAG-LC projection neurons, which are located in the ventral region, but not in PAG-RVM projection neurons (Fig. 2A, B, and D and *SI Appendix*, Fig. S6).

In addition, double immunofluorescence experiments performed using tissue from GAD67-GFP mice showed that the majority of PLC β 4 was detected in neurons that expressed CamKII α (calcium/calmodulin-dependent protein kinase II), but not in neurons that expressed GAD67-GFP (*SI Appendix*, Fig. S7), indicating that PLC β 4 was primarily expressed in excitatory neurons at the PAG. Moreover, double immunostaining for




Fig. 1. Improved analgesia in PLC β 4 $^{-/-}$ and PAG-specific PLC β 4-knockdown mice under endogenous and exogenous opioid conditions. (A) Schematic depiction of the endogenous opioid condition and the exogenous opioid condition paradigm used in this study. (B) Test of basal thermal sensitivity using a hotplate assay. (C) Comparison of the analgesic effect after the SSIA behavior test. (D and E) Time course of the effects of morphine-induced analgesia compared with saline control at the systemic and PAG level. Values are means \pm SEM. (*, # $P < 0.05$; **, ## $P < 0.01$; ***, ### $P < 0.001$; n.s., not significant).




Fig. 2. PLC β 4 is expressed in ventral PAG-LC but not in PAG-RVM projection neurons. (A) Representative image of cells in the PAG retrogradely labeled with two different-color tracers. Schematic illustration of dual retrograde tracing from both the LC and RVM (*Inset*). (B–D) Zoomed view of the subcellular localization of CTB, fluorospheres, and PLC β 4 in the dorsal PAG (dPAG) (B), lateral PAG (lPAG) (C), and ventral PAG (vPAG) (D). Arrows indicate PLC β 4 immunoreactivity at LC retrogradely labeled neurons. (Scale bars: A, 100 μ m; B–D, 25 μ m.)

tyrosine hydroxylase (TH), a marker for the LC (33, 34), showed no PLC β 4 signal in LC neurons (*SI Appendix, Fig. S8*). These findings raised the possibility that, because it is selectively expressed in PAG-LC projections, PLC β 4 may modulate the LC-mediated analgesia circuit.

PLC β 4-Expressing PAG Neurons Make a Direct Synaptic Input to LC→SC Projection Neurons. Although it has been well known that LC neurons receive dense projection from the PAG (29, 30), and that LC neurons innervate the SC to modulate pain transmission (22), confirmation of the direct connectivity of these three brain regions has not been achieved. To address this issue, we used the cTRIO method (29) to examine whether PLC β 4-expressing PAG neurons make monosynaptic connections to LC-SC projection neurons (Fig. 3A). We injected a Cre-dependent CAV2 vector carrying the flippase (Flp) recombinase [CAV2-FLEX (loxP)-Flp] into the SC of TH-Cre mice. Because CAV2 infects axons and is retrogradely transported to cell bodies, this combination allows us to express Flp specifically in Cre-expressing LC-SC projection neurons. We then injected the Flp-activatable AAV vectors, AAV10-CAG-FLEX (Frt)-TVA-mCherry and AAV10-CAG-FLEX (Frt)-RG, into the LC to express TVA and rabies glycoprotein (RG) in these cells (Fig. 3B). Finally, we injected SADAG-EGFP (EnvA) (35) in the same region (LC) to label upstream neurons that make direct synaptic contact with LC-SC projection neurons. As expected, we found that some PLC β 4-expressing PAG neurons expressed SADAG-EGFP (Fig. 3C), indicating that PLC β 4-expressing PAG neurons have monosynaptic connectivity to LC neurons and suggesting that these PAG neurons directly regulate the LC-SC projection neurons.

Deletion of PLC β 4 Eliminates Opioid-Induced Suppression of Firing in PAG-LC Projection Neurons. To define the physiological mechanism underlying the enhanced opioid-mediated analgesia associated with a PLC β 4 deficiency, we explored the physiological response of PAG-LC projection neurons to opioids in brain slices. We focused on the caudal portion of the ventral region of the PAG because it has been shown that cells in this region are sensitive to morphine *in vivo* (36–38) and because retrogradely labeled PAG-LC projection neurons were confined to this region (Fig. 4A). Patch-clamp recording showed no difference in the resting membrane potential of PAG-LC projection neurons between wild-type (-42.82 ± 2.99 mV) and PLC β 4 $^{-/-}$ (-41.11 ± 1.2 mV) mice. Moreover, neither wild-type nor PLC β 4 $^{-/-}$ PAG-LC projection neurons showed low-threshold spikes (LTSS) (Fig. 4B and D). Next, we recorded the discharge activity of PAG-LC

projection neurons at -35 mV for 1 min. The discharge activity of PAG-LC projection neurons under these conditions was about ~ 4 Hz, and there was no significant difference in this activity between wild-type and PLC β 4 $^{-/-}$ mice (Fig. 4C and E). However, there was a significant difference in sensitivity to the endogenous opioid neurotransmitter, [met⁵]-enkephalin (ME), between wild-type and PLC β 4-deficient neurons. ME is known to be released by stress *in vivo* and inhibits neurons mainly through μ ORs (4, 5, 39, 40). Wild-type PAG-LC projection neurons quickly responded to application of ME (30 μ M) with a reduction in discharge activity and became progressively hyperpolarized (14.72 ± 2.84 mV) compared with untreated controls ($t = 4.389$, $P = 0.0070$, paired two-tailed Student's *t* test; Fig. 4C and F). In contrast, PLC β 4 $^{-/-}$ PAG-LC projection neurons exhibited no change in discharge activity or in the membrane potential in response to the ME treatment ($t = 0.197$, $P = 0.848$, paired two-tailed Student's *t* test; Fig. 4E and F). Consistent with these results, treatment with the μ OR agonist, [D-Ala(2),N-Me-Phe(4),Gly(5)-ol]-enkephaline (DAMGO), suppressed the neuronal activity of wild-type PAG-LC projection neurons but showed no effect on PLC β 4 $^{-/-}$ PAG-LC projection neurons (*SI Appendix, Fig. S9*). To confirm that these results could not be attributed to a change in μ ORs caused by deletion of PLC β 4, we immunostained PAG-LC projection neurons for μ ORs. These experiments show that expression of μ ORs in PAG-LC projection neuron is not altered in PLC β 4 $^{-/-}$ mice ($t = 0.748$, $P = 0.456$, unpaired two-tailed Student's *t* test; *SI Appendix, Fig. S10*). However, the response of the PAG-RVM projection neurons to the ME treatment was not changed in both groups (*SI Appendix, Fig. S11*). These findings suggest that PLC β 4 is required for the μ OR-dependent signal transduction that mediates suppression of PAG-LC projection neurons. This, in turn, would lead to a decrease in the activity of downstream target noradrenergic neurons in the LC, resulting in a decrease in norepinephrine-dependent analgesia. In the absence of PLC β 4, noradrenergic neurons in the LC would remain active even in the presence of opioids, thus supporting norepinephrine-dependent analgesia. The net result is an enhancement in opioid-induced analgesia at the systems level in PLC β 4 $^{-/-}$ mice compared with wild-type mice.

Enhanced Opioid-Dependent Analgesia in PLC β 4 $^{-/-}$ Mice Is Reversed by Antagonism of α 2-Adrenergic Receptors. RVM-SC projection neurons innervated the laminae I and II of the lumbar spinal cord (41). Also, the LC is the principal noradrenergic nucleus in the central nervous system and the main source of noradrenergic innervation to the spinal dorsal horn, forming a well-described




Fig. 3. Transsynaptic tracing of PAG → LC to LC → SC circuit by cTRIO. (A) Schematic depiction of experimental procedure. CAV2-FLEX (loxP)-Flp was injected into the SC of TH-Cre mice to express Flp in LC neurons that send projections to the PAG (day 1). Flp-dependent AAV carrying TVA, and RG were then injected into the LC (day 1). SADΔG-EGFP (EnvA) was injected into the same site (day 15). (B, Left) Distribution of the starter cells (TVA⁺, SADΔG⁺ cells) in the LC region. (B, Right) Magnifications of regions in white rectangles in images (yellow arrows). (C, Left) PLCβ4-expressing PAG neurons merged with the SADΔG+EGFP signals. (C, Right) Magnifications of regions in white rectangles in images (white arrows).

analgesic circuit (42, 43). We made the output of LC neurons projecting to the SC to express fluorescence in a Cre-dependent manner to substantiate that RVM and LC neurons engage SC in different pathways (SI Appendix, Fig. S12A). Axonal terminals from the LC were observed throughout the laminae I ~ X of the lumbar spinal cord, which made us propose that the targeting of LC-SC neurons is different from that of the RVM-SC pathway (SI Appendix, Fig. S12B).

In the SC, norepinephrine suppresses pain signals through inhibitory actions on α2A-adrenoreceptors on central terminals of primary afferent nociceptors (presynaptic inhibition), as well as through direct α2-adrenergic actions on pain-relay neurons (postsynaptic inhibition) (19, 20, 23, 24, 27, 44). If the enhanced analgesic effect of locally or systemically infused morphine in

PLCβ4^{-/-} mice were a direct result of the continuous LC excitability cause by the persistent activity of PAG-LC projection neurons, blocking the noradrenergic inputs in this mutant should reverse the enhanced analgesic effect. First, we compared the c-Fos expression level in the LC between the wild-type and the PLCβ4^{-/-} mice after the SSIA. As we expected, the PLCβ4^{-/-} mice showed more c-Fos signals in the LC than the wild-type mice, implying hyperexcitability of the LC neurons in the mutant mice. ($t = 10.19$, $P < 0.001$, unpaired two-tailed Student's t test; SI Appendix, Fig. S13 A and B) Next, we used idazoxan, an α2-adrenergic receptor antagonist, that does not interfere with the morphine effect (45). i.p. administration of idazoxan (20 mg/kg) did not affect the basal thermal sensitivity in wild-type ($t = 0.103$, $P = 0.919$, unpaired two-tailed Student's t test) or PLCβ4^{-/-}




Fig. 4. Discharge activities at depolarized membrane potential in the LC-projecting neurons of the ventral PAG. (A, Left) Schematic depiction of retrograde labeling in the LC and patch-clamp recordings of cells in the PAG. (A, Right three images) DIC image (Left and Middle) and retrograde dye image of a patched neuron in the ventral PAG (vPAG; Right). Red circle indicates an example of PAG-LC projection neurons. (Scale bars: A, Middle Left, 500 μm; A, Middle Right and Right, 25 μm.) (B and D) Electrophysiological analysis of LTS in PAG-LC projection neurons from wild-type (B) and PLCβ4^{-/-} (D) mice in response to hyperpolarizing current pulses. (C and E) Changes in firing patterns for two periods from PAG-LC projection neurons from wild-type (C) and PLCβ4^{-/-} (E) mice. (F) Comparison of discharge activity before and after treatment with ME between in wild-type and PLCβ4^{-/-} PAG-LC projection neurons. Values are means ± SEM. (***) $P < 0.001$; n.s., not significant).




Fig. 5. The reversed opioidergic analgesia by antagonizing systemic α 2-adrenergic receptors in PLC β 4 mice. (A) Time course of the analgesic effects of idazoxan and morphine in the wild-type and in the PLC β 4 mutant mice by systemic treatment. (B) Time course of the analgesic effects of i.p. injected idazoxan with the PAG-specific delivery of morphine in both the wild-type and the PLC β 4 mutant mice. Values are means \pm SEM (* P < 0.05; ** P < 0.01; *** P < 0.001).

mice ($t = 0.137$, $P = 0.893$, unpaired two-tailed Student's t test) (SI Appendix, Fig. S14A). We subsequently infused morphine (10 mg/kg) in combination with a systemic injection of idazoxan to block α 2-adrenergic receptor activation. As expected, idazoxan did not influence the analgesic effect of morphine in the wild-type mice, as measured by RM ANOVA ($F_{1,25} = 2.735$, $P = 0.1107$, Fig. 5A) or unpaired Student's t test ($t = 1.696$, $P = 0.102$, SI Appendix, Fig. S14B). Notably, in mutant mice, a systemic injection of idazoxan significantly reduced morphine-induced analgesia compared with morphine alone, resulting in a nearly complete reversal of the potentiated analgesia to wild-type levels ($F_{1,16} = 126.2$, $P < 0.001$, RM ANOVA, Fig. 5A; $t = 14.78$, $P < 0.001$, unpaired two-tailed Student's t test; SI Appendix, Fig. S14B). Next, we systemically injected idazoxan together with PAG-specific morphine infusion in both wild-type and PLC β 4-deficient mouse groups. Systemic idazoxan injection did not affect the analgesia induced by PAG-specific morphine infusion in wild-type mice ($F_{1,26} = 0.3768$, $P = 0.5447$, RM ANOVA, Fig. 5B; $t = 0.424$, $P = 0.674$, unpaired two-tailed Student's t test, SI Appendix, Fig. S14C). However, the increased PAG-specific morphine-induced analgesia observed in the PLC β 4 $^{-/-}$ mice was returned to wild-type levels by i.p. injected idazoxan ($F_{1,23} = 7.678$, $P = 0.0109$, RM ANOVA, Fig. 4B; $t = 5.066$, $P < 0.001$, unpaired two-tailed Student's t test, SI Appendix, Fig. S14C).

Taken together, these results indicate that the persistent activity of LC neurons is responsible for the augmented opioid-induced analgesia in PLC β 4 $^{-/-}$ mice.

Discussion

Yin-and-Yang PAG Bifurcation Model. Our data suggest that the stress-induced, opioid-dependent, descending analgesia circuits bifurcate at the PAG into two mutually exclusive pathways that exert opposite effects: The PAG-RVM projection enhances, and the PAG-LC projection suppresses, the descending analgesia (Fig. 6, Left and Middle).

The most investigated form of the physiological activation of opioid-linked pain-modulating circuits originates under the general framework of stress-induced analgesia (7, 46). The midbrain PAG was the first discovered descending pain-modulating site in the brain, and the PAG-RVM circuits project to the dorsal horn of the SC and the trigeminal nucleus caudalis, where pain-transmitting neurons are located. In the disinhibition model, opioids activate the PAG-RVM descending pathway via suppression of the inhibitory influence of local GABAergic interneurons, thereby enhancing the antinociceptive drive of the neuronal output to the SC (9, 10). Now the results in this paper reveal the complex

organization of the PAG with regard to the control of the stress-induced opioid-dependent descending analgesia. Thus, opioid can both enhance and suppress the descending analgesia.

The Opioid Circuits in the PAG, a Safeguard Against the Adverse Effect of Stress.

The LC is activated by physiological and environmental stressors and is thought to play a role in cognitive aspects of the stress response (47). Corticotropin-releasing factor (CRF) acts through activation of the LC-norepinephrine (NE) system to increase the LC neuronal firing rate, thereby contributing to coordination of physiological and behavioral responses to stress (48, 49). Apart from the well-known descending LC-spinal pathway, which is important for pain control, an increase in LC discharge under stress conditions could potentially lead to a serious autonomic failure such as hypertensive challenge (50, 51). Our results suggest that endogenous opioids may serve to counterbalance the adverse effect of stress on the LC-NE system, while achieving analgesia.

Phospholipase C Signaling in the Opioid-Induced Analgesia.

Previously, we have reported that the thalamic PLC β 4 regulates the firing properties of thalamocortical neurons through a mechanism mediated by T-type calcium channels (15). Interestingly, PAG-LC projection neurons showed no LTS responses, and their firing was not attenuated by the opioids in the absence of PLC β 4 (Fig. 3). These observations demonstrate that PLC β 4 is required for fine control of opioid responses in the descending pain pathway. Indeed, numerous studies have reported that the PLC β isoforms are activated by the G $\beta\gamma$, which is liberated by μ ORs activation, and induce IP $_3$ and diacylglycerol formation. This, in turn, results in intracellular Ca $^{2+}$ release from IP $_3$ -sensitive Ca $^{2+}$ stores, PKC activation, and VDCC inhibition (52–57). These conclusions are consistent with prevailing views regarding the importance of the PLC system in opioid-signaling pathways (58, 59).

Additional studies will be required to define the antinociceptive function of other PLC systems in the PAG during the pain transmission. Importantly, our findings suggest that the




Fig. 6. Working model of the PAG descending pain-gating pathway for the opioid analgesic circuitry. (Left) The two distinct pathways in the PAG bifurcation model: PAG-RVM projection neurons and PAG-LC projection neurons. (Middle) The physiological role of the PAG-RVM pathway in positive control of analgesia (\oplus) proposed in the disinhibition model, and the new proposed model of the PAG-LC pathway for negative control of antinociceptive effects (\ominus) in wild-type mice. (Right) PAG-LC excitatory transmission showed persistent activity under opioidergic conditions in PLC β 4 $^{-/-}$ mice, leading to an enhanced analgesic effect. The thickness of the arrows indicates the relative activity of individual transmissions.

current “Yin-and-Yang” model at the PAG provides a therapeutic target in pain control (Fig. 6, *Right*).

Materials and Methods

Animal care and all experiments were conducted in accordance with the Institutional Review Board of Institute for Basic Science (IBS), Korea for the

ethical guidelines of Animal Care and Use. Detailed descriptions of study methods are provided in *SI Appendix, Materials and Methods*.

ACKNOWLEDGMENTS. We thanks to Dr. C. Park, Dr. S. Keum, and Dr. B. Lee for their valuable discussion with technical support. This work was supported by IBS Grant IBS-R001-D1G.

- Yaksh TL, Yeung JC, Rudy TA (1976) Systematic examination in the rat of brain sites sensitive to the direct application of morphine: Observation of differential effects within the periaqueductal gray. *Brain Res* 114:83–103.
- Reichling DB, Kwiat GC, Basbaum AI (1988) Chapter 2 Anatomy, physiology and pharmacology of the periaqueductal gray contribution to antinociceptive controls. *Prog Brain Res* 77:31–46.
- Dostrovsky JO, Deakin JFW (1977) Periaqueductal grey lesions reduce morphine analgesia in the rat. *Neurosci Lett* 4:99–103.
- Lewis VA, Gebhart GF (1977) Evaluation of the periaqueductal central gray (PAG) as a morphine-specific locus of action and examination of morphine-induced and stimulation-produced analgesia at coincident PAG loci. *Brain Res* 124:283–303.
- Basbaum AI, Fields HL (1984) Endogenous pain control systems: Brainstem spinal pathways and endorphin circuitry. *Annu Rev Neurosci* 7:309–338.
- Bandler R, Shipley MT (1994) Columnar organization in the midbrain periaqueductal gray: Modules for emotional expression? *Trends Neurosci* 17:379–389.
- Fields H (2004) State-dependent opioid control of pain. *Nat Rev Neurosci* 5:565–575.
- Terman GW, Shavit Y, Lewis JW, Cannon JT, Liebeskind JC (1984) Intrinsic mechanisms of pain inhibition: Activation by stress. *Science* 226:1270–1277.
- Lau BK, Vaughan CW (2014) Descending modulation of pain: The GABA disinhibition hypothesis of analgesia. *Curr Opin Neurobiol* 29:159–164.
- Park C, et al. (2010) T-type channels control the opioidergic descending analgesia at the low threshold-spiking GABAergic neurons in the periaqueductal gray. *Proc Natl Acad Sci USA* 107:14857–14862.
- North RA, Williams JT (1983) Opiate activation of potassium conductance inhibits calcium action potentials in rat locus coeruleus neurons. *Br J Pharmacol* 80:225–228.
- Pan ZZ, Williams JT, Osborne PB (1990) Opioid actions on single nucleus raphe magnus neurons from rat and guinea-pig in vitro. *J Physiol* 427:519–532.
- Belcheva MM, et al. (2005) μ and κ opioid receptors activate ERK/MAPK via different protein kinase C isoforms and secondary messengers in astrocytes. *J Biol Chem* 280:27662–27669.
- Rubovitch V, Gafni M, Sarne Y (2003) The μ opioid agonist DAMGO stimulates cAMP production in SK-N-SH cells through a PLC-PKC-Ca⁺⁺ pathway. *Brain Res Mol Brain Res* 110:261–266.
- Cheong E, et al. (2008) Tuning thalamic firing modes via simultaneous modulation of T- and L-type Ca²⁺ channels controls pain sensory gating in the thalamus. *J Neurosci* 28:13331–13340.
- Miyata M, et al. (2003) Role of thalamic phospholipase C[β]4 mediated by metabotropic glutamate receptor type 1 in inflammatory pain. *J Neurosci* 23:8098–8108.
- Kim D, et al. (2003) Thalamic control of visceral nociception mediated by T-type Ca²⁺ channels. *Science* 302:117–119.
- Shin H-S, Cheong E-J, Choi S, Lee J, Na HS (2008) T-type Ca²⁺ channels as therapeutic targets in the nervous system. *Curr Opin Pharmacol* 8:33–41.
- Jones SL, Gebhart GF (1986) Quantitative characterization of cereulospinal inhibition of nociceptive transmission in the rat. *J Neurophysiol* 56:1397–1410.
- Miller JF, Proudfit HK (1990) Antagonism of stimulation-produced antinociception from ventrolateral pontine sites by intrathecal administration of α -adrenergic antagonists and naloxone. *Brain Res* 530:20–34.
- West WL, Yeomans DC, Proudfit HK (1993) The function of noradrenergic neurons in mediating antinociception induced by electrical stimulation of the locus coeruleus in two different sources of Sprague-Dawley rats. *Brain Res* 626:127–135.
- Hickey L, et al. (2014) Optoactivation of locus ceruleus neurons evokes bidirectional changes in thermal nociception in rats. *J Neurosci* 34:4148–4160.
- Reddy SV, Maderdrut JL, Yaksh TL (1980) Spinal cord pharmacology of adrenergic agonist-mediated antinociception. *J Pharmacol Exp Ther* 213:525–533.
- Hammond DL, Yaksh TL (1984) Antagonism of stimulation-produced antinociception by intrathecal administration of methysergide or phentolamine. *Brain Res* 298:329–337.
- North RA, Yoshimura M (1984) The actions of noradrenaline on neurones of the rat substantia gelatinosa in vitro. *J Physiol* 349:43–55.
- Jones SL, Gebhart GF (1986) Characterization of coeruleospinal inhibition of the nociceptive tail-flick reflex in the rat: Mediation by spinal α 2-adrenoceptors. *Brain Res* 364:315–330.
- Kawasaki Y, Kumamoto E, Furue H, Yoshimura M (2003) α 2 adrenoceptor-mediated presynaptic inhibition of primary afferent glutamatergic transmission in rat substantia gelatinosa neurons. *Anesthesiology* 98:682–689.
- Sonohata M, et al. (2004) Actions of noradrenaline on substantia gelatinosa neurones in the rat spinal cord revealed by in vivo patch recording. *J Physiol* 555:515–526.
- Schwarz LA, et al. (2015) Viral-genetic tracing of the input-output organization of a central noradrenaline circuit. *Nature* 524:88–92.
- Lee HS, Kim M-A, Waterhouse BD (2005) Retrograde double-labeling study of common afferent projections to the dorsal raphe and the nuclear core of the locus coeruleus in the rat. *J Comp Neurol* 481:179–193.
- Jacquet YF (1988) The NMDA receptor: Central role in pain inhibition in rat periaqueductal gray. *Eur J Pharmacol* 154:271–276.
- Mantyh PW (1983) Connections of midbrain periaqueductal gray in the monkey. II. Descending efferent projections. *J Neurophysiol* 49:582–594.
- Sands SA, Guerra V, Morilak DA (2000) Changes in tyrosine hydroxylase mRNA expression in the rat locus coeruleus following acute or chronic treatment with valproic acid. *Neuropsychopharmacology* 22:27–35.
- Takeuchi T, et al. (2016) Locus coeruleus and dopaminergic consolidation of everyday memory. *Nature* 537:357–362.
- Wickersham IR, Finke S, Conzelmann K-K, Callaway EM (2007) Retrograde neuronal tracing with a deletion-mutant rabies virus. *Nat Methods* 4:47–49.
- Barr GA, Wang S (2013) Analgesia induced by localized injection of opiate peptides into the brain of infant rats. *Eur J Pain* 17:676–691.
- Bellgowan PSF, Helmstetter FJ (1998) The role of μ and κ opioid receptors within the periaqueductal gray in the expression of conditional hypoalgesia. *Brain Res* 791:83–89.
- Lovick TA (1991) Interactions between descending pathways From the dorsal and ventrolateral periaqueductal gray matter in the rat. *The Midbrain Periaqueductal Gray Matter*, NATO ASI Series (Springer, Boston), pp 101–120.
- Chiang B, Christie MJ (1994) Hyperpolarization by opioids acting on μ -receptors of a sub-population of rat periaqueductal gray neurones in vitro. *Br J Pharmacol* 113:121–128.
- Behbehani MM, Jiang M, Chandler SD (1990) The effect of [Met]enkephalin on the periaqueductal gray neurons of the rat: An in vitro study. *Neuroscience* 38:373–380.
- Aicher SA, Hermes SM, Whittier KL, Hegarty DM (2012) Descending projections from the rostral ventromedial medulla (RVM) to trigeminal and spinal dorsal horns are morphologically and neurochemically distinct. *J Chem Neuroanat* 43:103–111.
- Dahlstroem A, Fuxe K (1965) Evidence for the existence of monoamine neurons in the central nervous system. II. Experimentally induced changes in the intraneuronal amine levels of bulbospinal neuron systems. *Acta Physiol Scand Suppl* (Suppl 247), 1–36.
- Millan MJ (2002) Descending control of pain. *Prog Neurobiol* 66:355–474.
- Pertovaara A (2006) Noradrenergic pain modulation. *Prog Neurobiol* 80:53–83.
- Bhalla S, Rapolaviciute V, Gulati A (2011) Determination of α (2)-adrenoceptor and imidazoline receptor involvement in augmentation of morphine and oxycodone analgesia by agmatine and BMS182874. *Eur J Pharmacol* 651:109–121.
- Fields HL (2000) Pain modulation: Expectation, opioid analgesia and virtual pain. *Prog Brain Res* 122:245–253.
- Valentino RJ, Foote SL, Page ME (1993) The locus coeruleus as a site for integrating corticotropin-releasing factor and noradrenergic mediation of stress responses. *Ann N Y Acad Sci* 697:173–188.
- Snyder K, Wang W-W, Han R, McFadden K, Valentino RJ (2012) Corticotropin-releasing factor in the norepinephrine nucleus, locus coeruleus, facilitates behavioral flexibility. *Neuropsychopharmacology* 37:520–530.
- Homberg JR, Contet C (2009) Deciphering the interaction of the corticotropin-releasing factor and serotonin brain systems in anxiety-related disorders. *J Neurosci* 29:13743–13745.
- Gurtu S, Pant KK, Sinha JN, Bhargava KP (1984) An investigation into the mechanism of cardiovascular responses elicited by electrical stimulation of locus coeruleus and subcoeruleus in the cat. *Brain Res* 301:59–64.
- Drolet G, Gauthier P (1985) Peripheral and central mechanisms of the pressor response elicited by stimulation of the locus coeruleus in the rat. *Can J Physiol Pharmacol* 63:599–605.
- Galeotti N, Stefano GB, Guarna M, Bianchi E, Ghelardini C (2006) Signaling pathway of morphine induced acute thermal hyperalgesia in mice. *Pain* 123:294–305.
- Xie W, et al. (1999) Genetic alteration of phospholipase C β 3 expression modulates behavioral and cellular responses to μ opioids. *Proc Natl Acad Sci USA* 96:10385–10390.
- Perroy J, et al. (2000) Selective blockade of P/Q-type calcium channels by the metabotropic glutamate receptor type 7 involves a phospholipase C pathway in neurons. *J Neurosci* 20:7896–7904.
- Mathews JL, Smrcka AV, Bidlack JM (2008) A novel Gbetagamma-subunit inhibitor selectively modulates μ -opioid-dependent antinociception and attenuates acute morphine-induced antinociceptive tolerance and dependence. *J Neurosci* 28:12183–12189.
- Bourinet E, Soong TW, Stea A, Snutch TP (1996) Determinants of the G protein-dependent opioid modulation of neuronal calcium channels. *Proc Natl Acad Sci USA* 93:1486–1491.
- Al-Hasani R, Bruchas MR (2011) Molecular mechanisms of opioid receptor-dependent signaling and behavior. *Anesthesiology* 115:1363–1381.
- Wilding TJ, Womack MD, McCleskey EW (1995) Fast, local signal transduction between the μ opioid receptor and Ca²⁺ channels. *J Neurosci* 15:4124–4132.
- Taddese A, Nah S-Y, McCleskey EW (1995) Selective opioid inhibition of small nociceptive neurons. *Science* 270:1366–1369.