



Published in final edited form as:

Acc Chem Res. 2014 February 18; 47(2): 560–578. doi:10.1021/ar400193g.

Ynamides in Ring Forming Transformations

XIAO-NA WANG^a, HYUN-SUK YEOM^a, LI-CHAO FANG^a, SHUZHONG HE^a, ZHI-XIONG MA^a, BRANT L. KEDROWSKI^{b,*}, and RICHARD P. HSUNG^{a,*}

^aDivision of Pharmaceutical Sciences and Department of Chemistry, University of Wisconsin, Madison, WI 53705

^bDepartment of Chemistry, University of Wisconsin, Oshkosh, WI 54901


Conspectus

The ynamide functional group activates carbon-carbon triple bonds through an attached nitrogen atom that bears an electron-withdrawing group. As a result, the alkyne has both electrophilic and nucleophilic properties. Through the selection of the electron-withdrawing group attached to nitrogen chemists can modulate the electronic properties and reactivity of ynamides, making these groups versatile synthetic building blocks. The reactions of ynamides also lead directly to nitrogen-containing products, which provides access to important structural motifs found in natural products and molecules of medicinal interest. Therefore, researchers have invested increasing time and research in the chemistry of ynamides in recent years.

This Account surveys and assesses new organic transformations involving ynamides developed in our laboratory and in others around the world. We showcase the synthetic power of ynamides for rapid assembly of complex molecular structures. Among the recent reports of ynamide transformations, ring-forming reactions provide a powerful tool for generating molecular complexity quickly. In addition to their synthetic utility, such reactions are mechanistically interesting. Therefore, we focus primarily on the cyclization chemistry of ynamides.


This Account highlights ynamide reactions that are useful in the rapid synthesis of cyclic and polycyclic structural manifolds. We discuss the mechanisms active in the ring formations and describe representative examples that demonstrate the scope of these reactions and provide mechanistic insights. In this discussion we feature examples of ynamide reactions involving radical cyclizations, ring-closing metathesis, transition metal and non-transition metal mediated cyclizations, cycloaddition reactions, and rearrangements. The transformations presented rapidly introduce structural complexity and include nitrogen within, or in close proximity to, a newly formed ring (or rings). Thus, ynamides have emerged as powerful synthons for nitrogen-containing heterocycles and nitrogen-substituted rings, and we hope this Account will promote continued interest in the chemistry of ynamides.

*kedrowsk@uwosh.edu, rhsung@wisc.edu.



1. Introduction

Over the past 15 years, ynamides have become a modern functional group that has been prominently featured in a variety of synthetic transformations including natural product total syntheses.¹⁻³ Fueled by preparative access that is efficient and atom economical,^{4,5} the field of ynamide chemistry has rapidly expanded. Ynamides provide a means of activating carbon-carbon triple bonds, giving them both electrophilic and nucleophilic properties. The electronic properties of ynamides are tunable based on selection of the electron-withdrawing group attached to nitrogen, thereby rendering them highly versatile synthons. Furthermore, among all heteroatom-substituted alkynes, ynamides are special because nitrogen is one of the most privileged elements in nature. Consequently, many transformations involving ynamides offer a diverse array of novel structural entities that are not only powerful platforms for further transformations, but also prevalent among important pharmacophores. These properties have contributed to a continued and dramatic increase in the number of publications in the last few years.



This Account aims to examine the literature from late 2009 through early 2013 related to the use of ynamides in synthetic transformations that form cyclic and polycyclic manifolds. Consequently, the Account is organized by reaction types used in the ring formation, and representative examples are selected to showcase scope and mechanistic insight of each transformation. The intention here is not to comprehensively review the ynamide chemistry that appeared during this period, but to highlight major advancements in ynamide chemistry through revelation of its utility in rapid assembly of structural complexity. As a result, some beautiful recent works are not presented here. This includes improved ynamide preparations and syntheses of de novo structural analogs of ynamides. Both topics have been the subject of a thorough review published recently by Evano.⁴ In addition, new advances related to in situ generation of ketenimines or metallo-ynamides via Huisgen's azide-[3 + 2] cycloaddition/retro-[3 + 2] are not covered here.⁶

2. Radical Cyclizations

Balieu and Courillon⁷ reported the formation of six-membered rings **5** and **6**, and the eight-membered ring **3** via the radical cyclization of ynamides **1** (Scheme 1). A tandem radical cyclization also took place when R was 3,5-dimethoxyphenyl through vinyl radical **4**.

Chemla and Perez-Luna⁸ reported synthesis of 3-alkylidenetetrahydrofurans **11** from ynamides **7** through a 1,4-addition/alkyne carbozincation sequence based on a radical zinc-atom transfer process (Scheme 2). The addition of ethyl radical gave enoxy radical **8**, which underwent 5-*exo-dig* cyclization with the ynamide leading to vinyl radical **9** of *E*-geometry.

3. Ring-Closing Metathesis

Wakamatsu and Mori⁹ reported further development of their ring-closing metathesis of eneynamides, featuring syntheses of 7-membered heterocycles **13**, **15** and 8-membered heterocycles **17** using 2nd-generation Grubbs catalyst (Scheme 3).

4. Non-Transition Metal Mediated Cyclizations

Popik¹⁰ reported acid catalyzed cycloaromatizations of cyclic ynamides **18** (Scheme 4). Protonation of the ynamide followed by addition of the alkyne onto the resulting keteniminium ion **19** provided cation **20**, which underwent Friedel-Crafts additions. Alcoholic solvent trapping of **19** competed with cycloaromatization, especially with increasing in ring size.

Evano¹¹ developed a general and efficient approach toward 1,4-dihydropyridines **29** and pyridines **30** from readily available *N*-allyl-ynamides **24** via a tandem lithiation–isomerization–6-*endo-dig* intramolecular carbolithiation sequence (Scheme 5).

Flynn¹² reported a highly torquoselective Nazarov cyclization of 2-amido divinyl ketones **34** derived from chiral oxazolidinone-substituted ynamides **31** (Scheme 6). The diastereoselectivity can be very high, and Nazarov arrested products such as **37a** could also be obtained.

Cao¹³ described an efficient approach toward 3-allenyl-2-amidobenzofurans **40** and 3-alkyl-2-amidobenzofurans **44** via a novel carbocation-induced electrophilic cyclization of *o*-anisole-substituted ynamides with 1,1-diaryl-prop-2-yn-1-ol **39** and diarylmethanol **41**, respectively (Scheme 7).

Later, Cao¹⁴ developed a novel synthesis of 2-amidobenzofurans and 2-amidobenzothiophenes via electrophilic cyclization of *o*-anisole- and *o*-thioanisole-substituted ynamides **45** with I₂, NBS, and NCS. This strategy was also used to construct 1-amidonaphthalenes **50** and 1-amidobenzopyrans **52** from ynamides (Scheme 8).

Hsung¹⁵ unveiled a novel acid promoted 5-*endo-dig* cyclization of chiral γ -amino-ynamide **53** concomitant with the loss of the *t*-Bu group that led to the formation of isothiazole **54** and dihydroisothiazole *S*-oxide **55**. An inversion at the “S” center occurred in **55** (Scheme 9).

Hsung¹⁶ featured an *aza*-variant of a Meyer-Schuster rearrangement of γ -amino-ynamides **56**, which involved the formation of azetene intermediate **58** and pericyclic ring-opening (Scheme 10).

5. Transition Metal Mediated Cyclizations

5.1. Rhodium

Nishimura and Hayashi¹⁷ disclosed a rhodium-catalyzed asymmetric cycloisomerization of heteroatom-bridged 1,6-enamides such as **60** to afford 3-*aza* and oxabicyclo[4.1.0]heptene derivatives such as **61a** (Scheme 11). 2-oxazolidinone and 2-azetidinone substituents of the ynamides were critical for high enantioselectivities, as the carbonyl oxygen might coordinate to the metal during the transformation.

Tang¹⁸ developed an efficient method for the generation of *α-oxo* Rh(I) carbenes **63** and **69** from ynamides with 3,5-dichloropyridine *N*-oxide. The resulting Rh(I) carbenes then react intramolecularly with various alkynes or alkenes affording 2-oxopyrrolidines **66** and 3-azabicyclo[3.1.0]hexanes **71**, respectively (Scheme 12).

5.2. Palladium

Anderson¹⁹ reported a palladium-catalyzed tandem cascade of cyclization–cross-coupling–6π-electron electrocyclization using bromoenynamides **72** (Scheme 13).

Anderson²⁰ subsequently reported a related palladium-catalyzed cascade using bromoenynamides **75** affording cyclic 2-amidodienes **77** that could be used in ensuing Diels-Alder cycloadditions (Scheme 14). An alcohol served as a hydride source to terminate the carbopalladation process.

5.3. Platinum

Liu²¹ reported an equivalent of Pt(II)-catalyzed *oxo*-arylations of ynamide **79** (Scheme 15). This process employs nitrones **80** and provides imines **83** via intermediates **81** and **82**, and under reductive conditions using NaBH₃CN, 2-indolones **85** were obtained.

5.4. Copper

Chen²² reported a Cu(I)-catalyzed 1,2-aminothiolation of 1,1-dibromoalkenes **86** with 2-thiobenzoimidazole, leading to thiazolines **90** and **91** (Scheme 16). The regiochemistry of the 1,2-aminothiolation depends on whether it is 5-*endo dig* cyclization of *S*- or *N*-alkynylation intermediates (**88** or **89**).

Hashmi²³ reported a copper-mediated domino reaction of three simple components that included propargylcarboxamide **92**, protected amine **93**, and a chloride source. This cascade provides an efficient construction of highly functionalized oxazines **97** (Scheme 17).

Neuville²⁴ developed an efficient and regioselective approach to 1,2,4-trisubstituted imidazoles **103** via a copper-catalyzed oxidative diamination of terminal alkynes **99** by amidines **98** (Scheme 18). This transformation employs oxygen as the co-oxidant and proceeds through a direct *N*-alkynylation of the terminal acetylenes, thereby rendering the process atom economical.

Evano²⁵ reported a modular synthesis of polysubstituted indoles **105** from *N*-aryl-ynamides **104**. A bromine/lithium exchange of *N*-(2-bromoaryl)ynamides **104**, followed by a 5-*endo-dig* carbocupration afforded the substituted indoles efficiently (Scheme 19).

5.5. Silver

Malacria, Fentsterbank, and Aubert²⁶ reported a silver-catalyzed cycloisomerization of de novo allene-ynamides **106**, leading to amide-substituted cross-conjugated trienes **108** that are useful for tandem Diels-Alder cycloadditions (Scheme 20).

5.6. Gold

Hashimi²⁷ reported a gold-catalyzed cyclization of furanyl-ynamides **109** (Scheme 21). The course of the cyclization depended upon the length of the tether. Benzoannellated heterocycles **114** were produced when $n = 1$, while cyclopentadiene fused piperidines **118** were obtained when $n = 2$.

Skrydstrup²⁸ reported Au(I)-catalyzed hydroaminations or hydrations of diynamides **119** to access 2,5-diamidopyrroles **121** and 2,5-diamidofurans **122**, respectively (Scheme 22). This development represents a clever application of diynamides.

Liu²⁹ reported highly regioselective Au(I)-catalyzed oxidative ring-expansions of cyclopropyl-substituted ynamides **123** using Ph₂SO (Scheme 23). The ring expansion is not believed to proceed through an α -keto gold carbenoid intermediate but through **124**. Subsequently, Li³⁰ independently reported a similar gold-catalyzed oxidative ring-expansion (details not shown here).

Liu³¹ then explored 1,5-enynamides **127**, developing a Au(I)-catalyzed oxidative cyclization using 8-methylquinoline *N*-oxide **128** as the external oxidant to construct 3-carboxyamidoindenes **131** (Scheme 24). The transformation is believed to proceed through α -ketocarbenoid **129**.

Li³² reported a clever synthesis of 3-*aza*-bicyclo[3.1.0]hexan-2-one derivatives **135** via Au(I)-catalyzed oxidative cyclopropanations of *N*-allylynamides **132** using pyridine *N*-oxide as the external oxidant (Scheme 25).

Sueda³³ reported both Ag(I)- and/or Au(I)-catalyzed cyclizations of de novo ynamides **136**, which could be accessed for the first time. These cyclizations gave β -ketoimides **140** when using both Au(I) and Ag(I), while providing oxazoles **143** when using only Ag(I) (Scheme 26).

Hashmi³⁴ reported synthesis of highly functionalized cyclopentadienes **152** in moderate to good yields via Au(I)-catalyzed intermolecular cyclization of propargylic carboxylates **144** and ynamides **145** (Scheme 27).

Sahoo³⁵ developed Au(I)-catalyzed hydrative cyclization of easily accessible 5-yne-ynamides **153**, giving substituted 1,6-dihydropyridin-2(3*H*)ones **157** in good to excellent yields (Scheme 28). A mechanism involving a 6-*endo-dig* cyclization of intermediate **155** was proposed.

Bertrand³⁶ reported that deprotonation of oxazolium salt **159** initiated an interesting ring opening process giving ynamide **160** (Scheme 29). The acyclic ynamide readily reacts with various transition metals affording robust mesoionic carbene complexes **161**.

6. Cycloadditions and Formal Cycloadditions

6.1 [2 + 1]

Hsung³⁷ reported the first examples of stereoselective intramolecular cyclopropanations via a de novo class of push-pull carbenes derived from DMDO-epoxidations of chiral ynamides

162 (Scheme 30). This tandem epoxidation-cyclopropanation afforded a series of structurally unique amido-cyclopropanes **165**.

Buono³⁸ reported an unusual palladium catalyzed [2 + 1] cycloaddition of ynamides **166** with norbornene derivatives **167** giving various substituted aminomethylenecyclopropanes **170** (Scheme 31). Based on their previous study in alkyne system,³⁹ this process may involve a [2 + 2] cycloaddition of palladium vinylidene species with the double bonds of norbornene derivatives, followed by reductive elimination to furnish cyclopropanes.

6.2 [2 + 2]

Hsung⁴⁰ reported the first successful example of Ficini's [2 + 2] cycloaddition of ynamides **171** with enones **172** using CuCl₂ and AgSbF₆ as catalysts (Scheme 32).

Mezzetti⁴¹ subsequently reported Cu(OTf)₂ promoted Ficini [2 + 2] cycloadditions of ynamides with unsaturated β -keto esters in addition to a beautiful asymmetric variant using dicationic ruthenium(II)/PNNP complex (Scheme 33).

Danheiser⁴² reported the first examples of thermal [2 + 2] cycloadditions of 2-iodoynamides **181** with ketenes **182**, leading to 3-amido-2-iodocyclobutenones **183** (Scheme 34).

Lam⁴³ reported a [2 + 2] cycloaddition of ynamides **184** with nitroalkenes **185** catalyzed by a dirhodium complex and sodium tetraphenylborate, leading to nitro-substituted cyclobutenamides **189** and **190** (Scheme 35).

Takasu and Takemoto⁴⁴ reported selective syntheses of either *syn* or *anti* isomers of α,β -unsaturated amidines **195** and **198** through a tandem cascade of *aza*-[2 + 2] cycloaddition–4 π -electron pericyclic ring-opening by using Tf₂NH or CSA as catalyst, respectively (Scheme 36). The torquoselectivity of ring-opening was controlled by the Brønsted acidity of the catalyst and the polarity of the solvent.

Mikami⁴⁵ reported an asymmetric [2 + 2] cycloaddition of ynamide **199** with ethyl trifluoropyruvate **200** using a dicationic (*S*)-BINAP-Pd catalyst with excellent yield and enantioselectivity. This is also the first enantioselective synthesis of a stable oxetene derivative (Scheme 37).

6.3. [3 + 2]

Lam⁴⁶ reported a Rh-catalyzed formal [3 + 2] cycloaddition of ynamides **202** with arylboronic acids or esters **203** containing an electrophilic functional group at the *ortho*-position. This transformation effectively provides 2-amido-indenols **205** or 2-amido-indenes **206** in good regioselectivities (Scheme 38).

Sueda⁴⁷ applied their ynamides **207** to the copper-mediated Huisgen's azide-[3 + 2] cycloaddition giving 4-phthalimido-1-benzyl-1,2,3-triazole **208** (Scheme 39).

Davies⁴⁸ reported that 2,4,5-trisubstituted oxazoles **211** could be synthesized from ynamides **210** and 1,3-*N,O*-dipole equivalents **209** in a gold (III) catalyzed process (Scheme 40).

Gagosz and Skrydstrup⁴⁹ reported syntheses of cyclopentadienes **214** or tricycles **215** from dimerizations of ynamides **212** in the presence of a Au(I) complex (Scheme 41). While the divergence in this dimerization depends upon the substitution pattern, its efficiency is directly dependent on the electronic properties of the ynamide, which acts both as the electrophile and the nucleophile in the process.

Lai⁵⁰ reported the synthesis of δ -carbolines **220** from 2-iodoanilines **216** and *N*-tosyl-enynamides **217** via a Pd(0)-catalyzed cascade (Scheme 42). This cascade involves Larock's heteroannulation giving indoles **218** and an electrocyclization of 2-*aza*-trienes **219** after loss of TsOH.

Peng⁵¹ reported a AgOTf/Pd(OAc)₂ co-catalyzed [3 + 2]-cycloaddition of *N*-allyl-*N*-sulfonyl ynamides **221** with *N'*-(2-alkynyl-benzylidene)hydrazides **222** giving 2-amino-*H*-pyrazolo[5,1-*a*]isoquinolines **228** (Scheme 43). The transformation proceeds through 6-*endo-dig* cyclization of **222**, [3 + 2] cycloaddition between **224** and **225**, 3,3-sigmatropic rearrangement, and aromatization.

Batey⁵² reported a series of 1,3-dipole cycloadditions using ynehydrazides **229** that were first synthesized in these authors' lab. This exercise demonstrates the synthetic potential of these novel ynamides (Scheme 44).

6.4. [4 + 1]

Liu⁵³ reported a gold-catalyzed formal [4 + 1]-cycloaddition of ynamides **231** with 8-methylquinoline oxide **128**, leading to a series of substituted 2-amido-furans **237** (Scheme 45). Mechanistically, this formal cycloaddition likely proceeded through α -keto carbenoid **234** via an initial gold-catalyzed addition of 8-methylquinoline oxide **128** to ynamides **231** followed by an *oxa*-Nazarov cyclization.

6.5. [4 + 2]

Hoye⁵⁴ disclosed an intramolecular hexadehydro-Diels-Alder cycloaddition reaction of ynamide **238** (Scheme 46). The hexadehydro-Diels-Alder reaction of **238** led to the key benzyne intermediate **240**, which was trapped by the pendant silyloxy group, giving the tricycle **242** in 80% yield after an *O*-to-*C* silyl migration of the zwitter ionic intermediate **241**.

Lee⁵⁵ independently reported a similar hexadehydro-Diels-Alder cycloaddition of ynamide **243** catalyzed with silver (Scheme 47). What distinguishes this beautiful work from Hoye's is the alkane C-H insertion of the silver complex aryne to form carbon-carbon bonds.

Lee⁵⁶ subsequently reported a clever use of the Alder-ene process to trap the aryne intermediates derived from hexadehydro-Diels-Alder cycloaddition of ynamide **249** (Scheme 48). The metal catalyst was not essential, but increased the reaction's rate.

Danheiser⁵⁷ developed an ynamide-benzannulation using cyclobutenones **252** to synthesize highly substituted anilides **257** (Scheme 49). This benzannulation proceeds beautifully via four consecutive pericyclic processes, thereby constituting a formal [4 + 2] annulation. With olefin substitutions in R⁴ and on the nitrogen atom, these anilides could undergo ring-closing metathesis to generate complex *N*-heterocycles such as **258**.

6.6. [2 + 2 + 2]

Witulski⁵⁸ reported a ruthenium-catalyzed *hetero*-[2 + 2 + 2] cycloaddition of yne-ynamides **259** with Mander's reagent giving β - and γ -carbolines **260** and **261** (Scheme 50). The regioselectivity could be controlled by the steric hindrance of substitutions to the alkynes and ynamides. A total synthesis of eudistomin U **263** was achieved using regioselective β -carboline synthesis.

Nissen⁵⁹ subsequently reported the total synthesis of lavendamycin by Ru(II)-catalyzed *hetero*-[2 + 2 + 2] cycloaddition of ynamide **266** and an electron deficient nitrile to prepare the carboline scaffold (Scheme 51).

Witulski and Detert⁶⁰ later reported the total syntheses of perlolyrine **275** and isoperlolyrine **276**, featuring this Ru(II)-catalyzed *hetero*-[2 + 2 + 2] cycloaddition using yne-ynamides **271** and **273** (Scheme 52).

Malacria, Aubert and Gandon⁶¹ reported a Co(I)-catalyzed regioselective [2 + 2 + 2]-cycloaddition between ynamides **277** and nitriles **278** (Scheme 53). Through adjusting the substituent on the ynamides, regioselectivity of this cycloaddition could be tuned to favor either 3-aminopyridines **279** or 4-aminopyridines **280**.

Saito and Sato⁶² reported divergent total syntheses of (-)-herbindoles A-C through intramolecular [2 + 2 + 2] cycloaddition of ynamide **281** catalyzed by Wilkinson's catalyst (Scheme 54). All three herbindoles could be constructed from the common indoline intermediate **282**.

Liu⁶³ reported a Au(I)-catalyzed formal [2 + 2 + 2] cycloaddition of ynamides **286** with two equivalents of enol ethers (Scheme 55). Under activation by gold, two consecutive nucleophilic attacks by the enol ether followed by Prins-type cyclization furnished cyclic enamides **290** stereoselectively.

6.7. [3 + 2 + 2]

Saito⁶⁴ unveiled a novel Ni(0)-catalyzed [3 + 2 + 2] cycloaddition of ethyl cyclopropylideneacetate **291** with ynamides **292** (Scheme 56). The desired cycloadducts **293** were obtained in moderate yields with significant amounts of trisubstituted benzenes **294** resulting from trimerizations of the corresponding ynamides.

7. Rearrangements

Hsung⁶⁵ communicated the synthesis of azapin-2-one **299** via a sequence of *aza*-Claisen rearrangement, Pd(0)-catalyzed Overman rearrangement after trapping of ketenimine **296** with allyl alcohol, and ring-closing metathesis (Scheme 57).

Hsung⁶⁶ reported a novel synthesis of α,β -unsaturated cyclopentenimine **305** via a Pd-catalyzed *aza*-Rautenstrauch-type cyclization⁶⁷ (Scheme 58). It was proposed that the ynamido- π -allyl complex **301** derived from the oxidative addition of TIPS-terminated ynamide **295** underwent a Pd-[3,3] sigmatropic rearrangement giving the α -imino palladium carbenoid **302**, which is related to the key intermediates proposed in the Rautenstrauch cyclization.

Later, Hsung and DeKorver⁶⁸ discovered that the substrate scope for the Pd-catalyzed carbocyclization was quite broad. A variety of functionalized *N*-allyl- γ -branched ynamides were employed in cyclopentenimine synthesis. With *N*-sulfonyl ynamides **306**, palladium catalysis is required, as facile 1,3-sulfonyl shifts dominate under thermal conditions. However, since no analogous 1,3-phosphoryl shift is operational, *N*-phosphoryl ynamides **309** were used to prepare similar cyclopentenimines under thermal conditions through zwitter ionic intermediates **311** that underwent *N*-promoted *H*-shifts (Scheme 59).

DeKorver and Hsung⁶⁹ described a moderately stereoselective Staudinger-type ketenimine-imine [2 + 2] cycloaddition using *N*-phosphoryl ynamide **313** giving azetidin-2-imine **315**

bearing a quaternary carbon center (Scheme 60). The ketenimine intermediate **314** was generated in situ via an *aza*-Claisen reaction.

DeKorver and Hsung⁷⁰ showcased a tandem *aza*-Claisen carbocyclization of *N*-phosphoryl-*N*-allyl-ynamides that included possibilities such as ring-expansion via Meerwein-Wagner rearrangement and polyene-type cyclizations, thereby rapidly building structural complexity leading to fused bi- and tricyclic scaffolds (Scheme 61).

Hsung⁷¹ communicated a stereoselective synthesis of bridged or fused bicycloimines through a crossed or fused intramolecular [2 + 2] cycloaddition of ketenimines via palladium-catalyzed *aza*-Claisen rearrangements of *N*-allylynamides **330** and **333** (Scheme 62). Preference of cycloaddition pathways depended upon alkene substitutions.

Meyer and Cossy⁷² described an interesting Saucey-Marbet rearrangement of ynamide **336** containing an *N*-Bocglycinate motif, providing stereoselective access to functionalized allenamide **339**^{73,74} (Scheme 63). These de novo allenamides underwent silver catalyzed cyclization affording 3-pyrrolidine derivatives **340** having 2,5-*syn* relative stereochemistry.

8. Conclusion

This manuscript has highlighted recent advances in ynamide cyclization reactions that are useful in the rapid synthesis of cyclic and polycyclic structural motifs. The transformations presented are significant due to their rapid assembly of structural complexity, and inclusion of nitrogen within, or in close proximity to, the newly formed ring or rings. Thus, ynamides have emerged as powerful synthons for nitrogen-containing heterocycles and nitrogen-substituted rings. We hope this Account will help promote continued interest in the chemistry of ynamides.

Acknowledgments

We wish to thank NIH [GM066055] for funding. BLK also acknowledges the UW-Oshkosh Faculty Development Program for financial support.

Biographies

Xiao-Na Wang received a B.S. in chemistry from Nanyang Normal University in 2006, and her Ph.D. degree in 2011 at Institute of Chemistry, Chinese Academy of Sciences with Professor Song Ye. Currently, she is conducting postdoctoral research with Professor Richard Hsung at the University of Wisconsin.

Hyu-Suk Yeom received a B.S in chemistry from Hanyang University in 2007 and his Ph.D. degree in 2012 with Professor Shin at Hanyang University. Currently, he is conducting postdoctoral research with Professor Richard Hsung at the University of Wisconsin.

Lichao Fang received a B.S. in chemistry from Nankai University in 2007. He carried out doctoral research with Professor Zhen Yang at Peking University and obtained his Ph.D. degree in 2012. Currently, he is conducting postdoctoral research with Professor Richard Hsung at the University of Wisconsin.

Shuzhong He received his B.S. degree from Peking University in 2004. He then joined Professor Chi-Sing Lee's group in Peking University Shenzhen Graduate School and obtained his Ph.D. degree in 2011. Currently, he is a postdoctoral research scholar with Professor Richard Hsung at the University of Wisconsin.

Zhi-Xiong Ma received a B.S. in chemistry from University of Science and Technology of China in 2005, and his Ph.D. degree in 2010 at Shanghai Institute of Organic Chemistry with Professor Gang Zhao. Currently, he is conducting postdoctoral research with Professor Richard Hsung at the University of Wisconsin.

Brant L. Kedrowski obtained a B.S. in chemistry in 1994, and Ph.D. in organic chemistry in 2000 with Professor Wayland Noland at the University of Minnesota. He did postdoctoral work with Professor Clayton Heathcock at the University of California Berkeley before his appointment as an assistant professor at the University of Wisconsin Oshkosh in 2002. He was promoted to associate professor in 2008 and to full professor in 2013.

Richard P. Hsung obtained his B.S. in chemistry and mathematics from Calvin College and attended The University of Chicago for his M.S. and Ph.D. degrees in organic chemistry, respectively, with Professors Jeff Winkler and Bill Wulff. After postdoctoral stays with Professor Larry Sita in Chicago and Professor Gilbert Stork at Columbia University, he moved to University of Minnesota as an assistant professor in 1997. He was promoted to associate professor in 2002 and to full professor after moving to University of Wisconsin in 2006. He has coauthored over 200 publications and supervised over 150 students and postdoctoral fellows with research interests in developing stereoselective methods using allenamides, ynamides, enamides, and cyclic acetals, and applications in natural product syntheses.

References


1. DeKorver KA, Li H, Lohse AG, Hayashi R, Lu Z, Zhang Y, Hsung RP. Ynamides: A Modern Functional Group for the New Millennium. *Chem. Rev.* 2010; 110:5064–5106. [PubMed: 20429503]
2. Evano G, Coste A, Jouvin K. Ynamides: Versatile Tools in Organic Synthesis. *Angew. Chem. Int. Ed.* 2010; 49:2840–2859.
3. For Partial Reviews, see: Ackermann L, Potukuchi HK. Regioselective Syntheses of Fully-Substituted 1,2,3-Triazoles: The CuAAC/C - H Bond Functionalization Nexus. *Org. Biomol. Chem.* 2010; 8:4503–4513. [PubMed: 20733972] Domínguez G, Perez-Castells J. Recent Advances in [2 + 2] Cycloaddition Reactions. *Chem. Soc. Rev.* 2011; 40:3430–3444. [PubMed: 21431173] Weding N, Hapke M. Preparation and Synthetic Applications of Alkene Complexes of Group 9 Transition Metals in [2 + 2 + 2] Cycloaddition Reactions. *Chem. Soc. Rev.* 2011; 40:4525–4538. [PubMed: 21594299] Madelaine C, Valerio V, Maulide N. Revisiting Keteniminium Salts: More than the Nitrogen Analogs of Ketenes. *Chem. Asian J.* 2011; 6:2224–2239. [PubMed: 21721132]
4. Evano G, Jouvin K, Coste A. General Amination Reactions for the Synthesis of Ynamides. *Synthesis.* 2013; 45:17–26.
5. Also see: Mulder JA, Kurtz KCM, Hsung RP. In Search of an Atom-Economical Synthesis of Chiral Ynamides. *Synlett.* 2003; 10:1379–1390. Dehli JR, Legros J, Bolm C. Synthesis of Enamines, Enol Ethers and Related Compounds by Cross-coupling Reactions. *Chem. Commun.* 2005; 43:973–986. Tracey MR, Hsung RP, Antoline JA, Kurtz KCM, Shen L, Slafer BW, Zhang Y, Weinreb SM. Three Carbon-Heteroatom Bonds: Amides and Derivatives; Peptides; Lactams. Product Class 4: NArylalkanamides, Ynamides, Enamides, Dienamides, and Allenamides. *Science of Synthesis, Houben-Weyl Methods of Molecular Transformations.* 2005 Georg Thieme Verlag KG Stuttgart, Germany Chapter 21.4 Evano G, Blanchard N, Toumi M. Copper-Mediated Coupling Reactions and Their Applications in Natural Products and Designed Biomolecules Synthesis. *Chem. Rev.* 2008; 108:3054–3131. [PubMed: 18698737]
6. a Chen Z, Zheng D, Wu J. A Facile Route to Polysubstituted Indoles via Three-Component Reaction of 2-Ethynylaniline, Sulfonyl Azide, and Nitroolefin. *Org. Lett.* 2011; 13:848–851. [PubMed: 21299243] b Yavari I, Nematpour M. Copper-Catalyzed One-Pot Synthesis of Functionalized 1,4-Dihydroazete Derivatives from Sulfonyl Azides, Terminal Alkynes, and Tetramethylguanidine. *Synlett.* 2012; 23:2215–2218. c Jiang Z, Lu P, Wang Y. Three-Component Reaction of Propargyl

- Amines, Sulfonyl Azides, and Alkynes: One-Pot Synthesis of Tetrasubstituted Imidazoles. *Org. Lett.* 2012; 14:6266–6269. [PubMed: 23193963] d Yavari I, Nematpour M. Copper-Catalyzed Tandem Synthesis of Tetrasubstituted Pyrimidines from Alkynes, Sulfonyl Azides, Trichloroacetonitrile, and Tetramethylguanidine. *Synlett.* 2013; 24:165–168. e Cheng D, Ling F, Li Z, Yao W, Ma C. Three-Component Assembly of Conjugated Enyne Scaffolds via E-Selective Olefination of Ynals. *Org. Lett.* 2012; 14:3146–3149. [PubMed: 22671022] f Li B-S, Yang B-M, Wang S-H, Zhang Y-Q, Cao X-P, Tu Y-Q. Copper(I)-Catalyzed Intramolecular [2 + 2] Cycloaddition of 1,6-Enyne-Derived Ketenimine: An Efficient Construction of Strained and Bridged 7-Substituted-3-heterobicyclo[3.1.1]heptan-6-one. *Chem. Sci.* 2012; 3:1975–1979.
7. Balieu S, Toutah K, Carro L, Chamoreau L-M, Rousselière H, Courillon C. Radical Cyclization of Ynamides into Six- or Eight-membered Rings. Application to the Synthesis of a Protoberberine Analog. *Tetrahedron Lett.* 2011; 52:2876–2880.
 8. Chemla F, Dulong F, Ferreira F, Nüllen MP, Pérez-Luna A. Radical Zinc-Atom Transfer Based Multicomponent Approaches to 3-Alkylidene-Substituted Tetrahydrofurans. *Synthesis.* 2011; 9:1347–1360.
 9. Wakamatsu H, Sakagami M, Hanata M, Takeshita M, Mori M. Ring-Closing Metathesis of Ene-Ynamide: Application to the Synthesis of Medium-Sized Cyclic Dienamide. *Macromol. Symp.* 2010; 293:5–9.
 10. Poloukhine A, Rassadin V, Kuzmin A, Popik VV. Nucleophilic Cycloaromatization of Ynamide-Terminated Eneynes. *J. Org. Chem.* 2010; 75:5953–5962. [PubMed: 20684502]
 11. a Gati W, Rammah MM, Rammah MB, Couty F, Evano G. De Novo Synthesis of 1,4-Dihydropyridines and Pyridines. *J. Am. Chem. Soc.* 2012; 134:9078–9081. [PubMed: 22583001]
b Gati W, Rammah MM, Rammah MB, Evano G. Intramolecular Carbolithiation of N-Allylynamides: An Efficient Entry to 1,4-Dihydropyridines and Pyridines – Application to a Formal Synthesis of Sarizotan. *Beilstein J. Org. Chem.* 2012; 8:2214–2222. [PubMed: 23365632]
 12. Kerr DJ, Miletic M, Chaplin JH, White JM, Flynn BL. Oxazolidinone-Promoted, Torquoselective Nazarov Cyclizations. *Org. Lett.* 2012; 14:1732–1735. [PubMed: 22455473]
 13. Kong Y, Jiang K, Cao J, Fu L, Yu L, Lai G, Cui Y, Hu Z, Wang G. Synthesis of 3-Alkyl- or 3-Allenyl-2-amidobenzofurans via Electrophilic Cyclization of *o*-Anisole-Substituted Ynamides with Carbocations. *Org. Lett.* 2013; 15:422–425. [PubMed: 23301862]
 14. Kong Y, Yu L, Fu L, Cao J, Lai G, Cui Y, Hu Z, Wang G. Electrophilic Cyclization of *o*-Anisole- and *o*-Thioanisole-Substituted Ynamides: Synthesis of 2-Amidobenzofurans and 2-Amidobenzothiophenes. *Synthesis.* 2013; 45:1975–1982.
 15. Wang X-N, Hsung RP, Qi R, Fox SK, Lv M-C. A Highly Stereoselective Addition of Lithiated Ynamides to Ellman-Davis Chiral *N*-tert-Butanesulfinyl Imines. *Org. Lett.* 2013; 15:2514–2517. [PubMed: 23646900]
 16. Rui Q, Wang X-N, DeKorver KA, Tang Y, Wang C-C, Li Q, Li H, Lv M-C, Yu Q, Hsung RP. A Convenient Synthesis of γ -Amino-Ynamides via Additions of Lithiated Ynamides to Aryl Imines; Observation of an *Aza*-Meyer-Schuster Rearrangement. *Synthesis.* 2013; 45:1749–1758. [PubMed: 23976795]
 17. Nishimura T, Takiguchi Y, Maeda Y, Hayashi T. Rhodium-Catalyzed Asymmetric Cycloisomerization of 1,6-Eneynamides. *Adv. Synth. Catal.* 2013; 355:1374–1382.
 18. Liu R, Winston-McPherson GN, Yang Z-Y, Zhou X, Song W, Guzei IA, Xu X, Tang W. Generation of Rhodium(I) Carbenes from Ynamides and Their Reactions with Alkynes and Alkenes. *J. Am. Chem. Soc.* 2013; 135:8201–8204. [PubMed: 23701315]
 19. Greenaway RL, Campbell CD, Holton OT, Russell CA, Anderson EA. Palladium-Catalyzed Cascade Cyclization of Ynamides to Azabicycles. *Chem.-Eur. J.* 2011; 17:14366–14370. [PubMed: 22113952]
 20. Greenaway RL, Campbell CD, Chapman HA, Anderson EA. Reductive Cyclization of Bromoenynamides with Alcohols as Hydride Source: Synthesis and Reactions of 2-Amidodienes. *Adv. Synth. Catal.* 2012; 354:3187–3194.
 21. Bhunia S, Chang C-J, Liu R-S. Platinum-Catalyzed Oxoarylations of Ynamides with Nitrones. *Org. Lett.* 2012; 14:5522–5525. [PubMed: 23088158]


22. Xu H, Zhang Y, Huang J, Chen W. Copper-Catalyzed Synthesis of N-Fused Heterocycles through Regioselective 1,2-Aminothioloation of 1,1-Dibromoalkenes. *Org. Lett.* 2010; 12:3704–3707. [PubMed: 20669979]
23. Hashmi ASK, Schuster AM, Zimmer M, Rominger F. Synthesis of 5-Halo-4*H*-1,3-oxazine-6-amines by a Copper-Mediated Domino Reaction. *Chem.-Eur. J.* 2011; 17:5511–5515. [PubMed: 21491523]
24. Li J, Neuville L. Copper-Catalyzed Oxidative Diamination of Terminal Alkynes by Amidines: Synthesis of 1,2,4-Trisubstituted Imidazoles. *Org. Lett.* 2013; 15:1752–1755. [PubMed: 23528104]
25. Gati W, Couty F, Boubaker T, Rammah MM, Rammah MB, Evano G. Intramolecular Carbocupration of N-Aryl-ynamides: A Modular Indole Synthesis. *Org. Lett.* 2013; 15:3122–3125. [PubMed: 23734901]
26. Garcia P, Harrak Y, Diab L, Cordier P, Ollivier C, Gandon V, Malacria M, Fensterbank L, Aubert C. Silver-Catalyzed Cycloisomerization of 1,*n*-Allenynamides. *Org. Lett.* 2011; 13:2952–2955. [PubMed: 21534621]
27. Hashmi ASK, Pankajakshan S, Rudolph M, Enns E, Bander T, Rominger F, Frey W. Gold Catalysis: Anellated Heterocycles and Dependency of the Reaction Pathway on the Tether Length. *Adv. Synth. Catal.* 2009; 351:2855–2875.
28. Kramer S, Madsen JLH, Rottländer M, Skrydstrup T. Access to 2,5-Diamidopyrroles and 2,5-Diamidofurans by Au(I)-Catalyzed Double Hydroamination or Hydration of 1,3-Diynes. *Org. Lett.* 2010; 12:2758–2761. [PubMed: 20462239]
29. Li C-W, Pati K, Lin G-Y, Abu Sohel SM, Hung H-H, Liu R-S. Gold-Catalyzed Oxidative Ring Expansions and Ring Cleavages of Alkynylcyclopropanes by Intermolecular Reactions Oxidized by Diphenylsulfoxide. *Angew. Chem., Int. Ed.* 2010; 49:9891–9894.
30. Xu C-F, Xu M, Jia Y-X, Li C-Y. Gold-Catalyzed Synthesis of Benzil Derivatives and α -Keto Imides via Oxidation of Alkynes. *Org. Lett.* 2011; 13:1556–1559. [PubMed: 21332143]
31. Vasu D, Hung H-H, Bhunia S, Gawade SA, Das A, Liu R-S. Gold-Catalyzed Oxidative Cyclization of 1,5-Enynes Using External Oxidants. *Angew. Chem., Int. Ed.* 2011; 50:6911–6914.
32. Wang K-B, Ran R-Q, Xiu S-D, Li C-Y. Synthesis of 3-Azabicyclo[3.1.0]hexan-2-one Derivatives via Gold-Catalyzed Oxidative Cyclopropanation of *N*-Allylynamides. *Org. Lett.* 2013; 15:2374–2377. [PubMed: 23651134]
33. Sueda T, Kawada A, Urashi Y, Teno N. Ag- and Au-Catalyzed Addition of Alcohols to Ynimides: β -Regioselective Carbonylation and Production of Oxazoles. *Org. Lett.* 2013; 15:1560–1563. [PubMed: 23496249]
34. Rettenmeier E, Schuster AM, Rudolph M, Rominger F, Gade CA, Hashmi ASK. Gold Catalysis: Highly Functionalized Cyclopentadienes Prepared by Intermolecular Cyclization of Ynamides and Propargylic Carboxylates. *Angew. Chem. Int. Ed.* 2013; 52:5880–5884.
35. Ghosh N, Nayak S, Sahoo AK. Gold(I)-Catalyzed 6-Endo-Dig Hydrative Cyclization of an Alkyne-Tethered Ynamide: Access to 1,6-Dihydropyridin-2(3*H*)ones. *Chem.-Eur. J.* 2013; 19:9428–9433. [PubMed: 23788463]
36. Ung G, Mendoza-Espinosa D, Bertrand G. Ynamides: Stable Ligand Equivalents of Unstable Oxazol-4-ylidenes (Novel Mesoionic Carbenes). *Chem. Commun.* 2012; 48:7088–7090.
37. Li H, Antoline JE, Yang J-H, Al-Rashid ZF, Hsung RP. A Stereoselective Intramolecular Cyclopropanation via a de novo Class of Push - Pull Carbenes Derived from DMDO-Epoxidations of Chiral Ynamides. *New J. Chem.* 2010; 34:1309–1316.
38. Clavier H, Lepronier A, Bengobesse-Mintsa N, Gatineau D, Pellissier H, Giordano L, Tenaglia A, Buono G. Palladium-Mediated [2 + 1] Cycloaddition of Norbornene Derivatives with Ynamides. *Adv. Synth. Catal.* 2013; 355:403–408.
39. Bigeault J, Giordano L, Buono G. [2 + 1] Cycloadditions of Terminal Alkynes to Norbornene Derivatives Catalyzed by Palladium Complexes with Phosphinous Acid Ligands. *Angew. Chem., Int. Ed.* 2005; 44:4753–4757.
40. Li H, Hsung RP, DeKorver KA, Wei Y. Copper-Catalyzed Ficini [2 + 2] Cycloaddition of Ynamides. *Org. Lett.* 2010; 12:3780–3783. [PubMed: 20690601]

41. a Schotes C, Bigler R, Mezzetti A. Bicyclo[3.2.0]heptane-Based Enamides by Ru/PNNP-Catalyzed Enantioselective Ficini Reactions: Scope and Application in Ligand Design. *Synthesis*. 2012; 44:513–526. b Schotes C, Althaus M, Aardoom R, Mezzetti A. Asymmetric Diels–Alder and Ficini Reactions with Alkylidene β -Ketoesters Catalyzed by Chiral Ruthenium PNNP Complexes: Mechanistic Insight. *J. Am. Chem. Soc.* 2012; 134:1331–1343. [PubMed: 22191539] c Schotes C, Mezzetti A. Enantioselective Ficini Reaction: Ruthenium/PNNP-Catalyzed [2 + 2] Cycloaddition of Ynamides with Cyclic Enones. *Angew. Chem., Int. Ed.* 2011; 50:3072–3074. d Schotes C, Mezzetti A. Cu(I)- and Cu(II)-Catalyzed Cyclo- and Michael Addition Reactions of Unsaturated β -Ketoesters. *J. Org. Chem.* 2011; 76:5862–5866. [PubMed: 21639117]
42. Wang Y-P, Danheiser RL. Synthesis of 2-Iodoynamides and Regioselective [2 + 2] Cycloadditions with Ketene. *Tetrahedron Lett.* 2011; 52:2111–2114. [PubMed: 21479121]
43. Smith DL, Chidipudi SR, Goundry WR, Lam HW. Rhodium-Catalyzed [2 + 2] Cycloaddition of Ynamides with Nitroalkenes. *Org. Lett.* 2012; 14:4934–4937. [PubMed: 22954424]
44. Shindoh N, Kitaura K, Takemoto Y, Takasu K. Catalyst-Controlled Torquoselectivity Switch in the 4π Ring-Opening Reaction of 2-Amino-2-azetines Giving β -Substituted α,β -Unsaturated Amidines. *J. Am. Chem. Soc.* 2011; 133:8470–8473. [PubMed: 21557577]
45. Aikawa K, Hioki Y, Shimizu N, Mikami K. Catalytic Asymmetric Synthesis of Stable Oxetenes via Lewis Acid-Promoted [2 + 2] Cycloaddition. *J. Am. Chem. Soc.* 2011; 133:20092–20095. [PubMed: 22070285]
46. Gourdet B, Rudkin ME, Lam HW. Rhodium-Catalyzed Annulation of Ynamides with Bifunctional Arylboron Reagents. *Org. Lett.* 2010; 12:2554–2557. [PubMed: 20446723]
47. Sueda T, Oshima A, Teno N. N-Alkynyl Imides (Ynimides): Synthesis and Use as a Variant of Highly Labile Ethynamine. *Org. Lett.* 2011; 13:3996–3999. [PubMed: 21710984]
48. Davies PW, Cremonesi A, Dumitrescu L. Intermolecular and Selective Synthesis of 2,4,5-Trisubstituted Oxazoles by a Gold-Catalyzed Formal [3 + 2] Cycloaddition. *Angew. Chem. Int. Ed.* 2011; 50:8931–8935.
49. Kramer S, Odabachian Y, Overgaard J, Rottländer M, Gagosz F, Skrydstrup T. Taking Advantage of the Ambivalent Reactivity of Ynamides in Gold Catalysis: A Rare Case of Alkyne Dimerization. *Angew. Chem. Int. Ed.* 2011; 50:5090–5094.
50. Cao J, Xu Y, Kong Y, Cui Y, Hu Z, Wang G, Deng Y, Lai G. Synthesis of δ -Carbolines via a Pd-Catalyzed Sequential Reaction from 2-Iodoanilines and *N*-Tosyl-enynamines. *Org. Lett.* 2012; 14:38–41. [PubMed: 22126429]
51. Huang P, Chen Z, Yang Q, Peng Y. Silver Triflate and Palladium Acetate Co-catalyzed Reaction of *N'*-(2-Alkynylbenzylidene)hydrazide with *N*-Allyl Ynamide. *Org. Lett.* 2012; 14:2790–2793. [PubMed: 22616888]
52. Beveridge RE, Batey RA. Terminal Alkyne Addition to Diazodicarboxylates: Synthesis of Hydrazide Linked Alkynes (Ynehydrazides). *Org. Lett.* 2012; 14:540–543. [PubMed: 22216967]
53. Dateer RB, Pati K, Liu R-S. Gold-Catalyzed Synthesis of Substituted 2-Aminofurans via Formal [4 + 1]-Cycloadditions on 3-En-1-ynamides. *Chem. Commun.* 2012; 48:7200–7202.
54. Hoye TR, Baire B, Niu D, Willoughby PH, Woods BP. The Hexadehydro-Diels-Alder Reaction. *Nature*. 2012; 490:208–212. [PubMed: 23060191]
55. Yun SY, Wang K-P, Lee N-K, Mamidipalli P, Lee D. Alkane C–H Insertion by Aryne Intermediates with a Silver Catalyst. *J. Am. Chem. Soc.* 2013; 135:4668–4671. [PubMed: 23477300]
56. Karmakar R, Mamidipalli P, Yun SY, Lee D. Alder-Ene Reactions of Arynes. *Org. Lett.* 2013; 15:1938–1941. [PubMed: 23540274]
57. Mak XY, Crombie AL, Danheiser RL. Synthesis of Polycyclic Benzofused Nitrogen Heterocycles via a Tandem Ynamide Benzannulation/Ring-Closing Metathesis Strategy. Application in a Formal Total Synthesis of (+)-FR900482. *J. Org. Chem.* 2011; 76:1852–1873. [PubMed: 21322545]
58. Nissen F, Richard V, Alayrac C, Witulski B. Synthesis of β - and γ -Carbolines via Ruthenium and Rhodium Catalysed [2 + 2 + 2] Cycloadditions of Yne-ynamides with Methylcyanoformate. *Chem. Commun.* 2011; 47:6656–6658.


59. Nissen F, Detert H. Total Synthesis of Lavendamycin by a [2 + 2 + 2] Cycloaddition. *Eur. J. Org. Chem.* 2011;2845–2853.
60. Dassonneville B, Witulski B, Detert H. [2 + 2 + 2] Cycloadditions of Alkynylynamides-A Total Synthesis of Perlolyrine and the First Total Synthesis of “Isoperlolyrine”. *Eur. J. Org. Chem.* 2011;2836–2844.
61. a Garcia P, Evanno Y, George P, Sevrin M, Ricci G, Malacria M, Aubert C, Gandon V. Regioselective Cobalt-Catalyzed Formation of Bicyclic 3- and 4-Aminopyridines. *Org. Lett.* 2011; 13:2030–2033. [PubMed: 21413688] b Garcia P, Evanno Y, George P, Sevrin M, Ricci G, Malacria M, Aubert C, Gandon V. Synthesis of Aminopyridines and Aminopyridones by Cobalt-Catalyzed [2 + 2 + 2] Cycloadditions Involving Yne-Ynamides: Scope, Limitations, and Mechanistic Insights. *Chem. Eur. J.* 2012; 18:4337–4344. [PubMed: 22383395]
62. Saito N, Ichimaru T, Sato Y. Total Synthesis of (–)-Herbindoles A, B, and C via Transition-Metal-Catalyzed Intramolecular [2 + 2 + 2] Cyclization between Ynamide and Diynes. *Org. Lett.* 2012; 14:1914–1917. [PubMed: 22452396]
63. Dateer RB, Shaibu BS, Liu R-S. Gold-Catalyzed Intermolecular [4 + 2] and [2 + 2 + 2] Cycloadditions of Ynamides with Alkenes. *Angew Chem., Int. Ed.* 2012; 51:113–117.
64. Yamasaki R, Terashima N, Sotome I, Komagawa S, Saito S. Nickel-Catalyzed [3 + 2 + 2] Cycloaddition of Ethyl Cyclopropylideneacetate and Heteroatom-Substituted Alkynes: Application to Selective Three-Component Reaction with 1,3-Diynes. *J. Org. Chem.* 2010; 75:480–483. [PubMed: 20014802]
65. DeKorver KA, North TD, Hsung RP. An Efficient Synthesis of de novo Imidates via Aza-Claisen Rearrangements of *N*-Allyl Ynamides. *Synlett.* 2010;2397–2402. [PubMed: 21278848]
66. DeKorver KA, Hsung RP, Lohse AG, Zhang Y. A Divergent Mechanistic Course of Pd(0)-Catalyzed Aza-Claisen Rearrangement and *Aza*-Rautenstrauch-Type Cyclization of *N*-Allyl Ynamides. *Org. Lett.* 2010; 12:1840–1843. [PubMed: 20337418]
67. Rautenstrauch V. 2-Cyclopentenones from 1-Ethynyl-2-propenyl Acetates. *J. Org. Chem.* 1984; 49:950–952.
68. Wang X-N, Winston-McPherson GN, Walton MC, Zhang Y, Hsung RP, DeKorver KA. Synthesis of Cyclopentenimines from *N*-Allyl Ynamides via a Tandem *Aza*-Claisen Rearrangement–Carbocyclization Sequence. *J. Org. Chem.* 2013; 78:6233–6244. [PubMed: 23718841]
69. DeKorver KA, Walton MC, North TD, Hsung RP. Introducing a New Class of *N*-Phosphoryl Ynamides via Cu(I)-Catalyzed Amidations of Alkynyl Bromides. *Org. Lett.* 2011; 13:4862–4865. [PubMed: 21848304]
70. DeKorver KA, Wang X-N, Walton MC, Hsung RP. Carbocyclization Cascades of Allyl Ketenimines via *Aza*-Claisen Rearrangements of *N*-Phosphoryl-*N*-allyl-ynamides. *Org. Lett.* 2012; 14:1768–1771. [PubMed: 22414252]
71. DeKorver KA, Hsung RP, Song W-Z, Wang X-N, Walton MC. An Intramolecular [2 + 2] Cycloaddition of Ketenimines via Palladium-Catalyzed Rearrangements of *N*-Allyl-Ynamides. *Org. Lett.* 2012; 14:3214–3217. [PubMed: 22667819]
72. Brioché J, Meyer C, Cossy J. Synthesis of Functionalized Allenamides from Ynamides by Enolate Claisen Rearrangement. *Org. Lett.* 2013; 15:1626–1629. [PubMed: 23496162]
73. Wei L-L, Xiong H, Hsung RP. The Emergence of Allenamides In Organic Synthesis. *Acc. Chem. Res.* 2003; 36:773–782. [PubMed: 14567711]
74. Lu T, Lu Z, Ma Z-X, Zhang Y, Hsung RP. Allenamides: A Versatile Synthetic Building Block in Organic Synthesis. *Chem. Rev.* 2013; 113:4862–4904. [PubMed: 23550917]




Scheme 1.




Scheme 2.




Scheme 3.




Scheme 4.




Scheme 5.




Scheme 6.




Scheme 7.




Scheme 8.




Scheme 9.




Scheme 10.




Scheme 11.




Scheme 12.




Scheme 13.




Scheme 14.




Scheme 15.




Scheme 16.




Scheme 17.




Scheme 18.




Scheme 19.




Scheme 20.




Scheme 21.




Scheme 22.




Scheme 23.




Scheme 24.




Scheme 25.




Scheme 26.




Scheme 27.




Scheme 28.




Scheme 29.




Scheme 30.




Scheme 31.




Scheme 32.




Scheme 33.




Scheme 34.




Scheme 35.




Scheme 36.




Scheme 37.




Scheme 38.




Scheme 39.




Scheme 40.




Scheme 41.




Scheme 42.




Scheme 43.




Scheme 44.




Scheme 45.




Scheme 46.




Scheme 47.




Scheme 48.




Scheme 49.




Scheme 50.




Scheme 51.




Scheme 52.




Scheme 53.




Scheme 54.




Scheme 55.




Scheme 56.




Scheme 57.




Scheme 58.




Scheme 59.




Scheme 60.



Scheme 61.



Scheme 62.



Scheme 63.