

ORIGINAL ARTICLE

**Zinc toxicity on antioxidative response in (*Zea mays* L.) at two  
different pH**

Zahra Hosseini and Latifeh Poorakbar\*

*Department of Biology, Faculty of Sciences, Urmia University, Iran*

Tel.: +989144400987; fax: +98 (441) 2776707

\*E-Mail: [l.pourakbar@urmia.ac.ir](mailto:l.pourakbar@urmia.ac.ir)

Received September 22, 2012

Zn is the second most abundant transition metal after iron (Fe). Excess Zn can have negative effects on plants. The effect of Zn at two different pH on lipid peroxidation (MDA), membrane permeability (EC), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), non-protein thiols (NPT) and the activities of major antioxidant enzymes *Zea mays* were investigated under controlled growth conditions. Zn-excess conditions increased the EC, MDA, H<sub>2</sub>O<sub>2</sub> content and non-protein thiols and also activities of antioxidant enzymes were increased. Also zinc toxicity was higher in 4.5 pH than 7.5 pH.

*Key words: Zinc, pH, Antioxidant enzymes, MDA, H<sub>2</sub>O<sub>2</sub>, EC, NPT*

## ORIGINAL ARTICLE

## Zinc toxicity on antioxidative response in (*Zea mays* L.) at two different pH

Zahra Hosseini and Latifeh Poorakbar\*

Department of Biology, Faculty of Sciences, Urmia University, Iran

Tel.: +989144400987; fax: +98 (441) 2776707

\*E-Mail: [l.pourakbar@urmia.ac.ir](mailto:l.pourakbar@urmia.ac.ir)

Received September 22, 2012

Zn is the second most abundant transition metal after iron (Fe). Excess Zn can have negative effects on plants. The effect of Zn at two different pH on lipid peroxidation (MDA), membrane permeability (EC), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), non-protein thiols (NPT) and the activities of major antioxidant enzymes *Zea mays* were investigated under controlled growth conditions. Zn-excess conditions increased the EC, MDA, H<sub>2</sub>O<sub>2</sub> content and non-protein thiols and also activities of antioxidant enzymes were increased. Also zinc toxicity was higher in 4.5 pH than 7.5 pH.

**Key words:** Zinc, pH, Antioxidant enzymes, MDA, H<sub>2</sub>O<sub>2</sub>, EC, NPT

Soil pollution by toxic metals is a serious problem for the environment and is also one of the environmental stresses for higher plants. Evidence has suggested that heavy metal toxicity is manifested as oxidative stress, which is caused by the stimulation of free oxygen radical production (Gao *et al.*, 2011). Zinc is a heavy metal. It acts as a plant nutrient and plays an important role in many metabolic reactions but it is toxic at higher concentrations and causes growth inhibition (Vaillant *et al.*, 2005) and induction of oxidative damage in various plant species (Panda *et al.*, 2003). At the cellular level, excess Zn can significantly alter mitotic activity (Rout and Das,

2003), affect membrane integrity and permeability (Stoyanova and Doncheva, 2002), and even kill cells (Chang *et al.*, 2005). Various studies indicated that the availability of elements for leaching and plant uptake depend on their solubility in the soil solution (Gao *et al.*, 2010). As a result of the strong influence of pH on metal solubility (McBride *et al.*, 1997), anthropogenic processes which result in the lowering of substrate pH can cause metal toxicities, even if no extra metal has been added to the system (Robinson *et al.*, 1995). The goals of the recent study were to investigate the effect of zinc toxicity on antioxidative response and role of pH in toxicity on *Zea mays* plants.

## MATERIALS AND METHODS

Seeds were surface sterilized in 5% sodium hypochlorite solution for 10 minutes before use, to avoid fungal contamination. Seeds were then washed thoroughly with deionized, imbibed overnight in distilled water and germinated on moistened filter paper in trays for 5 days in darkness at 25°C. After 5 days, uniformly germinated seedlings were transferred to plastic cups. The plants were irrigated over a period 10 days with nutrient solution containing zinc between 0, 400 and 600  $\mu\text{M}$  as sulfate ( $\text{ZnSO}_4$ ) and two pH 4.5 and 7.5. After 12 days of Zn treatment, 9 plants from each group (control, Zn 400-pH4.5, Zn 400-pH7.5, Zn 600-pH4.5, Zn 600-pH7.5) were taken and divided into shoot and root fractions and analyzed.

Concentration of  $\text{H}_2\text{O}_2$  was determined following Jana and Choudhuri (1981). Lipid peroxidation was determined by measuring the amount of malondialdehyde (MDA) content according Heath and Packer (1968). Ascorbate peroxidase activity was measured according to Nakano and Asada (1981) by monitoring the rate of ascorbate oxidation at 290 nm. Catalase activity was assayed according to Aebi (1983). Guaiacol peroxidase activity was measured according to Upadhyaya *et al* (1985). GR Activity was assayed according to Foyer (1976). Percentage electrolyte leakage was assayed according to Lutts *et al* (1996). Non-protein thiols were measured with Edreva *et al* (1984).

### Statistical analysis

Results were based on three replicates and statistical analysis was performed using ANOVA (for  $P < 0.05$ ). Based on the ANOVA results, a Tukey test for mean comparison was performed, for a 95% confidential level, to test for significant differences among treatments.


## RESULTS

In the *Zea mays*, lipid peroxidation, which is measured as Malondialdehyde (MDA) content, increased at all Zn treatments than control and maximum increase was observed at 600  $\mu\text{M}$  (pH 4.5) (Fig. 1, A and B).  $\text{H}_2\text{O}_2$  content increased at all Zn treatments than those in the control while the maximum increase was observed at 600  $\mu\text{M}$  (pH 7.5). But no significant difference was observed between treatments and control (Fig. 1, C and D). Effect of zinc on APX and GR activities is shown in Fig. 2. APX and GR activity increased in all Zn treatments in both shoot and root compared with control. The highest value was occurred in the 600  $\mu\text{M}$  (pH 4.5) Zn treatment. APX and GR activity were higher in 600  $\mu\text{M}$  than 400  $\mu\text{M}$  concentration of zinc. But no significant difference was observed between pH 7.5 and pH 4.5 in APX activity while GR activity significantly was higher at pH 4.5 than pH 7.5 in both 400  $\mu\text{M}$  and 600  $\mu\text{M}$  concentration. CAT and GPX activity increased in all Zn treatments in both shoot and root compared with control (Fig. 3) and maximum increase was observed at 600  $\mu\text{M}$  (pH 4.5) treatment of zinc. CAT activity significantly was higher at pH 4.5 than pH 7.5 in shoot and there was significant difference between pH 7.5 and pH 4.5 of 600  $\mu\text{M}$  concentration in root. GPX activity significantly was higher at pH 4.5 than pH7.5 in 600  $\mu\text{M}$  concentration in root. But no significant difference was observed between pH 7.5 and pH 4.5 in shoot.


Non-protein thiol compounds are involved in heavy metal detoxification. Non-protein thiols increased in all Zn treatments than those in the control and this increase was higher in 600  $\mu\text{M}$  than 400  $\mu\text{M}$  of zinc (Fig. 4). Also Non-protein thiol compounds were significantly higher in pH 4.5 than pH 7.5 in shoot. Electrolyte leakage (Fig. 4, C)

reflected the plasmalemma damage caused by the Zn. Electrolyte leakage percentage in all Zn treatments increased as compared to the control.


But no significant difference was observed between Zn 400  $\mu$ M and Zn 600  $\mu$ M also between pH 7.5 and pH 4.5.




**Figure 1.** The effects of Zinc on Malondialdehyde content in shoot (A) and root (B) and  $H_2O_2$  content in shoot (C) and root (D) of *Zea mays*. Values are means of three replicates  $\pm$  S.E.



**Figure 2.** The effects of Zinc on APX activities in shoot (A) and root (B) and GR activities in shoot (C) and root (D) of *Zea mays*. Values are means of three replicates  $\pm$  S.E.



**Figure 3.** The effects of Zinc on CAT activities in shoot (A) and root (B) and GPX activities in shoot (C) and root (D) of *Zea mays*. Values are means of three replicates  $\pm$  S.E.



**Figure 4.** The content of non-protein -SH (NPT) in shoot (A) and root (B) and ELP (C) of *Zea mays* under Zn treatment. Values are means of three replicates  $\pm$  S.E.

## DISCUSSION

Heavy metal accumulation in plant lead to some physiological and biochemical changes. Production of free radicals in plants under heavy metal stress has been documented (McGeer *et al.*, 2000). Like other metals zinc also induces free radical generation in plants (Rout and Das, 2003). During present study zinc stimulated MDA production in *Zea mays*. As a product of membrane lipid peroxidation, MDA is used to assess the extent of oxidative stress in plants, and its level is increased under stress conditions (Liu *et al.*, 2009). In plants, heavy metals induce oxidative stress by generation of superoxide radical ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), hydroxyl radical ( $HO^*$ ), and singlet oxygen ( $^1O_2$ ), collectively termed ROS (Jain *et al.*, 2010). These ROS are highly reactive and cause the death of plants by damaging membrane lipids, proteins, pigments and nucleic acids. To cope up with the damages caused by the ROS, cells possess their own comprehensive and integrated endogenous antioxidant defense system composed of both enzymatic as well as non-enzymatic components (Miller *et al.*, 2008). Similar to our results an increase in MDA and  $H_2O_2$  content under excess zinc has been reported in sugarcane (Jain *et al.*, 2010). In the present study increased electrolyte leakage under zinc stress. Generation of free radicals and increased lipid peroxidation under zinc stress may have resulted in an increase in membrane permeability and loss of membrane integrity (Rout and Das, 2003). Jain *et al.* (2010) reported that elevated levels of  $H_2O_2$  and  $O_2^{2-}$  caused by stress facilitate the formation of highly active hydroxyl radicals ( $OH^*$ ). These hydroxyl radicals are generally considered to be the most likely ROS to initiate the peroxidation destruction of lipids that lead to membrane damage. Non-protein thiols, including

glutathione, thiol-rich peptides and other SH compounds, fulfill an important role in the detoxification of heavy metals in plants (Tiryakioglu *et al.*, 2006). In our study, an increase in the level of non-protein thiols observed with the increase Zn. Similarly, a marked increase was seen in non-protein thiols as a response to different metal stress in a number of plants (Tiryakioglu *et al.*, 2006). The induction of certain enzymes that detoxify ROS is considered to play an important role in the defense against oxidative stress caused by toxic metal concentrations (Miller *et al.*, 2008). In plants, a number of enzymes regulate  $H_2O_2$  intracellular levels, but CAT, APX and GPX are considered the most important (Liu *et al.*, 2009). These enzymes work independently in different parts of the plants to break up  $H_2O_2$ . The results of the present study show that the plants treated with Zn had high GPX (Fig. 3), CAT (Fig. 3), APX (Fig. 2) and GR activity (Fig. 2), indicating that there was efficient ROS scavenging activity in the system. Previous studies have also shown that induction of GPX activity has been reported in many plant species, when exposed to zinc stress (McGeer *et al.*, 2000).

The results also showed that at the low pH zinc toxicity higher than that at high pH in *Zea mays* because MDA content, electrolyte leakage, non-protein thiol and antioxidant enzymes activity was higher at pH 4.5 in comparison with that at pH 7.5. High toxicity in low pH might be the resulting in enhance zinc bioavailability in *Zea mays*. Lowering pH increased easily available Zn concentrations and enhanced metal uptake. Zn deficiency is a common problem in plants grown in high pH, calcareous soils (Cakmak *et al.* 1996), whereas in low pH soils Zn availability is generally high (Chaney, 1993). Autumn *et al.* (2006), reported that lowering pH is

an effective method to enhance metal bioavailability and *Thlaspi caerulescens* uptake for both Cd and Zn. The most obvious method for reducing plant Zn toxicity is to raise soil pH. Havlin and co-authors (2005) reported that available Zn and Mn concentrations decreased with raising the soil pH.

In conclusion, our results have demonstrated Zn phytotoxicity in *Zea mays*. Higher H<sub>2</sub>O<sub>2</sub>, MDA contents and increased activity of antioxidant enzymes in *Zea mays* indicating oxidative damage by zinc toxicity. Also at the low pH zinc toxicity was higher in *Zea mays*.

## REFERENCES

- Aebi, H.E. (1983) Methods of Enzymatic Analysis Verlag Chemic: Deer\_eld Beach, FL, 3rd ed. 3, pp. 273-286 .
- Autumn, S., Wang, J., Scott, A., Rufus, L., Chaney Thiery, A., Dolorme and Ronger, D. (2006) Reeves soil pH effect on uptake of Cd and Zn by *Thlaspi caerulescens*. *Plant and soil.*, **281**, 325-337.
- Cakmak, I., Yilmaz, A., Ekiz, H., Torun, B., Erenoglu, B. and Braun, H.J. (1996) Zinc deficiency as a critical nutritional problem in wheat production in Central Anatolia. *Plant and Soil.*, **180**, 165-172.
- Chang, H.B., Lin, C.W. and Huang, H.J. (2005) Zinc-induced cell death in rice (*Oryza sativa* L.) roots. *Plant Growth Regul.*, **46**, 261-266.
- Chaney, R.L. (1993) Zinc phytotoxicity. In: Robson A.D. (Ed.), Zinc in soil and plants. Kluwer Academic Publishers, Dordrecht, the Netherlands: 135-150.
- Edreva, A. and Hadjiiska, E. (1984) About the determination of sulfhydryl(thiol) group content in plant material. *Plant Physiol. (Bulg.)*, **10**, 73-83.
- Foyer, C.H. and Halliwell, B. (1976) The Presence of glutathione and glutathione reductase in chloroplasts: a proposed role in ascorbic acid metabolism. *Planta.*, **133**, 21-25.
- Gao, Y., Miao, C.Y., Xia, J., Mao, L., Wang, Y.F. and Zhou, P. (2011) Effects of coexisting plant species on soil enzyme activities and microbial community structure in multiple heavy metal contaminated soils: PCR-RAPD method. *Front Environ Sci Eng China*.
- Gao, Y., Zhou, P., Mao, L., Shi, W.J. and Zhi, Y.E. (2010) Phytoextraction of Cd and physiological changes in *Solanum nigrum* as a novel hyperaccumulator. *Russ J Plant Physiol.*, **57(4)**, 538-546.
- Havlin, J.L., Beaton, J.D., Tisdale, S.L. and Nelson, W.L. (2005) Soil fertility and fertilizer: An Introduction to Nutrient Management. Upper Saddle River, Newjersey. 515 pp.
- Heath, R.L. and Pacher, L. (1968) Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics.*, **125**, 189-198.
- Jana, S. and Choudhuri, M.A. (1981) Glycolate metabolism of three submerged aquatic anagiosperms during aging. *Aquatic Botany.*, **12**, 345-354.
- Jain, R., Srivastava, S., Solomon, S., Shrivastava, A.K., Chandra, A. (2010) Impact of excess zinc on growth parameters, cell division, nutrient accumulation, photosynthetic pigments and oxidative stress of sugarcane (*Saccharum* spp.). *Acta Physiol. Plant.*, **32**, 979-986.
- Liu, Z.J., Zhang, X.L., Bai, J.G., Suo, B.X., Xu, P.L. and

- Wang, L. (2009) Exogenous paraquat changes antioxidant enzyme activities and lipid peroxidation in droughtstressed cucumber leaves. *Sci. Hortic.*, **121**, 138–143.
- Lutts, S., Kinet, J.M., and Bouharmont, J. (1996) NaCl-induced senescence in leaves of rice (*Oriza sativa* L.) cultivar differing in salinity resistance. *Annal of Botany.*, **78**, 389–398.
- McBride, M., Sauve, S. and Hendershot, W. (1997) Solubility control of Cu, Zn, Cd and Pb in contaminated soils. *European Journal of Soil Science.*, **48**, 337-346.
- McGeer, J.C., Szebedinszky, C., McDonald, D.G. and Wood, C.M. (2000) Effects of chronic sublethal exposure to waterborne Cu, Cd or Zn in rainbow trout. 1: Lono-regulatory disturbance and metabolic costs. *Aqua. Toxicol.*, **50**, 231–243.
- Miller, G., Shulaev, V. and Mitter, R. (2008) Reactive oxygen signaling and abiotic stress. *Physiol. Plant.*, **133**, 481–489.
- Nakano, Y. and Asada, K. (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.*, **22**, 867-880.
- Panda, S.K., Chaudhury, I. and Khan, M.H. (2003) Heavy metals induced lipid peroxidation and effect antioxidants in wheat leaves. *Biol. Plant.*, **46**, 289-294.
- Rout, G.R. and Das, P. (2003) Effect of metal toxicity on plant growth and metabolism: I. Zinc. *Agronomie.*, **23**, 3-11.
- Robinson, B., Malfroy, H., Chartres, C., Helyar, K. and Ayers, G. (1995) The sensitivity of ecosystems to acid inputs in the Hunter Valley. Australia. *Water, Air and Soil Pollution.*, **85**, 1721-1726.
- Stoyanova, Z. and Doncheva, S. (2002) The effect of zinc supply and succinate treatment on plant growth and mineral uptake in pea plant. *Braz. J. Plant Physiol.*, **14**, 111-116.
- Tiryakioglu, M., Eker, S., Ozkutlu, F., Husted, S. and Cakmak, I. (2006) Antioxidant defence system and cadmium uptake in barley genotypes differing in cadmium tolerance. *J Trace Elem Med Biol.*, **20(3)**, 181–189.
- Updhyaya, A., Sankhla, D., Davis, T.D., Sankhla, N. and Smidth, B.N. (1985) Effect of paclobutrazol on the activities of some enzymes of activated oxygen metabolism and lipid peroxidation in senescing soybean leaves. *Plant Physiol.*, **121**, 453-461.
- Vaillant, N., Monnet, F., Hitmi, A., Sallanon, H. and Coudret, A. (2005) Comparative study of responses in four *Datura* species to a zinc stress. *Chemosphere.*, **59**, 1005–1013.